



UNIVERSITY OF LEEDS

This is a repository copy of *(3+1)-dimensional topological quantum field theory from a tight-binding model of interacting spinless fermions*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/99993/>

Version: Accepted Version

Article:

Cirio, M, Palumbo, G and Pachos, JK (2014) (3+1)-dimensional topological quantum field theory from a tight-binding model of interacting spinless fermions. *Physical Review B*, 90 (8). 085114. ISSN 2469-9950

<https://doi.org/10.1103/PhysRevB.90.085114>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

$(3 + 1)$ -dimensional topological quantum field theory from a tight-binding model of interacting spinless fermions

Mauro Cirio,¹ Giandomenico Palumbo,² and Jiannis K. Pachos²

¹*Centre for Engineered Quantum Systems, Department of Physics and Astronomy, Macquarie University, North Ryde, NSW 2109, Australia*

²*School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT, United Kingdom*

(Dated: July 15, 2014)

Currently, there is much interest in discovering analytically tractable $(3 + 1)$ -dimensional models that describe interacting fermions with emerging topological properties. Towards that end we present a three-dimensional tight-binding model of spinless interacting fermions that reproduces, in the low energy limit, a $(3 + 1)$ -dimensional Abelian topological quantum field theory called BF model. By employing a mechanism equivalent to the Haldane's Chern insulator, we can turn the non-interacting model into a three-dimensional chiral topological insulator. We then isolate energetically one of the two Fermi points of the lattice model. In the presence of suitable fermionic interactions, the system, in the continuum limit, is equivalent to a generalised $(3 + 1)$ -dimensional Thirring model. The low energy limit of this model is faithfully described by the BF theory. Our approach directly establishes the presence of $(2 + 1)$ -dimensional BF theory at the boundary of the lattice and it provides a way to detect the topological order of the model through fermionic density measurements.

PACS numbers: 11.15.Yc, 71.10.Fd

I. INTRODUCTION

The interest in strongly interacting fermionic systems has recently found new applications related to topological phases of matter. In the non-interacting case a complete classification [1, 2] of standard topological insulators [3] of free fermions exists. Unfortunately, it is not possible to straightforwardly extend these results to the interacting case. For example, it is not possible to generalise the band theory approach to topological invariants, so more flexible approaches have to be invented [4]. The introduction of interactions in a free fermion system can either connect different phases of matter [5] or give access to new ones [6]. Examples of the latter are the two-dimensional topological Mott insulators [7], where interactions can open an insulating gap and drive the system to topological phases not accessible in the non-interacting case.

Much progress in the study of interacting fermionic systems has already been made in $1+1$ and $2+1$ dimensions [8, 9]. In three spatial dimensions the situation is somehow less clear, though some analysis has been already carried out [6, 10]. Complications arise already in the effective description, where the Chern-Simons theory [11] only holds in even spatial dimensions with broken time reversal symmetry. A natural generalization of Chern-Simons theory is the topological BF theory, which is well defined in any dimensions [12]. In two spatial dimensions BF theories can be interpreted as double Chern-Simons theories, allowing for the description of time-reversal symmetric topological insulators [13]. BF theories have also been proposed as effective theories for describing topological insulators in any dimension [14–18]. Nevertheless, very few interacting fermionic models that give rise to BF theory are available.

Here we make another step into the exploration of

interactions-driven phases of matter. Our starting point is a cubic lattice of spinless fermions. For particular values of the couplings and in the absence of interactions the system becomes a chiral topological insulator [19]. Our approach is similar in spirit to Haldane's Chern insulator [20], which gives us the ability to arbitrarily tune the asymmetry in the energy spectrum of the theory. This allows us to enter a regime where the dynamics, associated with one of the two Dirac fermions present in the model, is adiabatically eliminated [21, 22]. Subsequently, we introduce interactions between the tight-binding fermions to obtain a generalization of the $(3+1)$ -dimensional massive Thirring model [23] with a tensorial current. By applying a series of transformations [24] we show that our system simulates a $(3 + 1)$ -dimensional topological massive gauge theory [25, 26]. The short distance behaviour of this theory is dominated by a Maxwell term. The large distance behaviour is characterised by an Abelian BF term which is topological in nature and it gives mass to the gauge field. The connection of the fermionic tight-binding model to the BF theory allows us to directly obtain that the boundary of the lattice is described by the $(2 + 1)$ -dimensional BF theory. Finally, we identify analytical expressions for topological invariants associated with the model and relate them to physical local fermionic observables. This method allows us to probe the topological properties of our three-dimensional system and provides a possible platform for simulating $(3 + 1)$ -dimensional gauge theories in the laboratory with cold atoms [27] in optical lattices [28, 29].

This article is organized as follows. In section II we introduce a free fermion tight binding model. We then (section II A) focus on the kinematic sector by analysing the (gapless) energy spectrum, the symmetry properties (introduced in Appendix A), and the low energy limit of the model. In section II B we consider the effect of ad-

ditional mass terms which open a gap in the spectrum and allows us (section II C) to show the existence of a chiral topological insulating phase (and its formal analogies with the Haldane model). In section III we leave the free fermion description by introducing 4-bodies interactions in the tight binding model (section III A). We then show that in the low energy limit the model is described by bosonic degrees of freedom (section III B) and we find (section III C) the corresponding effective theory through a duality operation. Interestingly, the effective theory contains a purely topological term. We then connect the effective and microscopic theory by proposing opportune bosonization rules (section III D). In section III E we explore two features of the theory in its purely topological regime. We find that the boundary of the model is described by a topological theory III E 1 (more details can be found in Appendix B). Finally, we describe microscopic fermionic observables (section III E 2) which can be used to test the topological features of the model (more details can be found in Appendix C).

II. FREE FERMION MODEL




FIG. 1: The tight-binding model, where spinless fermions (yellow) reside on the vertices of a cubic lattice. The plaquette unit cell has four fermions labelled a, b, c, d . The fermions tunnel along the lattice via Hamiltonian (1). Tunnelling takes place along the edges of the cubic lattice (black) with coupling t and a phase that is determined by the black arrows, i.e. $\chi^t = i$ for positive and $\chi^t = -i$ for negative direction. A purely imaginary staggering term in the y -direction (orange) has coupling δt and a phase $\chi^{\delta t}$ with the same phase convention as χ^t . Tunnelling along the diagonals of the cube (green) have coupling \bar{t} and phase $\chi^{\bar{t}} = i\bar{t}e^{\pm i\phi}$ with $\phi \in [0, \pi/2]$, where the negative (positive) sign is chosen for full (dashed) lines.

Let us begin with an overview of the model. We

consider spinless fermions, localised on the vertices of a three-dimensional cubic lattice Λ , as shown in Fig. 1. The tight-binding Hamiltonian is given by

$$H = t \sum_{\langle \mathbf{i}, \mathbf{j} \rangle} \chi_{\mathbf{i}\mathbf{j}}^t f_{\mathbf{i}}^\dagger f_{\mathbf{j}} + \delta t \sum_{\langle \mathbf{i}, \mathbf{j} \rangle_y} \chi_{\mathbf{i}\mathbf{j}}^{\delta t} f_{\mathbf{i}}^\dagger f_{\mathbf{j}} - \frac{\bar{t}}{2} \sum_{\langle\langle \mathbf{i}, \mathbf{j} \rangle\rangle} \chi_{\mathbf{i}\mathbf{j}}^{\bar{t}} f_{\mathbf{i}}^\dagger f_{\mathbf{j}}, \quad (1)$$

where $\mathbf{i}, \mathbf{j} \in \Lambda$ and $f_{\mathbf{i}}^\dagger$ and $f_{\mathbf{i}}$ are the creation and annihilation fermion operators at position \mathbf{i} of the lattice. We define planar unit cells populated by four fermion flavours $f \in \{a, b, c, d\}$, as shown in Fig. 1. Let us analyse each term of the Hamiltonian. The first term, which we call kinematic, has coupling t and corresponds to nearest-neighbour $\langle \mathbf{i}, \mathbf{j} \rangle$ hopping. The phases χ^t are such to create a net π flux through each plaquette. The term proportional to δt describes a staggering between sites along the y -direction indicated by $\langle \mathbf{i}, \mathbf{j} \rangle_y$. The last term corresponds to tunnelling between the next-next-nearest neighbouring sites, $\langle\langle \mathbf{i}, \mathbf{j} \rangle\rangle$, with coupling \bar{t} . The phase factors χ^t , $\chi^{\delta t}$ and $\chi^{\bar{t}}$ are defined in Fig. 1. Let us now study this model more explicitly.

The lattice of the unit cells (in blue in Fig. (1)) is given by: $\bar{\Lambda} = \{\mathbf{i} \in \mathbb{R}^3 : \mathbf{i} = n_1 \mathbf{s}_1 + n_2 \mathbf{s}_2 + n_3 \mathbf{s}_3\}$, with $n_i \in \mathbb{N}$ and $\mathbf{s}_1 = (2, 0, 0)$, $\mathbf{s}_2 = (0, 2, 0)$, $\mathbf{s}_3 = (1, 0, 1)$ written in units of a fixed reference length. The Hamiltonian in Eq. (1) can be written as

$$H = H_0 + H_m, \quad (2)$$

where $H_0 = t \sum_{\langle \mathbf{i}, \mathbf{j} \rangle} \chi_{\mathbf{i}\mathbf{j}}^t f_{\mathbf{i}}^\dagger f_{\mathbf{j}}$ is a kinematic Hamiltonian (defined through the black links in Fig. (1)) which has gapless spectrum. In order to open a gap in the model we introduce the Hamiltonian $H_m = \delta t \sum_{\langle \mathbf{i}, \mathbf{j} \rangle_y} \chi_{\mathbf{i}\mathbf{j}}^{\delta t} f_{\mathbf{i}}^\dagger f_{\mathbf{j}} - \frac{\bar{t}}{2} \sum_{\langle\langle \mathbf{i}, \mathbf{j} \rangle\rangle} \chi_{\mathbf{i}\mathbf{j}}^{\bar{t}} f_{\mathbf{i}}^\dagger f_{\mathbf{j}}$ which is defined along the red and green links in Fig. (1). Let us now define the two terms of the Hamiltonian one by one.

A. The kinematic model

As can be seen by inspecting Eq. (2) and Fig. (1), the kinematic Hamiltonian of the model can be written as

$$H_0 = it \sum_{\mathbf{i}} \left[(-a_{\mathbf{i}}^\dagger b_{\mathbf{i}} + b_{\mathbf{i}}^\dagger d_{\mathbf{i}} + d_{\mathbf{i}}^\dagger c_{\mathbf{i}} + c_{\mathbf{i}}^\dagger a_{\mathbf{i}}) \right. \\ + (a_{\mathbf{i}+\bar{\mathbf{s}}_1}^\dagger b_{\mathbf{i}} + c_{\mathbf{i}+\bar{\mathbf{s}}_1}^\dagger d_{\mathbf{i}} + d_{\mathbf{i}}^\dagger b_{\mathbf{i}+\bar{\mathbf{s}}_2} + a_{\mathbf{i}+\bar{\mathbf{s}}_2}^\dagger c_{\mathbf{i}}) \\ + (b_{\mathbf{i}+\bar{\mathbf{s}}_3-\bar{\mathbf{s}}_1}^\dagger a_{\mathbf{i}} + b_{\mathbf{i}}^\dagger a_{\mathbf{i}+\bar{\mathbf{s}}_3} + d_{\mathbf{i}}^\dagger c_{\mathbf{i}+\bar{\mathbf{s}}_3} + d_{\mathbf{i}+\bar{\mathbf{s}}_3-\bar{\mathbf{s}}_1}^\dagger c_{\mathbf{i}}) \\ \left. + \text{h.c.} \right], \quad (3)$$

where $\mathbf{i} \in \bar{\Lambda}$ is intended and where t is an energy scale. This Hamiltonian is known [30] to give rise in the continuum limit to two massless Dirac fermions. Let us now calculate the spectrum explicitly.

The reciprocal lattice is defined as $\bar{\Lambda}_p = \{\mathbf{p} \in \mathbb{R}^3 : \mathbf{p} = \sum_i n_i \mathbf{p}_i\}$ where the vectors \mathbf{p}_i satisfy $\mathbf{p}_i \cdot \mathbf{s}_j = 2\pi \delta_{ij}$ and are explicitly defined to be $\mathbf{p}_1 = \pi(1, 0, -1)$,

$\mathbf{p}_2 = \pi(0, 1, 0)$, $\mathbf{p}_3 = 2\pi(0, 0, 1)$. The Brillouin zone BZ is defined as the elementary cell in the reciprocal lattice $\text{BZ} = \{\mathbf{p} \in \bar{\Lambda}_p : \mathbf{p} = p_i \mathbf{p}_i\}$ with $p_i \in [0, 1)$. A generic vector in the Brillouin zone can be written as $\mathbf{p} \equiv (p_x, p_y, p_z) = \pi(p_1, p_2, 2p_3 - p_1)$ so that $p_x = \pi p_1$, $p_y = \pi p_2$, $p_z = \pi(2p_3 - p_1) = \pi 2p_3 - p_x$. The periodic invariance of the phase space allows us to parametrize the Brillouin zone in a different and somehow more convenient way. We can in fact define it as $\text{BZ} = \{\mathbf{p} \in \bar{\Lambda}_p : \mathbf{p} = (p_x, p_y, p_z)$ with $p_x \in [0, \pi)$, $p_y \in [0, \pi)$, $p_z \in [0, 2\pi)$ where the volume of the Brillouin zone is $2\pi^3$. We now have all the ingredients to define the Fourier transform $a_{\mathbf{r}} = \sum_{\mathbf{p} \in \text{BZ}} e^{-i\mathbf{p}\cdot\mathbf{r}} a_{\mathbf{p}}$ and analogously for b, c, d . By introducing the Fourier transformed operators inside the Hamiltonian in Eq. (3) we find

$$\begin{aligned} H_0 &= it \sum_{\mathbf{p}} [(-1 + e^{2\pi i p_1} - e^{-2\pi i(p_3 - p_1)} - e^{2\pi i p_3}) a_{\mathbf{p}}^\dagger b_{\mathbf{p}} \\ &\quad + (-1 + e^{2\pi i p_2}) a_{\mathbf{p}}^\dagger c_{\mathbf{p}} + i(1 - e^{2\pi i p_2}) b_{\mathbf{p}}^\dagger d_{\mathbf{p}} \\ &\quad + (-1 + e^{2\pi i p_1} - e^{2\pi i p_3} - e^{-2\pi i(p_3 - p_1)}) c_{\mathbf{p}}^\dagger d_{\mathbf{p}}] \\ &\quad + \text{h.c.} \\ &= it \sum_{\mathbf{p}} [(-1 + e^{2i p_x} - e^{-i(p_z - p_x)} - e^{i(p_z + p_x)}) a_{\mathbf{p}}^\dagger b_{\mathbf{p}} \\ &\quad + (e^{2i p_y} - 1) a_{\mathbf{p}}^\dagger c_{\mathbf{p}} + i(1 - e^{2i p_y}) b_{\mathbf{p}}^\dagger d_{\mathbf{p}} \\ &\quad + (-1 + e^{2i p_x} - e^{i(p_z + p_x)} - e^{-i(p_z - p_x)}) c_{\mathbf{p}}^\dagger d_{\mathbf{p}}] \\ &\quad + \text{h.c.} \\ &= \sum_{\mathbf{p}} \Psi'^\dagger \bar{H}'_0 \Psi' , \end{aligned} \quad (4)$$

with

$$\Psi' = \begin{pmatrix} a_{\mathbf{p}} \\ b_{\mathbf{p}} \\ c_{\mathbf{p}} \\ d_{\mathbf{p}} \end{pmatrix} \quad (5)$$

and with the kernel \bar{H}'_0 given by

$$\bar{H}'_0 = t \begin{pmatrix} 0 & B & C & 0 \\ B^* & 0 & 0 & -C \\ C^* & 0 & 0 & B \\ 0 & -C^* & B^* & 0 \end{pmatrix} , \quad (6)$$

where we have defined $B = i(-1 + e^{2i p_x} - e^{-i(p_z - p_x)} - e^{i(p_z + p_x)})$ and $C = i(e^{2i p_y} - 1)$. From the explicit expression of H_0 we can easily see that the set of vertices a and d only interacts with the set b and c . This condition defines a chiral symmetry. In fact, such a symmetry describes the existence of a bipartition of the lattice “broken” by *all* couplings (see Appendix A for more details). The existence of chiral symmetry allows to cast the Hamiltonian in an off-block diagonal form. In our case this is easily seen after the definition of a new basis

$$\Psi = \begin{pmatrix} a_{\mathbf{p}} \\ d_{\mathbf{p}} \\ c_{\mathbf{p}} \\ b_{\mathbf{p}} \end{pmatrix} , \quad (7)$$

the Hamiltonian takes the form

$$H_0 = \Psi^\dagger \bar{H}_0 \Psi , \quad (8)$$

with

$$\bar{H}_0 = t \begin{pmatrix} 0 & 0 & C & B \\ 0 & 0 & B^* & -C^* \\ C^* & B & 0 & 0 \\ B^* & -C & 0 & 0 \end{pmatrix} . \quad (9)$$

This Hamiltonian has eigenvalues (with degeneracy 2) given by

$$E_0 = \pm t \sqrt{6 - 2 \cos 2p_x - 2 \cos 2p_y + 2 \cos 2p_z} . \quad (10)$$

The spectrum has then two double degenerate bands and it is gapped everywhere in the Brillouin zone except for two Fermi points where the two bands touch each other. The two independent Fermi points are given by

$$\begin{cases} \mathbf{P}_+ &= (0, 0, \frac{\pi}{2}) \\ \mathbf{P}_- &= (0, 0, \frac{\pi}{2} + \pi) . \end{cases} \quad (11)$$

In order to study the behaviour around the Fermi points we now define the following matrices

$$\alpha^x = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} , \quad \alpha^y = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} ,$$

$$\alpha^z = \begin{pmatrix} 0 & 0 & 0 & +i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix} , \quad (12)$$

which satisfy the algebra

$$\{\alpha^i, \alpha^j\} = 2\delta^{ij} . \quad (13)$$

We now introduce coordinates around the Fermi points $\hbar\mathbf{p} = \hbar\mathbf{P}_\pm + (k_x, k_y, k_z)$ for small k_x , k_y and k_z , so that the Hamiltonian around the Fermi points looks like

$$\bar{\mathcal{H}}_\pm^0 = c(k_x \alpha_x + k_y \alpha_y \pm k_z \alpha_z) , \quad (14)$$

where $c = 2t/\hbar$. The Hamiltonians in Eq. (14) represent two (massless) Dirac fermions.

1. Symmetries

The symmetries of the kinematic model can be studied by analyzing the Hamiltonian kernel (9). In particular we are interested in checking the behaviour of the model under time-reversal, particle-hole and chiral symmetry. For an introduction to the definitions of these symmetries we refer to Appendix A. In the table below we express the conditions on the Hamiltonian kernel under which these symmetries are satisfied.

Inspection of the Hamiltonian kernel given in (9) shows us that $\bar{H}_0^*(-p) = -\bar{H}_0(p)$. This condition means that the system breaks time-reversal symmetry and preserves

Symmetry	Condition
Time Reversal	$\bar{H}(\mathbf{p}) = \bar{H}^*(-\mathbf{p})$
Particle-Hole	$\bar{H}(\mathbf{p}) = -\bar{H}^*(-\mathbf{p})$
Chiral	$\exists \bar{C}_s : \bar{C}_s^\dagger = \bar{C}_s^{-1} : \bar{C}_s \bar{H}(\mathbf{p}) = -\bar{H}(\mathbf{p}) \bar{C}_s$




FIG. 2: Energy bands for the full model described in Eq. (1) as a function of the momentum variable p_z for fixed $p_x = p_y = 0$, in arbitrary units. The parameters δt and \bar{t} are tuned to open a gap. The model is in fact a description of a chiral topological insulator. The dashed line represents the Fermi energy and highlights the insulating properties of the material.

particle-hole symmetry. We also have an explicit chiral symmetry since the Hamiltonian anticommutes with the matrix \bar{C}_s defined as

$$\bar{C}_s = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (15)$$

which is hermitian and unitary, as expected from the block structure of the Hamiltonian in (9).

B. Gapped model

The kinematic model introduced in the previous section is gapless. In order to study the topological features of the model, we now introduce a gap term. In this way the low energy physics of the model is described by a massive Dirac fermion. Such a mass term has to anticommute with all the α matrices (Eq. (12)), square to the identity and satisfy chiral symmetry. As can be checked, the mass term has to be proportional to $\beta = \bar{C}_s \alpha^x \alpha^y \alpha^z$.

In the chosen representation, we have

$$\beta = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}. \quad (16)$$

The implementation of such a mass term requires the introduction of additional couplings between the sites a and c and between b and d (as can be seen by inspecting the explicit form of β in the basis given by Eq. (7)). We introduce a staggering of the a, c and b, d couplings along the y axis and a next-next nearest neighbor (NNN) interactions as shown in Fig. (1). The staggering NNN interactions give an equal (opposite) mass term to the two Dirac fermions defined in Eq. (17). Explicitly we define

$$H_m = \sum_{\bar{r}} i \delta t a_{\bar{r}}^\dagger c_{\bar{r}} + i \delta t d_{\bar{r}}^\dagger b_{\bar{r}} + \frac{\bar{t}}{2} \sum_{\bar{r}} i e^{-i\phi} a_{\bar{r}}^\dagger c_{\bar{r}+\bar{s}_3} + i e^{i\phi} a_{\bar{r}}^\dagger c_{\bar{r}-\bar{s}_3} + \frac{\bar{t}}{2} \sum_{\bar{r}} i e^{-i\phi} d_{\bar{r}}^\dagger b_{\bar{r}+\bar{s}_3} + i e^{i\phi} d_{\bar{r}}^\dagger b_{\bar{r}-\bar{s}_3} + \text{h.c.}, \quad (17)$$

where we have introduced difference energy scales δt and \bar{t} for the staggering and NNN term respectively. We also introduced a phase ϕ associated with the NNN couplings. In momentum space this Hamiltonian becomes

$$H_m = \sum_{\mathbf{p}} i(\delta t + \bar{t} \cos(p_z + p_x + \phi)) a_{\mathbf{p}}^\dagger c_{\mathbf{p}} + i(\delta t + \bar{t} \cos(p_z + p_x + \phi)) d_{\mathbf{p}}^\dagger b_{\mathbf{p}} + \text{h.c.} \quad (18)$$

In the basis of Eq. (7) the kernel in momentum space of the interaction Hamiltonian H_1 reads

$$\bar{H}_m = (\delta t + \bar{t} \cos(p_z + p_x + \phi)) \beta, \quad (19)$$

which is exactly what we are looking for since it gives a contribution proportional to the matrix β which allows the interpretation of a fermion mass as discussed above. Now, we first notice that, when $\bar{t} \neq 0$ and $\phi \neq 0, \pi$ the particle-hole symmetry is broken: $\bar{H}_m^*(-k) \neq -\bar{H}_m(k)$. Incidentally, it is important to notice that by adding these interactions we did not restore time reversal symmetry (already broken in the kinematic model) in the full model.

The results of this section imply that the full Hamiltonian $H = H_0 + H_m$ breaks time-reversal and particle-hole symmetry while it is symmetric under chiral symmetry. We also note that the joint presence of staggering and NNN interactions allows to arbitrarily tune the fermion masses at the two Fermi points. This is easily seen by evaluating Eq. (19) at the two Fermi points to get two independent masses. More precisely, let us define

$$\begin{cases} m_+ c^2 = \delta t + \bar{t} \cos(\frac{\pi}{2} + \phi) \\ m_- c^2 = \delta t + \bar{t} \cos(\frac{3\pi}{2} + \phi) \end{cases}, \quad (20)$$

which, for the choice $\phi = \frac{\pi}{2}$ becomes

$$\begin{cases} m_+ c^2 = \delta t - \bar{t} \\ m_- c^2 = \delta t + \bar{t} \end{cases}. \quad (21)$$

With these definitions we get the expression for the full Hamiltonian around the two Fermi points (to be compared with Eq. (14))

$$\mathcal{H}_{\pm}(\mathbf{k}) = \bar{\Psi}_{\pm}(c\boldsymbol{\alpha} \cdot \mathbf{k} + m_{\pm}c^2\beta)\Psi_{\pm} , \quad (22)$$

where $\boldsymbol{\alpha} = \{\alpha_x, \alpha_y, \alpha_z\}$ and $\mathbf{k} = \{k_x, k_y, k_z\}$. Notice that when $\delta t = 0$ or $\phi = 0, \pi$ such an arbitrary tuning would not be possible and we would get $m_+ = m_-$.

We end up this section with the book-keeping explicit expression for the total Hamiltonian of the model of Eq. (1). From Eq. (9) and (19) and with the definitions (12), (16) and the ones below Eq. (6) the kernel \bar{H} of the total Hamiltonian reads

$$\begin{aligned} \bar{H} &= \bar{H}_0 + \bar{H}_m \\ &= t(\sin 2p_x - \sin(p_x + p_z) - \sin(p_x - p_z))\alpha^x \\ &\quad + t \sin 2p_y \alpha^y \\ &\quad + t(\cos 2p_x - \cos(p_x + p_z) - \cos(p_x - p_z) - 1)\alpha^z \\ &\quad + (t \cos 2p_y - t + \delta t + \bar{t} \cos(p_x + p_z + \phi))\beta . \end{aligned} \quad (23)$$

The spectrum of the total Hamiltonian has two double generate bands

$$E = \pm \sqrt{(4 - 2 \cos 2p_x + 2 \cos 2p_z + |M|^2)} , \quad (24)$$

where $M = (e^{2ip_y} - 1 + \delta t + t \cos(p_x + p_y + \phi))$.

C. Chiral Topological Insulator

Symmetry protected phases of matter for models described by a free fermion model are completely classified [1]. This classification characterizes phases of matter within 10 different symmetry classes determined by the symmetry properties under time-reversal, particle-hole, and chiral symmetry. More specifically, one starts by continuously deforming the Hamiltonian H that describes a free fermion model to a “reference” Hamiltonian Q [31]. This Hamiltonian has all occupied (empty) bands “flatten” with energy +1 (-1) in the whole Brillouin zone. This can be done by defining the operator $Q(k)$ as

$$Q(k) = 2P(k) - \mathbb{I}_{n+m} , \quad (25)$$

with

$$P(k) = \sum_{i=1}^m |u_i(k)\rangle \langle u_i(k)| , \quad (26)$$

where $u_i(k)$ $i = 1, \dots, m$ are the eigenvalues of the occupied bands for the total Hamiltonian and $m(n)$ is the number of occupied (empty) bands. The operator Q is such that $Q^\dagger = Q$, $Q^2 = \mathbb{I}$ and $\text{tr}(Q) = m - n$. This operator has eigenvalues +1 and -1 corresponding to occupied and empty bands. Each of the 10 symmetry classes mentioned above determines a manifold \mathcal{B} such that $Q : \text{BZ} \rightarrow \mathcal{B}$. Within each symmetry class (and hence for each manifold \mathcal{B}), we want to classify the phases of matter

described by the reference Hamiltonians Q . Two Hamiltonians belong to the same phase if they can be continuously deformed one into the other without encountering a critical point. As shown in [1] one can classify such phases through the d -th homotopy group π_d of the manifold \mathcal{B} where d is the spatial dimension of the model. For example, for the symmetry class A (all symmetries broken), we have that \mathcal{B} is isomorphic to the Grassmannian: $\mathcal{B} \simeq G_{n, n+m}(\mathbb{C}) \equiv U(n+m)/U(n) \times U(m)$. In fact, the collection of all energy eigenvectors describes an element of $U(n+m)$ modulo the “gauge” symmetry relabeling the eigenvectors corresponding to occupied and empty bands. Now, for two spatial dimensions we have an infinite number of different phases as implied by $\pi_2(G_{n, n+m}(\mathbb{C})) = \mathbb{Z}$ (specifying, for example, the number of edge states for the quantum Hall effect, which in fact, being a Chern insulator, belongs to the A symmetry class). In three spatial dimensions we have $\pi_3(G_{n, n+m}(\mathbb{C})) = e$ (where e represent the group trivial element) so that only the trivial phase is allowed. In three dimensions things can loose their trivial nature when other symmetry classes are considered. Specifically, we are interested in the symmetry class AIII where only chiral symmetry is preserved. In this case $n = m$ (positive and energy eigenstates come in pairs, see Appendix A), and one can write Q in the following block form [1]

$$Q(\mathbf{p}) = \begin{pmatrix} 0 & q(\mathbf{p}) \\ q^\dagger(\mathbf{p}) & 0 \end{pmatrix} , \quad (27)$$

with $q(\mathbf{p}) \in U(n)$. Our model is then described by the function $q : \text{BZ} \rightarrow \mathcal{B}$ where $\mathcal{B} \simeq U(n)$. Contrary to the class A example, we now find that $\pi_3(U(n)) = \mathbb{Z}$ allowing for non-trivial phases in three spatial dimensions. In fact, we can define [1] a winding number ν (associated with the map q) labelling all the possible phases as

$$\nu = \frac{1}{24\pi^3} \int d^3k \epsilon^{abc} \text{tr}[(q^{-1} \partial_a q)(q^{-1} \partial_b q)(q^{-1} \partial_c q)] , \quad (28)$$

where $a, b, c = 1, 2, 3$.

1. Phase Diagram

We now want to see under which conditions on the parameters of the full Hamiltonian of our model (Eq. (1)) we can get non-zero winding number (Eq. (28)). Let us first resume what we learned so far. First of all, from the results of section II B we know that non-trivial topological properties are forbidden for $\delta t = 0$ or $\phi = \{0, \pi\}$. In fact, in such a regime, the particle-hole symmetry is not broken leading to the impossibility to define a winding number as shown in [1]. Second, we know that the conditions $\delta t = \bar{t} \cos(\frac{\pi}{2} + \phi)$ and $\delta t = \bar{t} \cos(\frac{3\pi}{2} + \phi)$ imply (see Eq. (20)) that either $m_+ = 0$ or $m_- = 0$ meaning that the system is critical. Then, in these cases we expect the winding number in Eq. (28) to be not well defined.

Let us now draw the phase diagram of the model in the




FIG. 3: Phase diagram of Hamiltonian (1) as a function of the couplings δt and \bar{t} parameterised by the phase ϕ . The gapped regions with non-trivial winding number $\nu = \pm 1$ (yellow) are separated by phase transition (blue lines) from the topologically trivial regions with $\nu = 0$ (white). The winding number, ν , is correlated with the sign of $m_+ \cdot m_-$, in the same way as in Haldane's model. The critical line for $\phi = \pi/2$ (dashed blue) and for generic value of $\phi \in [0, \pi/2]$ (solid blue) are depicted.

parameter space $\{\delta t, \bar{t}\}$ (see Fig. (3)). A gapless system is described by the imposing the equations $m_+ = 0$ and $m_- = 0$ (see Eq. (20)). Pictorially, these equations are two straight lines in the $\delta t, \bar{t}$ plane, parametrized by the phase ϕ . They divide the parameter space $\delta t, \bar{t}$ in 4 disconnected parts. We studied the behaviour of the winding number in these 4 phases and we found that two of them are in fact non-trivial (see Fig. 3). When the parameter ϕ tends to 0 the two non-trivial phases disappear as the two critical lines merge together. This result is consistent to the fact that $\phi = 0$ corresponds to a system where particle hole symmetry is not broken (see the analysis following Eq. (19)) which is a sufficient condition for the absence of topological order [1]. The non-triviality of the winding number Eq. (28) (for a certain parameters regime) shows that the system is a chiral topological insulator.

To summarize, the introduction of NNN neighbour interactions ($\bar{t} \neq 0$) breaks particle-hole symmetry (provided that $\phi \neq 0$) and gives an opposite contribution to the masses in Eq. (20). On the contrary, the staggered interactions ($\delta t \neq 0$) give an equal contribution to the fermion masses. As a consequence, the simultaneous presence of both interactions allow to arbitrarily tune the masses m_{\pm} .

Intuitively, this model presents several formal analogies with the Haldane model [20] where spinless electrons hop on the vertices of a honeycomb lattice. Such a model has a kinematic term which preserves time-reversal and inversion symmetries (and breaks particle-hole symmetry)

that gives rise to two gapless Fermi points. In addition, next-nearest neighbour interactions (mimicking a nested magnetic field) and a staggered chemical potential break, respectively, time reversal and inversion symmetry. The breaking of each symmetry allows for a non-zero energy gap to appear. More precisely, the fermions at the two Fermi points acquire the same (opposite) mass due to the breaking of time-reversal (inversion) symmetry. In this sense, our NNN neighbour and staggering terms mimic, respectively, the staggered magnetic field and the chemical potential of the Haldane model. In the light of the classification given in [1], the key-feature to build a non-trivial topological phase on top of our (Haldane) kinematic theory is to break the particle-hole (time-reversal) symmetry. Despite these similarities, the Haldane model breaks all symmetries (it describes the physics of the quantum Hall effect without magnetic field) while we have to pay extra attention to preserve chiral symmetry in order to protect the topological phase.

III. INTERACTING FERMIONS MODEL

We now turn to the case of interacting fermions. The starting point is the effective theory described in Eq. (22) with $\phi = \pi/2$. This model has enough flexibility to allow us to arbitrarily tune the masses around the two Fermi points shown in Fig. 4. Following the approach in [21] we can define a hierarchy in the energy scales, given by $|m_+ c^2| \ll |m_- c^2|$, and adiabatically eliminate the physics around the second Fermi point, \mathbf{P}_- . We now introduce four-body fermionic interactions with coupling U that is small compared to the energy scale of \mathbf{P}_- , i.e. $\sqrt{(\hbar c)^3/U} \ll m_- c^2$, and comparable to $|m_+ c^2|$. These interactions are particularly designed so that they give rise to self-interacting current-current terms in the single Dirac fermion description corresponding to \mathbf{P}_+ . The resulting effective physics is encoded in the Hamiltonian

$$\mathcal{H}(\mathbf{p}) = \Psi^\dagger (c\boldsymbol{\alpha} \cdot \mathbf{p} + mc^2\beta) \Psi + \frac{g^2}{2m} (2J_{\mu\nu} J^{\mu\nu} - J_\mu J^\mu), \quad (29)$$

where $m \equiv m_+ = (\delta t - \bar{t})/c^2$ and $g^2 = 2mU$. There are two types of currents given by $J^\mu = \bar{\Psi} \gamma^\mu \Psi$ and $J^{\mu\nu} = \bar{\Psi} \gamma_5 [\gamma^\mu, \gamma^\nu] \Psi$, for $\bar{\Psi} = \Psi^\dagger \gamma^0$ with the gamma matrices γ^μ defined as $\boldsymbol{\gamma} = \beta \boldsymbol{\alpha}$, $\gamma^0 = \beta$ and $\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$. A dimensional analysis shows that the four-components Dirac field has dimensions $[\Psi] = (\text{Length})^{-3/2}$ compatible with the units of the Hamiltonian density above. This fixes the dimensions of the current-current interaction terms (to $(\text{Length})^{-6}$) which, in fact, implies that $[U] = \text{Energy} \cdot (\text{Length})^3$ has dimensions of Energy times Volume. The Hamiltonian in Eq. (29) is the tensorial generalization of the Thirring model [23] in $3+1$ dimensions. This generalization of the Thirring model is not renormalizable (at least by means of perturbative methods). It is analogous in spirit to the Nambu-Jona-Lasinio model [32] and the Fermi effective




FIG. 4: Adiabatic elimination of a Fermi point. The energy spectrum (in arbitrary units) of the model described by the Hamiltonian in Eq. (1) (as a function of the momentum variable p_z for fixed $p_x = p_y = 0$) allows to arbitrary tune the masses around the two Fermi points (see Eq. (20)). In particular, by opportunely choosing the parameters δt and \bar{t} , it is possible to work in a regime where $m_- \gg m_+$. If we suppose not to have any perturbation with energy scale bigger than $|m_-|$, then the low energy physics of the system is completely described around the P_+ Fermi point in the Brillouin zone.

model [33] for weak interactions which involves the non renormalizable point-like interaction between two currents. The Thirring interaction terms which appear in the Hamiltonian (29) have a coupling parameter $U = \frac{g^2}{m}$ which has dimensions Length \times Energy. The energy scale associated with such parameter is given by $\tilde{U} = \sqrt{\frac{(\hbar c)^3}{U}}$ (which is the only energy scale we can define from \hbar , c and U). This gives a dimensional analysis justification to the adiabaticity condition $\tilde{U} = \sqrt{\frac{(\hbar c)^3}{U}} \ll m_- c^2$ which allows us to restrict the physics around one Fermi point. From now on we will use units where $c = \hbar = 1$.

A. Microscopic prescription

We now want to find the microscopic description for the Hamiltonian in Eq. (29). To achieve this, we proceed backwards and substitute in J^μ and $J^{\mu\nu}$ the expressions for the spinor $\Psi = (a_{\mathbf{p}} \ d_{\mathbf{p}} \ c_{\mathbf{p}} \ b_{\mathbf{p}})^T$ and the gamma matrices as given by $\gamma_0 = \beta$ and $\gamma_i = \beta\alpha_i$. After some

tedious calculations one gets

$$\begin{aligned} J_\mu J^\mu &= (\bar{\Psi} \gamma_\mu \Psi)(\bar{\Psi} \gamma^\mu \Psi) \\ &= -2(a^\dagger a + b^\dagger b + c^\dagger c + d^\dagger d) \\ &\quad + 2(a^\dagger a)(d^\dagger d) + 4(a^\dagger a)(c^\dagger c) + 6(c^\dagger c)(d^\dagger d) \\ &\quad + 6(a^\dagger a)(b^\dagger b) + 4(b^\dagger b)(d^\dagger d) + 2(c^\dagger c)(b^\dagger b) \\ &\quad + 6a^\dagger b^\dagger c d + 6c^\dagger d^\dagger a b - 2a d b^\dagger c^\dagger - 2b c a^\dagger d^\dagger, \end{aligned} \quad (30)$$

and

$$\begin{aligned} J_{\mu\nu} J^{\mu\nu} &= (\bar{\Psi} \gamma^5 [\gamma_\mu, \gamma_\nu] \Psi)(\bar{\Psi} \gamma^5 [\gamma^\mu, \gamma^\nu] \Psi) \\ &= 48 [(a^\dagger a)(b^\dagger b) + (c^\dagger c)(d^\dagger d) \\ &\quad - (a^\dagger a)(d^\dagger d) - (b^\dagger b)(c^\dagger c)] \\ &\quad - 48 [a^\dagger b^\dagger c d + b^\dagger c^\dagger a d + c^\dagger d^\dagger a b + a^\dagger d^\dagger b c]. \end{aligned} \quad (31)$$

The terms involving only two fermions can be omitted since they give a contribution to the Hamiltonian kernel which is proportional to the identity and can be seen as a constant chemical potential on every site of the lattice. The other terms either interactions between two sites populations or between four sites.

This shows the explicit form of the microscopic interaction needed to simulate the Tensorial Thirring model at low energies given by Eq. (29). We note that some of these interactions are attractive while other are repulsive.

B. Bosonization

Throughout the rest of the section we assume that $g \neq 0$ in Eq. (29). The Thirring model describes a relativistic fermion with a potential given by self-interactions. In order to get a more accessible theory it is possible to linearize the interaction by introducing new degrees of freedom. Following this approach, we now show how to describe the low energy physics of the Tensorial Thirring model with a pure bosonic theory.

As it can be seen from Eq. (29), the effective theory of our model is described, in Euclidean space, by the action

$$Z_{\text{TTh}} = \int \mathcal{D}[\bar{\Psi}] \mathcal{D}[\Psi] e^{-\mathcal{S}_D - \mathcal{S}_J}, \quad (32)$$

where the Dirac action, $\mathcal{S}_D = \int d^4x \bar{\Psi}(\not{\partial} - m)\Psi$, and the action for the currents, $\mathcal{S}_J = \int d^4x \frac{g^2}{2m} (2J_{\mu\nu} J^{\mu\nu} - J_\mu J^\mu)$, are given in Euclidean space. Clearly, \mathcal{S}_J involves products of four spinors. To analytically treat this model we linearise the action in terms of the currents by introducing the Hubbard-Stratonovich transformation [34]. Indeed, we employ the bosonic degrees of freedom $\mathcal{F} \equiv (a^\mu, b^{\mu\nu})$ (in terms of a 4-vector field a^μ and an anti-symmetric tensor' field $b^{\mu\nu}$) to write

$$e^{-\mathcal{S}_J} = \int \mathcal{D}[\mathcal{F}] e^{\int d^4x \frac{1}{2} (\frac{1}{2} b_{\mu\nu} b^{\mu\nu} - a_\mu a^\mu) + \frac{g}{\sqrt{m}} (b_{\mu\nu} J^{\mu\nu} - a_\mu J^\mu)}. \quad (33)$$

Following [34] we can integrate out the Dirac fermions to find an effective bosonic theory

$$Z_{\text{TTh}} = \int D[a] D[b] e^{-\mathcal{S}_{\text{eff}} + \frac{1}{2} \int d^4x (\frac{1}{2} b_{\mu\nu} b^{\mu\nu} - a_\mu a^\mu)}, \quad (34)$$

where the effective action is defined as

$$S_{\text{eff}}[a_\mu, b_{\mu\nu}] = -\log \det \left(\not{\partial} - m + \frac{g}{\sqrt{m}} \not{\mathcal{F}} \right), \quad (35)$$

where $\not{\partial} = \gamma^\mu \partial_\mu$ and $\not{\mathcal{F}} = \gamma^\mu a_\mu + \gamma^5 [\gamma^\mu, \gamma^\nu] b_{\mu\nu}$. Up to terms of order ∂/m (p/m in momentum space) [34] this effective action can be written as

$$S_{\text{eff}}[a_\mu, b_{\mu\nu}] = -8 \frac{g^2}{(4\pi)^2} \int d^4x \epsilon^{\mu\nu\lambda\sigma} a_\mu \partial_\nu b_{\lambda\sigma}, \quad (36)$$

where $\epsilon^{\mu\nu\lambda\sigma}$ is the Levi-Civita symbol. The correction terms are insignificant in the large-wavelength/low-energy regime we are interested in. By neglecting the irrelevant constants, the partition function for the final theory can be written as

$$\tilde{Z} = \int \mathcal{D}[\mathcal{F}] e^{-\int d^4x \frac{1}{2} (a_\mu a^\mu - \frac{1}{2} b_{\mu\nu} b^{\mu\nu}) + \frac{g^2}{2} \epsilon^{\mu\nu\rho\alpha} a_\mu \partial_\nu b_{\rho\alpha}}. \quad (37)$$

We note that the fields a and b have dimension (Length)⁻². We could be tempted to consider the field a as a sort of electromagnetic field and b as a sort of curvature field. Unfortunately, such an interpretation is not obvious at this stage. In fact, the theory is not invariant under the gauge-like transformation [35]

$$\begin{cases} a_\mu & \rightarrow a_\mu + \partial_\mu \chi \\ b_{\mu\nu} & \rightarrow b_{\mu\nu} + \partial_\mu \xi_\nu - \partial_\nu \xi_\mu, \end{cases} \quad (38)$$

where χ and ξ_μ are a scalar and a vector field respectively. In fact, the kinetic terms $a_\mu a^\mu$ and $b_{\mu\nu} b^{\mu\nu}$ explicitly break invariance under these transformations (as, for example, the vector potential appears explicitly). Hence, the partition function \tilde{Z} describes a massive spin-1 theory that does not allow easy interpretations. We would like to recast this theory in a more suitable form given in terms of a “vector potential” and a “curvature” field, which naturally leads to next sections’ topic. This process is analogous to the (2+1)-dimensional one where a duality between a selfdual free massive field theory and a topologically massive theory [36] has been demonstrated. As a final note, it is important to stress that it is not possible to apply the bosonization procedure proposed here to free Dirac fermions, i.e. without the presence of the current-current interactions in Eq. (29). This means that we cannot tune the parameter g^2 to zero without encountering non-analytical points, which justifies the presence of the factor $1/g^2$ in Eq. (40). Physically, the naive replacement $g = 0$ into the initial (Eq. (29)) and final (Eq. (37)) theories would lead to a mapping between *free* fermionic degrees of freedom into *free* bosonic ones, which is clearly forbidden by the statistics of the fields involved. The presented “transmutation” of degrees of freedom holds only for interacting theories ($g \neq 0$), as for example happens in superconductivity where the interaction between electrons in a metal leads to a physics described by bosonic degrees of freedom in terms of Cooper pairs [37].

C. Duality

In order to recast the theory defined in Eq. (37) in a more suitable form, one can employ a BFT quantization procedure [38, 39] to show the equivalence of the massive spin-1 theory

$$L = -\frac{1}{4} b_{\mu\nu} b^{\mu\nu} + \frac{1}{2} a_\mu a^\mu + \frac{g^2}{2} \epsilon_{\mu\nu\lambda\sigma} b^{\mu\nu} \partial^\lambda a^\sigma, \quad (39)$$

to one involving an “electromagnetic” field A^μ and a so called Kalb-Ramond field $B^{\mu\nu}$ [40]. The two theories can in fact be embedded in the same enlarged theory from which they descend as different choices of gauge fixing [41, 42]. The resulting theory, in the Lorentzian signature, is described by the Cremmer-Scherk Lagrangian [25]

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{12} H_{\mu\nu\lambda} H^{\mu\nu\lambda} + \frac{1}{4g^2} \epsilon_{\mu\nu\lambda\sigma} B^{\mu\nu} F^{\lambda\sigma}, \quad (40)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and $H_{\mu\nu\lambda} = \partial_\mu B_{\nu\lambda} + \partial_\nu B_{\lambda\mu} + \partial_\lambda B_{\mu\nu}$. The field A is an electromagnetic field (with dimension (Length)⁻¹) while the field B (with dimension (Length)⁻¹) is the so called Kalb-Ramond field. As above, we can define the symmetry transformation

$$\begin{cases} A_\mu & \rightarrow A_\mu + \partial_\mu \chi \\ B_{\mu\nu} & \rightarrow B_{\mu\nu} + \partial_\mu \xi_\nu - \partial_\nu \xi_\mu. \end{cases} \quad (41)$$

Contrary to the theory described in Eq. (37), this one is explicitly invariant under this “gauge” transformation. Let us now take some time to analyze the terms appearing in the action in Eq. (40). The first two kinematic terms are geometric (the metric appears explicitly) while the last one has a topological nature and it is the standard BF term. This theory is a topological massive gauge theory in 3 + 1 dimensions [25, 26] and represents the natural abelian generalization of Chern Simons-Maxwell theory in 2 + 1 dimensions as the Chern Simons theory cannot exist in 3 + 1 dimensions. This way of generating mass for the electromagnetic field (through a topological interaction) is an alternative to the Higgs mechanism and can in fact be connected to superconducting phenomena [43, 44]. The theory is renormalizable [26] and explicitly gauge invariant in the bulk.

D. Bosonization Rules

The possibility to map the Tensorial Thirring model to a massive gauge theory does not come as a surprise. In fact, in the (1 + 1)-dimensional case the massive Thirring model is equivalent to the sine-Gordon massive scalar theory [45], while in (2 + 1) dimensions is equivalent to the Maxwell-Chern-Simons theory [24], where, the Maxwell field acquire a mass through a topological mechanism. Motivated by the analogies with the low-dimensional cases we propose the natural generalization

of the *bosonization rules* to the three-dimensional case. These rules connect the degrees of freedom of the equivalent fermionic and bosonic theories (up to multiplicative factors) in the following way

Dimensions	Theory	Bosonization Rules
1+1	sine-Gordon	$J^\mu \rightarrow \epsilon^{\mu\nu} \partial_\nu \phi$
2+1	Maxwell-CS	$J^\mu \rightarrow \epsilon^{\mu\nu\lambda} \partial_\nu A_\lambda$
3+1	Cremmer-Scherk	$\begin{cases} J^\mu \rightarrow \epsilon^{\mu\nu\lambda\gamma} \partial_\nu B_{\lambda\gamma} \\ J^{\mu\nu} \rightarrow \epsilon^{\mu\nu\lambda\gamma} \partial_\lambda A_\gamma \end{cases}$

Let us take a little more time to emphasize the analogies with the lower dimensional cases and get some more intuitions on the bosonization procedure. We can note that the Thirring model is always equivalent to some massive theory. In the (1+1) dimensional case the equivalent theory is a sine-Gordon massive scalar theory. The equivalence with the Thirring model has been shown by Coleman [45] (see also [46] for extension of the proof to the finite temperature case). In (2+1) dimensions the Thirring model has been proven by Fradkin and Schaposnik to be equivalent to a Maxwell-Chern-Simons theory. The proof relies on a dualization procedure first showed by Deser and Jackiw [36] and the equivalent theory corresponds to a massive gauge theory where the mass of the photon is given thanks to the interaction with a topological Chern-Simons term (an alternative to the Higgs procedure to give mass to a gauge theory). In 3+1 dimensions the Tensorial Thirring model we are studying is going to be equivalent to a so called Cremmer-Scherk model [25] where a Maxwell theory is coupled to a Kalb-Ramond field [40] thanks to a BF term. This is, in analogy with the (2+1)-dimensional case, a massive gauge theory where the mass comes from the topological interaction with the Kalb-Ramond field. Note that, in this case, we need two fields since we are considering two different types of current-current interactions in the fermionic model. These two fields are necessary to generate mass for the gauge theory in a topological fashion [59]. We also note that similar bosonization rules in $d+1$ dimensions were proposed in [47].

E. Pure Topological Regime

From now on we work in a regime where the contribution of the topological BF term in (40) is dominant, i.e. we want to work with energy scales much smaller than $1/g^2$. Intuitively, this suggests that the Maxwell and Kalb-Ramond field have small kinetic energy compared to their (topologically) acquired mass. In such a regime we are sufficiently close to the ground state and the important contributions to the effective theory come from the topological BF term

$$S_{\text{BF}} = \frac{1}{4g^2} \int_{\mathcal{M}} d^4x \epsilon_{\mu\nu\lambda\sigma} B^{\mu\nu} F^{\lambda\sigma}, \quad (42)$$

where \mathcal{M} is the spacetime manifold associated with our theory.

1. Boundary Behaviour

We now consider the behaviour of our lattice model at its physical boundary. Several approaches are possible based, for example, on the Symanzik method [48] or on gauge invariance analysis [14, 49]. Focusing on the latter at the bosonic level a BF theory defined on a non-compact space, \mathcal{M} , is not manifestly gauge invariant due to contributions from the boundary, $\partial\mathcal{M}$. Restoring gauge invariance generates a (2+1)-dimensional BF theory on the boundary, while leaving the bulk theory unchanged [14, 49]. Here, we show that the bosonization rules give rise to exactly the same theory on the boundary of the (3+1)-dimensional fermionic lattice model (for more detail we refer to Appendix B). We start by introducing a minimal coupling between the tight-binding fermions and a pure gauge $U(1)$ field $\hat{A}_\mu = \partial_\mu \varphi$ parameterised by φ . This coupling extends (42), in the continuum limit, by

$$S_\varphi = \int_{\mathcal{M}} d^4x J^\mu \partial_\mu \varphi, \quad (43)$$

but it leaves the physics of the model unchanged. We can now employ the bosonization rule $J^\mu \rightarrow \epsilon^{\mu\nu\lambda\gamma} \partial_\nu B_{\lambda\gamma}$, together with Stokes' theorem and an integration by parts to show that

$$S_\varphi = \int_{\partial\mathcal{M}} d^3x \phi \epsilon_{\mu\nu\lambda} \partial^\mu B^{\lambda\nu}, \quad (44)$$

where here (and throughout the rest of the paper for integrations on the boundary) the indices run through the coordinates that parameterise $\partial\mathcal{M}$. The field ϕ can now be interpreted as a Lagrange multiplier enforcing the condition $dB = 0$ on $\partial\mathcal{M}$. This implies that, locally, $B_{\mu\nu} = \partial_\mu \eta_\nu - \partial_\nu \eta_\mu$, which conveniently implies $S_\varphi = 0$. This means that the possibility to add S_φ to our action is equivalent to the constraint $dB = 0$ on $\partial\mathcal{M}$. We are now ready to find the effective action on the boundary. In fact, we can rewrite the right hand-side of Eq. (42) as

$$\frac{1}{4g^2} \left(\int_{\partial\mathcal{M}} d^3x \epsilon_{\mu\nu\lambda} B^{\mu\nu} A^\lambda - \int_{\mathcal{M}} d^4x \epsilon_{\mu\nu\lambda} \partial^\mu B^{\nu\lambda} A^\sigma \right), \quad (45)$$

which, by restriction on the boundary, implies the following form for the theory on the boundary

$$S_{\partial\mathcal{M}} = \frac{1}{4g^2} \int_{\partial\mathcal{M}} d^3x \epsilon_{\mu\nu\lambda} \eta^\mu \partial^\nu A^\lambda, \quad (46)$$

This is indeed a (2+1)-dimensional BF theory. It is equivalent to a double Chern-Simons theory that describes time-reversal symmetric physics on the boundary.

2. Physical Observables

We now want to identify physical observables associated with the purely topological part, S_{BF} . Gauge invariant observables of the $(3+1)$ -dimensional BF theory are given by expectation values of Wilson *surface* operators [50, 51], which are a generalization of the $(2+1)$ -dimensional Wilson loop operators. These observables, $W_B = \langle e^{\frac{i}{g^2} \int_{\partial\Sigma} B} \rangle$, are defined for any two-dimensional boundary $\partial\Sigma$ of a three-dimensional volume Σ , where B is the Kalb-Ramond field [40]. The corresponding fermionic observables are given by $W_\Psi = \langle e^{iq \int_\Sigma d^3x \Psi^\dagger \Psi} \rangle$, where q is a generic charge of the (string-like) excitations associated with the field B . The correspondence is easily proven by an opportunely manipulation of the Noether charge $Q = \int_\Sigma J^0 d^3x$ (where $J^0 = q\Psi^\dagger \Psi$). The joint use of the bosonization rule $J^0 = \frac{1}{g^2} \epsilon^{ijk0} \partial_\nu B_{ij}$ (where the constant g^2 has been introduced for dimensional reasons) and Stokes' theorem immediately leads to $Q = \frac{1}{g^2} \int_{\partial\Sigma} B$. This proves that $q \int_\Sigma d^3x \Psi^\dagger \Psi = \frac{1}{g^2} \int_{\partial\Sigma} B$ (where any proportionality constant implicit in definition of the bosonization rules is absorbed inside the charge q) or, in other words

$$W_B = W_\Psi. \quad (47)$$

For more details on this proof we refer to Appendix C 1. It is a well known fact that $W_B = 1$, identically [12]. Indeed, one can explicitly confirm (see Appendix C 2) that

$$\frac{1}{g^2} \int_{\partial\Sigma} B = 2\pi n, \quad n \in \mathbb{Z}, \quad (48)$$

for all permissible configurations of B . This implies that the charge $q \int_\Sigma d^3x \Psi^\dagger \Psi$ inside a volume Σ takes discrete values. While this condition gives, as expected, trivial values for the observable W_Ψ it can be employed to distinguish between trivial (product) states and topologically ordered ones [21]. Indeed, product states correspond to a fixed value of n for a given Σ , while the highly correlated ones can give different values at each measurement. These values of n are experimentally accessible by measuring fermion populations on the vertices of the tight-binding model that are inside Σ .

IV. CONCLUSIONS

In summary, we presented a tight-binding model of spinless fermions that has a variety of behaviours. In the absence of interactions it generalises the methodology employed in the $(2+1)$ -dimensional Haldane model to the $(3+1)$ -dimensional case giving a chiral topological insulator. In the presence of interactions it gives rise, in the continuum limit, to the $(3+1)$ -dimensional BF theory accompanied by a Maxwell term. Our model can be tuned to be in the topological (BF) or the non-topological

(Maxwell) regimes, thus being of relevance to both condensed matter and high energy physics. The versatile method we presented for detecting the topological character of the model can become a powerful diagnostic tool for experimentally probing the topological properties of three-dimensional systems.

Acknowledgements.— This work was supported by EPSRC and the ARC via the Centre of Excellence in Engineered Quantum Systems (EQuS), project number CE110001013. We would like to thank G.K. Brennen for useful discussions and support.

APPENDIX A: DISCRETE SYMMETRIES

In this appendix we briefly analyze the definitions of the three symmetries used to study the model given in Eq. (1). Following the main text, throughout this appendix we restrict to translationally invariant spinless fermionic systems.

1. Time Reversal Transformation

Time reversal transformations are associated with the inversion of time. From a physical point of view we want to address whether the system distinguishes a time direction or not. More precisely, given a certain Hamiltonian H and a solution $\Psi(t)$ of the Schrödinger equation $H\Psi(t) = i\partial_t\Psi(t)$ we want to know if there also exist a solution $\Psi'(-t)$ of the equation $H\Psi'(-t) = i\partial_{-t}\Psi'(-t)$. Let us define the (antiunitary) time reversal operator T by its action on $\Psi(t)$ as $T\Psi(t) = \Psi'(-t)$.

We define a system to be time reversal symmetric if such an operator T exists and it satisfies the equation $HT\Psi(t) = i\partial_{-t}T\Psi(t)$, where $\Psi(t)$ is known to be a solution of the Schrödinger equation $H\Psi(t) = i\partial_t\Psi(t)$. This condition is equivalent to impose $T^{-1}HT\Psi(t) = T^{-1}iT\partial_{-t}\Psi(t)$ which is satisfied if

$$\begin{cases} T^{-1}iT = -i \\ T^{-1}HT = H \end{cases}. \quad (A1)$$

The first condition tells us that the operator T must be antiunitary while the second can be viewed as a restriction for the Hamiltonian. Given these two conditions, a solution $\Psi(t)$ of the Schrödinger equation $H\Psi(t) = i\partial_t\Psi(t)$ implies that $T^{-1}HT\Psi(t) = T^{-1}iT\partial_t\Psi(t)$ which in turn implies $HT\Psi(t) = i\partial_{-t}\Psi(t)$, that is $T\Psi(t)$ satisfies the Schrödinger equation with reversed time. The operator T can be written as $T = T_U K$ where K is the complex conjugation operator and T_U a generic unitary operator. From this, it is easy to show that

$$\begin{cases} TT^\dagger = \mathbb{I} \\ T^T = T \end{cases}, \quad (A2)$$

since we have $TT^\dagger = T_U K U^T K = U U^\dagger = \mathbb{I}$ and $T^T = K U^T = K U^* = U K = T$.

In the case of spinless fermions, the operator T_U can be chosen to be the identity, so that $T^{-1}f_{\vec{r}}^\dagger T = f_{\vec{r}}^\dagger$ for every generic fermion operator $f_{\vec{r}}$ labeled by its position \vec{r} . The action on the Fourier transformed fermion operator $a_{\mathbf{p}}$ is easily found to be $T^{-1}a_{\mathbf{p}}T = \sum_{\vec{r}} T^{-1}e^{i\mathbf{p}\cdot\vec{r}}Ta_{\vec{r}} = \sum_{\vec{r}} e^{-i\mathbf{p}\cdot\vec{r}}a_{\vec{r}} = a_{-\mathbf{p}}$. Basically, the time reversal operator maps a particle with momentum \mathbf{p} to a particle with momentum $-\mathbf{p}$. The time reversal action on the Hamiltonian kernel $\bar{H}(\mathbf{p})$ in momentum space follows from $T^{-1}HT = \sum_{\mathbf{p}} f_{-\mathbf{p}}^\dagger \bar{H}^*(\mathbf{p})f_{-\mathbf{p}}$, where $*$ denotes the complex conjugation which is introduced accordingly to the first of Eqs. (A1). We then just saw that time reversal induces an action \bar{T} on the Hamiltonian kernel given by $\bar{T}\bar{H}(\mathbf{p})\bar{T}^\dagger = \bar{H}^*(-\mathbf{p})$ (with \bar{T} unitary such that $\bar{T} = \bar{T}^\dagger$). Invariance under time reversal is then equivalent of the request

$$\bar{H}^*(-\mathbf{p}) = \bar{H}(\mathbf{p}) . \quad (\text{A3})$$

Note that the time invariance operator for spinless particles is just complex conjugation so that $T^2 = \mathbb{I}$.

2. Charge Conjugation Transformation

The charge conjugation transformation does not involve any action on spatial or temporal coordinates. Let us define the charge conjugation operator in momentum space following the analysis in [52]. We impose that, under the action of the (unitary) charge conjugation operator C the operator that annihilates a particle f transforms in the operator that annihilates an antiparticle f' as $C^{-1}f_{\mathbf{p}}C = f'_{\mathbf{p}}$. Incidentally, from this definition and from the linearity of the operator we can derive the action in real space: $C^{-1}f_{\vec{r}}C = C^{-1}\sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\vec{r}}f_{\mathbf{p}}C = \sum_{\vec{r}} e^{i\mathbf{p}\cdot\vec{r}}f'_{\vec{r}} = f'_{\vec{r}}$. In our case we identify the antiparticle with a hole imposing that $f'_{\mathbf{p}} = f_{-\mathbf{p}}^\dagger$, to finally get: $C^{-1}f_{\mathbf{p}}C = f_{-\mathbf{p}}^\dagger$, where the transpose operator has been introduced to match the notation used so far where creation (annihilation) operators are accommodated in a row (column) vector. The action on the Hamiltonian kernel is given by $:C^{-1}HC: = :C^{-1}\sum_{\mathbf{p}} f_{\mathbf{p}}^\dagger \bar{H}(\mathbf{p})f_{\mathbf{p}}C: = \sum_{\mathbf{p}} :f_{-\mathbf{p}}^\dagger C^{-1}\bar{H}(\mathbf{p})C f_{-\mathbf{p}}^\dagger: = -\sum_{\mathbf{p}} f_{-\mathbf{p}}^\dagger \bar{H}(\mathbf{p})^T f_{-\mathbf{p}}$, where $:$ indicates the normal ordering operator (which imposes creation operators to be on the left of annihilation ones) and where the minus sign takes into account the fermionic statistics. Charge conjugation then implies an action \bar{C} on the Hamiltonian kernel given by $\bar{C}\bar{H}(\mathbf{p})\bar{C}^\dagger = \bar{H}(-\mathbf{p})^*$ (with \bar{C} unitary and such that $\bar{C} = \bar{C}^\dagger$).

A system is defined to be invariant under charge conjugation if $:C^{-1}HC := :H:$ which implies

$$\bar{H}(-\mathbf{p})^* = -\bar{H}(\mathbf{p}) , \quad (\text{A4})$$

as one can see by comparing the expression given above for $:C^{-1}HC:$ and the expression for the Hamiltonian

in momentum space (and taking into account that the Hamiltonian is hermitian). This is in accordance with the intuition that, if we want to treat particles and holes in a symmetric fashion the Hamiltonian has to change sign when transforming all the particles into holes and viceversa.

We also note that the charge conjugation operator for spinless particles is just complex conjugation so that $C^2 = \mathbb{I}$. This condition, together with the unitarity one implies

$$\begin{cases} CC^\dagger = \mathbb{I} \\ C^\dagger = C \end{cases} , \quad (\text{A5})$$

It is worth noticing that the nomenclature for the charge conjugation symmetry (and its relation with particle-hole symmetry) is not uniform in the literature, as pointed out in, for example, [53].

3. Chiral Symmetry

We define a system to have chiral symmetry if there exist a unitary matrix \bar{C}_s that anticommutes with the Hamiltonian kernel in momentum space [1, 31]

$$\bar{C}_s \bar{H}(\mathbf{p}) = -\bar{H}(\mathbf{p})\bar{C}_s , \quad (\text{A6})$$

and such that $\bar{C}_s^2 = \mathbb{I}$. This immediately implies that, for each eigenfunction $\Psi_{\mathbf{p}}$ with energy $E_{\mathbf{p}}$ there exist an eigenfunction $\bar{C}_s \Psi_{\mathbf{p}}$ with energy $-E_{\mathbf{p}}$, since $\bar{H}(\mathbf{p})\bar{C}_s \Psi_{\mathbf{p}} = -\bar{C}_s \bar{H}(\mathbf{p})\Psi_{\mathbf{p}} = -E_{\mathbf{p}}\bar{C}_s \Psi_{\mathbf{p}}$. In the context of this thesis, chiral symmetry reflects a particular structure of the lattice. In fact, a sufficient condition for the existence of chiral symmetry is the possibility to colour the lattice such that two vertices of the same colour do not have a common link. This property is known as bi-colourability. In this case it is clear that the Hamiltonian kernel can be written in a block off-diagonal form $\begin{pmatrix} 0 & \cdot \\ \cdot & 0 \end{pmatrix}$ which implies the anticommutation with σ_z .

Another sufficient condition for the presence of chiral symmetry is the existence of both time reversal and charge conjugation symmetries. In this case we can define an (antiunitary) operator $C_s = TC$ whose action on the Hamiltonian kernel is given by (see sections above) $\bar{H} \rightarrow \bar{C}_s \bar{H} \bar{C}_s^\dagger$, where $\bar{C}_s = \bar{T} \cdot \bar{C}$. The existence of both time and charge conjugation symmetries implies that the operator \bar{C}_s anticommutes with the Hamiltonian since $\bar{T} \cdot \bar{C}\bar{H}(\mathbf{p}) = -\bar{T}\bar{H}^*(-\mathbf{p})\bar{C} = -\bar{H}(\mathbf{p})\bar{T} \cdot \bar{C}$. One can take this as a definition of the symmetry (which, since it combines time reversal and charge conjugation, it can be named particle-hole symmetry as it maps a particle with momentum \mathbf{p} to a hole with momentum $-\mathbf{p}$).

APPENDIX B: BEHAVIOUR ON THE BOUNDARY

In this appendix we study in detail how to obtain the effective theory describing the boundary of our material. Throughout this appendix we will use the differential forms formalism [54]. In this language, the bosonic fields introduced in section IIIB consist of a 1-form A and a 2-form B .

Given their importance for what follows we re-write here the bosonization rules connecting the fermionic microscopic degrees of freedom and the bosonic effective ones

$$\begin{cases} J^\mu & \rightarrow \frac{1}{g^2} \epsilon^{\mu \nu \lambda \gamma} \partial^\nu B^{\lambda \gamma} \rightarrow \frac{1}{g^2} * dB \\ J^{\mu\nu} & \rightarrow \frac{1}{g^2} \epsilon^{\mu\nu \lambda \gamma} \partial^\lambda A^\gamma \rightarrow \frac{1}{g^2} * dA \end{cases}, \quad (\text{B1})$$

where $*$ denotes the Hodge operator [54] and where we retain the correct dimensions through the coupling g^2 .

We now introduce an example of a procedure to obtain the theory on the boundary as proposed in [14, 49]. This approach relies on restoring the gauge invariance for the theory on the boundary. We then propose a procedure specific to the model presented here. The advantage of this procedure is that it does not require any physical hypothesis on the system.

1. Example: how to restore gauge invariance on the boundary

We start from the BF theory defined in Eq. (42)

$$S_{\text{BF}} = \frac{1}{4g^2} \int_{\mathcal{M}} B \wedge F, \quad (\text{B2})$$

Let us begin by showing that the theory is not gauge invariant on the boundary. The gauge transformation considered here is the one in Eq. (41) that is

$$\begin{cases} A & \rightarrow A + d\chi \\ B & \rightarrow B + d\xi \end{cases}, \quad (\text{B3})$$

where χ is a function and ξ is a 1-form. It is easy to see that, when we add a boundary ∂M to the manifold M , the theory keeps the invariance under the gauge transformations in Eq. (41) of A alone as only the gauge invariant quantity F (because of the properties of the exterior derivative [54] which imply that $d^2\chi = 0$) appears in the action. Unfortunately, it loses the invariance under the generalised gauge transformations for the B term (see Eq. (41)). In fact under such a transformation the action changes as

$$S \rightarrow S + \Delta S, \quad (\text{B4})$$

where

$$\Delta S = \int_M d\xi \wedge F = \int_{\partial M} \xi \wedge F, \quad (\text{B5})$$

where in the last step we used integration by parts, Stokes theorem and the abelian Bianchi identity $dF = d^2A = 0$. We now want to modify the original action to restore gauge invariance on ∂M .

Following [49] and [14] (see also [44]) we now add a boundary term $\int_{\partial M} B \wedge A$ to the action so that $S' = S + \int_{\partial M} B \wedge A$. This solves the gauge invariance problem for the field B as easily shown with the following

$$\begin{aligned} \Delta S' &= \Delta S + \int_{\partial M} d\xi \wedge A \\ &= \int_{\partial M} \xi \wedge F - \int_{\partial M} \xi \wedge F + \int_{\partial M} d(\xi \wedge A) \\ &= 0, \end{aligned} \quad (\text{B6})$$

where in the last equality we used the Stokes theorem together with the fact that $\partial\partial M = 0$. Note that we now have broken the gauge invariance under the transformation on A as can be easily seen by simple inspection of the additional term $\int_{\partial M} B \wedge A$ which is explicitly dependent on the (gauge) field A . In order to restore full gauge invariance, we introduce a new scalar field ϕ with the following transformation properties

$$\phi \rightarrow \phi - \chi, \quad (\text{B7})$$

where χ is the same function appearing in the transformation rule for A . We now notice that if we redefine $A \rightarrow A' = A + d\phi = D\phi$ we get $\Delta A' = d\chi - d\chi = 0$. We then can define a final gauge invariant action as

$$S_{\text{tot}} = \int_M B \wedge F' + \int_{\partial M} B \wedge A'. \quad (\text{B8})$$

Explicitly, the total action is

$$\begin{aligned} S_{\text{tot}} &= \int_M B \wedge F' + \int_{\partial M} B \wedge A' \\ &= \int_M B \wedge dA' + \int_{\partial M} B \wedge A' \\ &= \int_M B \wedge (dA + d^2\phi) + \int_{\partial M} B \wedge (A + d\phi). \end{aligned} \quad (\text{B9})$$

Notice that we have modified the action only on the boundary and that the additional term breaks time reversal symmetry (B is even for time reversal since it is a sort of electric field, while A is odd) and it is in fact odd under such symmetry if we impose that $\phi \rightarrow -\phi$ under time reversal.

What is the role of the new field ϕ in our theory? This field is actually not a dynamical one. We can see this by using its equation of motion. Let us start by computing $\delta_\phi S_{\text{tot}}$. We have

$$\begin{aligned} \delta_\phi S_{\text{tot}} &= \int_{\partial M} B \wedge \delta d\phi \\ &= \int_{\partial\partial M} d(B \wedge \delta\phi) - \int dB \wedge \delta\phi \\ &= - \int dB \wedge \delta\phi \end{aligned} \quad (\text{B10})$$

The equation of motion for the field ϕ is given by $\delta_\phi S_{\text{tot}} = 0$ which (from Eq. (B10)) is fulfilled if $dB = 0$ on the boundary ∂M . In fact, the field ϕ is nothing but a Lagrange multiplier enforcing the constraint

$$dB = 0 \quad \text{on } \partial M. \quad (\text{B11})$$

We can now suppose that the boundary for our system is $\partial M = \mathbb{R} \times \Sigma$ where the spatial manifold Σ is topologically equivalent to S^2 . Otherwise stated, our boundary is a sphere embedded in the space. The constraint in Eq. (B11) is telling us that B is a closed 2 form on this boundary. Since the second de Rham cohomology class on ∂M is non-trivial we can say that $B = d\eta$ only locally on ∂M (i.e. B is a pure gauge there) so that the total action is

$$S_{\text{tot}} = \int_M B \wedge F + \int_{\partial M} d\eta \wedge A' , \quad (\text{B12})$$

where the boundary term is local on the boundary. We can then conclude that we have (locally) a BF theory on the boundary ∂M . The kinetic terms $F \wedge *F$ and $H \wedge *H$ on the boundary are not topological and they are of no interest in this matter (see also [44]).

Incidentally, we notice that this method of imposing gauge invariance on the boundary is not the only possible one. In fact one can as well change the transformation rules of the B field imposing that it has to transform trivially $B \mapsto B$ on the boundary (that is: B is a true ‘‘curvature’’ form on the boundary). In this case we can avoid adding the term $\int B \wedge A$.

2. BF theory on the boundary

In the previous example, the existence of a BF theory on the boundary was proved by invoking additional terms on the boundary (which also involve a new scalar field), justified by the requirement of gauge invariance. In this section we want to closely follow this procedure. Specifically, we want to add a scalar field in order to impose $dB = 0$ on the boundary ∂M . The main question we want to address is: can we justify the addition of such a field? We will find a positive answer as a consequence of the bosonization rules given in Eq. (B1).

Let us begin by introducing a pure gauge electromagnetic field $A_\phi = d\phi$ in the fermionic tight binding model (we stress that we do not actually require the field A_ϕ in the system but we introduce it as a pure gauge only to prove that $dB = 0$ on ∂M). The interaction can be imposed through minimal coupling of the fermionic current to the field A_ϕ . This implies the addition of a term $J_\mu A_\phi^\mu$ to the microscopic action. Such a term, manipulated through the bosonization rules (thanks to Stokes theorem) gives

$$\begin{aligned} \int_M J_\mu A_\phi^\mu &= \int_M *J \wedge A_\phi \\ &= \int_M dB \wedge d\phi \\ &= \int_M d(B \wedge d\phi) \\ &= \int_{\partial M} B \wedge d\phi . \end{aligned} \quad (\text{B13})$$

This shows the expression for the additional term to be added to the effective theory as a consequence of the minimal coupling. We can notice that this additional term contains a scalar field ϕ . Compared to the example given in the previous section the introduction of this term is

now naturally arising from the minimal coupling of the microscopic theory with a pure gauge degree of freedom. The possibility of this result is given by the bosonization rules present in our analysis. This pure gauge is totally arbitrary and does not change the physics of the model. We can then treat this field as being a Lagrange multiplier enforcing the condition $dB = 0$ on the boundary as explained in the example above. We then have $B = d\eta$ on ∂M and we can write the total action for our theory as

$$S_{\text{tot}} = \int_M B \wedge F + \int_{\partial M} d\eta \wedge d\phi . \quad (\text{B14})$$

We can now notice that the second term in this expression (the one coming from the minimal coupling) is actually zero since $\int_{\partial M} d\eta \wedge d\phi = \int_{\partial M} d(\eta \wedge d\phi) = \int_{\partial \partial M} \eta \wedge d\phi = 0$ (where we used Stokes theorem and the fact that $\partial \partial M = 0$). Basically, the gauge field ϕ ‘‘lives’’ just enough to impose the constraint $dB = 0$ on ∂M before dying without leaving any trace! The action is then given by

$$\begin{aligned} S_{\text{tot}} &= \int_M B \wedge F \\ &= \int_M B \wedge dA \\ &= \int_M d(B \wedge A) - \int_M dB \wedge A \\ &= \int_{\partial M} B \wedge A - \int_M dB \wedge A . \end{aligned} \quad (\text{B15})$$

Since $dB = 0$ on the boundary we have that on ∂M the value of the action is just

$$S_{\partial M} = \int_{\partial M} B \wedge A , \quad (\text{B16})$$

and, locally

$$S_{\partial M} = \int_{\partial M} d\eta \wedge A . \quad (\text{B17})$$

This is indeed a $(2 + 1)$ -dimensional BF theory and proves Eq. (46) in the main text. It is equivalent to a double Chern-Simons theory that describes time-reversal symmetric physics on the boundary.

In summary, we have seen that our model is equivalent to one which has a theory on the boundary with a topological BF term. Notice that, without the term $\int B \wedge A$ (described in section B1) the introduction of the scalar field done in this section is not enough to restore the gauge invariance on the boundary. More precisely, the theory in Eq. (B17) is invariant for gauge transformations involving the field A alone but not for ones involving also the field B .

APPENDIX C: OBSERVABLES

It is possible [50, 55] to define observables which witness the topological nature of the BF theory describing the low energy physics of the model. These observables are defined as expectation values of Wilson surface operators

$$\langle e^{i \frac{q}{g^2} \int_{\partial \Sigma} B} \rangle , \quad (\text{C1})$$

where the surface $\partial\Sigma$ is defined as the boundary of a generic three-dimensional spatial manifold Σ and where q is a generic charge of the string-like (since the two-form field $B_{\mu\nu}$ couples with tensorial currents $J^{\mu\nu}$ and not to the usual vector currents J^μ as the electromagnetic field A^μ) excitations associated with the field B . In [55] it is shown that such an expectation value is trivial and equal to 1, for the (3+1)-dimensional case considered here. In fact, it represents the trivial case in which the surface $\partial\Sigma$ does not intersect any other surface leading to a null linking number. Explicitly we have

$$\langle e^{i\frac{q}{g^2} \int_{\partial M} B} \rangle = 1 . \quad (C2)$$

In this appendix we will check this result for our specific model and we will describe how to map this observable to a microscopic fermionic observable.

1. Effective Noether charge as a topological number

We begin by noticing that the Wilson surface observable can be written as an *effective* Noether charge. The effective Noether charge can be written as a function of the zeroth component of the Noether fermionic current as

$$Q = \int_{\Sigma} J^0 d^3x . \quad (C3)$$

The general expression for the current is obtained by the bosonization rule as

$$\begin{aligned} J &= \frac{1}{g^2} *dB \\ &= \frac{1}{g^2} \partial_\lambda B_{\mu\nu} * [dx^\lambda \wedge dx^\mu \wedge dx^\nu] \\ &= \frac{1}{g^2} \partial_\lambda B_{\mu\nu} \epsilon^{\lambda\mu\nu} dx^\rho , \end{aligned} \quad (C4)$$

where we used

$$\begin{cases} B &= B_{\mu\nu} dx^\mu \wedge dx^\nu \\ dB &= \partial_\lambda B_{\mu\nu} dx^\lambda \wedge dx^\mu \wedge dx^\nu . \end{cases} \quad (C5)$$

Eq. (C4) directly leads to the expression for the zeroth component of the Noether current

$$\begin{aligned} J_0 &= \frac{1}{g^2} \partial_\lambda B_{\mu\nu} \epsilon^{\lambda\mu\nu} \\ &= \frac{1}{g^2} \partial^i B^{jk} \epsilon_{ijk0} . \end{aligned} \quad (C6)$$

Any proportionality constant left implicit in the bosonization rules can simply be used to rescale the parameter g^2 . We now note that this is the same expression in coordinates as the exterior differential in the space dimensions of the form $\varphi_t^*(B_{\mu\nu} dx^\mu dx^\nu)$ where φ_t^* denotes the pull back [54] of the form B in a constant time slice of spacetime under the map $\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ given by $\{\vec{x}\} \mapsto \{\vec{x}, t\}$. We in fact simply have

$$\begin{aligned} d(\varphi_t^*(B_{\mu\nu} dx^\mu dx^\nu)) &= d(B_{ij} dx^i dx^j) \\ &= \partial_k B_{ij} dx^i dx^j dx^k \\ &= \partial_k B_{ij} \epsilon^{kij} dx^1 dx^2 dx^3 \\ &= g^2 J_0 d^3x . \end{aligned} \quad (C7)$$

We can finally write the expression for the effective Noether charge as

$$\begin{aligned} Q &= \int_{\Sigma} J^0 d^3x \\ &= \frac{1}{g^2} \int_{\Sigma} *dB \\ &= \frac{1}{g^2} \int_{\Sigma} d(\varphi_t^*(B_{\mu\nu} dx^\mu dx^\nu)) . \end{aligned} \quad (C8)$$

Since $J^0 = q\Psi^\dagger\Psi$ for a given charge q , we can now use these results to identify observables for the effective topological theory with fermionic physical observables as

$$e^{\frac{i}{g^2} \int_{\partial\Sigma} B} = e^{iq \int_{\Sigma} \Psi^\dagger \Psi} . \quad (C9)$$

2. Check of quantization of the effective Noether charge

In this subsection we further analyze the left hand side of Eq. (C9) in order to check the validity of Eq. (C2). For simplicity, we start by rescaling the field $\frac{1}{g^2}B \rightarrow B$. We can consider the embedding $\varphi_\Sigma : \partial\Sigma \rightarrow \mathbb{R}^3$ of the two dimensional manifold $\partial\Sigma$ in \mathbb{R}^3 and include it in the definition of Q . In fact, such an embedding induces a pull-back map φ_Σ^* [54] which takes differential forms defined in \mathbb{R}^3 to differential forms defined in $\partial\Sigma$. We get

$$Q = \int_{\partial\Sigma} B_{jk} \varphi_\Sigma^* [dx^j \wedge dx^k] . \quad (C10)$$

If we introduce coordinates θ^1, θ^2 on $\partial\Sigma$ and write the pull-back function in coordinates [54], we find

$$\begin{aligned} Q &= \int_{\partial\Sigma} B_{ij} \frac{\partial\varphi_\Sigma^i}{\partial\theta^a} \frac{\partial\varphi_\Sigma^j}{\partial\theta^b} d\theta^a \wedge d\theta^b \\ &= \int_{\partial\Sigma} \tilde{B}_{ab} d\theta^a \wedge d\theta^b , \end{aligned} \quad (C11)$$

with $i, j = 1, 2, 3$ and $a, b = 1, 2$ and where, for simplicity, we defined \tilde{B} as a 2-form living on $\partial\Sigma$ (notice that the above integral can be written as a flux $\int \vec{B} \cdot d\vec{S}$ with $\vec{B}_i = \epsilon_{ijk} B_{jk}$, $d\vec{S}^i = \epsilon_{ijk} \frac{\partial\varphi_\Sigma^j}{\partial\theta^a} \frac{\partial\varphi_\Sigma^k}{\partial\theta^b} d\theta^a \wedge d\theta^b$). Tracking down the definition of \tilde{B} we find that \tilde{B} is obtained by after applying two pullback on B : $\tilde{B} = \varphi_\Sigma^* \varphi_t^* B$.

Eq. (C11) tells us that we have to compute the surface integral of a two form. In a two-dimensional space every spatial two-form is always closed, that is $d_{\text{space}} \tilde{B} = 0$. In our case we are interested in $\partial\Sigma = S^2$ so that the second de Rham cohomology group is not trivial allowing us to conclude that B is exact only locally on the sphere. We can define two patches of the sphere called N and S respectively around the north and south pole. We can suppose that the two patches intersect on a closed loop γ (let us say the equator). We can define $\tilde{B} = dA_N$ and $\tilde{B} = dA_S$ on the two patches (let us say: North (N) and South (S)) and write

$$\begin{aligned} Q &= \int_N dA^N + \int_S dA^S \\ &= \int_N (\partial_1 A_2^N + \partial_2 A_1^N) d\theta^1 d\theta^2 \\ &\quad + \int_S (\partial_1 A_2^S + \partial_2 A_1^S) d\theta^1 d\theta^2 . \end{aligned} \quad (C12)$$

We can now use Stokes theorem and write

$$Q = \int_{\gamma} A_N - A_S , \quad (\text{C13})$$

where, as defined above, γ is the common line where the surfaces N and S intersect. The origin of the minus sign lies in the fact that $\partial\Sigma$ has no boundary so that γ has to be taken with different orientations depending if we are integrating on N or S .

How are the two “potentials” A^N and A^S related on γ ? We know that, in general, our theory is invariant under the symmetry $B \mapsto B + d\xi$ where ξ is a 2-form, which in coordinates reads: $B_{\mu\nu} \mapsto B_{\mu\nu} + \partial_{\mu}\xi_{\nu} + \partial_{\nu}\xi_{\mu}$. We now remember, from the analysis given above, that $\tilde{B} = \varphi_{\Sigma}^* \varphi_t^* B$. Since the exterior derivative d commutes with the pullback [54] we have that $\tilde{B} \mapsto B + d\tilde{\xi}$, where $\tilde{\xi} = \varphi_{\Sigma}^* \varphi_t^* \xi$. This transformation has been studied before [56] but its connection with a gauge group is not clear and we will in fact do not suppose any association with a gauge group. Since $B = dA$ the transformation has to act on the potentials A as $A \mapsto A + \xi$, or in coordinates $A_{\mu} \mapsto A_{\mu} + \xi_{\mu}$. We also want the field B to be single valued on γ . This can be imposed by writing the simple looking relation: $B_N = B_S$ (on γ) which leads to

$$dA_N = dA_S = dA_N + d\xi , \quad (\text{C14})$$

so that $d\xi = 0$ on γ . This means that ξ is closed and locally exact ($\xi = d\chi$) on γ and also allows to write

$$A_N - A_S = d\chi , \quad (\text{C15})$$

where χ is a function defined on γ everywhere except for a point. Since we can take a point out of the integral over γ without affecting the value of the integral we can write, from Eq. (C13)

$$Q = \oint_{\gamma} d\chi = \oint_{\gamma} (\vec{\nabla}\chi) \cdot d\vec{\gamma} . \quad (\text{C16})$$

Unfortunately what was done so far did not give us any constraints on the value of the discontinuity in χ around γ . This is a reflection of the fact that we decided (in all generality) not to associate a gauge group to the transformation properties of the field B . Nevertheless, we can still obtain such constraint by invoking the observable nature of the Wilson surface operators $\langle e^{i \int_{\partial\Sigma} B} \rangle$. As such, we do not want these observables to be dependent on some “gauge” choice. In particular, we can always use the arguments given above to show that every transformation of the fields implies: $\langle e^{i \int_{\partial\Sigma} B} \rangle \mapsto \langle e^{i \int_{\partial\Sigma} B} \rangle e^{i \oint_{\gamma} d\chi}$. Since we do not want the value of the observable to be affected by a (generalised) gauge transformation, we have to impose the condition $\int_{\gamma} d\chi = 2\pi n$ (see also [57, 58]) which leads to the final result

$$Q = 2\pi n . \quad (\text{C17})$$

-
- [1] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
 - [2] A. Kitaev, AIP Conf. Proc. **1134**, 22 (2009).
 - [3] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
 - [4] V. Gurarie, Phys. Rev. B **83**, 085426 (2011).
 - [5] L. Fidkowski and A. Kitaev, Phys. Rev. B **81**, 134509 (2010).
 - [6] C. Wang, A.C. Potter and T. Senthil, arXiv:1306.3238 (2013).
 - [7] S. Raghu, X.-L. Qi, C. Honerkamp and S.-C. Zhang, Phys. Rev. Lett. **100**, 156401 (2008).
 - [8] L. Fidkowski and A. Kitaev, Phys. Rev. B **83**, 075103 (2011).
 - [9] Y.-M. Lu and A. Vishwanath, Phys. Rev. B **86**, 125119 (2012).
 - [10] Y. Zhang, Y. Ran and A. Vishwanath, Phys. Rev. B **79**, 245331 (2009).
 - [11] X.-G. Wen, *Quantum Field Theory of Many-body Systems* (Oxford University Press, 2004).
 - [12] D. Birmingham, M. Blau, M. Rakowski and G. Thompson, Phys. Rep. **209**, 129 (1991).
 - [13] B.A. Bernevig and S.-C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006).
 - [14] G. Y. Cho and J. E. Moore, Ann. Phys. **326**, 1515 (2011).
 - [15] A. M. Chan, T. L. Hughes, S. Ryu, and E. Fradkin, Phys. Rev. B **87**, 085132 (2013).
 - [16] G. Palumbo, R. Catenacci and A. Marzuoli, Int. J. Mod. Phys. B **28**, 1350193 (2014).
 - [17] M. C. Diamantini, P. Sodano and C.A. Trugenberger, New. J. Phys. **14**, 063013 (2012).
 - [18] J. Maciejko, V. Chua and G.A. Fiete, arXiv:1307.5566v1 (2013).
 - [19] P. Hosur, S. Ryu, and A. Vishwanath, Phys. Rev. B **81**, 045120 (2010).
 - [20] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
 - [21] G. Palumbo and J. K. Pachos, Phys. Rev. Lett. **110**, 211603 (2013).
 - [22] G. Palumbo and J. K. Pachos, arXiv:1311.2871v1 (2013).
 - [23] W. E. Thirring, Ann. Phys. **3**, 91 (1958).
 - [24] E. Fradkin and F. A. Schaposnik, Phys. Lett. B **338**, 253 (1994).
 - [25] E. Cremmer and J. Scherk, Nucl. Phys. B **72**, 117 (1974).
 - [26] T. Allen, M. J. Bowick, and A. Lahiri, Mod. Phys. Lett. A **6**, 559 (1991).
 - [27] E. Zohar, J. I. Cirac and B. Reznik, Phys. Rev. A **88**, 023617 (2013).
 - [28] D. Banerjee, M. Bögli, M. Dalmonte, E. Rico, P. Stebler, U.-J. Wiese and P. Zoller, Phys. Rev. Lett. **110**, 125303 (2013).

- [29] M. Lewenstein, A. Sanpera and V. Ahufinger, *Ultra-cold Atoms in Optical Lattices* (Oxford University Press, 2012).
- [30] L. Susskind, Phys. Rev. D **16**, 3031 (1977).
- [31] A. P. Schnyder, S. Ryu, A. Furusaki and A. W. W. Ludwig, AIP Conf. Proc. **1134**, 10 (2009).
- [32] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
- [33] E. Fermi. Z. Physik **88** (1934).
- [34] M. Botta Cantcheff and J. A. Helayel-Neto, Phys. Rev. D **67**, 025016 (2003).
- [35] M. Blau and G. Thompson. Annals of Physics **205**(1), 130 (1991).
- [36] S. Deser, R. Jackiw and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982).
- [37] L.N. Cooper, Phys. Rev. **104**, 4 (1956).
- [38] I. A. Batalin and E. S. Fradkin, Nucl. Phys. B **279**, 514 (1987).
- [39] I. A. Batalin and I. V. Tyutin, Int. J. Mod. Phys. A **6**, 3255 (1991).
- [40] M. Kalb and P. Ramond, Phys. Rev. D **9**, 2273 (1974).
- [41] E. Harikumar and M. Sivakumar, Nucl. Phys. B **565**, 385 (2000).
- [42] Y.-W. Kim, S.-K. Kim and Y.-J. Park, Mod. Phys. Lett. A **18**, 2287 (2003).
- [43] T.H. Hansson, V. Oganesyan and S.L. Sondhi, Ann. Phys. **313**, 497 (2004).
- [44] A. Balachandran and P. Teotonio-Sobrinho, Int. J. Mod. Phys. A **08**(04), 723 (1993).
- [45] S. Coleman, Phys. Rev. D **11**, 2088 (1975).
- [46] D. Delpine, R. G. Felipe, and J. Weyers, Phys. Lett. B **419**(14), 296 (1998).
- [47] F.A. Schaposnik, Phys. Lett. B **356**, 39 (1995).
- [48] A. Amoretti, A. Blasi, N. Maggiore and N. Magnoli, New J. Phys. **14** 113014 (2012).
- [49] A. Momen, Phys. Lett. B **394**, 269 (1997).
- [50] G. Horowitz and M. Srednicki, Comm. Math. Phys. **130**, 83 (1990).
- [51] I. Oda and S. Yahikozawa, Phys. Lett. B **238**, 272 (1990).
- [52] S. Weinberg. *The Quantum Theory of Fields, Volume 1: Foundations* (Cambridge University Press, 1995).
- [53] P. V. Sriluckshmy, A. Mishra, S. R. Hassan, and R. Shankar. arXiv:1310.0791v1 (2013).
- [54] M. Nakahara, Taylor & Francis, 1989.
- [55] D. Birmingham, M. Blau, M. Rakowski, and G. Thompson, Physics Reports **209**(45), 129 (1991).
- [56] A. S. Cattaneo, P. Cotta-Ramusino, J. Frohlich, and M. Martellini, J. Math. Phys. **36**(11), 6137 (1995).
- [57] R. J. Szabo, Annals of Physics **280**(1), 163 (2000).
- [58] M. Bergeron, G. W. Semenoff, and R. J. Szabo, Nuclear Physics B **437**(3), 695 (1995).
- [59] Without them, it would be impossible to generate such a mass with just one field. In the abelian case this is very easily seen. In fact, with the presence of one field the only possibilities for a gauge invariant 4-form are $F \wedge F$ and $F \wedge *F$. The first term has a topological nature while the second term is geometric. The topological term can be rewritten as $F \wedge F = d(A \wedge dA)$. This allows to conclude that the addition of such a differential to the lagrangian would not change the equation of motion (since it adds a total derivative to the Lagrangian) and for this reason it cannot possibly add a mass to the gauge field.