

This is a repository copy of *The antimicrobial activity of a carbon monoxide releasing molecule (EBOR-CORM-1) is shaped by intraspecific variation within3 Pseudomonas aeruginosa populations.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/127017/>

Version: Accepted Version

---

## Article:

Flanagan, Lindsey Anne, Steen, Rachel Rosemary, Saxby, Karinna Isobel et al. (7 more authors) (2018) The antimicrobial activity of a carbon monoxide releasing molecule (EBOR-CORM-1) is shaped by intraspecific variation within3 Pseudomonas aeruginosa populations. *Frontiers in Microbiology*. 195.

<https://doi.org/10.3389/fmicb.2018.00195>

---

## Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

## Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# The antimicrobial activity of a carbon monoxide releasing molecule (EBOR-CORM-1) is shaped by intraspecific variation within *Pseudomonas aeruginosa* populations

Lindsey Flanagan<sup>1,2</sup>, Rachel Steen<sup>2</sup>, Karinna Saxby<sup>1,2</sup>, Mirre Klatter<sup>1</sup>, Benjamin J. Aucott<sup>2</sup>, Craig Winstanley<sup>3</sup>, Ian J. S. Fairlamb<sup>2</sup>, Jason M. Lynam<sup>2</sup>, Alison Parkin<sup>2</sup> and Ville-Petri Friman<sup>1</sup>

<sup>1</sup> The University of York, Department of Biology, Wentworth Way, York, YO10 5DD, UK

<sup>2</sup> Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK

<sup>3</sup> Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, Ronald Ross Building, University of Liverpool, 8 West Derby Street, Liverpool, L69 7BE, UK

Correspondence: Ville-Petri Friman, The University of York, Department of Biology, Wentworth Way, York, YO10 5DD, UK, tel: 01904 328675, e-mail: ville.friman@york.ac.uk

## ABSTRACT

Carbon monoxide releasing molecules (CORMs) have been suggested as a new synthetic class of antimicrobials to treat bacterial infections. Here we utilised a novel EBOR-CORM-1 ( $[\text{NEt}_4][\text{MnBr}_2(\text{CO})_4]$ ) capable of water-triggered CO-release, and tested its efficacy against a collection of clinical *Pseudomonas aeruginosa* strains that differ in infection-related virulence traits. We found that while EBOR-CORM-1 was effective in clearing planktonic and biofilm cells of *P. aeruginosa* strain PAO1 in a concentration dependent manner, this effect was less clear and varied considerably between different *P. aeruginosa* cystic fibrosis (CF) lung isolates. While a reduction in cell growth was observed after 8 hours of CORM application, either no effect or even a slight increase in cell densities and the amount of biofilm was observed after 24 hours. This variation could be partly explained by differences in bacterial virulence traits: while CF isolates showed attenuated *in vivo* virulence and growth compared to strain PAO1, they formed much more biofilm, which could have potentially protected them from the CORM. Even though no clear therapeutic benefits against a subset of isolates was observed in an *in vivo* wax moth acute infection model, EBOR-CORM-1 was more efficient at reducing the growth of CF isolate co-culture populations harbouring intraspecific variation, in comparison with efficacy against more uniform single isolate culture populations. Together these results suggest that CORMs could be effective at controlling genetically diverse *P. aeruginosa* populations typical for natural chronic CF infections and that the potential benefits of some antibiotics might not be observed if tested only against clonal bacterial populations.

**Keywords:** Biofilms, Carbon monoxide releasing molecules, CORM, Cystic fibrosis, Polymicrobial infections, *Pseudomonas aeruginosa*, Synthetic chemistry, Virulence

## 1. INTRODUCTION

The rapid emergence of multidrug-resistant bacteria is a global problem that is predicted to cause ten million deaths per year by 2050 (O'Neill, 2014). Antibiotic resistance often evolves very quickly via *de novo* mutations and horizontal gene transfer (Normark and Normark, 2002), and as a result, antibiotic discovery has not been able to replace all of the antibiotics that have now become ineffective (Brown and Wright, 2016). New methods and approaches for treating bacterial infections are thus urgently required.

In recent years, carbon monoxide has emerged as a new potential therapeutic due to its properties as a homeostatic and cytoprotective molecule with important signalling capabilities (Motterlini and Otterbein, 2010). Carbon monoxide can be delivered via carbon monoxide releasing molecules (CORMs), which are small molecules that release carbon monoxide in response to certain environmental triggers such as enzymes (Stamellou et al., 2014) or light (Jimenez et al., 2016). Nobre *et al.* first investigated the effect of CORMs on bacteria (Nobre et al., 2007) and found that CORM-2 and CORM-3 reduced the number of colony-forming units of *Escherichia coli* in minimal salts media and *Staphylococcus aureus* in Luria Broth (LB) media (Nobre et al., 2007). The CORM effects were stronger in near-anaerobic conditions and the activation of CORM required direct contact between the molecule and its cellular targets (Nobre et al., 2007). Moreover, the effect of CORM-3 on *Pseudomonas aeruginosa* wild type strain PAO1 was investigated by Desmard *et al.* (Desmard et al., 2009), who found that treatment with the CORM reduced bacterial densities and increased the survival of immunocompromised mice during an infection. It has also previously been found that CORM-2 effectively reduces the densities of *P. aeruginosa* planktonic and biofilm cultures with wild type and clinical strains (Murray et al., 2012). Another study found that manganese-based Trypto-CORM is able to inhibit the growth of *E. coli* when exposed to photochemical stimulus (Ward et al., 2014), while in the dark it is active against *Neisseria gonorrhoeae* (Ward et al., 2017). In both cases, control experiments indicate that the CO liberated from the metal is responsible for the observed behaviour. However, most of the studies thus far have concentrated on exploring CORM effects on relatively short time span (less than 24 hours). Furthermore, although it has been established that many infections are polymicrobial, and that clinical bacterial pathogens can respond differently to CORMs than laboratory strains, no studies have explored CORM effects on bacterial co-cultures.

Cystic fibrosis (CF) is a genetically inherited disease which affects 1 in 2000 to 3000 newborn infants in the EU (who.int, 2010). Patients with CF often develop a thick mucus in the lungs which they are unable to clear (Flume et al., 2010). This mucus makes patients susceptible to frequent and recurring bacterial chest infections and the presence of *P. aeruginosa* is often associated with increasing morbidity and loss of lung function (Pritt et al., 2007). One of the key features of *P. aeruginosa* is its capability to rapidly adapt to the lung environment and to become highly resistant to the antibiotics that are used to treat infections (Smith et al., 2006; Poole, 2011; Folkesson et al., 2012; Winstanley et al., 2016). As a result, *P. aeruginosa* populations show high levels of genetic variation within and between CF patients (Marvig et al., 2013; Williams et al., 2015; O'Brien et al., 2017). This includes phenotypic and genomic heterogeneity within genetically-related populations of *P. aeruginosa* derived from the same clonal lineage (Mowat et al., 2011; Workentine et al., 2013; Williams et al., 2015). This variation might also affect the applicability of potential alternative therapies if it is linked with bacterial life-history traits that relate to potential resistance mechanisms.

Here we synthesised and characterised a water-soluble CORM (EBOR-CORM-1),  $[\text{NEt}_4][\text{MnBr}_2(\text{CO})_4]$ , and tested its effectiveness against *P. aeruginosa* strain PAO1 and a selection of *P. aeruginosa* CF isolates originating from a single sputum sample from the lungs of a CF patient, namely patient CF03 from previously published studies (Mowat et al., 2011; Williams et al., 2015). Based on genome sequence data presented in a previous study, these CF isolates were classified into two genetically distinct Liverpool Epidemic Strain (LES) lineages, A and B (Williams et al., 2015; Williams et al., 2016), that differ regarding their virulence traits (O'Brien et al., 2017). These genetically diverged lineages have been shown to commonly coexist within individual patients and to share mutations via homologous recombination that potentially help strains to adapt to the airway during chronic infection (Williams et al., 2015). However, the implications of within-patient genetic variation have been seldom considered in the context of antimicrobial therapies. We hypothesised that effects of EBOR-CORM-1 could vary between different clinical isolates and lineages, and that the susceptibility of isolates could be linked to expression of some other bacterial virulence factors. We found that the CORM was effective in reducing both planktonic and biofilm cells of strain PAO1 in a density-dependent manner. However, CORM effects were more varied and generally weaker against clinical CF isolates. Regardless, CORM efficiently reduced the growth of CF strain lineage co-cultures, which suggest that CORMs could be effective at controlling genetically diverse *P. aeruginosa* infections.

## 2. MATERIALS AND METHODS

### ***Synthesis and properties of [NEt<sub>4</sub>][MnBr<sub>2</sub>(CO)<sub>4</sub>], EBOR-CORM-1***

EBOR-CORM-1 was synthesised as described previously (Angelici, 1964): Mn(CO)<sub>5</sub>Br (466 mg, 1.69 mmol) and 330 mg (1.57 mmol) of [(C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>N]Br were heated in 18 mL of absolute methanol under a nitrogen atmosphere at 50 °C for 1 hour. The methanol was then evaporated from the orange solution at the above temperature. The remaining yellow solid was dissolved in 40 mL of chloroform, and the solution was filtered under nitrogen. After adding 200 mL of hexane to the filtrate, the cloudy solution was allowed to stand under nitrogen for 2 hours. The air-stable yellow crystals were separated by filtration, washed with hexane, and dried under vacuum giving a yield of 88 % (636 mg). The compound was characterised *via* solid state IR spectroscopy recorded using a KBr disk. Four main bands were seen at 2090, 2001, 1984 and 1942 cm<sup>-1</sup> and a small shoulder was seen at 1897 cm<sup>-1</sup>. This is consistent with the literature values (Angelici, 1964). In a chloroform solution of CORM four distinct bands were observed at 2092, 2015, 1987 and 1943 cm<sup>-1</sup>, again this is similar to previously reported literature values (Angelici, 1964). The change in the number of carbonyl bands between the solid and solution phase measurements typically reflects that different orientations are present in the solid state. The stability of the CORM in the solid state was tested by heating a sample to 50 °C and running ATR IR spectra at 1 hour intervals.

Infrared detection of CO release from EBOR-CORM-1 following dissolution in different solvents was conducted by dissolving 12 mg of CORM in 4 mL of solvent in a 25 mL round bottomed flask attached to vacuum evacuated gas IR cell via a closed tap. After 1 h of stirring the flask, the tap was opened to enable gas from the headspace of the flask to enter the IR cell. Carbon monoxide could then be identified via the distinctive gaseous IR signature of a double band, with fine rotational splitting, centred at 2150 cm<sup>-1</sup> (Klein et al., 2014). The impact of different solvents can be quantified by comparison of the intensity of the CO bands to those from CO<sub>2</sub>, which is assumed to act as an effective internal standard.

The release of CO from EBOR-CORM-1 following dissolution in water was also followed via solution phase monitoring of the metal complex's IR bands. In contrast to chloroform, when EBOR-CORM-1 was first dissolved in water only two main IR bands were observed at 2050 and 1943 cm<sup>-1</sup>. In order to investigate activity of EBOR-CORM-1 in liquid culture media, we compared the effects of active and 'inactivated' CORM on the growth of PAO1 strain in LB media as described previously (Murray et al., 2012). Briefly, CORM was inactivated by storing a 2 mM CORM stock LB solution (10% v/v of standard LB concentration, i.e., the same that was used in all the experiments; see below) at room temperature for 24 hours. To

estimate the effect of CORM inactivation on PAO1 growth, we added 50  $\mu$ L of freshly prepared 2 mM CORM, 50  $\mu$ L of inactivated 2 mM CORM or 50  $\mu$ L 10% v/v LB (control) to 150  $\mu$ L of PAO1 starter culture on 96-well microplate. All treatments were replicated five times and PAO1 growth monitored for 8 hours at 37 °C with spectrophotometer (OD 600 nm; Tecan Infinite).

### **Bacterial strains and culture media**

In this study we used *P. aeruginosa* strain PAO1 (ATCC 15692), the earliest archived isolate of the Liverpool Epidemic strain, LESB58 (Winstanley et al., 2009), and 19 clinical *P. aeruginosa* LES isolates from the same sputum sample of a chronically infected CF patient (Williams et al., 2015). The CF lung LES isolates originate from the sputum sample of one patient, identified as patient CF03 in previous studies, and consist of two genetically separate lineages A and B (Williams et al., 2015). Lineage A was represented by six isolates, namely isolates: 2, 5, 10, 19, 23 and 25. Lineage B was represented by 13 isolates, namely isolates: 1, 6, 8, 17, 24, 26, 28, 32, 33, 34, 35, 36 and 37. Clinical isolates were collected with the consent of the patient and under institutional human investigation approval. All strains and isolates of *P. aeruginosa* were routinely cultured in liquid or solid LB media containing 10.0 g tryptone, 5.0 g yeast extract and 10.0 g NaCl in 1 L of ultra-pure water (final pH adjusted to 7.0 and 15 g of agar was used for solid media). For all experiments, starter cultures were prepared from cryofrozen stocks by streaking frozen stock culture onto LB plates. After 24 hours growth, a single colony was selected and inoculated into 5 mL of liquid LB and grown overnight in a shaking incubator at 37 °C in 50 mL centrifuge tubes. Overnight cultures were centrifuged at 4000 rpm (11.5 g) for 15 min (Eppendorf), the resultant pellets were suspended in 10% LB and bacterial densities adjusted to optical density at 600 nm of 0.066 before use (OD 600nm), equalling  $\sim 1 \times 10^8$  cells mL<sup>-1</sup>.

### **Measuring the effects of EBOR-CORM-1 concentration on *P. aeruginosa* PAO1 strain**

We measured the effect of CORM concentration on *P. aeruginosa* PAO1 in four different ways. First, we examined how EBOR-CORM-1 affects PAO1 growth after both 8 and 24 hours of inoculation in 10% LB media (bacteria and CORM inoculated at the same time). Additionally, we measured how effective EBOR-CORM-1 is at clearing both established planktonic and biofilm PAO1 cultures (bacteria pre-grown before adding EBOR-CORM-1). All measurements were conducted on 96-well microplates and each treatment was replicated 5 times. A variety of EBOR-CORM-1 concentrations were tested by first preparing a 4 mM CORM stock solution (dissolving EBOR-CORM-1 in 10% LB media by vortexing for 30 s

and sonicating for 1.5 min). The stock solution was then sterilised with syringe filtration and serially diluted to result in 1 mM, 0.5 mM, 0.25 mM, 0.125 mM and 0 mM (control) EBOR-CORM-1 concentrations and  $1 \times 10^8$  PAO1 cells mL<sup>-1</sup> with final volume of 200 µl of media. The microplate was then incubated at 37 °C for 24 hours.

All replicate populations were sampled at 8 and 24 hours after the start of the experiment (20 µl of samples) and serially diluted in sterile PBS on microplates to quantify the number of living versus dead cells by flow cytometry. Briefly, DAPI (4',6-diamidino-2-phenylindole for dead and living cells) and PI (Propidium iodide for dead cells) fluorescent stains (both from Sigma-Aldrich) were added to microplate wells with diluted bacterial samples at concentrations of 1 µg/mL and 50 µM, respectively. Plates were then incubated at room temperature for 1 hour before measuring cell densities with a Cytoflex flow cytometer and the CytExpert program. Every well was sampled for 60 s at fast speed setting. Gating of live and dead cells was performed by monitoring DAPI staining on the PB450 channel with the 405 nm laser, and PI staining on the ECD channel of the 488 nm laser. Number of living cells was determined as total cells (DAPI) – dead cells (PI).

To quantify the effects of EBOR-CORM-1 on established planktonic and biofilm cultures, PAO1 was first grown in the absence of CORM at 37 °C for 48 hours. Cell cultures were then inoculated with stock CORM solution to reach the same final concentrations as above: 1 mM, 0.5 mM, 0.25 mM, 0.125 mM and 0 mM (control) of CORM. The plate was incubated for four more hours at 37 °C before sampling (20 µL), serial dilution and flow cytometry as described above. To quantify effects of EBOR-CORM-1 on biofilm, crystal violet was added to the remaining cell cultures at 10% v/v. After 15 min of incubation, the plate was rinsed with deionised water and solubilised with 228 µL ethanol per well. The biofilm was quantified by measuring absorbance at 600 nm.

### ***Measuring the effects of EBOR-CORM-1 on clinical P. aeruginosa isolates in mono- and co-cultures***

Similar to the PAO1 strain experiments, we measured the effect of EBOR-CORM-1 on clinical *P. aeruginosa* isolates after 8 and 24 hours of inoculation in 10% LB media. We also measured the impact of growing the isolates in the absence of EBOR-CORM-1 for 48 hours and then applying EBOR-CORM-1 for 4 hours using both flow cytometry and crystal violet staining. We used only one EBOR-CORM-1 concentration, 0.5 mM, which resulted in clear reduction of PAO1 cultures (see results) alongside control treatment (no CORM).

In addition to measuring the effects of EBOR-CORM-1 in monocultures of each clinical isolate, we also quantified the effect of the CORM on mixtures of the CF clinical isolates from patient CF03. First, we prepared the clinical isolate starter cultures as described above, then we mixed the standardised monocultures together in three different ways: as a whole mix (all isolates mixed together in equal proportions), lineage A mix (all isolates classified as lineage A mixed together in equal proportions) and lineage B mix (all isolates classified as lineage B mixed together in equal proportions). All final mixes contained approximately  $1 \times 10^8$  cells mL<sup>-1</sup> before the application of 0.5 mM of EBOR-CORM-1. Each experiment was replicated 5 times. After 24 hours growth at 37 °C, bacterial densities were measured by using a Tecan infinite spectrophotometer: optical density measurements correlate well with the proportion of living cells measured with flow cytometer (Supplementary figure 1).

### ***Characterising bacterial virulence and growth***

To characterise production of the virulence factors pyocyanin and pyoverdine, all clinical isolates were grown in 200 µL of 10% LB media in round-bottomed 96-well microplates for 48 hours at 37 °C (no shaking). After incubation, we measured the bacterial densities (OD 600 nm) and centrifuged the microplate for 10 min. at 4000 rpm (11.5 g) in a swing rotor Eppendorf centrifuge. To measure pyocyanin and pyoverdine production, 150 µL of the supernatant of each well was transferred to flat-bottomed 96-well microplates and the absorbance spectrum measured with a spectrophotometer (Tecan infinite). Per capita pyocyanin production was measured for each isolate by measuring the absorbance of supernatant at 691 nm, and then standardizing by bacterial OD (Reszka et al., 2004). Per capita production of the iron-chelating siderophore, pyoverdine, was measured by using excitation-emission assay (O'Brien et al., 2017) where the fluorescence of each supernatant well was measured at 470 nm following excitation at 380 nm, using a Tecan infinite M200 pro spectrophotometer. Also, OD was measured at 600 nm to quantify the ratio fluorescence/OD as a quantitative measure of per capita pyoverdine production (O'Brien et al., 2017). The isolate biofilm production was measured as described previously and growth as maximum density and growth rate h<sup>-1</sup> during 24-hour growth period. Lastly, we also measured the *in vivo* virulence of each isolate by using wax moth model as described previously (O'Brien et al., 2017).

### ***Testing EBOR-CORM-1 antimicrobial activity in wax moth model in vivo***



To test the efficacy of EBOR-CORM-1 to constrain bacterial infections *in vivo*, we used a wax moth larvae model (*Galleria mellonella* [Lepidoptera: Pyralidae], Livefood UK Ltd) and followed the infection methodology described previously (O'Brien et al., 2017). We chose three strains for infection experiments: PAO1, LESB58 and isolate 36 (Lineage B) from the clinical sample collection. Before infection, we first grew the selected *P. aeruginosa* isolates for 24 hours at 37 °C and subsequently diluted all cultures to approximately similar densities (equalling approximately  $1 \times 10^6$  cells mL<sup>-1</sup> in 0.8% w:v NaCl). The virulence of every isolate was then tested in 16 independent wax moth larvae. We also infected 16 larvae with 0.8% w/v NaCl salt solution to control for the damage caused by the injection itself. The larvae were injected with either 20 µL of one bacterial solution or NaCl buffer (“non-infected”) between the abdominal segments six and seven with a 1 mL Terumo syringe. After 2 hours, 8 larvae from each bacterial infection or non-infection group were treated with 20 µL injection of 500 µM EBOR-CORM-1, and the other 8 were injected with 0.8% w:v NaCl salt solution (control placebo) in the same location where the bacteria were originally injected. After infection, larvae were placed on individual wells of 24-well cell culture plates and the survival was monitored at 2-hour intervals for 3 days at 37 °C. Larvae were scored as dead when they did not respond to touch with forceps. Larvae that were still alive after 7 days from the infection were given a time of death of 168 hours. Every bacterial isolate was tested for three times. It was concluded that the EBOR-CORM-1 injection alone did not affect larval survival in the absence of bacteria (mortality similar between non-infected CORM-injected larvae and non-infected CORM-free larvae: 5-10%).

### **Statistical analysis**

All data were analysed with Generalized Mixed Models (factorial ANOVA) or regression analysis where bacterial densities (Figs. 2, 3 and 4b) or trait values (Fig. 4a; Supplementary figure 4) were explained with the presence and/or concentration of EBOR-CORM-1, CF isolate identity (isolate number) or CF lineage (A or B). All proportional data (%) were arcsine transformed before the analysis to meet the assumptions of parametric models.

## **3. RESULTS**

### **Chemistry of EBOR-CORM-1**

The stability of EBOR-CORM-1 in the solid state was demonstrated by heating a sample of solid to 50 °C in air, and showing that there is very little difference in the carbonyl bands observed in ATR IR spectra measured at 1 hour intervals over a 3-hour period (Fig. 1A). In contrast, gas phase infrared analysis proved that CO release from EBOR-CORM-1 can be

triggered by dissolution in water, phosphate buffer or LB media, or addition of water to a solution of the compound in an organic solvent (Fig. 1B).

Solution phase monitoring of the CO stretches of the compound showed that there was no reaction with water over short periods of time, since dissolving EBOR-CORM-1 in water, immediately re-drying it on a vacuum line and then re-dissolving the resultant solid in chloroform yielded an IR spectra which matched that of the as-purified compound in chloroform (Fig. 1C). The only two observed IR bands in the CORM spectrum in water (2050 and 1943  $\text{cm}^{-1}$ ) were therefore attributed to the molecular symmetry of the hydrated complex, rather than an immediate loss of CO upon contact with water. However, after 90 min in water, a loss of these carbonyl bands was observed, and this was attributed to the release of all the CO from the complex (Fig. 1D).

### **EBOR-CORM-1 activity against planktonic and biofilm cells of *P. aeruginosa* PAO1**

We found that applying EBOR-CORM-1 had generally negative effects on *P. aeruginosa* PAO1 growth both after 8 and 24 hours of application ( $F_{4, 25} = 50.9$ ,  $p < 0.001$  and  $F_{4, 25} = 31.8$ ,  $p < 0.001$  for proportion of living cells after 8 and 24 hours, respectively, Fig. 2A) and that these negative effects increased along with the increasing concentration of applied EBOR-CORM-1 (regression analysis:  $F_{1, 24} = 43$ ,  $p < 0.001$  and  $F_{1, 24} = 35$ ,  $p < 0.001$  for proportion of living cells after 8 and 24 hours, respectively, Fig. 2A). Similarly, EBOR-CORM-1 was highly effective against both established planktonic and biofilm *P. aeruginosa* PAO1 cultures ( $F_{4, 25} = 77.5$ ,  $p < 0.001$  and  $F_{4, 25} = 39.5$ ,  $p < 0.001$ , respectively, Fig. 2A-B) and the antimicrobial activity of CORM increased in a density-dependent manner (regression analysis:  $F_{1, 24} = 92$ ,  $p < 0.001$  and  $F_{1, 24} = 54$ ,  $p < 0.001$ , respectively, Fig. 2A-B).

### **EBOR-CORM-1 activity against planktonic and biofilm cells of clinical *P. aeruginosa* cystic fibrosis isolates**

Similar to strain PAO1, we found that EBOR-CORM-1 had inhibitory effects on all tested clinical *P. aeruginosa* isolates after 8 hours of application of CORM ( $F_{1, 152} = 11969$ ,  $p < 0.001$ , Fig. 3A). While this effect did not depend on the lineage (CORM  $\times$  lineage:  $F_{1, 152} = 1.4$ ,  $p < 0.001$ ), it varied between different clinical isolates (CORM  $\times$  isolate:  $F_{18, 152} = 11969$ ,  $p < 0.001$ , Fig. 3A). In contrast, EBOR-CORM-1 had slightly positive effects on *P. aeruginosa* growth after 24 hours of application ( $F_{1, 152} = 256$ ,  $p < 0.001$ , Fig. 3B) and this effect varied between different isolates (CORM  $\times$  strain:  $F_{18, 152} = 2.8$ ,  $p = 0.001$ ) being slightly stronger (i.e. positive) with isolates belonging to a lineage B (CORM  $\times$  lineage:  $F_{1, 152} = 11969$ ,  $p < 0.001$ , Fig. 3B).

$_{152} = 24.9, p < 0.001$ , Fig. 3B). EBOR-CORM-1 also had negative effects when applied to established *P. aeruginosa* cell cultures ( $F_{1, 152} = 222, p < 0.001$ , Fig. 3C). However, these effects depended on the isolate (CORM  $\times$  isolate:  $F_{18, 152} = 2.8, p = 0.001$ ) and the lineage ( $F_{1, 152} = 65.2, p = 0.001$ ), reduction being relatively larger with isolates belonging to lineage A (Fig. 3C). In the case of established biofilms, EBOR-CORM-1 had a slightly positive effect ( $F_{1, 152} = 9.6, p = 0.002$ , Fig. 3D) and while this effect varied between different isolates ( $F_{18, 152} = 2.0, p = 0.01$ ) it did not differ between the lineages ( $F_{1, 152} = 1.2, p = 0.265$ , respectively, Fig. 3D). Together these results suggest that compared to strain PAO1, EBOR-CORM-1 effects varied more with the clinical *P. aeruginosa* isolates having negative, neutral or positive effects on bacterial growth depending on the isolate identity, lineage and the timing of CORM application.

### **Linking EBOR-CORM-1 antimicrobial activity with clinical *P. aeruginosa* isolate virulence and growth**

We found that all the isolates belonging to a lineage A formed non-mucoid colonies (6 out of 6), while most of the isolates belonging to a lineage B formed mucoid (i.e., mucus-like) colonies (11 out of 13) on LB plates (typical mucoid and non-mucoid colonies shown in supplementary figure 3). All clinical isolates differed from the non-mucoid PAO1 strain respective of their virulence and growth (Fig. 4A). More specifically, clinical isolates produced less pyoverdine ( $F_{1, 23} = 286, p < 0.001$ ) and pyocyanin ( $F_{1, 23} = 170, p < 0.001$ ) and grew slower ( $F_{1, 23} = 91, p < 0.001$ ) and reached lower maximum densities in LB medium ( $F_{1, 23} = 15.5, p = 0.001$ , Fig. 4A). However, clinical isolates produced a considerably larger amount of biofilm ( $F_{1, 23} = 21.7, p < 0.001$ ) and showed very low virulence (high time to death) in wax moth larvae *in vivo* ( $F_{1, 23} = 1296, p < 0.001$ , Fig. 4A).

When comparing the two CF lineages, we found that isolates belonging to a lineage B consistently outperformed the isolates belonging to a lineage A by producing more pyoverdine ( $F_{1, 18} = 6.06, p = 0.025$ ), biofilm ( $F_{1, 18} = 15.08, p = 0.001$ ) and by growing faster ( $F_{1, 18} = 22.35, p < 0.001$ ) and to higher maximum densities ( $F_{1, 18} = 6.27, p = 0.023$ ) in LB medium (Fig. 4A; Supplementary figure 4). However, lineages did not differ in pyocyanin production ( $F_{1, 18} = 1.99, p = 0.176$ ) or virulence ( $F_{1, 18} = 1.03, p = 0.324$ ; Fig. 4A; Supplementary figure 4). Across all clinical isolates, density reduction by CORM correlated negatively with biofilm formation ( $F_{1, 18} = 4.8, p = 0.042$ ). Together these results suggest that clinical isolates differed from PAO1 and from each other respective to various life-history traits important for establishing an infection.

### **EBOR-CORM-1 activity against clinical *P. aeruginosa* CF isolate co-cultures**

Despite the observed isolate-specific variation in *P. aeruginosa* monocultures, EBOR-CORM-1 was effective in reducing the growth of *P. aeruginosa* co-cultures after 24 hours of application (CORM:  $F_{1, 24} = 132$ ,  $p < 0.001$ , Fig. 4B). Moreover, this reduction was the same regardless of whether the mix contained only one lineage or both lineages (CORM  $\times$  co-culture:  $F_{2, 24} = 0.5$ ,  $p = 0.612$ ). These results suggest that intraspecific *P. aeruginosa* population heterogeneity makes the bacteria more susceptible to EBOR-CORM-1 treatment.

#### **EBOR-CORM-1 activity against *P. aeruginosa* strains in wax moth model**

We found that *P. aeruginosa* isolates differed in their virulence (time to death) from each other ( $F_{2, 24} = 12.2$ ,  $p < 0.001$ ): PAO1 and LESB58 strains were equally virulent, and both exhibited higher virulence than the clinical isolate 36 (killing larvae approximately in 17 hours [PAO1], 36 hours [LESB58] and 92 hours [clinical isolate 36]; values averaged over both non-CORM and CORM treatments, Fig. 5). In contrast to *in vitro* results, application of EBOR-CORM-1 did not increase the survival of infected larvae ( $F_{1, 24} = 1.3$ ,  $p = 0.257$ ) with any of the infected strains (CORM  $\times$  strain:  $F_{2, 24} = 1.4$ ,  $p = 0.273$ , Fig. 5). All larvae became highly pigmented (black throughout) during the infection regardless of the *P. aeruginosa* isolate.

## **4. DISCUSSION**

Here we set out to study the antimicrobial activity of  $[\text{NEt}_4][\text{MnBr}_2(\text{CO})_4]$ , EBOR-CORM-1, against clinical *P. aeruginosa* isolates *in vitro*. This CORM was chosen as a suitable representative of this class of molecule based on the aqueous solubility, facile synthesis (Angelici, 1964), content of a non-toxic metal core, and simple architecture which makes it akin to a “parent compound” for CORMs that have been engineered to possess sophisticated CO release mechanisms. In contrast to more complex CORMs, the molecule was shown to have a water activated mechanism of CO release, as seen in previous studies of  $[\text{MX}(\text{CO})_5]^-$  species, where X is a halide (Zhang et al., 2009). Such water induced degradations are believed to proceed via a two-step pathway whereby water causes loss of the halide followed by formation of a dimer species; from which the CO is released. This may explain the changes in the IR spectra recorded in water when compared to chloroform, although the data do not directly match those for  $[\text{Mn}_2\text{Br}_2(\text{CO})_8]$  (El-Sayed and Kaesz, 1963), the product expect on loss of  $\text{Br}^-$  from EBOR-CORM-1. We found that while EBOR-CORM-1 showed density-dependent antimicrobial activity against both planktonic and biofilm cells of the widely studied laboratory-adapted strain PAO1, these effects were more varied and weaker against clinical CF lung isolates. Regardless, EBOR-CORM-1 was efficient at reducing the

growth of CF isolate lineage mixes, which suggests that it could have therapeutic potential in controlling heterogeneous *P. aeruginosa* infections. Solutions of inactivated EBOR-CORM-1 were essentially inactive against *P. aeruginosa* strain PAO1 (Supplementary figure 2) implying that, at least in this case, the observed activity was due to CO released from the complex rather than the residual metal salts (or indeed  $[\text{NEt}_4]^+$ ).

Similar to a study published by Murray *et al.* (2012), we found considerable variation in CORM antimicrobial activity between different clinical CF isolates, which depended whether we explored EBOR-CORM-1 effects on relatively short (8 hours) or long timescales (24 hours) and if we compared CORM antimicrobial activity on actively growing and established cell cultures (after 48 hours of bacterial growth). Our results after 8 hours of EBOR-CORM-1 application are very similar to a previous study (Murray *et al.*, 2012) showing clear reduction in bacterial densities. However, this effect vanished by the 24 hour time point, and surprisingly, some bacterial isolate cultures reached higher optical densities in the presence compared to absence of CORM, which could have been due to increase in number of cells or expression of exoproducts that were picked up by OD600 nm (e.g. pyocyanin or alginate). The most likely explanation for this is that CORM effects were short-lived (Fig. 1), which allowed bacteria to recover and grow to high densities during 24 hours after application of CORM. However, when CORM effects were measured after 4 hours of application to established cell cultures, we could still observe clear reduction in mean bacterial densities. Together these results suggest that CORM effects could be seen up to 4 hours post application and that CORM could eradicate bacterial cells whether they are at exponential or stationary phase of their growth. Interestingly, CORM effects varied between clinical isolates and were clearer with the isolates belonging to lineage A. While Murray *et al.* (2012) did not observe clear variation in CORM effects against planktonic cell cultures, they found differences in CORM efficiency in eradicating bacterial biofilms. This is also consistent with our data and reinforces the hypothesis that *P. aeruginosa* clinical isolates are likely to respond differently to CORM therapies.

To explore clinical isolate variation in more detail, we compared differences in bacterial virulence and growth traits between the PAO1 and clinical CF lung isolates. We found that relative to strain PAO1, clinical CF isolates grew slower, had lowered virulence and produced lower amounts of pyoverdine and pyocyanin, which are important virulence factors (O'Brien *et al.*, 2017). This is consistent with previous research and typical for *P. aeruginosa* isolates retrieved from chronic lung infections (Smith *et al.*, 2006; Folkesson *et al.*, 2012; Marvig *et al.*, 2013; Williams *et al.*, 2015). The clinical isolates produced much more

biofilm compared to strain PAO1 and biofilm formation was the highest in the isolates belonging to lineage B. Biofilms could potentially provide a protective function against CORMs. Biofilms often have much higher antibiotic resistance than their aquatic counterparts (Stewart and William Costerton, 2001) and there are multiple reasons for this. First, antibiotics might be ineffective because the biofilm acts as a diffusion barrier (de Beer et al., 1997). Second, subpopulations within the biofilm can sometimes differentiate into a highly protected phenotypes that can repopulate the biofilms (Cochran et al., 2000). Third, the biofilm might change the chemical microenvironment, forming zones of nutrient and oxygen depletion or waste accumulation that prevents the antibiotics from functioning optimally (de Beer et al., 1994). Although we did not explore this specifically, clinical isolates belonging to a lineage A were more susceptible to CORMs and produced relatively less biofilm compared to strains belonging to a lineage B. Thus, overall a negative correlation was found between density reduction by CORM and biofilm formation. Our results therefore suggest that biofilm might provide a protective function against the CORM.

Despite the isolate variations observed in bacterial monocultures, EBOR-CORM-1 was effective at reducing the growth of *P. aeruginosa* clinical isolate mixed cultures. One explanation for this is that, in addition to CORM, *P. aeruginosa* growth was limited by antagonistic intraspecific species interactions in co-cultures. *P. aeruginosa* has been shown to exert both facilitative and antagonistic effects on each other via siderophore (Harrison et al., 2008) and bacteriocin (Ghoul et al., 2015) production. In our case, all the clinical isolates were derived from the same Liverpool Epidemic Strain clonal lineage and therefore likely carried the same siderophore and bacteriocin genes. Additionally, resource competition is likely to further limit *P. aeruginosa* growth both in CF lungs and simplified laboratory microcosms. As a result, even though some clinical strains were relatively insensitive to EBOR-CORM-1, their growth could have been constrained by competition with the other strains in co-cultures. We found that this was the case for all strain mixes regardless if the strains belonged to a lineage A, B or them both. This suggests that the susceptibility of the lineages measured in monocultures did not predict the susceptibility of isolate mixes within or between lineages. However, such antagonism was not observed in the absence of EBOR-CORM-1, which suggests that CORM-triggered antagonistic intraspecific interactions in *P. aeruginosa* co-cultures. Mechanistically, this could have been driven by competition sensing in response to CORM-mediated cell damage in *P. aeruginosa* populations (Cornforth and Foster, 2013). However, this needs to be confirmed in future experiments. Interestingly, all the clinical strains we used originated from a single CF patient and interactions between them thus reflect the realistic ecology of CF lungs. In the future, it would be useful to determine

pairwise interactions between these CF strains and look at CORM effects on other coexisting bacterial species observed in CF infections (Folkesson et al., 2012).

We found that EBOR-CORM-1 had no clear therapeutic benefits in the wax moth infection model. There are several potential explanations for this. First, EBOR-CORM-1 had limited long-term activity when in contact with water. As a result, the bactericidal effect may only have elicited lag in the initial phase of bacterial growth and proliferation within the wax moths. Second, insect tissue is not homogeneous and it is possible that we failed to deliver the CORM to the specific area of infection, or that bacteria were able to colonise new areas that were not exposed to the CORM. Third, insects differ from laboratory media (such as LB) as a bacterial growth environment, which could also affect pathogen virulence. For example, it has been recently demonstrated that plant versus animal based growth media can have physiological effects on bacterial virulence (Ketola et al., 2016) and that LB media does not adequately reflect *P. aeruginosa* growth on lung tissue (Harrison et al., 2014; Harrison and Diggle, 2016). Hence, the wax moth injection model might not reliably reflect the virulence of CF isolates derived from chronic infections. However, it is also the case that many of the affordable and available CF infection animal models do not truly reflect the real CF lung disease environment. It remains to be established whether CORM therapy could be applied in the context of CF lung infections. It is possible, for example, that it might be more suitable for treating topical infections such as burn wounds, for which better animal models are available (Rumbaugh et al., 2012).

Further work is also needed to understand the mode of action of EBOR-CORM-1. While respiratory oxidases and globins at heme targets are generally considered the prime targets of CO and CORMs (Wareham et al., 2015), it has been demonstrated that CORMs can have multiple different other targets (Wilson et al., 2015). For example, CO also binds to the di-iron site in bacterial NO reductases and to iron, copper, and nickel sites in certain microbial proteins such as CO dehydrogenase (Lu et al., 2004; Wasser et al., 2005). In some cases, CORMs might have intracellular targets but their accumulation within the cells can be very weak (Tinajero-Trejo et al., 2016). Moreover, in the future it would be important to test if EBOR-CORM-1 is cytotoxic to eukaryotic cells. The concentration we used are in line with previously published work where no, or very mild, cytotoxic effects were observed (Murray et al., 2012). We are currently conducting experiments to validate this independently and to understand how EBOR-CORM-1 interacts with bacterial cells. While, our wax moth assays show that the concentrations we used had no negative effects on short-term insect viability, more detailed cytotoxicity assays are needed in the future. Lastly, the low solubility of

EBOR-CORM-1 in water, and its activation in this medium, is problematic for delivery and activation at specific sites within patients. In addition to chemically increasing the molecule stability, CORMs could be enclosed in microvesicles (van Dommelen et al., 2012) to ensure more efficient antimicrobial activity and drug delivery.


In conclusion, our results show that EBOR-CORM-1 shows antimicrobial activity against both planktonic and biofilm cells of *P. aeruginosa* strain PAO1 but that these effects are more varied and less pronounced against clinical CF lung isolates in monocultures. In contrast, more heterogeneous *P. aeruginosa* populations comprising intraspecific phenotypic variants were more susceptible to CORM treatment. This potentially has wider implications in the testing of novel therapeutics. At present, this is done almost exclusively using clonal *P. aeruginosa* populations. Our observations suggest that testing carried out on more heterogeneous populations of *P. aeruginosa*, more closely resembling those found in the CF lung, may give different and sometimes more promising results.

## ACKNOWLEDGEMENTS

We thank James Pitt for carrying out an initial EBOR-CORM-1 synthesis and TARGeTED Antimicrobial Resistance (AMR) Project and EPSRC council for funding (EP/M027538/1). Ville-Petri Friman is also supported by the Wellcome Trust [reference no. 105624] through the Centre for Chronic Diseases and Disorders (C2D2) at the University of York.


## FIGURES AND FIGURE LEGENDS






**Figure 1.** The stability of EBOR-CORM-1. Panel A shows the IR spectra of EBOR-CORM-1 upon heating at 50° C for 0 hours (black line), for 1 hours (red line), for 2 hours (blue) and for 3 hours (pink). The structure of the EBOR-CORM-1 is shown in inset on the left. Panel B shows the gas phase IR spectra of EBOR-CORM-1 in chloroform with added water where the \* indicates a band from chloroform. Panel C shows the IR spectra of EBOR-CORM-1 in chloroform (top) and water (bottom) and panel D the IR spectra of CO in water after 1 min (top) and 90 min (bottom) dissolution. All frequencies given are in cm<sup>-1</sup>.

501




502


503 **Figure 2.** The EBOR-CORM-1 effects on planktonic and biofilm cells of *P. aeruginosa*  
504 PAO1. In panel A, different lines denote for cell densities after 8 hours (blue line) and 24  
505 hours (green line) of EBOR-CORM-1 application and EBOR-CORM-1 effects on established  
506 cell cultures (red line) in different CORM concentrations. Panel B shows EBOR-CORM-1  
507 effects on PAO1 biofilms in different CORM concentrations. The  $R^2$  denotes for the fit of  
508 regression with our data, and in panel B, bars denote for  $\pm 1$  s.e.m.



**Figure 3.** The EBOR-CORM-1 effects on planktonic and biofilm cells of clinical *P. aeruginosa* CF isolates. Panels A and B show the proportion of living cells after 8 hours and 24 hours of EBOR-CORM-1 application, respectively. Panel C and D show the EBOR-CORM-1 effects on established cell cultures and biofilms, respectively. In all panels, bars denote for  $\pm 1$  s.e.m.



**Figure 4.** Differences in *P. aeruginosa* growth and virulence trait variation between PAO1 and clinical CF isolates (panel A) and EBOR-CORM-1 effects on clinical CF isolate lineage mixes (panel B). In panel A, different colours denote for pyocyanin (blue) and pyoverdine (green) production, time to death (black), growth rate (purple), maximum density (yellow) and biofilm production for clinical isolates belonging to lineages A and B. The dashed line shows the mean performance of PAO1 strain. Panel B shows EBOR-CORM-1 effect on clinical CF isolate mixes after 24 hours of CORM application. In panel A, bars denote for  $\pm 1$  s.e.m., and in panel B, extreme values around lower and upper quartile (black line shows the median).



528

529 **Figure 5.** The EBOR-CORM-1 activity against three *P. aeruginosa* strains in wax moth  
 530 model. Boxplots show larval survival in the absence (light grey) and presence (blue) of  
 531 EBOR-CORM-1 for PAO1, LESB58 and clinical isolate #36 (lineage B). Bars show extreme  
 532 values around lower and upper quartile and black lines show the median.

533 **REFERENCES**

- 534 Angelici, R.J. (1964). Preparation, Characterization, and Reactions of the cis-  
535 Dihalotetracarbonylmanganate (I) Anions. *Inorganic Chemistry* 3, 1099-1102.
- 536 Brown, E.D., and Wright, G.D. (2016). Antibacterial drug discovery in the resistance era.  
537 *Nature* 529, 336-343.
- 538 Cochran, W., Mcfeters, G., and Stewart, P. (2000). Reduced susceptibility of thin  
539 *Pseudomonas aeruginosa* biofilms to hydrogen peroxide and monochloramine.  
540 *Journal of applied microbiology* 88, 22-30.
- 541 Cornforth, D.M., and Foster, K.R. (2013). Competition sensing: the social side of bacterial  
542 stress responses. *Nature Reviews Microbiology* 11, 285-293.
- 543 De Beer, D., Stoodley, P., and Lewandowski, Z. (1997). Measurement of local diffusion  
544 coefficients in biofilms by microinjection and confocal microscopy. *Biotechnology*  
545 *and Bioengineering* 53, 151-158.
- 546 De Beer, D., Stoodley, P., Roe, F., and Lewandowski, Z. (1994). Effects of biofilm structures  
547 on oxygen distribution and mass transport. *Biotechnology and bioengineering* 43,  
548 1131-1138.
- 549 Desmard, M., Davidge, K.S., Bouvet, O., Morin, D., Roux, D., Foresti, R., Ricard, J.D.,  
550 Denamur, E., Poole, R.K., Montravers, P., Motterlini, R., and Boczkowski, J. (2009).  
551 A carbon monoxide-releasing molecule (CORM-3) exerts bactericidal activity against  
552 *Pseudomonas aeruginosa* and improves survival in an animal model of bacteraemia.  
553 *Faseb j* 23, 1023-1031.
- 554 El-Sayed, M., and Kaesz, H. (1963). Infrared Spectra and Structure of the Tetracarbonyl  
555 Halide Dimers of Manganese, Technetium, and Rhenium. *Inorganic Chemistry* 2,  
556 158-162.
- 557 Flume, P.A., Mogayzel, P.J., Robinson, K.A., Rosenblatt, R.L., Quittell, L., and Marshall,  
558 B.C. (2010). Cystic Fibrosis Pulmonary Guidelines. *American Journal of Respiratory*  
559 *and Critical Care Medicine* 182, 298-306.
- 560 Folkesson, A., Jelsbak, L., Yang, L., Johansen, H.K., Ciofu, O., Hoiby, N., and Molin, S.  
561 (2012). Adaptation of *Pseudomonas aeruginosa* to the cystic fibrosis airway: an  
562 evolutionary perspective. *Nature Reviews Microbiology* 10, 841-851.
- 563 Ghouli, M., West, S.A., Johansen, H.K., Molin, S., Harrison, O.B., Maiden, M.C., Jelsbak, L.,  
564 Bruce, J.B., and Griffin, A.S. (2015). Bacteriocin-mediated competition in cystic  
565 fibrosis lung infections. *Proc Biol Sci* 282.
- 566 Harrison, F., and Diggle, S.P. (2016). An ex vivo lung model to study bronchioles infected  
567 with *Pseudomonas aeruginosa* biofilms. *Microbiology* 162, 1755-1760.
- 568 Harrison, F., Muruli, A., Higgins, S., and Diggle, S.P. (2014). Development of an ex vivo  
569 porcine lung model for studying growth, virulence, and signaling of *Pseudomonas*  
570 *aeruginosa*. *Infection and immunity* 82, 3312-3323.
- 571 Harrison, F., Paul, J., Massey, R.C., and Buckling, A. (2008). Interspecific competition and  
572 siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *ISME J* 2, 49-55.
- 573 Jimenez, J., Chakraborty, I., Carrington, S.J., and Mascharak, P.K. (2016). Light-triggered  
574 CO delivery by a water-soluble and biocompatible manganese photoCORM. *Dalton*  
575 *Trans* 45, 13204-13213.
- 576 Ketola, T., Mikonranta, L., Laakso, J., and Mappes, J. (2016). Different food sources elicit  
577 fast changes to bacterial virulence. *Biol Lett* 12, 20150660.
- 578 Klein, M., Neugebauer, U., Gheisari, A., Malassa, A., Jazzazi, T.M., Froehlich, F.,  
579 Westerhausen, M., Schmitt, M., and Popp, J.R. (2014). IR spectroscopic methods for  
580 the investigation of the CO release from CORMs. *The Journal of Physical Chemistry*  
581 *A* 118, 5381-5390.

- Lu, S., Suharti, De Vries, S., and Moënne-Loccoz, P. (2004). Two CO Molecules Can Bind Concomitantly at the Diiron Site of NO Reductase from *Bacillus a zotoformans*. *Journal of the American Chemical Society* 126, 15332-15333.
- Marvig, R.L., Johansen, H.K., Molin, S., and Jelsbak, L. (2013). Genome analysis of a transmissible lineage of *Pseudomonas aeruginosa* reveals pathoadaptive mutations and distinct evolutionary paths of hypermutators. *PLoS Genet* 9, e1003741.
- Motterlini, R., and Otterbein, L.E. (2010). The therapeutic potential of carbon monoxide. *Nat Rev Drug Discov* 9, 728-743.
- Mowat, E., Paterson, S., Fothergill, J.L., Wright, E.A., Ledson, M.J., Walshaw, M.J., Brockhurst, M.A., and Winstanley, C. (2011). *Pseudomonas aeruginosa* population diversity and turnover in cystic fibrosis chronic infections. *Am J Respir Crit Care Med* 183, 1674-1679.
- Murray, T.S., Okegbe, C., Gao, Y., Kazmierczak, B.I., Motterlini, R., Dietrich, L.E., and Bruscia, E.M. (2012). The carbon monoxide releasing molecule CORM-2 attenuates *Pseudomonas aeruginosa* biofilm formation. *PLoS One* 7, e35499.
- Nobre, L.S., Seixas, J.D., Romao, C.C., and Saraiva, L.M. (2007). Antimicrobial action of carbon monoxide-releasing compounds. *Antimicrob Agents Chemother* 51, 4303-4307.
- Normark, B.H., and Normark, S. (2002). Evolution and spread of antibiotic resistance. *Journal of internal medicine* 252, 91-106.
- O'brien, S., Williams, D., Fothergill, J.L., Paterson, S., Winstanley, C., and Brockhurst, M.A. (2017). High virulence sub-populations in *Pseudomonas aeruginosa* long-term cystic fibrosis airway infections. *BMC Microbiol* 17, 30.
- O'Neill, J. (2014). "Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations", in: *Review on Antimicrobial Resistance*. HM Government & Wellcome Trust ).
- Poole, K. (2011). *Pseudomonas aeruginosa*: resistance to the max. *Front Microbiol* 2, 65.
- Pritt, B., O'brien, L., and Winn, W. (2007). Mucoid *Pseudomonas* in cystic fibrosis. *Am J Clin Pathol* 128, 32-34.
- Reszka, K.J., O'malley, Y., McCormick, M.L., Denning, G.M., and Britigan, B.E. (2004). Oxidation of pyocyanin, a cytotoxic product from *Pseudomonas aeruginosa*, by microperoxidase 11 and hydrogen peroxide. *Free Radic Biol Med* 36, 1448-1459.
- Rumbaugh, K.P., Trivedi, U., Watters, C., Burton-Chellew, M.N., Diggle, S.P., and West, S.A. (2012). Kin selection, quorum sensing and virulence in pathogenic bacteria. *Proc Biol Sci* 279, 3584-3588.
- Smith, E.E., Buckley, D.G., Wu, Z., Saenphimmachak, C., Hoffman, L.R., D'argenio, D.A., Miller, S.I., Ramsey, B.W., Speert, D.P., Moskowitz, S.M., Burns, J.L., Kaul, R., and Olson, M.V. (2006). Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proceedings of the National Academy of Sciences of the United States of America* 103, 8487-8492.
- Stamellou, E., Storz, D., Botov, S., Ntasis, E., Wedel, J., Sollazzo, S., Kramer, B.K., Van Son, W., Seelen, M., Schmalz, H.G., Schmidt, A., Hafner, M., and Yard, B.A. (2014). Different design of enzyme-triggered CO-releasing molecules (ET-CORMs) reveals quantitative differences in biological activities in terms of toxicity and inflammation. *Redox Biol* 2, 739-748.
- Stewart, P.S., and William Costerton, J. (2001). Antibiotic resistance of bacteria in biofilms. *The Lancet* 358, 135-138.
- Tinajero-Trejo, M., Rana, N., Nagel, C., Jesse, H.E., Smith, T.W., Wareham, L.K., Hippler, M., Schatzschneider, U., and Poole, R.K. (2016). Antimicrobial Activity of the Manganese Photoactivated Carbon Monoxide-Releasing Molecule [Mn (CO) <sub>3</sub> (tpa-κ<sup>3</sup> N)]<sup>+</sup> Against a Pathogenic *Escherichia coli* that Causes Urinary Infections. *Antioxidants & redox signaling* 24, 765-780.

- Van Dommelen, S.M., Vader, P., Lakhal, S., Kooijmans, S., Van Solinge, W.W., Wood, M.J., and Schiffelers, R.M. (2012). Microvesicles and exosomes: opportunities for cell-derived membrane vesicles in drug delivery. *Journal of Controlled Release* 161, 635-644.
- Ward, J.S., Lynam, J.M., Moir, J., and Fairlamb, I.J. (2014). Visible - Light - Induced CO Release from a Therapeutically Viable Tryptophan - Derived Manganese (I) Carbonyl (TryptoCORM) Exhibiting Potent Inhibition against E. coli. *Chemistry-A European Journal* 20, 15061-15068.
- Ward, J.S., Morgan, R., Lynam, J.M., Fairlamb, I.J., and Moir, J.W. (2017). Toxicity of tryptophan manganese (I) carbonyl (Trypto-CORM), against *Neisseria gonorrhoeae*. *MedChemComm* 8, 346-352.
- Wareham, L.K., Poole, R.K., and Tinajero-Trejo, M. (2015). CO-releasing Metal Carbonyl Compounds as Antimicrobial Agents in the Post-antibiotic Era. *J Biol Chem* 290, 18999-19007.
- Wasser, I.M., Huang, H.-W., Moënné-Loccoz, P., and Karlin, K.D. (2005). Heme/non-heme diiron (II) complexes and O<sub>2</sub>, CO, and NO adducts as reduced and substrate-bound models for the active site of bacterial nitric oxide reductase. *Journal of the American Chemical Society* 127, 3310-3320.
- Who.Int (2010). *WHO | Genes and human disease* [Online]. Available: <http://www.who.int/genomics/public/geneticdiseases/en/index2.html> [Accessed 16-02 2017].
- Williams, D., Evans, B., Haldenby, S., Walshaw, M.J., Brockhurst, M.A., Winstanley, C., and Paterson, S. (2015). Divergent, coexisting *Pseudomonas aeruginosa* lineages in chronic cystic fibrosis lung infections. *Am J Respir Crit Care Med* 191, 775-785.
- Williams, D., Paterson, S., Brockhurst, M.A., and Winstanley, C. (2016). Refined analyses suggest that recombination is a minor source of genomic diversity in *Pseudomonas aeruginosa* chronic cystic fibrosis infections. *Microbial Genomics* 2.
- Wilson, J.L., Wareham, L.K., Mclean, S., Begg, R., Greaves, S., Mann, B.E., Sanguinetti, G., and Poole, R.K. (2015). CO-Releasing Molecules Have Nonheme Targets in Bacteria: Transcriptomic, Mathematical Modeling and Biochemical Analyses of CORM-3 [Ru(CO)<sub>3</sub>Cl(glycinate)] Actions on a Heme-Deficient Mutant of *Escherichia coli*. *Antioxid Redox Signal* 23, 148-162.
- Winstanley, C., Langille, M.G., Fothergill, J.L., Kukavica-Ibrulj, I., Paradis-Bleau, C., Sanschagrin, F., Thomson, N.R., Winsor, G.L., Quail, M.A., Lennard, N., Bignell, A., Clarke, L., Seeger, K., Saunders, D., Harris, D., Parkhill, J., Hancock, R.E., Brinkman, F.S., and Levesque, R.C. (2009). Newly introduced genomic prophage islands are critical determinants of in vivo competitiveness in the Liverpool Epidemic Strain of *Pseudomonas aeruginosa*. *Genome Res* 19, 12-23.
- Winstanley, C., O'brien, S., and Brockhurst, M.A. (2016). *Pseudomonas aeruginosa* Evolutionary Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. *Trends Microbiol* 24, 327-337.
- Workentine, M.L., Sibley, C.D., Glezerson, B., Purighalla, S., Norgaard-Gron, J.C., Parkins, M.D., Rabin, H.R., and Surette, M.G. (2013). Phenotypic heterogeneity of *Pseudomonas aeruginosa* populations in a cystic fibrosis patient. *PLoS One* 8, e60225.
- Zhang, W.-Q., Atkin, A.J., Thatcher, R.J., Whitwood, A.C., Fairlamb, I.J., and Lynam, J.M. (2009). Diversity and design of metal-based carbon monoxide-releasing molecules (CO-RMs) in aqueous systems: revealing the essential trends. *Dalton Transactions*, 4351-4358.