

This is a repository copy of *Manganese(II)-Catalyzed C–H Activation: The Key Role of a 7-Membered Manganacycle in H-Transfer and Reductive Elimination*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/105600/>

Version: Published Version

Article:

Yahaya, Nasiru P., Appleby, Kate M., Teh, Magdalene et al. (11 more authors) (2016) Manganese(II)-Catalyzed C–H Activation: The Key Role of a 7-Membered Manganacycle in H-Transfer and Reductive Elimination. *Angewandte Chemie International Edition*. pp. 12455-12459. ISSN 1433-7851

<https://doi.org/10.1002/anie.201606236>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

C–H Activation

International Edition: DOI: 10.1002/anie.201606236
German Edition: DOI: 10.1002/ange.201606236

Manganese(I)-Catalyzed C–H Activation: The Key Role of a 7-Membered Manganacycle in H-Transfer and Reductive Elimination

Nasiru P. Yahaya, Kate M. Appleby, Magdalene Teh, Conrad Wagner, Erik Troschke, Joshua T. W. Bray, Simon B. Duckett, L. Anders Hammarback, Jonathan S. Ward, Jessica Milani, Natalie E. Pridmore, Adrian C. Whitwood, Jason M. Lynam,* and Ian J. S. Fairlamb*

Dedicated to Professors Michael Bruce and Robin N. Perutz

Abstract: Manganese-catalyzed C–H bond activation chemistry is emerging as a powerful and complementary method for molecular functionalization. A highly reactive seven-membered Mn^I intermediate is detected and characterized that is effective for H-transfer or reductive elimination to deliver alkenylated or pyridinium products, respectively. The two pathways are determined at Mn^I by judicious choice of an electron-deficient 2-pyrone substrate containing a 2-pyridyl directing group, which undergoes regioselective C–H bond activation, serving as a valuable system for probing the mechanistic features of Mn C–H bond activation chemistry.

C–H bond activation–functionalization chemistry is a central arena for catalyst development and synthetic application.^[1] Transition metals mediate the efficient and selective activation of C–H bonds, with recent attention focusing on environmentally benign and sustainable metals, for example, Mn, Co, Fe, and Cu.^[2] Mn^I promotes C–H activation of substrates containing nitrogen-directing groups.^[3] For example, **1** gives cyclomanganated complex **2**, with subsequent reaction with alkyne **3** forming a proposed 7-membered ring intermediate **4** (Scheme 1).^[4] Formation of either **5**, **6**, or **7** results from reductive elimination, H-transfer, or dehydrogenative annulation, respectively.

Processes utilizing Mn^I , particularly $[Mn(C^{\wedge}N)(CO)_4]$ **2**,^[5,6] have been of broad interest. The mechanistic features of the remarkable synthetic work of Ackermann and Wang,^[3,4] where intermediates **4a–c** have been proposed,

prompted us to examine whether they could be detected and characterized and then subsequently be shown to deliver organic products such as **5–7**. Complexes **4d–f**, formed by insertion of internal alkynes are known,^[6,7] but their competence in terms of a fully connected reaction system, affording organic products, has not been examined. As 18-electron species containing four CO ligands, possessing high thermodynamic stability, they are unlikely to be directly involved in the catalytic cycle.^[8]

Herein we describe a suitable reaction system (**1g**→**4g**→**5g** or **6g**, Scheme 1) that takes advantage of the exquisite reactivity of an electron-deficient 2-pyrone ring system containing a 2-pyridyl directing group (**1g**). We recognized that the 2-pyrone could act as a hemilabile ligand in 7-membered manganacycle **4g**, potentially providing sufficient stabilisation for observation of this key intermediate. Our findings demonstrate that **4g** acts as a central manifold to reductive elimination and H-transfer, giving products **5g** and **6g**, respectively, with details described herein.


Our study began with the reaction of 2-pyrone **1g** with $BnMn(CO)_5$ in hexane at 75 °C, which gave cyclometalated **2g** cleanly and in quantitative yield (Scheme 2). Complex **2g** was fully characterized (see the Supporting Information); a single crystal X-ray structure confirmed that regioselective C–H activation occurred at C3, in keeping with Pd^{II} -direct arylations of 2-pyrones,^[9] albeit most likely by a σ -CAM-type process.^[10]

We hypothesized that UV irradiation^[11] of **2g** would lead to solvated intermediate I_{Pyr} (Scheme 2, middle inset).^[12] Subsequent alkyne trapping via intermediate II_{Pyr} , would then convert into the alkyne insertion manganacycle **4g**. UV irradiation (Hg/Xe Arc lamp, 200–2500 nm) of a mixture of **2g** and **3** (1.1 equiv) in $[D_8]THF$ at 240 K (at 5 min intervals), and reaction monitoring by 1H NMR spectroscopy between intervals, revealed the formation of a new intermediate that grows up to 9.6 % conversion. Further irradiation resulted in spectral broadening (paramagnetic species), but crucially, full NMR analysis of manganacycle **4g** was possible, with HMQC/HMBC correlation methods/n.O.e. experiments. Analysis shows that **4g** formed regioselectively at C3 (Scheme 2, bottom inset). MS analysis also confirmed the presence of **4g** (LIFDI m/z 427 for $[M]^+$ and ESI m/z 428 for $[MH]^+$) in solution.

[*] Dr. N. P. Yahaya, K. M. Appleby, M. Teh, C. Wagner, E. Troschke, J. T. W. Bray, Prof. S. B. Duckett, L. A. Hammarback, Dr. J. S. Ward, J. Milani, Dr. N. E. Pridmore, Dr. A. C. Whitwood, Dr. J. M. Lynam, Prof. I. J. S. Fairlamb
Department of Chemistry, University of York
York, YO10 5DD (UK)
E-mail: jason.lynam@york.ac.uk
ian.fairlamb@york.ac.uk

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<http://dx.doi.org/10.1002/anie.201606236>.

© 2016 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Scheme 1. Manganese(I)-catalyzed C–H activation, and potential products and intermediates.


Experimentally there is evidence in **4g** of an interaction between the 2-pyrone olefinic bond (C6–C11) and the Mn^I center at $\delta = 159.7$ ppm (C6) and $\delta = 145.9$ ppm (C11), which stabilizes the tricarbonyl complex. Computational studies (DFT methods) confirm that HOMO–4 within **4g** has 2-pyrone–Mn bonding character (see the Supporting Information), confirming **4g** as a feasible structure. The small coordination shifts in the ¹³C{¹H} NMR spectrum imply this interaction is weak, although generation of a vacant site at Mn (**4g'**) and subsequent alkyne coordination (**4g''**) ought to be feasible. The DFT studies for **III_{pyr}** (**4g**) and **III_{ph}** (**4a**) indicate no low-lying vacant orbitals (HOMO–LUMO gap = 1.70154–1.97588 eV), consistent with Mn having an 18-electron count.

Warming of the [D₈]THF solution of **4g** to room temperature led to the formation of the reductive elimination product **5g** (Scheme 3). Complex **5g** was fully characterized (see the Supporting Information) and confirmed by X-ray analysis to possess a Mn(CO)₃ anion. **5g** was also formed in 87% yield on treatment of **2g** with **3** (1.1 equiv.) at 80 °C, Et₂O, 18 h (sealed tube). Thus, the same reaction pathway (**2g** + **3** → **5g**) results from either UV irradiation or thermal heating, validating our approach in utilizing UV irradiation to enable detection and characterization of intermediate **4g**.

Interestingly, catalytic reactions of **1g** with **3**, under the reaction conditions reported by Wang et al.^[4] for 2-phenyl-




Scheme 2. Cyclomanganation of **1g** gives **2g**, which upon photolysis with phenylacetylene **3** gives **4g**. The X-ray structure of **2g** is given (top right, ellipsoids set at 50% probability; H-atoms omitted and Mn atom labeled only for clarity). Insets: proposed transient intermediates on route to **4g** and the key NMR data for **4g**.



Scheme 3. Thermally controlled reductive elimination from either **2g** or **4g** to give **5g**. An X-ray structure of a single crystal of **5g** is also shown (ellipsoids set to 50% probability; H-atoms omitted and Mn atom labeled only, for clarity).


pyridine **1a** (conditions: BrMn(CO)₅, Cy₂NH, Et₂O, 100 °C for 6–24 h), do not lead to formation of alkenylated products (for example, **6g**). This indicates that the rate of reductive elimination from **4g** to give **5g** is faster than the rate for alkyne H-transfer to give **6g** (see above). We rationalized that reaction of **2g** in neat phenylacetylene **3** would enable H-transfer to become the dominant pathway (Scheme 4), but the reaction afforded three new products. Firstly, the H-transfer product **6g** was formed in 28% yield; an excess of **3** favors H-transfer over reductive elimination. Central to the success of the reaction is coordination of a second molecule of alkyne **3** and subsequent alkyne H-transfer of intermediate **4g**. The other products **8** and **9** were unexpected, resulting from a noteworthy Diels–Alder reaction (DAR) of **3** with the 2-pyridine ring,^[13] followed by ring fragmentation (single-



Scheme 4. Reaction of **2g** in neat phenylacetylene **3**. The green bonds show the newly formed bonds in the organic products, with red showing the insertion location of **3** (**5g** not observed under these reaction conditions).

crystal X-ray structures of **8** and **9** confirmed the molecular connectivity, correlating with NMR spectroscopy, see the Supporting Information). Compound **9** shows that the 2-pyrone participated in a secondary inverse electron demand DAR.^[14] Along with **6g**, both **8** and **9** derive from **4g**, where the DARs and 2-pyridyl fragmentation are secondary reactions.

To understand the steps leading to the formation of **5g** DFT methods were used (Scheme 5, see the Supporting




Scheme 5. DFT calculations showing the feasibility of reductive elimination from **5a** and **5g**, starting from intermediates **II_{ph}** and **II_{Pyr}** respectively. Energies are zero point energy-corrected electronic energies and Gibbs energies at 298.15 K in kJ mol⁻¹ relative to **II**.

Information for details of DFT calculations). Starting from **II_{Pyr}**, formed via loss of CO from **2g** and coordination of **3**, insertion of coordinated alkyne into the Mn–C(pyrone) bond proceeds through a low-energy transition state (**TS_{IIPyr-IIPyr}**) to give **III_{Pyr}**. The latter intermediate is equivalent to characterized **4g**. C–N reductive elimination from **III_{Pyr}** via transition state **TS_{IIPyr-5g-iso}**, results in the formation of the 2-methyl-4-oxo-6-phenyl-4H-3,7λ⁵-pyrano[4,3-a]quinolizin-7-ylum ring system (**5g**). A DRC analysis of **TS_{IIPyr-5g-iso}** revealed that the imaginary eigenvector led to **5g-iso** (the coordination isomer of **5g**); a π -slip then gives **5g**.

The corresponding potential energy surface for the phenyl-substituted system (giving the Chen and Wang product **5a**) revealed that the same reaction pathway was viable (pathway shown in gray in Scheme 5). The barrier to insertion of **3** (**TS_{II-III}**) was slightly greater (Gibbs energies at 298.15 K

relative to the respective compound **II** + 25 kJ mol⁻¹ for 2-pyrone versus + 34 kJ mol⁻¹ for phenyl) and that **III_{Pyr}** was relatively higher in energy than **III_{ph}** (–76 kJ mol⁻¹ versus –95 kJ mol⁻¹). To explain the different outcome from the phenyl and 2-pyrone substituents it is informative to consider the higher energy of **TS_{IIPh-5a-iso}** (+ 26 kJ mol⁻¹) against **TS_{IIPyr-5g-iso}** (–16 kJ mol⁻¹). Therefore, the energetic spans for reductive elimination are 60 kJ mol⁻¹ (2-pyrone) and 121 kJ mol⁻¹ (phenyl). When compared with the formation of **IV_{Pyr}** and **IV_{ph}**, which is the next step in forming H-transfer products **5g** and **5a**, respectively, it is evident that the reductive elimination to form **5g** is competitive, but in the case of **5a** the much larger energetic span to reductive elimination allows for productive catalysis via alkyne coordination to give **IV_{ph}**.^[4]


While no double alkyne insertion products were detected in reactions of **2g** with phenylacetylene **3**, the reaction of related derivative **2h** with **3** resulted in exclusive formation of double alkyne insertion product **10** (Scheme 6; the structure



Scheme 6. Double alkyne insertion into **2h**. Dotted lines show Mn coordination in complex **10** for clarity (ellipsoids set to 50% probability; H-atoms omitted and Mn atom labeled only, for clarity).

of **6h** is shown as an expected alkenylated product). This remarkable result shows the impact that a subtle change to the pyridyl directing group has on the barriers to these steps.

We rationalized the experimental observations by DFT calculations, which enabled a mechanism for this reaction and the differences between the phenyl- and 2-pyrone-substituted complexes to be proposed (Scheme 7). In the case of the pyrone derivative, coordination of alkyne to **III_{Pyr}** results in formation of **IV_{Pyr}** having two energetically accessible fates. H-transfer through **TS_{IVPyr-VIPyr}** (+ 3 kJ mol⁻¹) results in the formation of alkynyl complex **V_{Pyr}** which would liberate **6h**, however insertion of the alkyne into the Mn–C bond of **IV_{Pyr}** through **TS_{IVPyr-VIPyr}** (+ 4 kJ mol⁻¹) affords more energetically favourable **VI_{Pyr}**. The process seen for reactions of **2h** has



Scheme 7. DFT calculations showing the feasibility of a double alkyne insertion pathway to rationalize formation of double alkyne insertion product **10**. Energies are zero point energy-corrected electronic energies and Gibbs energies at 298.15 K in kJ mol^{-1} relative to **II**.^[17]

resulted in the formation of two C–C bonds. Preliminary investigations indicate that this proceeds through a “two-steps no intermediate” pathway^[15] with the initial insertion into the Mn–C bond, followed by cyclization giving a six-membered ring without an intermediate. However, in **VI_{Pyr}**, the Mn is η^3 -coordinated to the pendant pyridyl group and newly formed ring. To form **10_{Pyr}**, which is the lowest point on the potential energy surface at -320 kJ mol^{-1} , the Mn needs to migrate to the alternative ring-face. We postulate that this involves migration onto one of the phenyl rings in the ligand, for example, **VIIa_{Pyr}**. The ring rotates allowing the Mn to migrate to the other face of the pentadienyl system, giving **VIIb_{Pyr}**. It is reasonable to presume that this proceeds via a low energy ring-walking process.^[16]

In the case of the phenyl derivative, all of the states predicted for the 2-pyrone system are viable; however, $TS_{IVPh-VPh}$ is far higher in energy than $TS_{IVPh-VPh}$ ($+41 \text{ kJ mol}^{-1}$ versus -1 kJ mol^{-1}). Therefore, insertion of the second alkyne is non-competitive, with the H-transfer pathway leading to the alkenylated product, consistent with experimental observations.

In conclusion, we have detected and characterized a commonly proposed 7-membered mangana-cycle **4g** (of direct relevance to generic structure **4**, Scheme 1). Mangana-cycle **4g** sits at the selectivity junction to reductive elimination or H-transfer steps. Depending on the reaction conditions, **5g** or **6g** products form that correspond to reductive elimination and protonation pathways, respectively. Double alkyne insertion to give **10** has also been revealed in these studies. Our observations provide the first clear cut evidence that mangana-cycles such as **4** are key intermediates in Mn^I -mediated C–H bond activation processes involving substrates containing directing groups.^[3,4,7] More generally, such intermediates

may be considered as leading to side reactions, but here we have shown that it presents an opportunity to control product selectivity. Serendipitously we have uncovered a rare example of a DAR of a pyridine derivative, where the intermediate fragments to form products such as **8** and **9**. Taken together, our findings provide a unique insight into Mn^I -mediated C–H bond activation processes, especially how relatively minor changes in substrate structure influence product selection; Mn^I -based metallocycles clearly offer rich chemistry,^[3] much potential, and warrant further study more generally in organic and organometallic chemistry.

Acknowledgements

EPSRC funded the computational equipment used in this study (ref. EP/H011455/ and EPSRC “ENERGY” grant, ref. no. EP/K031589/1). EPSRC PhD studentships to KMA and LAR are gratefully acknowledged (EP/K503216/1 and EP/N509413/1). We acknowledge PTDF (Nigeria) for funding the Ph.D. studies of N.P.Y. We acknowledge the ERASMUS Scheme for supporting the research visits of C.W. and E.T. We are grateful to Christopher Morrell for assistance with this study.

Keywords: catalysis · C–H activation · functionalization · manganese · sustainability

How to cite: *Angew. Chem. Int. Ed.* **2016**, *55*, 12455–12459
Angew. Chem. **2016**, *128*, 12643–12647

- [1] a) *C–H and C–X Bond Functionalization. Transition Metal Mediation* (Ed.: X. Ribas), RSC Catalysis, No. 11, **2013**; b) *Metal Catalyzed Cross-Coupling Reactions and More* (Eds.: A. de Meijere, S. Bräse, M. Oestreich), Wiley-VCH, Weinheim, **2014**; c) “Syntheses via C–H Bond Functionalizations”: L. Ackermann, A. R. Kapdi, H. K. Potukuchi, S. I. Kozhushkov, in *Handbook of Green Chemistry* (Eds.: C.-J. Li), Wiley-VCH, Weinheim, **2012**, pp. 259–305.
- [2] Very recent examples, at Co, see: a) H. Wang, J. Koeller, W. Liu, L. Ackermann, *Chem. Eur. J.* **2015**, *21*, 15525; At Mn, see: b) W. Liu, J. Bang, Y. Zhang, L. Ackermann, *Angew. Chem. Int. Ed.* **2015**, *54*, 14137; *Angew. Chem.* **2015**, *127*, 14343; c) W. Liu, S. C. Richter, Y. Zhang, L. Ackermann, *Angew. Chem. Int. Ed.* **2016**, *55*, 7747; *Angew. Chem.* **2016**, *128*, 7878; at Fe, see: d) D. Gärtner, A. L. Stein, S. Grupe, J. Arp, A. J. von Wangelin, *Angew. Chem. Int. Ed.* **2015**, *54*, 10545; *Angew. Chem.* **2015**, *127*, 10691; e) T. Jia, C. Zhao, R. He, H. Chen, C. Wang, *Angew. Chem. Int. Ed.* **2016**, *55*, 5268; *Angew. Chem.* **2016**, *128*, 5354.
- [3] For recent reviews, see: a) C. Y. Wang, *Synlett* **2013**, *24*, 1606; b) W. Liu, L. Ackermann, *ACS Catal.* **2016**, *6*, 3743.
- [4] a) B. Zhou, H. Chen, C. Wang, *J. Am. Chem. Soc.* **2013**, *135*, 1264; for selected other catalytic C–H bond functionalization reactions mediated by Mn, see: b) Y. Kuninobu, Y. Nishina, T. Takeuchi, K. Takai, *Angew. Chem. Int. Ed.* **2007**, *46*, 6518; *Angew. Chem.* **2007**, *119*, 6638; c) M. Yoshikai, S. Zhang, K. Yamagata, H. Tsuji, E. Nakamura, *J. Am. Chem. Soc.* **2009**, *131*, 4099; d) Y. Kuninobu, A. Kawata, M. Nishi, S. S. Yudha, J.-J. Chen, *Chem. Asian J.* **2009**, *4*, 1424; e) L. Shi, X. Zhong, H. She, Z. Lei, F. Li, *Chem. Commun.* **2015**, *51*, 7136.
- [5] J. S. Ward, J. M. Lynam, J. W. B. Moir, D. E. Sanin, A. P. Mountford, I. J. S. Fairlamb, *Dalton Trans.* **2012**, *41*, 10514.

- [6] a) M. I. Bruce, B. L. Goodall, I. Matsuda, *Aust. J. Chem.* **1975**, 28, 1259; b) W. Tully, L. Main, B. K. Nicholson, *J. Organomet. Chem.* **1996**, 507, 103; c) W. Tully, L. Main, B. K. Nicholson, *J. Organomet. Chem.* **2001**, 633, 162; d) W. J. Grigsby, L. Main, B. K. Nicholson, *Organometallics* **1993**, 12, 397; e) A. Suárez, F. Faraldo, J. M. Vila, H. Adams, A. Fernández, M. López-Torres, J. J. Fernández, *J. Organomet. Chem.* **2002**, 656, 270.
- [7] Key references: a) R. He, Z.-T. Huang, Q.-Y. Zheng, C. Wang, *Angew. Chem. Int. Ed.* **2014**, 53, 4950; *Angew. Chem.* **2014**, 126, 5050; b) B. Zhou, P. Ma, H. Chen, C. Wang, *Chem. Commun.* **2014**, 50, 14558; c) W. Liu, D. Zell, M. John, L. Ackermann, *Angew. Chem. Int. Ed.* **2015**, 54, 4092; *Angew. Chem.* **2015**, 127, 4165.
- [8] J. Hartwig, *Organotransition Metal Chemistry From Bonding to Catalysis*, University Science Books, Sausalito, **2010**, p. 233.
- [9] M. J. Burns, R. J. Thatcher, R. J. K. Taylor, I. J. S. Fairlamb, *Dalton Trans.* **2010**, 39, 10391.
- [10] R. N. Perutz, S. Sabo-Etienne, *Angew. Chem. Int. Ed.* **2007**, 46, 2578; *Angew. Chem.* **2007**, 119, 2630.
- [11] A control experiment established no reaction of **2a** with phenylacetylene **3a** (1.1 equiv) in [D₈]THF at ambient temperature.
- [12] a) I. R. Farrell, P. Matousek, M. Towrie, A. W. Parker, D. C. Grills, M. W. George, A. Vlček, *Inorg. Chem.* **2002**, 41, 4318; b) A. Vlček, I. R. Farrell, D. J. Liard, P. Matousek, M. Towrie, A. W. Parker, D. C. Grills, M. W. George, *J. Chem. Soc. Dalton Trans.* **2002**, 701.
- [13] DARs of pyridines are rare. For a report showing how tungsten can aid DARs of pyridines with alkenes, see: a) P. M. Graham, D. A. Delafuente, W. Liu, W. H. Myers, M. Sabat, W. D. Harman, *J. Am. Chem. Soc.* **2005**, 127, 10568; b) B. Halford, *Chem. Eng. News* **2005**, 83, 11; in our case we believe that the Mn is necessary for the DARs affording both **8** and **9**—mass spectrometric evidence for coordination of a Mn(CO)₃ fragment to **8** has been gained (see the Supporting Information).
- [14] R. T. Kohl, T. Katto, J. N. Braham, J. K. Stille, *Macromolecules* **1978**, 11, 340.
- [15] D. A. Singleton, C. Hang, M. J. Szymanski, M. P. Meyer, A. G. Leach, K. T. Kuwata, J. S. Chen, A. Greer, C. S. Foote, K. N. Houk, *J. Am. Chem. Soc.* **2003**, 125, 1319.
- [16] a) M. Orbach, S. Shankar, O. V. Zenkina, P. Milko, Y. Diskin-Posner, M. E. van der Boom, *Organometallics* **2015**, 34, 1098; b) O. V. Zenkina, A. Karton, D. Freeman, L. J. W. Shimon, J. M. L. Martin, M. E. van der Boom, *Inorg. Chem.* **2008**, 47, 5114.
- [17] The calculations support the difference in behavior between the phenyl- and pyrone-substituted complexes; however, the potential energy surface of the reaction of the OMe-substituted pyridine complex, **2h**, shows no significant differences to the unsubstituted case. We postulate that the lack of DAR chemistry in the OMe case permits the observation of **10**.

Received: June 27, 2016

Published online: September 7, 2016