

This is a repository copy of *Efficacy and cost-effectiveness of a community-based smoke-free-home intervention with or without indoor-air quality feedback in Bangladesh (MCLASS II): a three-arm, cluster-randomised, controlled trial.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/170269/>

Version: Accepted Version

Article:

Mdege, Noreen Dadirai orcid.org/0000-0003-3189-3473, Fairhurst, Caroline Marie orcid.org/0000-0003-0547-462X, Wang, Han I orcid.org/0000-0002-3521-993X et al. (12 more authors) (2021) Efficacy and cost-effectiveness of a community-based smoke-free-home intervention with or without indoor-air quality feedback in Bangladesh (MCLASS II): a three-arm, cluster-randomised, controlled trial. *The Lancet Global Health*. e639-e650. ISSN 2214-109X

[https://doi.org/10.1016/S2214-109X\(21\)00040-1](https://doi.org/10.1016/S2214-109X(21)00040-1)

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Title: Muslim Communities Learning About Second-hand Smoke in Bangladesh (MCLASS II): a three-arm, cluster randomised controlled trial of the effectiveness and cost-effectiveness of a community-based smoke-free homes intervention, with or without indoor air quality feedback

Author names and highest degrees

Noreen Dadirai Mdege, PhD^{1*}

Caroline Fairhurst, MSc²

Han-I Wang, PhD¹

Tarana Ferdous, MPH³

Anna-Marie Marshall, PhD¹

Catherine Hewitt, PhD²

Rumana Huque, PhD^{3,4}

Cath Jackson, PhD⁵

Ian Kellar, DPhil⁶

Steve Parrott, MSc¹

Sean Semple, PhD⁷

Aziz Sheikh, MD⁸

Qi Wu, MSc¹

Zunayed Al Azdi, MPH³

Kamran Siddiqi, PhD^{1,9}

On behalf of the MCLASS II Trial Team

Author affiliations and addresses

¹Department of Health Sciences, Faculty of Sciences, University of York, York YO10 5DD, UK

²York Trials Unit, Department of Health Sciences, Faculty of Sciences, University of York, York YO10 5DD, UK

³ARK Foundation, Suite C-3, C-4, House number 06, Road 109, Dhaka - 1212, Bangladesh

⁴Department of Economics, Dhaka University, Dhaka, Bangladesh

⁵Valid Research Ltd, Sandown House, Sandbeck Way, Wetherby LS22 7DN, UK

⁶School of Psychology, Faculty of Medicine and Health, University of Leeds, Leeds LS2 9JT, UK

⁷Institute for Social Marketing and Health, University of Stirling, Stirling FK9 4LA, UK

⁸Usher Institute, University of Edinburgh, Edinburgh EH8 9AG, UK

⁹Hull York Medical School, University of York, Heslington, York YO10 5DD, UK

* Corresponding author: noreen.mdege@york.ac.uk; tel: +44 1904 321836; Seebohm Rowntree Building, Health Sciences, University of York, Heslington, York YO10 5DD, UK

Word count: 4491

SUMMARY

Background

Exposure to second-hand smoke (SHS) from tobacco is a major contributor to global morbidity and mortality. We evaluated the effectiveness and cost-effectiveness of a community-based smoke-free homes (SFH) intervention, with or without indoor air quality (IAQ) feedback, in reducing SHS exposure in homes in Bangladesh.

Methods

We conducted a pragmatic, three-arm, cluster randomised controlled trial in which we randomised mosques and consenting households from their congregations to SFH intervention plus IAQ feedback, SFH intervention only, or usual services. The SFH intervention consisted of health messages delivered within an Islamic discourse by religious leaders at mosques over 12 weeks. IAQ feedback comprised providing households with feedback on their 24-hour IAQ. The primary outcome was the 24-hour mean household fine particulate matter <2.5 microns in diameter (PM_{2.5}) concentration (a marker of SHS) at 12 months post-randomisation. Cost-effectiveness was estimated using incremental cost-effectiveness ratios (ICERs). Trial registration number: ISRCTN49975452.

Findings

45 mosques (1,801 households) were recruited between April and August 2018. At 12 months, the adjusted mean differences in the primary outcome were -1.0µg/m³ (95% confidence interval (CI) -12.8 to 10.9, p=0.88) for SFH intervention plus IAQ feedback versus usual services (primary comparison), 5.0µg/m³ (95% CI -7.9 to 18.0, p=0.45) for SFH intervention only versus usual services, and -6.0µg/m³ (95% CI -18.3 to 6.3, p=0.34) for SFH intervention plus IAQ feedback versus SFH intervention only. The ICER for the primary comparison was US\$653/QALY gained, which was above the US\$427/QALY threshold. The SFH intervention only incurred higher costs, but generated less QALYs compared to usual services.

Interpretation

The SFH intervention, with or without IAQ feedback, was neither effective nor cost-effective in reducing household SHS exposure compared to usual services. These interventions are therefore not recommended for Bangladesh.

Funding

Medical Research Council UK under the Global Alliance for Chronic Diseases research programme: MR/P008941/1

Research in context**Evidence before this study**

We reviewed relevant literature identified from four databases: MEDLINE, Embase, CINAHL, and the Cochrane Controlled Register of Trials. We looked for randomised controlled trials of community-based interventions to reduce second-hand smoke (SHS) exposure conducted in low- and middle-income countries (LMICs). The studies had to have reported biochemically verified SHS exposure (e.g., through measuring air quality, air nicotine concentrations, or cotinine concentration in the blood, urine, saliva or hair) as an outcome. We searched for English language publications from the inception of the databases till July 2017 and updated the searches in December 2018. We found six studies that met our eligibility criteria. All six studies evaluated counselling or educational interventions targeted at reducing SHS exposure, particularly among children. Three studies evaluated interventions delivered to children within schools. Of these, two conducted in China found that the interventions were effective in reducing mean urine cotinine concentration, and the remaining study conducted in Bangladesh by our team did not find any statistically significant difference between groups on saliva cotinine concentrations. A study conducted in China where the intervention was delivered in people's homes found that the intervention was effective on reducing mean urine cotinine concentration among children. Another study conducted in Iran identified participants from health centres: it demonstrated that the community-based intervention was effective in reducing mean urine cotinine concentrations among children. The sixth study was conducted in Armenia and did not find any benefits from the intervention on hair cotinine concentrations. Our review concluded that, whilst there is some evidence of the effectiveness of community-based interventions in reducing SHS exposure, this evidence is very limited. Moreover, the potential of religion in promoting behaviours that are protective from SHS exposure had yet to be explored. We did not identify any studies evaluating the costs or cost-effectiveness of community-based interventions to reduce SHS exposure.

Added value of this study

Our trial is the first to investigate the effectiveness and cost-effectiveness of community-based interventions, delivered within a faith-based discourse by imams and other religious leaders in mosques with or without an individual-level indoor air quality feedback intervention, for reducing SHS exposure within the home. Our interventions were neither effective nor cost-effective when compared to usual services. However, we demonstrated that it is feasible to conduct large studies of such interventions within faith-based settings in low-income country contexts.

Implications of all the available evidence

Current evidence on the effectiveness and cost-effectiveness of community-based interventions to reduce SHS exposure in LMICs is limited and the findings are mixed. Unless future studies provide strong evidence demonstrating their effectiveness and cost-effectiveness, such interventions cannot be recommended for scale up within LMIC settings. There is a need for more studies exploring interventions that have shown promise in high-income countries such as those that combine smoke-free home interventions with smoking cessation advice and support for smokers within the home.

BACKGROUND

Every year approximately 1·2 million people die worldwide from exposure to second-hand smoke (SHS) from tobacco.¹ 47% and 28% of these deaths occur in women and children respectively, mostly in low- and middle-income countries (LMICs).² About 11 million disability-adjusted life years are lost due to SHS exposure worldwide every year, with children carrying approximately 61% of the burden of disease attributable to SHS.² In Bangladesh, 39% (40·8million) of adults and 31% of students aged 12 to 16 years (classes 7-9) are exposed to SHS in their homes.^{3,4} A survey of 12 schools in Dhaka, Bangladesh found that 95% of 9-11 year olds had saliva cotinine levels consistent with recent SHS exposure.⁵ The mean cotinine value of those living with a smoker was approximately double that of those not living with a smoker.⁵ Thus, homes remain a key source of SHS exposure in children in Bangladesh.

In Bangladesh, 88% of the total population are Muslims.⁶ Religion has an influence on both health-risk behaviour and health.⁷⁻⁹ It is a far-reaching conveyor of social norms, potentially through clear and direct precepts regarding the pursuit of a healthy life, or religious tenets that have an indirect effect on health.⁷ Religion, including the Islamic faith, can have a prohibitive influence against tobacco use and promote quitting among smokers.^{10,11} Reinforcing health messages in interventions using Islamic scripture to change smoking behaviours has been reported as acceptable.¹² Islamic faith-based teachings and teachers thus have a potential role in tobacco control, including in reducing SHS exposure in the home; but evidence on the effectiveness of faith-based interventions on changing smoking behaviours is lacking.¹³ Indoor air quality (IAQ) feedback based on markers of SHS such as the concentration of airborne particulate matter less than 2·5 microns (PM_{2·5}) in diameter can potentially motivate households to make their homes smoke-free.¹⁴ However, using IAQ feedback this way is under-researched, particularly in LMICs.

For the Muslim Communities Learning About Second-hand Smoke (MCLASS) II trial, we built our interventions on theoretical work on the role of faith-based interventions to reduce smoking and the potential motivational effects of IAQ feedback.^{10,13,14} We also built our methods on a pilot trial conducted in England that found that a Smoke-Free Homes (SFH) intervention was acceptable to Muslim communities and feasible to deliver through mosques.¹⁵ We designed a community-based SFH intervention in which religious leaders (i.e., imams and khatibs) encouraged their mosque congregations to change their smoking behaviours. In this paper, we report results from the evaluation of the effectiveness and cost-effectiveness of the community-based SFH intervention, with or without IAQ feedback, in reducing exposure to SHS in the home, the frequency and severity of respiratory symptoms, health service use and in improving quality of life.

METHODS

Study design and participants

We conducted a pragmatic, three-arm, open label, cluster randomised controlled trial (RCT) and economic evaluation in which mosques in Mirpur, Dhaka, Bangladesh were randomised and households from their catchment communities enrolled. The mosques were situated in residential areas of Dhaka, hosted regular communal prayers (including Friday Jumu'ah prayers) and had a non-smoking religious leader (imam/khatib). They were affiliated with the Islamic Foundation under the Ministry of Religious Affairs, Bangladesh. A household (i.e.,

a single housing unit shared by one or more people) was eligible if it had at least one resident attending one of the participating mosques, at least one adult resident who smoked cigarettes and/or other forms of smoked tobacco (e.g., bidi, waterpipe) regularly (at least 25 days/month), and at least one non-smoking resident of any age. Households were excluded if they were planning to move home in the next 12 months, or used coal or biomass fuel for domestic cooking or heating. A resident was defined as an adult or child who had been staying in the home for the preceding three months and planned to stay for at least one more year.

Written informed consent was obtained from imams/khatibs for their and their mosques' participation, household heads for recruiting their households, adults in respective households for data collection and if they are parents/guardians then also for collecting data on their children. Ethics approval was obtained from the Bangladesh Medical Research Council's National Research Ethics Committee (Ref: BMBC/NREC/2016–2019/358) and the University of York's Health Sciences Research Governance Committee.

Randomisation and masking

Once recruitment and baseline data collection were completed within a mosque, it was ready for central randomisation 1:1:1 to: SFH intervention plus IAQ feedback, SFH intervention only, or usual services. Minimisation, via MinimPY (<https://sourceforge.net/projects/minimpy/>), was used to balance on the average estimated size of the Friday Jumu'ah prayer congregation (≤ 1500 / >1500) and geographical location (wards within Mirpur). Randomisation was performed by a statistician at the University of York who was not involved in recruiting mosques or households. Mosques were input into MinimPY in a random order unknown to anyone but the statistician. Thus, even if the minimisation factors for the mosques were known, the allocations could not be predicted in advance and allocation concealment was assured. Blinding of participants or imams/khatibs was not possible. Outcome data collection and statistical analyses were also not conducted blind to allocation.

Procedures

The SFH intervention consisted of health messages relating to smoking and SHS exposure, each supported by at least one Qur'an verse (ayah), or an Islamic faith-based decree. The messages were developed through iterative workshops involving Islamic scholars, public health professionals and behavioural scientists. The messages were delivered by imams/khatibs to those attending Friday Jumu'ah prayer in mosques over 12 weeks (one message per week- see examples in Supplementary Table S1). The full intervention manual is available at <https://www.york.ac.uk/healthsciences/research/public-health/projects/mclass11/#tab-3>. The messages addressed key determinants of current smoking behaviours including: lack of knowledge on, and attitudes towards, smoking and SHS exposure by providing information on health consequences of smoking and SHS exposure including addressing misconceptions; and perceptions about social norms by providing general information on others' approval. The messages also targeted prompting intentions, goal setting (both for behaviour, e.g., quit attempt, and the desired outcome of SFH), self-efficacy, commitment, action planning, coping planning, and sources of social support. The intervention logic model is provided as Supplementary Figure S1. Imams/khatibs in mosques randomised to deliver the SFH intervention received an

intervention booklet-based half-day training on the intervention and its delivery. They also received copies of the SFH intervention booklet to distribute to their congregation members after Friday Jumu'ah prayers or in study circles. Intervention delivery started immediately after training and continued for 12 weeks.

IAQ feedback comprised providing households with personalised information on the PM_{2.5} concentration measured within their home at baseline, in the form of a two-page bespoke leaflet, aimed at motivating changes in smoking behaviour in households. The leaflet is available at <https://www.york.ac.uk/healthsciences/research/public-health/projects/mclass11/#tab-3>. PM_{2.5} concentration was measured in homes using the Dylos DC 1700 (Dylos, California, USA), an optical particle counter validated for use in domestic settings.¹⁶ Feedback included a comparison of the household's 24-hour mean PM_{2.5} concentration to the World Health Organization (WHO) guidance limit of 25µg/m³,¹⁷ the total time the IAQ was above this guidance limit, and the maximum concentration measured. Feedback also included pictorial information about the household PM_{2.5} concentration (with classifications: hazardous if >150µg/m³, unhealthy if 36-150µg/m³, moderate if 12-35µg/m³, and good if <12µg/m³), information about adverse effects of SHS exposure, recommendations to reduce SHS exposure in the home, and a target that was achievable by implementing SFH rules within the home. The leaflet was designed in consultation with lay community members. Trial field investigators delivered and discussed the IAQ feedback with household members in person in approximately 10 minutes. All followed-up households in the three groups received feedback on their 12-month IAQ measurements after the final follow-up.

No intervention was offered to households in mosques randomised to receive usual services; however, following trial completion, mosques in the usual service group were offered the SFH toolkit free of charge.

Field investigators recruited mosques by providing their leaders with trial information and screening the mosques for eligibility. They also approached household heads living in the catchment area and attending prayers at any of the participating mosques, either at the mosque or through a home visit, and provided them with study information; those interested were screened for eligibility.

Data were collected at baseline, and three, six and 12 months post-randomisation (Supplementary Table S2) using paper-based questionnaires administered by 16 field investigators after receiving three-days training on trial procedures. The data was entered into a password-protected database on a secure web application, Research Electronic Data Capture (REDCap) (<https://projectredcap.org/software/>). More details on study design, participant and study procedures are provided in our published protocol.¹⁸

Outcomes

The primary outcome was 24-hour mean household PM_{2.5} concentration at 12 months post-randomisation. Household-level secondary outcomes were: 24-hour mean PM_{2.5} concentration at three months; and smoking restrictions at home assessed through a questionnaire directed at adults in the households. Participant-level secondary outcomes assessed at each follow-up were: frequency and severity of respiratory symptoms assessed

using Part 1 (eight questions) of the validated St. George's Respiratory Questionnaire (SGRQ)¹⁹ for participants 11 years and over, and the severity scale developed and validated by Chauhan et al²⁰ for participants younger than 11; health-related quality of life (HRQoL) assessed using the EQ-5D-5L²¹ for adults (≥ 18 years), EQ-5D-Y²² for adolescents (11-17 years inclusive) and PedsQoL²³ for children younger than 11 years. A questionnaire previously used in a pilot trial in England,¹⁵ and adapted to the Bangladesh context was used for healthcare resource utilisation.

Statistical analysis

We planned to recruit 45 mosques and 40 households per mosque ($n=1,800$), and to follow-up 30 households per mosque at three months ($n=1,350$) prioritising those with an average baseline $PM_{2.5} \geq 35\mu g/m^3$. Assuming an intra-cluster correlation coefficient (ICC) of 0.02 and 20% attrition at 12 months, this would provide 90% power to detect an effect size of 0.3 of a standard deviation (SD; equivalent to a difference of 13.5, from $76\mu g/m^3$ to $62.5\mu g/m^3$, assuming a SD of 45), for each pairwise comparison, using a two-sided alpha of 0.05.

Analyses followed a pre-specified analysis plan, approved by the Trial Steering Committee prior to the completion of 12-month data collection. No post-hoc analyses were conducted. Analyses were conducted using intention-to-treat and two-sided statistical tests at the 5% significance level in Stata v15. Baseline and outcome data were summarised by group.

The primary analysis compared household 24-hour mean $PM_{2.5}$ measurement between the groups using a covariance pattern, mixed-effect linear regression model incorporating the two post-randomisation time points (three and 12 months). The model included baseline $PM_{2.5}$ value (household-level), geographical area and size of Friday Jumu'ah prayer congregation in its continuous form (mosque-level), and time point, randomised group, and a time by group interaction as fixed effects. Household and mosque were specified as random effects. An unstructured covariance pattern for the correlation of observations within households over time was specified, based on minimising the Akaike information criterion. Visual inspection of model assumptions demonstrated substantial deviations (Supplementary material S1: Primary analysis model assumptions; and Supplementary Figures S2 and S3). Log-transformation of the outcome data improved model fit (Supplementary Figures S4 and S5) and was explored in sensitivity analyses. The pairwise mean difference, 95% confidence interval (CI) and p-value at three and 12 months was extracted from the model.

The primary comparison was between SFH intervention plus IAQ feedback and usual services at 12 months. All other comparisons served as secondary investigations. To account for non-compliance with randomised group, a complier average causal effect (CACE) analysis²⁴ for the primary outcome was conducted. A two-stage, least squares instrumental variable (IV) approach was used, with randomised group as the IV. Two analyses compared the 12-month outcome for each intervention with usual services. Within the SFH intervention only group, compliance was defined at the household-level as the lead adult reporting that they or another member of their household had received the SFH intervention from any mosque at any time point. Within the SFH intervention plus IAQ feedback group, compliance additionally included self-reported receipt of IAQ feedback by the three-month follow-up.

Calibration of the Dylos machines prior to the 12-month follow-up indicated that they were consistently underestimating PM_{2.5} concentrations, relative to a 'gold standard', factory calibrated device, due to degradation of the laser particulate counter caused by heavy use at the baseline and three-month assessments. This underestimation was corrected for in the primary analysis; details of sensitivity analyses assessing the impact of this correction are provided in Supplementary material S2: Sensitivity analyses.

A subgroup analysis considered whether the benefits of the interventions were greater among households with a baseline PM_{2.5} value $\geq 35\mu\text{g}/\text{m}^3$ than those with $< 35\mu\text{g}/\text{m}^3$, by including an interaction between dichotomised baseline PM_{2.5} value and group in the primary analysis.

Participant-level respiratory symptom scores were analysed in an analogous way to the primary outcome. Participant, household, and mosque were nested random effects. Analyses were performed separately for the SGRQ Symptoms component score for adults, for children aged 11-17 years, and for the total symptoms severity scale for children aged less than 11 years. Since both these instruments measure the same construct, respiratory symptoms, an additional analysis that included all participants was conducted using standardised scores. Model assumptions were assessed as for the primary analysis; no major deviations were observed so data transformation was unnecessary.

Cost-effectiveness analysis

We conducted a within-trial cost-effectiveness analysis comparing the SFH intervention with and without IAQ feedback to usual services. The analysis used a healthcare sector and intervention provider perspective to include healthcare resource use and intervention delivery costs. No discounting was applied as the follow-up period was 12 months.

All costs were calculated using a bottom-up approach. Costs for training and delivering the interventions (teaching materials, support, etc.) were estimated based on the cost incurred alongside the trial, while information on health care resources use (number of inpatient stays, outpatient visits, etc.) was collected from participants. The unit costs of home visits by doctors or nurses were obtained from the Bangladesh Bureau of Statistics,²⁵ those of inpatient stay and outpatient visit were derived from WHO's Bangladesh-specific unit costs,²⁶ and those of emergency department visits were extracted from the Bangladesh essential health service package (Table 1).²⁷ All cost results were expressed in 2018/19 US dollars (US\$) using the 2018 World Development Indicators exchange rates.²⁸

Due to the absence of unified and established tariffs from Bangladesh for the three instruments used to measure HRQoL (i.e. PedsQL, EQ-5D-Y and EQ-5D-5L), relevant UK value sets were used and mapped to the corresponding EQ-5D-3L values.^{23,29} This allowed us to obtain unified utility estimates across individuals. A sensitivity analysis using Thailand value sets³⁰ for EQ-5D-Y and EQ-5D-5L were conducted to test the robustness of the results. Quality-adjusted life-years (QALYs) of individuals were calculated using the area under the curve method over the trial period.³¹

Cost-effectiveness was evaluated using the pairwise incremental cost-effectiveness ratios (ICERs) method³¹ at a household level and assessed based on the Bangladesh willingness-to-pay (WTP) threshold: US\$30 to US\$427 per QALY gained.³² Seemingly unrelated regression was used to account for potential correlations between costs and QALYs, and adjust for prognostic baseline covariates.³³ Uncertainty was estimated using the non-parametric bootstrapping technique with 5,000 replications that were presented on a cost-effectiveness plane.³¹

Role of funding source

This trial was funded by the Medical Research Council UK under the Global Alliance for Chronic Diseases (GACD) research programme [MR/P008941/1]. The funder had no role in the design of the trial and collection, analysis, interpretation of data or in writing the manuscript.

RESULTS

Trial population

116 mosques were assessed for eligibility and 45 were recruited (Figure 1). Reasons for exclusion were: being less than half a kilometre from another mosque (n=54); imams/khatibs not providing consent to participate (n=7), mosque catchment area having entry restrictions (n=7), small catchment area (n=2), and not performing Friday Jumu'ah prayers (n=1). Between 11th April and 2nd August 2018, 4,430 households were screened for eligibility; 1,801 (40.7%) were eligible and recruited. Reasons for ineligibility are in Supplementary Table S3. Every mosque recruited 40 households except one (allocated to usual services), which recruited 41. 16 mosques were randomised to the SFH intervention plus IAQ feedback, 14 to SFH intervention only and 15 to usual services (Supplementary Table S4).

Of the 712 households with average baseline PM_{2.5} ≥35µg/m³, 614 (86.2%) were followed-up at three months; 98 had either moved away from the study area or did not wish to continue the study. To achieve the target of 1,350 households followed-up at three months, we randomly selected another 736 households with average baseline PM_{2.5} <35µg/m³ to follow-up as per our trial protocol. Household baseline data as randomised and as followed up are summarised in Table 2, and Supplementary Tables S5-S12. The 1,350 households were contacted for follow-up again at 12 months and 1,314 completed it (2.7% attrition rate; usual services 2.0%, SFH intervention only 3.6%, SFH intervention plus IAQ feedback 2.5%).

Primary outcome

At 12 months, the average 24-hour PM_{2.5} measurement was 65.2µg/m³ (SD 44.7), 68.9µg/m³ (SD 49.5) and 65.8µg/m³ (SD 39.6) for usual services, SFH intervention only and SFH intervention plus IAQ feedback, respectively (Table 3). No evidence of a difference was observed at 12 months for any pairwise comparison, including when the outcome data were log-transformed (Table 4; Supplementary Figure S6). The adjusted mean differences were: SFH intervention plus IAQ feedback vs. usual services (primary comparison) -1.0µg/m³ (95% CI -12.8 to 10.9, p=0.88); SFH intervention only vs. usual services 5.0µg/m³ (95% CI -7.9 to 18.0, p=0.45); and SFH intervention plus IAQ feedback vs. SFH intervention only -6.0µg/m³ (95% CI -18.3 to 6.3, p=0.34). The log-transformed sensitivity analysis indicated that the average PM_{2.5} concentrations in the SFH intervention plus IAQ feedback households were

expected to be 1.02 times larger (95% CI 0.86 to 1.21, $p=0.79$) than the usual services group. The mosque-level ICC was estimated at 0.08 (95% CI 0.05 to 0.14).

Secondary outcomes

There was no evidence of a difference in the average 24-hour $PM_{2.5}$ concentration at three months for any pairwise comparison, including when the outcome data were log-transformed (Table 4; Supplementary Figure S6). There was evidence of small differences in some secondary comparisons between SFH intervention only and SFH intervention plus IAQ feedback favouring SFH intervention plus IAQ feedback (SGRQ adults at six months: 2.4, 95% CI 0.3 to 4.6, $p=0.03$; respiratory symptoms <11 years at three months: 2.0, 95% CI 0.5 to 3.4, $p=0.01$; standardised respiratory scores all participants at six months: 0.14, 95% CI 0.00 to 0.29, $p=0.04$). No other differences were observed at three, six or 12 months (Supplementary Tables S13-S18).

Overall, 22.9% of households (usual services $n=110$, 24.9%; SFH intervention only $n=76$, 18.8%; SFH intervention plus IAQ feedback $n=115$, 26.6%) reported at 12 months that residents were permitted to smoke anywhere inside the home (Supplementary Table 18).

Sensitivity analyses

Within the SFH intervention only group, 331 (78.8%) households received the intervention; in addition, 91 (20.2%) of households in the usual services group reported receiving some element of the intervention. The CACE estimate of receipt of the SFH intervention was an increase in $PM_{2.5}$ concentration of $11.3\mu g/m^3$ (95% CI -6.7 to 29.2). In the SFH intervention plus IAQ feedback group, 351 (73.1%) households received the SFH intervention and IAQ feedback by the three-month follow-up: the CACE estimate of receipt of the SFH intervention and an IAQ report was a decrease in $PM_{2.5}$ concentration ($-3.0\mu g/m^3$, 95% CI -17.4 to 11.4).

No differences were observed at three or 12 months in the other sensitivity analyses (Supplementary Tables S19 and S20; Supplementary Figures S7 and S8).

Subgroup analysis

There was no evidence of an interaction with baseline $PM_{2.5}$ value ($<35/\geq 35\mu g/m^3$) in the subgroup analysis (Supplementary Table 21; Supplementary Figures S9 and S10).

Cost-effectiveness results

Based on the 1,350 households followed up at three months, 4,893 out of 5,143 participants (95.1%) had complete cost and QALY data at all follow-ups. After removing households with members showing incomplete data, a total of 1,237/1,350 (91.7%) households were included in the cost-effectiveness analysis. The SFH intervention plus IAQ feedback group incurred the highest mean total cost (\$32.8, SD 22.0) and generated the highest mean QALYs (3.31, SD 1.20) (Table 5). The SFH intervention only group incurred higher costs, but generated less QALYs, compared to the usual services group, and was therefore dominated. Due to high delivery cost of IAQ (\$16.1), intervention cost was the key cost driver for the SFH intervention plus IAQ feedback group but not for the SFH intervention only group (Supplementary Table S22). The SFH intervention plus IAQ feedback group was found not to be cost-effective as the ICER of US\$653/QALY against the usual services group was above

the threshold of US\$30 to 427/QALY gained. The results of the 5,000 bootstrapped SUR models are shown in Table 5 and Supplementary Figure S11. The bootstrapped ICERs fell within the top left quadrant of the cost-effectiveness plane and above the WTP threshold lines, indicating that both the SFH intervention only, and the combination of the SFH intervention plus IAQ feedback, were not cost-effective even when taking uncertainty into consideration. The results of sensitivity analysis were similar to the main findings (Supplementary Table S23 and Supplementary Figure S12).

DISCUSSION

Compared to usual services, the SFH intervention alone or in combination with IAQ feedback did not reduce exposure to SHS measured as the mean 24-hour $PM_{2.5}$ concentration within households. The economic evaluation suggests that both the SFH intervention only and the combination of the SFH intervention and IAQ feedback were not cost-effective due to high intervention costs and minimal QALY improvements.

To the best of our knowledge, our trial is the first to investigate the effectiveness and cost-effectiveness of community-based interventions, delivered within an Islamic discourse by imams/khatibs in mosques, for reducing SHS exposure within the home. We demonstrated that it is feasible and acceptable to conduct large studies of such interventions within mosques. The trial is also the largest of its kind to provide 24-hour household-level $PM_{2.5}$ concentration data, and explore the usefulness of using IAQ feedback as a motivational tool for reducing SHS exposure in the home, in a LMIC setting. Other strengths were the rigour and quality with which the trial was conducted, including a cluster RCT design, achieving the required sample size, high follow-up rates, high levels of data completeness, and the assessments of 12-month outcomes.

There are a number of potential explanations for the lack of effectiveness of our interventions. Sermons where the intervention messages were delivered are not mandatory and therefore some people may have joined the prayers, but not attended or paid the desired level of attention to the sermons. Although intervention compliance as defined in our trial was high, it is likely that individuals might have received some, but not all, messages. Because of the nature of the intervention, it was not possible to calculate the 'dose' of the intervention received by household members. The interventions targeted reducing SHS exposure in the home directly and did not offer smoking cessation support to smokers within the home. Aspirations to make SFH might have been constrained by limited social and environmental opportunities to change behaviour.^{34,35} Thus, a standalone community-based intervention delivered over a limited period might have been insufficient to change smoking behaviours in Bangladesh where regulatory and fiscal measures for tobacco control are weak, cigarettes and bidis are cheap, and smoking cessation services are scarce. In addition, the personalised feedback was delayed due to the need to take the IAQ machine back to the office to download the data and generate the graphical and numerical feedback, and was not targeted specifically at the smokers.

The intervention effects on $PM_{2.5}$ concentration in the home could have been diluted due to a Hawthorne effect across all trial groups during the baseline 24-hour measurement period when the Dylos devices were present in the home.³⁶ Measuring $PM_{2.5}$ concentrations over a longer period of time could have reduced this potential bias by making it more difficult to

sustain behaviour change over the whole measurement period.³⁶ However, this would have been more costly due to the need for more devices.

PM_{2.5} is not specific to tobacco smoke – it can also be generated by non-tobacco sources such as using solid fuels and vehicle and industrial emissions. Our trial addressed other PM_{2.5} influences by excluding households that used coal or biomass fuel for domestic use and restricting measurements to the period April-October when outdoor air pollution levels are lowest in Dhaka.³⁷ We also used a cluster RCT design in order to balance such confounders across the two arms. Therefore, any change observed in the primary outcome between the two arms would have been most likely due to change in smoking behaviour. Confidence in our findings is also enhanced by the fact that baseline PM_{2.5} concentrations were significantly lower in smoke-free homes when compared to smoking permitted homes despite high ambient air pollution.³⁸

Participants' HRQoL was measured using three different instruments (i.e. the EQ-5D-5L for adults,²¹ EQ-5D-Y for adolescents²² and PedsQoL for children²³) due to the absence of a universal instrument that could measure the HRQoL across all age groups. As HRQoL may differ depending on which instruments are being used, this approach can result in household QALY estimations being sensitive to the number of people and the age composition in each household. However, this is unlikely to have affected our conclusion as the household composition was controlled for in the analysis. With no established Bangladesh population tariffs, the UK population tariffs were used for QALY calculations as they are the only tariffs that can convert all three instrument measurements into consistent EQ-5D-3L values. Future studies on Bangladesh tariffs, and for other LMICs, across all age groups are required in order to obtain more precise estimates.

In relation to faith-based behaviour change intervention, our findings can be generalised to other community-based interventions delivered primarily through mosques. When considering IAQ feedback, our study findings can be generalized to other urban centres similar to Dhaka with high population density, high levels of ambient air pollution and limited opportunities to smoke outside.

Contrary to our findings, studies in other areas such as cardiovascular diseases, obesity, and breast cancer screening have suggested that health programmes in faith-based organisations can improve outcomes.^{8,9} Islamic faith-based smoking cessation interventions have also been found to be effective in encouraging Muslim smokers to stop smoking during Ramadan, although the sustainability of the behaviour change is unclear.^{12,39} Nevertheless, our findings are consistent with those from other studies targeting reduction of SHS exposure within the home using behavioural interventions and IAQ feedback. A recent review found that the effectiveness of several counselling and educational interventions that have been used to try to reduce SHS exposure has not been clearly demonstrated.⁴⁰ More successful interventions seem to be those that combine SFH interventions with smoking cessation advice and support for smokers within the home, or those that target smoking cessation as a pathway to reducing SHS exposure.⁴¹ In addition, a recent study showed that real-time particle feedback and coaching contingencies reduced indoor air pollution from behaviours such as smoking cigarettes or burning candles.⁴² Hence, future research in LMICs should investigate the effectiveness of interventions that include offering

smoking cessation to smokers within the household, and measures that offer real-time or immediate, rather than delayed, feedback on IAQ. Nevertheless, these technologies need to be low cost if they are to be cost-effective and scalable in LMICs.

In conclusion, the SFH intervention, alone or in combination with IAQ feedback, was not effective or cost-effective in reducing exposure to SHS in the home when compared to usual services and should therefore not be recommended in Bangladesh.

DATA SHARING STATEMENT

De-identified individual participant data on which summary statistics and tables are based will be made available from the point of, and up to five years after the, acceptance for publication of the main findings from the final dataset. These data can be requested from the Principal Investigator (Professor Kamran Siddiqi, kamran.siddiqi@york.ac.uk) and will be shared after the provision of a methodologically sound proposal, and only under a data-sharing agreement that provides for: (1) a commitment to using the data only for research purposes and not to identify any individual participant; (2) a commitment to securing the data using appropriate computer technology; and (3) a commitment to destroying or returning the data after analyses are completed. The proposals will be assessed and approved by members of the Programme Management Group.

The intervention manual and indoor air quality feedback leaflet are available on the study webpage: <https://www.york.ac.uk/healthsciences/research/public-health/projects/mclass11/#tab-3>

Other material such as participant information sheets, informed consent forms and questionnaires will be made widely and freely available to anyone who wishes to access them from the point of, and up to five years after the, acceptance for publication of the main findings from the trial. They can be requested from the Principal Investigator.

The study protocol is available in the public domain for free: Mdege N, Fairhurst C, Ferdous T, et al. Muslim Communities Learning About Second-hand Smoke in Bangladesh (MCLASS II): study protocol for a cluster randomised controlled trial of a community-based smoke-free homes intervention, with or without Indoor Air Quality feedback. *Trials* 2019; 20: 11. doi: 10.1186/s13063-018-3100-y.

CONTRIBUTORS

NDM drafted the manuscript, contributed to study design, conduct and interpretation of findings. CF contributed to the writing of the methods and results sections of the manuscript, and designed and conducted the statistical analysis. H-IW and QW conducted the health economic analysis contributed to the writing of the methods and results sections of the manuscript. TF coordinated the implementation of the study in Bangladesh including data collection and management. AMM coordinated the study. CH participated in study design and oversaw the statistical analysis. RH contributed to study design, oversaw the implementation and conduct of the study in Bangladesh, and provided critical inputs to interpretation of results. CJ contributed to study design, and particularly led on process evaluation. IK contributed to the design of the interventions including the intervention logic model. ZAA contributed to intervention design and process evaluation. SP contributed to

study design and specifically led the design of the health economic evaluation. SS contributed to study design and led the design of IAQ measurement and IAQ feedback. AS contributed to study design, interpretation of results and manuscript writing. KS conceived the study idea and contributed to the study design, conduct, interpretation of results and manuscript writing. All authors participated in manuscript revisions, and read and approved the final manuscript.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

ACKNOWLEDGEMENTS

This trial was funded by the Medical Research Council UK under the Global Alliance for Chronic Diseases (GACD) research programme [MR/P008941/1]. We would like to thank members of the independent Trial Steering Committee: Dr Andrew Fogarty (Chair), University of Nottingham; Professor Jo Leonardi-Bee, University of Nottingham; and Dr Saidur Rahman Mashreky, Bangladesh University of Health Sciences. Our gratitude also goes to members of our national-level reference group: the Joint Secretary of the Ministry of Religious Affairs, Director of Research of the Islamic Foundation, Head of the Imam Training Academy, and a representative from the ARK Foundation. We are also grateful to the project working group comprising an Islamic Foundation representative, an imam, a research fellow, a public health expert and a member of the public. We are grateful to the trial field investigators; the Islamic Foundation, Bangladesh; participating mosques, their mosque committee leaders, imams, and khatibs; all participants; professionals and other researchers who have contributed to the trial.

REFERENCES

1. World Health Organisation. Health topic: Tobacco. https://www.who.int/health-topics/tobacco#tab=tab_1 (Accessed 15 March 2020).
2. Öberg M, Jaakkola MS, Woodward A, Peruga A, Prüss-Ustün A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* 2011; **377**: 139-46.
3. Ministry of Health and Family Welfare. Global Adult Tobacco Survey: Bangladesh Factsheet 2017. Dhaka: World Health Organization, Country Office for Bangladesh, 2017.
4. WHO/SEARO/Country Office for Bangladesh, Ministry of Health and Family Welfare. Global Youth Tobacco Survey (GYTS) Bangladesh Report, 2013. Dhaka: World Health Organization, Country Office for Bangladesh, 2013.
5. Shah S, Kanaan M, Huque R, et al. Secondhand smoke exposure in primary school children: a survey in Dhaka, Bangladesh. *Nicotine Tob Res* 2017; **21**: 416-23.
6. Bangladesh Bureau of Statistics. Report on Bangladesh sample vital statistics 2018. Dhaka: Bangladesh Bureau of Statistics, 2019.
7. Nunziata L, Toffolutti V. "Thou Shalt not Smoke": Religion and smoking in a natural experiment of history. *SSM Popul Health* 2019; **8**: 100412. doi: 10.1016/j.ssmph.2019.100412
8. DeHaven MJ, Hunter IB, Wilder L, Walton JW, Berry J. Health programs in faith-based organizations: are they effective? *Am J Public Health* 2004; **94**: 1030-6.
9. Lancaster KJ, Carter-Edwards L, Grilo S, Shen C, Schoenthaler AM. Obesity interventions in African American faith-based organizations: a systematic review. *Obes Rev* 2014; **15**: 159-76.
10. Weaver AJ, Flannelly KJ, Strock AL. A review of research on the effects of religion on adolescent tobacco use published between 1990 and 2003. *Adolescence* 2005; **40**: 761-76.
11. Widyaningrum N, Yu J. Tobacco use among the adult Muslim population in Indonesia: a preliminary study on religion, cultural, and socioeconomic factors. *J Drug Issues* 2018; **48**: 676-88.
12. Pratt R, Ojo-Fati O, Adam A, et al. Text message support for smoking cessation during Ramadan: a focus group study with Somali immigrant Muslim men. *Nicotine Tob Res* 2020; **22**: 1636-9.
13. Jabbour S, Fouad FM. Religion-based tobacco control interventions: how should WHO proceed? *Bull World Health Organ* 2004; **82**: 923-7.
14. Marsh J, McNeill A, Lewis S, et al. Protecting children from secondhand smoke: a mixed-methods feasibility study of a novel smoke-free home intervention. *Pilot Feasibility Stud* 2016; **2**: 53. doi: 10.1186/s40814-016-0094-7.
15. Shah S, Ainsworth H, Fairhurst C, et al. Muslim communities learning about second-hand smoke: a pilot cluster randomised controlled trial and cost-effectiveness analysis. *NPJ Prim Care Respir Med* 2015; **25**: 15052. doi: 10.1038/npjpcrm.2015.52.
16. Semple S, Ibrahim AE, Apsley A, Steiner M, Turner S. Using a new, low-cost air quality sensor to quantify second-hand smoke (SHS) levels in homes. *Tob Control* 2015; **24**: 153-158.
17. World Health Organization. WHO guidelines for indoor air quality: selected pollutants. Copenhagen: World Health Organization, 2010.

18. Mdege N, Fairhurst C, Ferdous T, et al. Muslim Communities Learning About Second-hand Smoke in Bangladesh (MCLASS II): study protocol for a cluster randomised controlled trial of a community-based smoke-free homes intervention, with or without Indoor Air Quality feedback. *Trials* 2019; **20**: 11. doi: 10.1186/s13063-018-3100-y.
19. Jones PW, Quirk FH, Baveystock CM. The St George's Respiratory Questionnaire. *Respir Med* 1991; **85 Suppl B**: 25-31; Discussion 3-7.
20. Chauhan AJ, Inskip HM, Linaker CH, et al. Personal exposure to nitrogen dioxide (NO₂) and the severity of virus-induced asthma in children. *Lancet* 2003; **361**: 1939-44.
21. EuroQol Group. EQ-5D-5L User guide: basic information on how to use the EQ-5D-5L instrument (version 2.1). Rotterdam: EuroQol Research Foundation, 2015.
22. Ravens-Sieberer U, Wille N, Badia X, et al. Feasibility, reliability, and validity of the EQ-5D-Y: results from a multinational study. *Qual Life Res* 2010; **19**: 887-97.
23. Khan KA, Petrou S, Rivero-Arias O, Walters SJ, Boyle SE. Mapping EQ-5D utility scores from the PedsQL™ generic core scales. *Pharmacoeconomics* 2014; **32**: 693-706.
24. Dunn G, Maracy M, Dowrick C, Ayuso-Mateos JL, Dalgard OS, Page H, et al. Estimating psychological treatment effects from a randomised controlled trial with both non-compliance and loss to follow-up. *Br J Psychiatry* 2003; **183**: 323-31.
25. Bangladesh Bureau of Statistics. Bangladesh - Labour Force Survey 2016/17. Dhaka: Bangladesh Bureau of Statistics, 2017.
26. World Health Organization. Country-specific unit costs – Bangladesh 2008. https://www.who.int/choice/country/country_specific/en/ (Accessed 14 April 2020).
27. Islam Z, Ahmed S, Dorin FA, et al. Costs of the Bangladesh essential health service package: 2016- 2022. Dhaka: Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh, 2018.
28. The World Bank Group. World Development Indicators: Exchange rates and prices 2017. <http://wdi.worldbank.org/table/4.16> (Accessed 03 January 2020).
29. van Hout B, Janssen MF, Feng YS, et al. Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health* 2012; **15**: 708-15.
30. Pattanaphesaj J, Thavorncharoensap M, Ramos-Goñi JM, Tongsiri S, Ingsrisawang L, Teerawattananon Y. The EQ-5D-5L Valuation study in Thailand. *Expert Rev Pharmacoecon Outcomes Res* 2018; **18**: 551-8.
31. Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. Methods for the economic evaluation of health care. Fourth ed. Oxford: Oxford University Press; 2015.
32. Woods B, Revill P, Sculpher M, Claxton K. Country-level cost-effectiveness thresholds: initial estimates and the need for further research. *Value Health* 2016; **19**: 929-35.
33. Willan AR, Briggs AH, Hoch JS. Regression methods for covariate adjustment and subgroup analysis for non-censored cost-effectiveness data. *Health Econ* 2004; **13**: 461-75.
34. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011; **6**: 42. doi: 10.1186/1748-5908-6-42.

35. O'Donnell R, Amos A, Turner SW, et al. 'They only smoke in the house when I'm not in': understanding the limited effectiveness of a smoke-free homes intervention. *J Public Health* 2020; fdaa042. <https://doi.org/10.1093/pubmed/fdaa042>.
36. Semple S, Turner S, O'Donnell R, et al. Using air-quality feedback to encourage disadvantaged parents to create a smoke-free home: results from a randomised controlled trial. *Environ Int* 2018; **120**: 104-10.
37. U.S. Embassy in Bangladesh. Air quality data. <https://bd.usembassy.gov/embassy/air-quality-data/> (Accessed 21 October 2017).
38. Ferdous T, Siddiqi K, Semple S, et al. Smoking behaviours and indoor air quality: a comparative analysis of smoking-permitted versus smoke-free homes in Dhaka, Bangladesh. *Tob Control* 2020. doi: 10.1136/tobaccocontrol-2020-055969.
39. Ismail S, Abdul Rahman H, Abidin EZ, et al. The effect of faith-based smoking cessation intervention during Ramadan among Malay smokers. *Qatar Med J* 2016; **2016**. <https://doi.org/10.5339/qmj.2016.16>.
40. Behbod B, Sharma M, Baxi R, Roseby R, Webster P. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. *Cochrane Database Syst Rev* 2018; **1**: CD001746-CD.
41. Ratschen E, Thorley R, Jones L, et al. A randomised controlled trial of a complex intervention to reduce children's exposure to secondhand smoke in the home. *Tob Control* 2018; **27**: 155-62.
42. Hovell MF, Bellettiere J, Liles S, et al. Randomised controlled trial of real-time feedback and brief coaching to reduce indoor smoking. *Tob Control* 2020; **29**: 183-190.