UNIVERSITY of York

This is a repository copy of Application of stranded pelagic sargassum biomass as compost for seedling production in the context of mangrove restoration.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/id/eprint/190939/</u>

Version: Accepted Version

Article:

Trench, Camilo, Thomas, Shanna-Lee, Thorney, Delroy et al. (7 more authors) (2022) Application of stranded pelagic sargassum biomass as compost for seedling production in the context of mangrove restoration. Frontiers in Environmental Science. 932293. ISSN: 2296-665X

https://doi.org/10.3389/fenvs.2022.932293

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.





Application of stranded pelagic sargassum biomass as compost for seedling production in the 1 context of mangrove restoration 2 3 Camilo Trench¹, Shanna-Lee Thomas¹, Delroy Thorney¹, Gina-Marie Maddix², Patrice Francis², Hugh Small³, Carla Botelho 4 Machado⁴, Dale Webber⁵, Thierry Tonon^{4*}, Mona Webber² 5 ¹Discovery Bay Marine Laboratory- UWI, Queens Highway, Discovery Bay, St. Ann, Jamaica. 6 7 ²Centre for Marine Sciences, 1 Anguilla Close, University of the West Indies, Mona, Kingston 7, Jamaica. ³Port Royal Marine Laboratory- UWI, Port Royal, Kingston 1, Jamaica. 8 ⁴Centre for Novel Agricultural Products (CNAP), Department of Biology, University of York, Heslington, York YO10 5DD, United 9 10 Kingdom. ⁵Department of Life Sciences, 2 Anguilla Close, University of the West Indies, Mona, Kingston 7, Jamaica. 11 12 * Correspondence: 13 Thierry Tonon 14 15 thierry.tonon@york.ac.uk 16 Keywords: Sargassum compost, urbanised coastlines, mangrove seedlings, element analysis 17 18 Abstract 19

20 The Sargassum spp. inundations across the Atlantic and Caribbean that began in 2011 have continued unabated and new uses for the biomass are being continuously explored. Mangroves protect shorelines, store carbon, enhance water quality and promote biodiversity. 21 22 Their restoration can be hindered by poor soils associated with urbanized coastlines. *Sargassum* spp. application in the form of mulch, 23 compost and plant tonics has yielded positive results in a range of plants. As part of transforming the inundations to benefit communities, 24 Sargassum spp. compost (SC) was assessed in mangrove seedling production for restoration. Pure SC was mixed with soil/sand medium, as 25 different treatments, for the production of Rhizophora mangle seedlings in 'wet' and 'dry' nurseries. Plants in the 'wet' nursery performed 26 poorly, with 90-100% of plants in 50 and 100% SC, respectively, dying after six weeks. Seedlings in all SC treatments in the 'dry' nursery 27 survived with obvious and statistically significant treatment differences. Height and number of leaves indicated best growth in the 75% SC 28 treatment while the control (0% SC) had poorest growth. Seedling health, greatest in the control, was poorest in 50 and 100% SC. Elemental 29 analysis of SC, seedlings and soil/sand medium indicated that several elements (Na, K, Ca, As, Se) found in high concentrations in the SC, 30 were low in the plants. Overall, low sequestration of elements by mangrove seedlings, and the reported ability of mangrove soils to reduce 31 element mobilization through chelation, indicate potential for use of Sargassum spp. in soil amelioration for mangrove restoration without 32 proportional contamination of the ecosystem. We see potential for the use of nuisance Sargassum spp. blooms to support mangrove 33 restoration, leading to increased benefits to coastal communities who are being affected by the inundations.

34 1 Introduction

35 Large floating mats of pelagic sargassum seaweed, comprised of Sargassum natans I, S. natans VIII and S. fluitans III (Schell et al. 2015), 36 have been inundating shorelines across the Caribbean since 2011 (Wang et al., 2019) and there is evidence that these annual blooms will 37 continue into the foreseeable future; becoming the "new normal" (Wang et al., 2019; Salter et al., 2020 and Machado et al., 2022). Although 38 widely considered a nuisance (van Tussenbroek et al., 2017), many have come to regard the algal biomass as a resource (Milledge and Harvey, 2016; Chávez et al., 2020; Amador-Castro et al., 2021) with a plethora of projects exploring uses, which include soil amelioration 39 40 (Desrochers et al., 2020). The effectiveness of Sargassum spp. in improving plant production (applied as mulch, compost or liquid fertilizer) has been shown for several food crops (CARDI, 2015). However, there is a threat of increasing the salinity of the soil (CARDI, 2015) and 41 although this may be ameliorated by washing the Sargassum spp. before and during composting (Eyras et al., 1998), this measure can be too 42 43 expensive for some communities and also leaches nutrients (CARDI, 2015).

44 While there are examples of the use of *Sargassum* spp. seaweed for restoration of dunes (Desrochers et al., 2020) and to enhance the actual growth of dune plants (Williams and Feagin, 2010; Thompson et al., 2020), there is no previous research on the use of sargassum seaweed in 45 mangrove restoration or rehabilitation. It is hypothesized that mangroves and other coastal plant species growing in poor soil conditions will 46 47 benefit from the soil amelioration properties of Sargassum spp. (improved texture, water holding capacity and increased nutrients), and is 48 expected to also be able to tolerate increased soil salinity. The production of compost from pelagic Sargassum spp. by mixing with other plant material (wood chips and food waste) has been reported (Sembera et al., 2018). However, there is yet no example of composting pure 49 50 Sargassum spp., which would make use of larger quantities of the feedstock, and facilitate an assessment of the effect of the pure compost 51 on coastal plant species like mangroves.

52 Mangrove ecosystem services are well known and include shoreline protection, water quality improvement, biodiversity support through the

- 53 provision of a range of habitats and carbon sequestration. Since 2006, over 200 benefits of mangroves have been documented (Lee et al.,
- 54 2014) and later grouped into regulating, supporting, provisioning and cultural ecosystem services (e.g., Uddin et al., 2013; Webber et al.,
- 55 2016; Kathiresan, 2012; Mitra, 2018). There has also been increasing global recognition of mangrove importance in carbon sequestration
- 56 (Alongi, 2012; Sandeman et al., 2018), with mangroves showing high rates of carbon capture and almost permanent storage in the forest
- 57 sediment as well as living biomass of leaves, stems and roots (Donato et al., 2011; Alongi, 2014).

58 Despite the ecological, economic, cultural and climate regulation value of mangroves, these forests are being destroyed globally at rates of 1

-2% per year (Giri et al., 2008; Hogart, 2015); such that mangrove cover has declined by 35% over the last two decades due to human

60 activities (Krauss et al., 2008). Mangrove forests have been threatened by mariculture (primarily shrimp farming) (Giri et al., 2008),

- 61 excessive timber harvest (Kairo et al., 2001), deforestation for development (Valiela et al., 2001), hydrology changes (Turner and Lewis,
- 62 1997) and rising sea levels which force mangroves to recede to unsuitable land margins (Webber et al., 2016).

63 As is the case with many developing countries, Jamaica has lost almost 20% of mangroves in the last decade, due to coastal development

and land reclamation as well as extreme weather events (Trench, 2021). Mangrove rehabilitation efforts have had increasing success but in
 some cases require severe interventions like, sediment replacement and slope reconstruction to restore hydrology as well as replanting of

- 66 nursery-reared (hardened) seedlings (Nguyen et al., 2016 and Trench, 2021). In some cases the sediments used for mangrove rehabilitation
- 67 sites is riverine sand, which does not provide high levels of organic matter as is naturally found in mangrove forests (Reef et al., 2010).

68 Urbanised coastlines (like the shores of Kingston Harbour, Jamaica) provide a good example of mangrove areas that have experienced

69 severe habitat modification and have little chance of restoration being facilitated by natural recruitment of seedlings from adjacent forests.

Such shorelines are said to be "propagule limited" (Lewis, 2009) and often require extensive replanting using nursery reared seedlings for

- restoration efforts to succeed. Furthermore, soil conditions in such areas are often so altered that there is need for sediment replacement or substantial amelioration for the introduced plants to survive. It was hypothesized that nursery propagation and out-planting of seedlings for
- results and out-planting of second se
- 74 of the algal biomass associated with the pelagic *Sargassum* spp. inundations.

Rhizophora is the most common mangrove genus along Atlantic coasts (Cerón-Souza et al., 2020) and *R. mangle* is the most commonly used and easily propagated mangrove. *R. mangle* usually grows at the seaward edge of the forest, has been shown to be the most productive of the coastal mangrove species and has great ecosystem services due to the sheltered, nursery-habitat (Mandal et al., 2019) created by its proproots hanging into adjacent lagoon. In this context, our work aimed to determine whether *R. mangle* seedlings showed enhanced growth and optimal health with the use of pure *Sargassum* spp. compost (SC) under nursery conditions. The use of pure SC would also facilitate identification of contaminants reported to be associated with pelagic *Sargassum* spp. blooms (Chávez et al., 2020; Milledge et al., 2020); in particular arsenic (Davis et al., 2021; Tonon et al., 2022).

82 **2. Materials and Methods**

83 2.1 Compost production

84 Pure Sargassum spp. compost (SC) was produced by storage of fresh un-washed and unsorted Sargassum spp. from which all non-85 sargassum plants and debris had been removed. The sargassum biomass was collected from near shore waters using a surf net at two locations in Jamaica; Hellshire coastline, St Catherine, and Fort Rocky, Palisadoes coastline (Kingston) in the summer months of 2019 and 86 87 2020 respectively. Care was taken to ensure that the collected material was free of beach sand. The Sargassum spp., which consistently 88 comprised of a mix of S. fluitans III, S. natans I and S. natans VIII (Machado et al., 2022; Schell et al., 2015) in descending order of biomass, was placed in large black plastic bags which were stored unsecured (allowing air through but not rainwater) in an earthen pit for 89 90 three months. The material was periodically 'turned' to effect mixing. The highest temperature recorded from the composting material 91 during the day was ~ 36 °C.

92 **2.2. Propagule establishment**

93 Similar sized *R. mangle* propagules, collected from mangrove forests along the north coast of Jamaica were soaked in fresh water for 24

hours, measured and weighed, and then planted in a soil/sand mixture (in approximately 50:50 ratio) used as potting medium in the

95 mangrove nursery. Mangrove propagules (10 per treatment for wet nursery experiment and 3 per treatment for dry nursery experiment) were 96 planted to the same depth in potting bags containing SC, soil/sand or a mix based on the following five treatments: A- 100% SC; B- 75% SC

97 with 25% soil/sand; C- 50% SC and soil/sand; D- 25% SC and 75% soil/sand; E- 0% SC (a soil/sand only control).

98 The experiments were conducted at a mangrove nursery located at the Discovery Bay Marine Laboratory, north coast, Jamaica in successive

99 years for wet and dry nursery experiments. The wet nursery experiments were conducted in September 2019 and only maintained for

approximately 6 weeks due to death of all plants in selected treatments. The experiment was repeated under 'dry' nursery conditions

101 approximately one year later.

102 During 'wet' nursery conditions the plants were exposed to normal seawater (35 psu) or brackish water salinity of 20 psu. The water level in

103 the nursery enclosures was altered on 12 hour cycles and the enclosures in which the plants were kept allowed to completely drain for 12

hours before replacing the water for another 12 hours. Thus diurnal tides were simulated. Water levels were just enough to cover the soil

- during the 'high tide'. Treatments with different SC and soil/sand ratios were segregated, kept in different enclosures in the 'wet' nursery so
- 106 as not to have leachate from one treatment influencing another.
- For the 'dry' nursery experiment no 'tidal' fluctuations were introduced, however plants were watered from above with a sprinkler type set up twice per day (06:30 and 17:00). Treatments were also segregated, kept far enough apart in the nursery so as not to have the leachate mixing across treatments.
- 110 The dry nursery experiment was established in September 2020 and terminated in June 2021 (40 weeks) and plants in all treatments were
- alive up to the end of the experiment. Seedlings were assessed each week for number of leaves, number of nodes, height above soil level

- and health status. Height, nodal distances, leaves and biomass have proved to be useful mangrove seedling parameters (Gab-Alla et al.,
- 113 2003; Hoppe-Speer et al., 2011). The health index was developed for our assessments to facilitate having an objective indication of seedling
- 114 condition considering the potential deleterious effects of the *Sargassum* spp. The health index was based on a numeric scale of 0-5 as
- 115 follows:
- 116 0 Dead, no leaves, no green areas on stem, stem withered.
- 117 1- Surviving (plant has green colour on stem) but no leaves;
- 118 2- Surviving, plant with few withered leaves, pale or yellow with curling, scarring or spotting of leaves;
- 119 3- Poor health, possessing few leaves with evidence of thinning, and pale or yellow;
- 120 4- Good health, leaves not fully green but succulent (not thin or curled);
- 121 5- Best health possible, leaves fully green and succulent.

122 2.3 Elemental analysis using Inductively Coupled Plasma Mass Spectrometry (ICP-MS)

123 At the end of the dry nursery experiment *R. mangle* seedlings were removed from the soil, thoroughly washed with fresh water and sun-dried

- before roots and leaves were separated from the stems and ground separately for elemental analysis. Sun-drying was done over 48 hours at
- temperatures between 30 and 35 °C in the day and 20 to 24 °C at night. Stems were not analysed. Samples of the pure SC and the soil/sand
- potting medium (n = 3) were also sun-dried for elemental analysis, as indicated for the mangrove seedlings. All samples (\sim 0.2g) were digested in a CEM MARS6 microwave digestion system in 20 ml Xpress vessels using concentrated sub boiled nitric acid at 200 °C for 10
- digested in a CEM MARS6 microwave digestion system in 20 ml Xpress vessels using concentrated sub boiled nitric acid at 200 °C for 10 min (CEM standard Xpress procedure for plant material). Following digestion, the samples were diluted with Milli-Q water to form an ~ 20
- min (CEW standard Apress procedure for plant material). Following digestion, the samples were diluted with Wini-Q water to form an ~ 20 min mother solution then subsampled to give an $\sim 1000x$ total dilution. The daughter samples were spiked to give a final concentration of 5
- 130 ppb for In and Re to act as internal standards. Standards were made from the Inorganic Ventures Environmental standard (IV-Stock-50) and
- also spiked to give 5 ppb In and Re final concentrations. The samples were analysed on an Agilent 8900 QQQ-ICP-MS in standard, He and
- 132 O₂ modes depending on the element of interest.

133 2.4 Statistical tests

Statistical tests were conducted using SPSS Version 22 to assess if there was significant difference in plant parameters across the different treatments. P < 0.05 was accepted as significant for all parameters. The Shapiro-Wilks analysis was used to test normality amongst all the parameters, and due to the assumptions of the data, where normality was achieved (number of leaves and number of nodes) the one-way ANOVA test of variance was conducted. The Kruskal-Wallis analysis was used where normality was not achieved (Height and Health Ranking). Kruskal-Wallis was also used to assess significant differences in element concentration (in ppm) between treatments for roots and shoots, pure SC and soil/sand medium. Principal component analysis was used to group the 25 elements based on concentrations across

140 treatments, pure SC and soil/sand medium.

141 **3. Results**

142 **3.1 'Dry' nursery experiment**

143 After 40 weeks, median height (p < 0.001), number of leaves (p = 0.045) and seedling health (p < 0.001) were found to vary significantly

144 between treatments (Supplementary File S1). Height (Fig. 1) indicated best seedling growth occurred in the 75% Sargassum spp. compost

145 (SC) treatment while the poorest growth was from the control (0% SC); followed by seedlings in 100% SC.

146 Number of leaves was similarly greatest for seedlings in 75% SC treatments (Fig. 2) and by the end of the experiment lateral branching was

147 observed only in that treatment. Seedling health index (Fig. 3) was highest in the control and poorest in 100% SC, followed by the 50% SC

148 treatment which also showed poor health (comparable to 100% SC treatment). This was indicated by curling of the leaves and dark spots

along with yellowing/loss of colour (Fig. 4). Leaves also appeared thin (loss of succulence). Number of nodes did not vary significantly

150 between treatments (p = 0.860).

151 **3.2 'Wet' nursery experiments**

152 All seedling parameters assessed in the wet nursery varied significantly between SC treatments (height, p < 0.001; number of leaves, p <

153 0.001; number of nodes, p < 0.001; health index, p < 0.001) (Supplementary File S2). However, the most noteworthy finding was the rapid 154 demise of the plants under 'wet' nursery conditions. Seedling survival in the 'wet' nursery was especially poor for treatments of 100% and

154 demise of the plants under wet nursery conditions. Seeding survival in the wet nursery was especially poor for treatments of 100% and 155 50% SC. Complete loss of leaves and presumed death of the plants was seen after three weeks (Table 1). One plant of ten subsequently

recovered in the 50% treatment but there was no recovery of plants in the 100% treatment after 6 weeks. The control (0 % SC), by contrast

157 showed 90% seedling survival.

158 The growth parameters (height, number of leaves, number of nodes) and health rank assessed in the surviving 'wet' nursery plants showed a

159 similar pattern across treatments. Number of leaves, number of nodes and health were greatest in the control (0% SC) while the 50 and

160 100% SC treatments showed lowest values (Fig. 5). Height was again similar with 50% and 100% SC treatments showing poorest growth

161 while plants in 75% SC treatment showed greatest values (similar to the control and the 25% SC treatment).

162 **3.3 Elemental analysis**

163 Elemental analysis of seedlings after the dry nursery experiments indicated 25 elements (Table 2). Six elements (Na, Mg, K, Ca, As, Se),

164 found in high concentrations in the SC had minimal concentrations in the plants (roots or leaves) (Supplementary Fig. S1). Most of the

elements (Be, Al, V, Cr, Ni, Co, Mn, Fe, Ag, Ba, Pb, Th, U) had high concentrations in the soil/sand medium compared to the SC, however

these were still low in the plants (Supplementary File S.3). Most elements (19 of 25) displayed a general pattern of greater mean

167 concentrations in the roots than in the leaves while only four (Mg, Ca, Mn and Ti) were higher in the leaves.

168 Elements of interest, As and Na, had greatest mean concentrations in the pure SC (202.43 ppm and 50,676.21 ppm, respectively), however 169 concentrations of these were much lower in the *R. mangle* seedlings (Table 2). Arsenic (As) concentrations in the *R. mangle* seedlings were 170 less than 4% of concentrations in the SC for all except roots grown in 100% SC (which had 11.9% of As concentration in the SC). For Na,

171 mean concentration in the R. mangle seedlings grown in 100% SC were all lower than 17.7% of the SC concentration, except for roots in

172 100% SC which had mean Na concentrations of 21.5% of the SC concentration.

173 Principal Component Analysis (PCA) indicated four groups (Fig. 6). PCA also confirmed the groupings by the inclusion of elements found

174 high in the pure SC (Mg, Na, K, Ca, As, Se) in one cluster group. Ni which was high in the soil/sand medium, low in SC and with varied

175 concentrations across *R. mangle* roots and leaves was identified by the PCA as an outlier.

176 4. Discussion

177 Since the Sargassum spp. influx across the Caribbean began in 2011, these algae have been assessed for efficacy in crop production with

178 indication of success, in particular improved crop productivity as well as resistance against pests and harsh environmental conditions

(CARDI, 2015; Desrochers et al., 2020). Mohanty et al. (2013) and Walsh (2019) used Sargassum spp. in compost form and examined the 179

180 production of this for agriculture and horticulture applications. Liquid *Sargassum* spp. extract has also been used in foliar application to

181 achieve significant growth (Miranda et al., 2021). Pelagic Sargassum spp. have been shown to be rich in macro and micronutrients (Davis et

al., 2021; Machado et al., 2022; Tonon et al., 2022) and so the production of commercial fertiliser from Sargassum spp. is well established 182 183 (Desrochers et al., 2020). Interestingly, the use of *Sargassum* spp. for mangrove restoration has not been previously explored. This is

184 particularly relevant as we seek to achieve successful mangrove rehabilitation in urbanised areas which have less than ideal sediment

185 conditions, and we saw the need to explore the utility of sargassum for this application.

4.1 Impact of Sargassum spp. compost (SC) on mangrove seedlings under different nursery conditions 186

187 The wet nursery results clearly demonstrated that use of high concentrations of SC (100%) combined with waterlogged conditions result in

188 very poor outcomes, quickly affecting plant survival. Rotting of the roots was observed concomitantly with the presence of a strong smell of

189 hydrogen sulphide (H_2S) gas. Rotting *Sargassum* spp. has been reported to produce H_2S gas which has resulted in decline in natural

190 mangrove and seagrass habitats (Oxenford et al., 2021). The finding of survival in the 50% SC being similarly poor to that of the 100% SC

191 treatment is difficult to explain. The height of plants in 50% SC was also as low as the 100%, while 75% compost in the wet nursery yielded

- 192 the greatest heights. The value of the 'wet' nursery experiments however, was to demonstrate the deleterious effect of SC in water-logged
- 193 soils and a similar propensity for H₂S production as was observed with rotting of fresh Sargassum spp. in coastal areas.

194 The positive response of red mangrove seedlings to 75% SC treatment in the dry nursery was best indicated by changes in seedling height. 195 The gradual change in median height from maximum in the 75% SC along the decreasing treatment percentages (50%, followed by 25% and

then the control) indicated the positive effect of Sargassum spp. on R. mangle seedling growth. Growth in pure SC (100% treatment) was 196

197 low, being marginally higher than the control, and was accompanied by poorest health, showing that pure SC is deleterious to plant health

198 and growth. There were indications of high concentrations of growth inhibiting substances or other constituents that have deleterious effects 199

on growth and health of the plants in the undiluted compost. Several authors have reported occurrence of heavy metals and substances like

200 phenols (sometimes in high concentrations) from *Sargassum* spp. (Milledge and Harvey, 2015; Milledge et al., 2020). Arsenic in particular

- 201 has been measured from fresh and dried *Sargassum* spp. at concentrations above the accepted EPA values and with a dominance of the
- highly toxic inorganic form (Davis et al., 2021; Milledge et al., 2020; Tonon et al., 2022). The inhibitory effects from use of the 100% SC
- treatment would support the recommendation that the SC be mixed with other plant material (Milledge et al., 2020) or used in lower
- 204 concentrations with soil mixes. Use of 25 to 50% Sargassum spp. in compost trials by Walsh (2019) achieved good plant growth and low
- arsenic levels (4.2-7.2 ppm in the plants) which are within the EPA standards for such activity (Desrochers et al., 2020).
- 206 The experiment facilitated assessment of the growth (and health) parameters normally used to assess seedlings within a mangrove nursery or
- after out-planting (Trench 2021). The effect of *Sargassum* spp. on number of leaves was not as clear as seen for plant height in the dry
- 208 nursery experiments. Leaf production and retention was however noticeably best in the 75% SC treatment while the 50% treatment was the
- 209 poorest. We do not currently have any clear explanation for the observed responses in 50% SC. Mangroves, especially the *Rhizophora* spp.,
- 210 have been reported to show necrosis and leaf loss (as well as low leaf production) in response to increasing and above-normal salinity
- 211 (Aslam et al., 2011) as well as to other chemical stressors (Proffit and Develin, 1998). Other effects, like leaf thickening (succulence), have
- also been reported for *Rhizophora* spp. grown in high salt conditions (Parida and Jha, 2010). While the leaf damage observed in our
- 213 experiment was expected, the occurrence of thinning of the leaves (reduced thickness or loss of succulence) in especially the 100% treatment
- as well as in the 50% SC, would suggest that the plants were not responding to salt stress (Parida and Jha, 2010). The responses may be due
- to the effect of other (chemical) stressors. Heavy metals and other toxins, (e.g. insecticides like chlordecone), which are sequestered by
- 216 Sargassum spp. after prolonged periods in the environment (Devault et al., 2020) could be toxic to the plants in high compost concentrations.
- 217 Hence the need for elemental analysis of the plants, compost and soil used as potting medium.

218 **4.2** Partitioning of elements between soil/sand medium, *Sargassum* spp. compost and mangrove seedlings

- Our findings of high elemental concentrations associated with the soil/sand support the idea that mangrove sediments may act as an efficient metal trap (Silva et al., 1990; Saher and Asmat, 2017). These authors also indicated that mangrove sediments can be the main reservoir for heavy metals when compared to the living biomass; the later accounting for less than 1% of what was retained in the sediments. Others have reported that mangrove plants, especially *Rhizophora* spp., due to ion exchange at the roots, can actually regulate heavy metal uptake (Alongi et al., 2004; MacFarlane and Burchett, 2002; MacFarlane et al., 2007. The combined effect may account for the low elemental concentrations we observed in the plants.
- It was our expectation that seedling roots would have greater elemental concentrations than leaves based on direct contact with source, and our results did indicate that generally mangrove seedling roots had higher concentrations of elements than the leaves. Mangroves are reported to be resistant to heavy metals (Machado and Lacerda, 2004) and tolerance, in especially *R. mangle*, has been suggested by Walsh et al. (1979) to be due to formation of non-toxic sulfides in the root or at the root surface. The plants may also employ detoxification at the tissue level and ion-exclusion at the roots which is part of the same salt-exclusion mechanism (Scholander et al., 1962), or a combination of
- these factors. Metal exclusion is therefore strongly suggested but also detoxification at the tissue level due to high tissue residues without

proportional effect on the plant (Walsh et al., 1979). This was also seen in our experiments where for some elements in high concentrations in the soil/sand potting medium and with high root uptake in the control treatment, low concentrations were found in the leaves and there was no obvious compromise to plant health. Furthermore, we observed that the only elements found to be higher in leaves than roots tended to be those naturally associated with plant leaves (e.g. Mg, Ca, Mn).

Of the elements found to be high in pure SC (and low in plants), Na and As were deemed of concern because of their abundance in naturally occurring *Sargassum* spp., and their potential to cause poor outcomes and plant damage. While sodium and magnesium are common seawater elements (Ceron-Sousa et al., 2021), sodium was important to consider because salt stress can be very damaging even to coastal plants. According to Aslam et al. (2011), no toxic substance restricts plant growth more than excess salt. Mangrove seedlings have shown leaf damage in persistent hypersalinity (Krauss et al., 2008), condition under which stunting and poor growth (sometimes leading to death) of mangroves has also been observed (Barreto, 2008; Drexler and Ewel, 2001). Arsenic is another element of concern because of its toxicity, potential to cause environmental contamination (Rodríguez-Martínez et al., 2020), and the fact that it has been consistently reported in high

concentrations in pelagic *Sargassum* spp. associated with the blooms (Davis et al., 2021; Desrochers et al., 2020).

243 **4.2.1 Sodium contamination**

244 Mangrove plants are said to be facultative halophytes meaning they do not need to grow in saline soils and are found in areas with both high 245 and fluctuating salinities (Parida and Jha, 2010). They employ various mechanisms to tolerate saline environments, including exclusion of 246 salt at the roots, excretion of salts (usually by the leaves) and some tolerance of relatively high salt concentration in the tissues (Hogart, 247 2015). Rhizophora spp. have concentric layers of hypodermal and endodermal cells (called a double-endodermis) in the roots that act as a 248 barrier (Aslam et al., 2011). This genus is also known to move salt into older leaves which are then shed (Hoppe-Speer et al., 2011). Despite 249 these mechanisms, *Rhizophora* is not able to survive in hypersaline soils and prefers the margins of the forest where a range from normal 250 seawater (35 psu) to hyposaline and tidal flushing are experienced (Hoppe-Speer et al., 2011). The natural mangrove environment, however, 251 is dynamic and species therefore tend to be tolerant to a range of conditions even for a short time. In less than ideal situations, these species 252 may exhibit slower growth or appear less 'healthy' rather than be eliminated (Krauss et al., 2008). In the case of hypersalinity or high 253 sodium, mangroves may also expend a lot of energy to remove salt, using large amounts of "photosynthetic energy" (Medina, 1999) and so

exhibit slower growth.

255 4.2.2 Arsenic contamination

Holopelagic Sargassum spp., like many brown seaweeds, contain concerning concentrations of arsenic (Rodríguez-Martínez et al., 2020). In

257 our experiments the poor health of the seedlings in 100% SC treatment was thought to be due to increased levels of contaminants like

arsenic. The indicators of poor health which included spotting (brown) and curling as well as yellowing of the leaves were very evident in

- the 100% Sargassum spp. compost plants. However, while elemental analysis indicated greatest mean As concentration (202.43 ppm) in the
- 260 pure SC, the only plant tissue with concentrations greater than 20% (24.1 ppm) was roots of seedlings grown in the 100% pure SC treatment.

- 261 This indicates the lack of uptake and sequestration by the *R. mangle* seedlings and the possible utility of *Sargassum* spp. in nursery
- 262 propagation of this plant. Furthermore, application on a wider scale (out-planting) may have positive outcomes.

263 **5. Application**

Collecting and drying or composting of *Sargassum* spp. in tropical coastal communities with high insolation (and low rainfall) is a relatively low cost activity that produces material with potential for several applications. In this context, composting has the greatest potential for soil amelioration as both the properties (texture and water holding capacity) as well as nutrient content of the soil or sand is enhanced (Adugna, 2016; Pan et al., 2012). Desrochers et al. (2020) have indicated that using seaweed as compost in coastal areas is one of the simplest, cheapest and most practical methods of making use of the large algal biomass. In line with this, our experiments show that in appropriate proportions, pure SC does enhance mangrove seedling growth with height, and to a lesser extent leaf production, being good indicators.

270 While potentially toxic elements found naturally in or sequestered by Sargassum spp. remain a concern, for mangroves there is evidence that

especially *R. mangle* has developed strategies to minimize uptake. These attributes of mangroves have actually been proposed for

application to mitigate metal pollution in coastal areas (Machado and Lacerda, 2004). There is therefore potential for use of the algal

273 biomass in nuisance pelagic *Sargassum* spp. blooms to support mangrove rehabilitation under conditions where the application is in

relatively dry areas (e.g. along the urbanised shorelines like Kingston Harbour, Jamaica) and under controlled applications where the

ameliorant can be contained in the immediate planting area. This could ensure better success of rehabilitation efforts in such areas as well as

276 make use of the algal biomass; leading to the benefit of impoverished coastal communities who are ultimately most affected by *Sargassum*

spp. inundations and mangrove loss.

278 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

281 Author Contributions

282 CT: investigation, formal analysis, visualization, writing - review & editing; SLT: investigation, formal analysis, writing - review & editing;

283 DT: investigation, formal analysis, writing - review & editing; GM: investigation, formal analysis, writing - review & editing; PF:

284 investigation, formal analysis, writing - review & editing; HS: formal analysis, writing - review & editing; CBM: formal analysis, writing -

285 review & editing; DW: formal analysis, writing - review & editing; TT: conceptualization, funding acquisition, investigation, formal

analysis, visualization, writing - review & editing. MW: conceptualization, funding acquisition, investigation, formal analysis, visualization,
 writing - review & editing.

288 Funding

289 This work was supported by the Economic and Social Research Council [grant number ES/T002964/1].

290 Acknowledgments

291 The authors are grateful to the following DBML interns, Javel Noble and Kadian McCalla as well as colleagues of the International Centre

for Environment and Nuclear Sciences (ICENS) and Natural Products Institute (NPI) at the University of the West Indies (Mona) for

assistance with sample processing and storage.

294 Legend of Figures

- Fig. 1. Median seedling height (bar), 25-75% range (box) and non-outlier range (whisker) for red mangrove seedlings in each treatment indicating the % sargassum compost (SC).
- Fig. 2. Median number of leaves (bar), 25-75% range (box) and non-outlier range (whisker) for red mangrove seedlings in each treatment indicating the % sargassum compost (SC).
- Fig. 3. Median health rank (bar), 25-75% range (box) and non-outlier range (whisker) for red mangrove seedlings in each treatment indicating the % sargassum compost (SC).
- Fig. 4. *R. mangle* seedlings showing leaf damage after growth in 100% SC (A) and seedlings in 75% SC showing lateral branching (B). The lateral branching was observed at the end of the experiment.
- Fig. 5. Change in seedling parameters; number of leaves, number of nodes, height and health (means with \pm 95% CI) across SC treatment in the 'wet' nursery averaged for all weeks.
- 305 Fig. 6. Principal Component analysis (PCA) showing grouping of elements according to concentrations in ppm.
- 306

307 Supporting materials

- 308 S1. Supplementary File S1 (Excel File). Statistical tests and boxplots for dry nursery plant parameters.
- 309 S2. Supplementary File S2 (Excel File). Statistical tests and boxplots for wet nursery plant parameters.
- 310 S3. Supplementary File S3 (Excel File). Raw data and boxplots for all elements in plants, soil/sand and SC.

311 Fig. S1. Boxplots of six elements found high in SC and low in the mangrove plants.

312

313

314 **References**

- Adugna, G., 2016. A review on impact of compost on soil properties, water use and crop productivity. *Academic Research Journal of Agricultural Science and Research* 4, 93-104.
- 317 Alongi, D.M., 2012. Carbon sequestration in mangrove forests. *Carbon Management* 3, 313-322.
- 318 Alongi, D.M., 2014. Carbon cycling and storage in mangrove forests. *Annual Review of Marine Science* 6, 195-219.
- Alongi, D.M., Wattayakorn, G., Boyle, S., Tirendi, F., Payn, C. and Dixon, P., 2004. Influence of roots and climate on mineral and trace element storage and flux in tropical mangrove soils. *Biogeochemistry* 69, 105-123.
- Amador-Castro, F., García-Cayuela, T., Alper, H.S., Rodriguez-Martinez, V., and Carrillo-Nieves, D., 2021. Valorization of pelagic
 Sargassum sp. biomass into sustainable applications: Current trends and challenges. *Journal of Environmental Management* 283, 112013.
- Aslam, R., Bostan, N., Nabgha-e-Amen, M., and Safdar, W., 2011. A critical review on halophytes: Salt tolerant plants. *Journal of Medicinal Plants Research* 5, 7108-7118.
- Barreto, M.B., 2008. Diagnostics about the state of mangroves in Venezuela: case studies from the National Park Morrocoy and wildlife
 refuge Cuare. In: Lieth PH, Sucre DMG, Herzog B (eds) Mangroves and halophytes: restoration and utilisation. Springer, Netherlands, 51 64
- 328 CARDI, 2015. *Sargassum* seaweed and its use in crop and livestock production: Possible agri-business opportunities. Policy Brief 2015, 1.
 329 Available online: http://www.cardi.org/ Sargassum-Seaweed-and-its-use-incrop-and-livestock-production-CARDI-Policy-Brief.pdf
 330 (accessed on 23 February 2022).
- Cerón-Souza, I., Barreto, M.B., Barreto-Pittol, E., Silva, A., Feliner, G.N., and Medina, E., 2021. *Rhizophora* zonation, salinity, and nutrients in the western atlantic. *Biotropica* 53, 384-396.

333 Chávez, V., Uribe-Martínez, A., Cuevas, E., Rodríguez-Martínez, R.E., van Tussenbroek, B.I., Francisco, V., Estévez, M., Celis, L.B.,

- 334 Monroy-Velázquez, L.V., Leal-Bautista, R., and Álvarez-Filip, L., 2020. Massive influx of pelagic *Sargassum* spp. on the coasts of the
- 335 Mexican Caribbean 2014–2020: challenges and opportunities. *Water* 12, 2908.
- Davis, D., Simister, R., Campbell, S., Marston, M., Bose, S., McQueen-Mason, S.J., Gomez, L.D., Gallimore, W.A., and Tonon, T., 2021.
- 337 Biomass composition of the golden tide pelagic seaweeds Sargassum fluitans and S. natans (morphotypes I and VIII) to inform valorisation
- 338 pathways. *Science of the Total Environment* 762, 143134.
- Desrochers, A., Cox, S., Oxenford, H., and van Tussenbroek, B. 2020. *Sargassum* sp. uses guide: A resource for Caribbean researchers,
 entrepreneurs and policy makers. *CERMES Technical Reports* 97, 172.
- 341 Devault, D.A., Massat, F., Baylet, A., Dolique, F., and Lopez, P.J., 2022. Arsenic and chlordecone contamination and decontamination 342 toxicokinetics in *Sargassum* sp. *Environmental Science and Pollution Research* 29, 6-16.
- Donato, D., Kauffman, J., Murdiyarso, D., Kurnianto, S., Stidham, M., and Kanninen M., 2011. Mangroves Among the Most Carbon-rich
 Forests in the Tropics. *Nature Geoscience* 4, 293-297.
- Drexler, J.Z., and Ewel, K.C., 2001. Effect of the 1997–1998 ENSO-related drought on hydrology and salinity in a micronesian wetland
 complex. *Estuaries* 24, 347-356.
- 347 Eyras, M., Rostagn, C., and Defosse, G., 1998. Biological evaluation of Sargassum composting. Compost Science and Utilization 6, 74-81.
- Gab-Alla, A., Khafagi, I., Salama, W. and Fouda, M., 2003. Production of nursery-reared seedlings of the gray mangrove *Avicennia marina* under laboratory conditions. *Egyptian Journal of Biology* 5, 55-61.
- Giri, C., Zhu, Z., Tieszen, L., Singh, A., Gillette, S., and Kelmelis J., 2008. Mangrove Forest Distributions and Dynamics (1975–2005) of the Tsunami-Affected Region of Asia. *Journal of Biogeography* 35, 519-528.
- 352 Hogarth, P., 2015. The Biology of Mangroves and Seagrasses. 3rd Ed. United Kingdom: Oxford University Press.
- Hoppe-Speer, S.C., Adams, J.B., Rajkaran, A., and Bailey, D., 2011. The response of the red mangrove *Rhizophora mucronata* Lam. to salinity and inundation in South Africa. *Aquatic Botany* 95, 71-76.
- Kairo, J., Dahdouh-Guebas, F., Bosire, J., and Koedam N., 2001. Restoration and Management of Mangrove Systems A Lesson for and from the East African Region. *South African Journal of Botany* 67, 383-389.

- 357 Kathiresan, K., 2012. Importance of mangrove ecosystem. *International Journal of Marine Science* 2, 70-89.
- Krauss, K.W., Lovelock, C.E., McKee, K.L., López-Hoffman, L., Ewe, S.M., and Sousa, W.P., 2008. Environmental drivers in mangrove establishment and early development: a review. *Aquatic Botany* 89, 105-127.
- Hoppe-Speer, S.C., Adams, J.B., Rajkaran, A., and Bailey, D., 2011. The response of the red mangrove *Rhizophora mucronata* Lam. to salinity and inundation in South Africa. *Aquatic Botany* 95, 71-76.
- Lee, S.Y., Primavera, J.H., Dahdouh-Guebas, F., McKee, K., Bosire, J.O., Cannicci, S., Diele, K., Fromard, F., Koedam, N., Marchand, C.,
 and Mendelssohn, I., 2014. Ecological role and services of tropical mangrove ecosystems: a reassessment. *Global Ecology and Biogeography* 23, 726-743.
- Lewis, R.R., 2009. Methods and criteria for successful mangrove forest restoration. *Coastal Wetlands: An integrated ecosystem approach*,
 787-800.
- MacFarlane, G.R., and Burchett, M.D., 2002. Toxicity, growth and accumulation relationships of copper, lead and zinc in the grey mangrove
 Avicennia marina (Forsk.) Vierh. *Marine Environmental Research* 54, 65-84.
- MacFarlane, G.R., Koller, C.E., and Blomberg, S.P., 2007. Accumulation and partitioning of heavy metals in mangroves: a synthesis of field-based studies. *Chemosphere* 69, 1454-1464.
- 371 Machado, W., and Lacerda, L.D., 2004. Overview of the biogeochemical controls and concerns with trace metal accumulation in mangrove 372 sediments. In *Environmental geochemistry in tropical and subtropical environments* (pp. 319-334). Springer, Berlin, Heidelberg.
- 373 Machado, C. B., Maddix, G. Francis, Thomas, S., Burton, J., Langer, S., Larson, T., Marsh, R., Webber, M., and Tonon. T., 2022. Pelagic
- *Sargassum* events in Jamaica: morphotype abundance, provenance, and influence of sample preparation on biochemical composition.
 Science of the Total Environment 817, 152761.
- Mandal, A., Smith R-A., Edwards, T., Kinlocke, R., Mitchell, S., Webber M., Trench, C., Francis P., Spence, A., and Edwards, P., 2019.
 Forces of Nature: Assessment of the Economic Valuation of Coastal Protection Services Provided by Mangroves in Jamaica. Edited by The
 World Bank-PROFOR-World Bank Caribbean Office.
- Medina, E., 1999. Mangrove physiology: the challenge of salt, heat, and light stress under recurrent flooding. *Ecosistemas de manglar en América tropical* 109-126.

Milledge, J.J., and Harvey, P.J., 2016. Golden tides: problem or golden opportunity? The valorisation of *Sargassum* from beach inundations.
 Journal of Marine Science and Engineering 4, 60.

Milledge, J.J., Nielsen, B.V., and Bailey, D., 2016. High-value products from macroalgae: the potential uses of the invasive brown seaweed,
 Sargassum muticum. Reviews in Environmental Science and Bio/Technology 15, 67-88.

Miranda, J.L.L., Celis, L.B., Estévez, M., Chávez, V., Van Tussenbroek, B.I., Uribe-Martínez, A., Cuevas, E., Pantoja, I.R., Masia, L.,
 Cauich-Kantun, C., and Silva, R., 2021. Commercial Potential of Pelagic *Sargassum* spp. in Mexico. *Frontiers in Marine Science*.

Mitra, A., Zaman, S., and Gobato, R., 2018. Indian Sundarban Mangroves: A potential Carbon Scrubbing System. *Parana Journal of Science and Education* 4, 7-29.

389 McCalla, K. 2019. Assessing the effects of *Sargassum* sp. compost in different proportions on the growth of red mangrove seedlings.

BIOL3413 Research Project, Department of Life Sciences, University of the West Indies. Pp. 42.

391 Medina, E., 1999. Mangrove physiology: the challenge of salt, heat, and light stress under recurrent flooding, p. 109-126. In: A. Yáñez-

392 Arancibia y A. L. LaraDomínguez (eds.). Ecosistemas de Manglar en América Tropical. Instituto de Ecología A.C. México, UICN/ORMA,

393 Costa Rica, NOAA/NMFS Silver Spring MD USA.

Mohanty, D., Adhikary, S., Chattopadhyay, G., Prakasam, V., Gopamma, D., Ramoji, G., et al. (2013). Seaweed liquid fertilizer (SLF) and its role in agriculture productivity. The Ecoscan III 23–26, 1-10.

Nguyen, T.P., Van Tam, N., and Parnell, K.E., 2016. Community perspectives on an internationally funded mangrove restoration project:
 Kien Giang province, Vietnam. *Ocean & Coastal Management* 119, 146-154.

Oxenford, H.A., Cox, S.A., van Tussenbroek, B.I., and Desrochers, A., 2021. Challenges of turning the *Sargassum* crisis into gold: current
 constraints and implications for the Caribbean. *Phycology* 1, 27-48.

400 Pan, I., Dam, B., and Sen, S.K., 2012. Composting of common organic wastes using microbial inoculants. *3 Biotech* 2, 127-134.

401 Parida. A.K., and Jha, B. 2010. Salt tolerance mechanisms in mangroves: a review. *Trees* 24, 199-217.

402 Reef, R., Feller, I.C., and Lovelock, C.E., 2010. Nutrition of mangroves. *Tree Physiology* 30, 1148-1160.

Rodríguez-Martínez, R.E., Roy, P.D., Torrescano-Valle, N., Cabanillas-Terán, N., Carrillo-Domínguez, S., Collado-Vides, L., GarcíaSánchez, M., and van Tussenbroek, B.I., 2020. Element concentrations in pelagic *Sargassum* sp. along the Mexican Caribbean coast in
2018-2019. *PeerJ* 8, e8667.

Saher, N.U., and Asmat, S.S., 2017. Evaluation of heavy metals contamination in mangrove sediments and their allied fiddler crab species
 (*Austruca sindensis* (alcock, 1900) from Hawks bay, Karachi, Pakistan. *Pakistan International Journal of Biology and Biotechnology* 14,
 411-417.

Salter, M.A., Rodríguez-Martínez, R.E., Álvarez-Filip, L., Jordán-Dahlgren, E., and Perry, C.T., 2020. Pelagic *Sargassum* as an emerging
 vector of high rate carbonate sediment import to tropical Atlantic coastlines. *Global and Planetary Change* 195, 103332.

Sembera, J.A., Meier, E.J., and Waliczek, T.M., 2018. Composting as an alternative management strategy for sargassum drifts on coastlines.
 HortTechnology 28, 80-84.

Schell, J.M., Goodwin, D.S., and Siuda, A.N., 2015. Recent *Sargassum* inundation events in the Caribbean: shipboard observations reveal
 dominance of a previously rare form. *Oceanography* 28, 8-11.

415 Silva, C.A.R., Lacerda, L.D., and Rezende, C.E., 1990. Metals reservoir in a red mangrove forest. *Biotropica* 339-345.

Thompson, T.M., Young, B.R., and Baroutian, S., 2020. Pelagic *Sargassum* for energy and fertiliser production in the Caribbean: A case
 study on Barbados. *Renewable and Sustainable Energy Reviews* 118, 109564.

Tonon, T. Machado, C.B. Webber, M. Webber, D. Smith, J. Pilsbury, A. Cicéron, F. Herrera-Rodriguez, L. Jimenez, E.M. Suarez, J.V., et al., 2022. Biochemical and Elemental Composition of Pelagic Sargassum Biomass Harvested across the Caribbean. *Phycology* 2, 204-215.

- 420 Trench, C., 2021. Hydrological restoration approaches for degraded mangrove forests in Jamaica. PhD Thesis. Department of Life Sciences,
 421 UWI (Mona).
- 422 Turner, R., and Lewis, R. III. 1997. Hydrologic Restoration of Coastal Wetlands. Wetland Ecology Management 4, 65-72.

Uddin, M.S., van Steveninck, E.D.R., Stuip, M., and Shah, M.A.R., 2013. Economic valuation of provisioning and cultural services of a
 protected mangrove ecosystem: A case study on Sundarbans Reserve Forest, Bangladesh. *Ecosystem Services* 5, 88-93.

425 Valiela, I., Bowen, J., and York, J. 2001. Mangrove Forests: One of the World's Threatened Major Tropical Environments: At Least 35% of

426 the Area of Mangrove Forests Has Been Lost in the Past Two Decades, Losses that Exceed those for Tropical Rain Forests and Coral Reefs,

427 Two Other Well-Known Threatened Environments. *Bioscience* 51, 807-815.

van Tussenbroek, B.I., et al. (2017). Severe impacts of brown tides caused by *Sargassum* sp. spp. on near-shore Caribbean seagrass
 communities. *Marine Pollution Bulletin* 122, 272-281.

- Walsh, K.T., 2019. Examining the quality of a compost product derived from *Sargassum (Sargassum fluitans* and *Sargassum natans*). MS
 Thesis- Texas State University.
- Walsh, G.E., Ainsworth, K.A., and Rigby, R., 1979. Resistance of red mangrove (*Rhizophora mangle* L.) seedlings to lead, cadmium, and
 mercury. *Biotropica* 22-27.
- Wang, M., Hu, C., Barnes, B.B., Mitchum, G., Lapointe, B., and Montoya, J.P., 2019. The great Atlantic *Sargassum* sp. belt. *Science* 365, 83-87.

436 Webber, M., Calumpong, H., Ferreira, F., Granek, E., Green, S., Ruwa, R., and Soares, M., 2016. Mangroves. The First Global Integrated

- 437 Marine Assessment: World Ocean Assessment I, pp.877-886. Cambridge University Press-
- 438 http://www.un.org/depts/los/global_reporting/WOA_RPROC/Chapter_48.pdf.
- Williams, A., and Feagin, R. 2010. *Sargassum* sp. as a natural solution to enhance dune plant growth. *Environmental Management* 46, 738 747.
- 441
- 442
- 443
- 444
- . . .
- 445
- 446
- 447
- 448

Table 1. Mangrove seedling survival in 'wet' nursery conditions showing death of the plants (0% survival) by week 3 in 100% SC.
Experiment terminated after week 6 when one treatment had death of all plants.

Number of mangrove seedlings remaining alive in each treatment per									
SC Treatments	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6	% survival		
0% SC (Control)	10	10	9	9	9	9	90		
25% SC	10	10	9	7	6	6	60		
50% SC	10	10	0	0	1	1	10		
75% SC	5	5	5	3	3	3	60		
100% SC	10	10	0	0	0	0	0		

451



Table 2. Element concentration (mean with S.E.) in parts-per-million (ppm) for plant sections in all SC treatments, pure SC and soil/sand
 potting mix.

Element	Leaves in 0% SC	Leaves in 25% SC	Leaves in 50% SC	Leaves in 75% SC	Leaves in 100% SC	Roots in 0% SC	Roots in 25% SC	Roots in 50% SC	Roots in 75% SC	Roots in 100% SC	Soil/ Sand	Pure SC
Ве	0.01±.01	0.000±.000	0.008±.006	0.004±.002	0.003±.003	0.05±.014	0.025±.004	0.025±.01	0.013±.012	0.002±.002	0.20±.01	0.01±.003
Na	1560.78	2267.14	2966.58	2444.78	4726.84	6430.93	7247.74	8979.04	8087.39	10893.02	830.95	50676.21
	±64.27	±43.04	±44.07	±146.84	±80.28	±144.24	±150.194	±135.05	±205.03	±224.90	±67.48	±525.80
Mg	4537.53	3911.61	4618.84	4107.99	5903.27	2512.29	2101.96	2105.24	2145.72	3036.78	9172.54	20767.5
	±293.23	±56.70	±111.47	±86.21	±59.42	±180.17	±7.59	±89.43	±21.60	±68.12	±730.24	±623.27
Al	57.55	26.26	63.74	58.84	44.83	2344.14	1157.15	560.85	315.01	111.99	11496.95	160.21
	±10.24	±1.99	±5.82	±13.51	±2.92	±371.06	±43.12	±101.34	±50.73	±4.43	±500.80	±10.69
K	9648.36	5212.63	6346.68	4924.16	5747.46	7812.36	4625.87	4727.49	4851.39	3094.17	1041.63	76863.61
	±321.71	±179.92	±155.64	±118.76	±129.90	±861.39	±7.44	±363.06	±196.54	±12.65	±39.98	±1621.92
Ca	26119.82	26104.25	29395.10	27501.35	30795.27	12325.62	14735.22	15985.00	17131.54	22251.60	90140.80	143919.47
	±1137.61	±512.98	±585.52	±640.01	±554.75	±340.93	±247.82	±1076.33	±408.91	±328.42	±8746.77	±6981.48
V	0.23±.06	0.08±.01	0.34±.07	0.1±.05	0.13±.01	11.61±2.46	7.19±.31	6.01±.77	3.65±.41	2.99±.08	56.01±2.27	7.21±.73

Cr	1.35±.20	0.73±.19	0.63±.08	0.35±.10	0.27±.02	27.33±2.47	18.53±1.22	9.72±1.40	9.04±1.74	7.90±.06	22.68±4.40	0.99±.14
Mn	75.83	92.67	140.22	168.29	113.99	59.72	85.36	51.63	40.37	29.95	393.70	32.08
	±4.16	±3.06	±4.65	±6.80	±2.98	±20.11	±4.83	±5.46	±3.40	±.72	±12.55	±1.07
Fe	65.50	38.09	79.11	47.49	33.62	2619.25	1158.37	686.82	439.19±58.7	161.75±1.02	17077.63	114.98
	±14.24	±11.22	±13.67	±10.27	±1.43	±804.21	±47.85	±130.58	9		±522.80	±6.28
Со	0.08±.01	0.08±.01	0.19±.01	0.19±.01	0.17±.01	1.23±.30	1.00±.05	1.34±.24	1.17±.16	1.68±.07	8.39±.24	1.47±.06
Ni	4.80±1.99	1.03±.07	1.54±.37	0.77±.15	1.15±.38	7.93±.79	5.18±.85	2.47±.31	2.38±.64	5.33±.99	13.31±.71	4.52±.14
Cu	21.67	11.54	11.06	7.67	7.83	53.46	135.30	44.01	57.79	46.79	42.76	5.59
	±2.51	±.54	±1.41	±1.96	±1.31	±1.29	±76.39	±9.11	±7.98	±6.01	±2.05	±.22
Zn	11.32	7.42±.46	8.44±.64	7.49	8.60	47.83	78.05	44.48	37.28±5.91	37.47±1.05	72.44±.95	14.07±.46
	±1.46			±1.10	±.45	±1.59	±34.06	±10.26				
As	0.44±.04	0.63±.03	1.14±.04	0.99±.07	1.11±.02	1.89±.01	5.72±.47	14.85	10.00	24.11	27.09	202.43
								±2.65	±1.52	±.95	±2.38	±2.13
Se	0.17±.004	0.11±.01	0.09±.02	0.06±.01	0.04±.01	0.19±.03	0.22±.02	0.19±.04	0.11±.01	0.16±.02	0.68±.06	1.29±.03
Мо	0.39±.04	0.26±.002	0.25±.01	0.16±.02	0.10±.01	0.90±.09	0.80±.05	0.57±.06	0.48±.04	0.55±.04	0.70±.15	0.33±.03

Ag	0.004	$0.002 \pm .001$	0.004	$0.003 \pm .00$	0.003	$0.02 \pm .002$	$0.02 \pm .006$	0.01±.002	0.01±.002	0.01±.001	$0.05 \pm .002$	0.02±.002
	±.001		±.001		±.001							
Cd	0.01±.002	0.01±.001	0.02±.002	0.02±.003	0.02±.002	0.40±.01	0.47±.03	0.65±.16	0.31±.05	0.58±.02	1.02±.20	0.64±.01
Sb	0.01±.003	0.01±.003	0.01±.00	0.004	0.003±.00	0.03±.002	$0.04 \pm .004$	0.03±.01	0.04±.01	0.07±.006	0.03±.004	0.08±.003
				±.001								
Ba	44.86	21.35	14.02	8.29	3.08	51.17	17.26	12.26	7.24	3.51	148.91	38.98
	±1.61	±.48	±.25	±.11	±.002	±9.78	±.44	±.92	±.91	±.11	±14.53	±.48
Ti	0.02±.002	0.02±.00	0.02±.002	0.02±.00	0.02±.002	0.01±.001	0.01±.00	0.01±.001	0.01±.001	0.01±.001	0.04±.00	0.03±.004
Pb	0.76±.15	0.31±.03	0.41±.07	0.28±.09	0.29±.04	2.23±.06	4.45±2.09	2.10±.45	2.68±.39	2.66±.15	8.87±.64	0.34±.03
Th	0.01±.001	0.003±.00	$0.005 \pm .00$	0.01±.001	$0.004 \pm .00$	0.09±.01	0.05±.002	0.02±.01	0.02±.003	0.01±.00	0.86±.03	0.02±.001
U	0.01±.002	0.01±.001	0.01±.001	0.02±.004	0.01±.001	0.30±.002	0.24±.01	0.27±.03	0.20±.03	0.28±.01	1.75±.13	1.22±.08