

This is a repository copy of *Nuclear moments of indium isotopes reveal abrupt change at magic number 82*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/191938/>

Version: Accepted Version

Article:

Vernon, A. R., Garcia Ruiz, R. F., Miyagi, T. et al. (22 more authors) (2022) Nuclear moments of indium isotopes reveal abrupt change at magic number 82. *Nature*. pp. 260-265. ISSN 0028-0836

<https://doi.org/10.1038/s41586-022-04818-7>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Nuclear moments of indium isotopes reveal abrupt change at magic number 82

A.R. Vernon^{1,2,*}, R.F. Garcia Ruiz^{2,3,*}, T. Miyagi⁴, C.L. Binnersley¹, J. Billowes¹, M.L. Bissell¹, J. Bonnard⁵, T.E. Cocolios⁶, J. Dobaczewski^{5,7}, G.J. Farooq-Smith⁶, K.T. Flanagan^{1,8}, G. Georgiev⁹, W. Gins^{6,10}, R.P. de Groote^{6,10}, R. Heinke^{3,13}, J.D. Holt^{4,11}, J. Hustings⁶, Á. Koszorús⁶, D. Leimbach^{12,13,14}, K.M. Lynch³, G. Neyens^{6,3}, S.R. Stroberg¹⁵, S.G. Wilkins^{1,2}, X.F. Yang^{6,16}, and D.T. Yordanov^{3,9}

¹School of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, United Kingdom

²Massachusetts Institute of Technology, Cambridge, MA 02139, USA

³EP Department, CERN, CH-1211 Geneva 23, Switzerland

⁴TRIUMF, 4004 Wesbrook Mall, Vancouver, BC V6T 2A3, Canada

⁵Department of Physics, University of York, Heslington, York YO10 5DD, United Kingdom

⁶KU Leuven, Instituut voor Kern- en Stralingsfysica, B-3001 Leuven, Belgium

⁷Institute of Theoretical Physics, Faculty of Physics, University of Warsaw, ul. Pasteura 5, PL-02-093 Warsaw, Poland

⁸Photon Science Institute, The University of Manchester, Manchester M13 9PL, UK

⁹IJCLab, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France

¹⁰Department of Physics, University of Jyväskylä, PB 35(YFL) FIN-40351 Jyväskylä, Finland

¹¹Department of Physics, McGill University, 3600 Rue University, Montréal, QC H3A 2T8, Canada

¹²EN Department, CERN, CH-1211 Geneva 23, Switzerland

¹³Institut für Physik, Johannes Gutenberg-Universität Mainz, D-55128 Mainz, Germany

¹⁴Department of Physics, The University of Gothenburg, Gothenburg, Sweden

¹⁵Department of Physics, University of Washington, Seattle, Washington 98105, USA

¹⁶School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China

*Corresponding authors: vernona@mit.edu, rgarcia@mit.edu

In spite of the high-density and strongly correlated nature of the atomic nucleus, experimental and theoretical evidence suggests that around particular ‘magic’ numbers of nucleons, nuclear properties are governed by a single unpaired nucleon^{1,2}. A microscopic understanding of the extent of this behavior and its evolution in neutron-rich nuclei remains an open question in nuclear physics^{3–5}. The indium isotopes are considered a textbook example of this phenomenon⁶, where the constancy of their electromagnetic properties indicated a single unpaired proton hole can provide the identity of a complex many-nucleon system^{6,7}.

Here, we present precision laser spectroscopy measurements performed to investigate the validity of this simple single-particle picture. Observation of an abrupt change in the dipole moment at $N = 82$ indicates that while the single-particle picture indeed dominates at neutron magic number $N = 82^{2,8}$, it does not for previously studied isotopes.

To investigate the microscopic origin of these observations, our work provides a combined effort with developments in two complementary nuclear many-body methods: *ab-initio* valence space in-medium similarity normalization group and density functional theory. We find that the inclusion of time symmetry-breaking mean fields is essential for a correct description of nuclear magnetic properties, which were previously poorly constrained. These experimental and theoretical findings are key to understand how seemingly simple single-particle phenomena naturally emerge from complex interactions among protons and neutrons.

Introduction

The atomic nucleus is formed by strongly-interacting nucleons (protons, Z , and neutrons, N), packed tightly into a volume around a trillion times smaller than that of atoms. Hence, describing the atomic nuclei and predicting their properties at extreme values of mass and charge are major long-standing challenges for nuclear science. Similar to electrons in an atom, the nucleons (protons and neutrons) in the atomic nucleus occupy quantum ‘shells’. Thus, nuclei with a single valence particle or hole around a nuclear closed shell provide important foundations for our understanding of the atomic nucleus.

Their simpler structure vastly reduces the complexity of the quantum many-body problem, providing critical guidance for the development of nuclear theory.

Recent advances in our understanding of the strong interaction and the development of many-body methods, combined with escalation in computer power, have enabled theoretical descriptions of increasingly complex nuclei. Isotopes around the proton closed shell $Z = 50$, are now the frontier of *ab initio* calculations^{9,10}. The properties of these nuclei can be described by complementary many-body methods such as configuration-interaction methods⁴ and nuclear Density Func-




Figure 1. (color online) **Evolution of nuclear electromagnetic properties for the $9/2^+$ ground states of $^{105-131}\text{In}$ isotopes.** a) the electric quadrupole moments and b) magnetic dipole moments. The horizontal dotted line indicates the single-particle value (Schmidt limit). Experimental results are compared with theoretical calculations from *ab initio* (VS-IMSRG) and Density Functional Theory (DFT). Literature experimental values for $^{105-127}\text{In}$ were taken from Ref.⁷. The evolution of collective properties of these isotopes are illustrated at the bottom of the figure: left) quadrupole polarisation gradually reduces to a single proton hole value at $N = 82$. right) the magnetic dipole moments abruptly approach the value for a single-proton hole in a ^{132}Sn core at $N = 82$, as the dominant effect changes from charge to spin distribution.

tional Theory (DFT)¹¹. This has led to an increased focus on studying this region of the nuclear chart (around $Z = 50$, $N = 50, 82$) over the last decade^{2,8,12-14}.

Here we present measurements of two fundamental properties of indium isotopes using precision laser spectroscopy: the (spectroscopic) magnetic dipole moment, μ , and electric quadrupole moment, Q . Measurements were performed for the neutron-rich In ($Z = 49$) isotopes, reaching up to ^{131}In , which possesses a magic number of $N = 82$ neutrons (see Methods section for details). With a single proton hole configuration with respect to the well established^{2,8,14} proton closed-shell of $Z = 50$, the low-energy structure of the odd-mass indium isotopes is expected to be governed by a single-hole configuration in the proton orbit $\pi 1g_{9/2}$.

As we show here, the nuclear magnetic dipole moments of odd-mass indium isotopes are determined by the total spin distribution of the nucleus induced by the unpaired valence proton hole. This is schematically illustrated at the bottom right of Figure 1. The nuclear electric quadrupole moment provides a complementary measurement of the nuclear charge distribution, and is highly sensitive to the collective motion

of all nucleons¹⁵. These observables together therefore probe distinct aspects of the nucleon distribution and measuring them across a large range of neutron numbers allows a unique insight into the evolution of the interplay between single-particle and collective nuclear phenomena.

Previously, the magnetic moments of the ground state $I^\pi = 9/2^+$ of indium isotopes were known to exhibit remarkably little variation over 22 isotopes, from $A = 105$ to 127 (see Figure 1b) open symbols)⁷. The constant value of the magnetic moment over such a long range of isotopes has been presented as an archetypal example of the independent-particle behaviour of single-particle states near a proton shell closure^{6,7}. ‘How do these seemingly simple patterns emerge from complex interactions among protons and neutrons?’ and ‘Do they prevail at extreme number of neutrons?’ are two major open questions that we address in this work.

In addition to the ground state, the indium isotopes can also exist in excited nuclear configurations with relatively long lifetimes - isomers - with spin $I^\pi = 1/2^-$. These isomeric states provide additional insight and are expected to be described by a single-hole configuration based on a different proton orbital

($\pi 2p_{1/2}$). However, in contrast to the $I^\pi = 9/2^+$ states, the μ values of these isomeric states exhibit significant variations (see Figure 3), posing a three-decades-long puzzle in our description of these nuclei⁷.

To unravel the microscopic origin of the electromagnetic properties of these isotopes, we compare our experimental results with two complementary state-of-the-art theoretical methods: i. *ab initio* valence-space in-medium similarity renormalisation group (VS-IMSRG) calculations^{9,16}, which start from nucleon-nucleon interactions derived from chiral effective field theory¹⁷; and ii. symmetry-breaking nuclear Density Functional Theory (DFT)^{11,18}. The latter assumes nucleons moving within their own self-consistently-generated spin-dependent broken-symmetry-confining potential. DFT provides a satisfactory description of bulk nuclear properties such as radii and binding energies across the whole nuclear chart^{19–21}. Here, we have developed its symmetry-restored version²² to be able to provide accurate calculations of spectroscopic μ and Q moments.

Experimental and Theoretical developments

Our measurements were performed using the collinear resonance ionisation spectroscopy (CRIS) technique at the ISOLDE facility of CERN²³ (see Methods section for details). From the hyperfine structure, we extracted the magnetic dipole and electric quadrupole parameters, A_{hf} , B_{hf} of the probed atomic states, corresponding to the two long-lived nuclear states, $9/2^+$ and $1/2^-$, present in each isotope. Recent improvements in the sensitivity of the technique allowed us to achieve high-resolution spectroscopy measurements, despite production of the indium isotopes at rates below 1000 atoms/s in the presence of large isobaric contamination. In order to enable these measurements, laser ionisation spectroscopy schemes had to be developed which are both sensitive to low isotope rates from a large background and sensitive to nuclear properties through hyperfine structure, in addition to the development of increased accuracy atomic calculations for the extraction of their moments^{24–26}.

We performed *ab initio* VS-IMSRG calculations (see Methods for further details) using two different sets of initial two- (NN) and three-nucleon (3N) forces derived from chiral effective field theory^{27,28}; the 1.8/2.0(EM)¹⁷ and the more recent N²LOGO²⁹. The 1.8/2.0(EM) set is constrained only by fitting to properties of two-, three-, and four-nucleon systems. N²LOGO was recently developed to include Δ -isobar degrees of freedom and is additionally fit to reproduce saturation properties of infinite nuclear matter²⁹.

We performed DFT calculations using both Hartree-Fock (HF) and Hartree-Fock-Bogilbov (HFB) approaches, corresponding to configurations of the nucleus constructed using single-nucleon (HF), or nucleon-hole pair excitations (HFB) as basis states, with HFB calculations introducing pairing correlations. The electromagnetic moments of semi-magic ± 1 nucleon systems, such as the indium isotopes, are well suited to study DFT time-symmetry-breaking (time-odd) contribu-

tions to the mean field^{30,31}, which vary with the time-reversal operator. These fields are predominantly generated by the two-body spin-spin interaction terms, and up until now were poorly constrained within DFT theory. However, they are of particular interest as our understanding of time-reversal-violating mean fields is critical to tackle open problems of modern physics, e.g. the search for new physics^{32–34} and dark matter searches³⁵. Our experimental results presented an excellent opportunity to perform and test these developments. To investigate the relative importance of time-odd fields and pairing correlations, DFT calculations were performed by turning “on” and “off” of each effect. [See Methods for more details on the symmetry-restored calculations.](#) Contrary to shell-model calculations based on empirical interactions, where effective single-nucleon charges and g-factors are used to reproduce nuclear moments data, in our DFT and *ab-initio* calculations we use free single-nucleon g-factors and bare nucleon charges. In addition, the Landau parameter g'_0 was able to be constrained in the DFT calculations by the ¹³¹In magnetic moment.

The results of both *ab initio* and DFT types of calculation are shown alongside the experimental Q and μ moments in Figure 1a) and 1b), respectively, for the $9/2^+$ states of ^{105–131}In. The μ moments of the $1/2^-$ states are shown in Figure 3. All experimental data are presented in Tables 1 and 2 of Methods, and compared with literature values that exist for ^{105–127}In.

Results and discussion

In the single-particle picture, a proton hole configuration induces an intrinsic polarisation of the whole nucleus, as indicated schematically in Figure 1 (bottom left). A gradual decrease in the quadrupole moments of the $9/2^+$ states was previously observed up to $N = 78$. Our measurements reveal a notably larger decrease at $N = 82$, indicating a significant decrease in polarisation (Figure 1a), as the value expected for the single proton hole configuration^{36,37} is reached.

The VS-IMSRG calculations reproduce the experimental trends, i.e local variations in neutron number, a dip around $N = 64$, and a gradual decrease towards $N = 82$. However, the magnitude of the Q is underestimated. The reproduction of the magnitude of the quadrupole moments is a known challenge for *ab initio* nuclear theory, as the Q moments are a highly collective emergent property of the nucleus which can require the inclusion of extensive many-body correlations³⁸.

Conversely, our DFT calculations are able to closely reproduce the overall magnitude of the Q moments. As $N = 82$ is approached, the agreement with the calculations without pairing (HF), shows that describing individual neutron orbitals becomes important. However, due to effects induced by occupying individual neutron orbitals, an inaccurate staggering with neutron number is also produced, compared to HFB. As shown in Figure 1a), time-odd contributions have a negligible effect on Q moments.

In contrast to the Q moments, the μ moments of the $9/2^+$ states were known to exhibit little variation⁷, which is contin-

ued up to ^{129}In in our observation. However, we observe an abrupt increase at ^{131}In , see Figure 1b), and the atomic spectra in Figure 4. The extreme single-particle magnetic moment of a proton in the $\pi g_{9/2}$ orbit, the so-called ‘Schmidt’ limit³⁹, is also indicated in Figure 1b).

Our results now reveal a value much closer to the extreme single-particle limit at $N = 82$, in agreement with decay spectroscopy and moment measurements in the region^{40–44}. The μ value at ^{131}In reaches 93% of the Schmidt free particle value, and 81–84% for $N < 82$. However, the sudden change in the μ values from $N < 82$ to $N = 82$ indicates a departure from the single-particle picture for the former, which contradicts the generally accepted conclusions from the earlier magnetic moment^{6,7} and transfer reaction⁴⁵ studies. Extension of transfer reaction studies to isotopes in the $N = 82$ region would therefore be highly valuable, and should be possible in the near future with the development of experiments at radioactive beam facilities, such as SOLARIS at FRIB⁴⁶, and ISS at ISOLDE⁴⁷.

Owing to the standard angular-momentum coupling rules, the nuclear quadrupole moments can be measured only in states with $I > 1/2$. The $I = 9/2^+$ states of the indium isotopes thus provide access simultaneously to the nuclear quadrupole moments and nuclear magnetic dipole moments. The indium isotopes are therefore a unique case to study the effect of single-proton hole coupling to the nuclear core on both observables. This is in contrast to thallium ($Z = 81$), the heavier experimentally accessible proton-hole isotopic chain, which has a $1/2^+$ configuration.

We find that the inclusion of time-odd fields proves to be the primary missing component of the mean field for an accurate description of magnetic moments. We observe that without the time-odd mean fields, the DFT value obtained for ^{131}In is close to the single-particle value. However, the addition of a spin-spin interaction term corresponding to the isovector Landau parameter⁴⁸ of $g'_0 = 0.82$ generates time-odd mean fields that result in a perfect agreement between the DFT and the experimental ^{131}In magnetic moment. The same value of g'_0 was kept in lighter indium isotopes, so the reproduced abrupt change of μ between ^{129}In and ^{131}In was obtained without further adjustment of parameters. The value of g'_0 adopted here is about two standard deviations smaller than that of $g'_0 = 1.7(4)$ obtained in the recent global analysis of Ref.⁴⁹, which for other Skyrme functionals gave values of $g'_0 = 1.3(4)$ and $1.0(4)$. It is also somewhat smaller than that of $g'_0 = 1.2$ recommended in Refs.^{48,50} based on the analysis of the Gamow-Teller β decays and better aligned with that of $g'_0 = 0.6(1)$ extracted in Ref.⁵¹ from the Gamow-Teller response functions, see also discussions in Refs.^{52,53}. Further studies are needed to see if the time-odd mean fields can be consistently modelled by the single spin-spin term or a more involved description is needed.

In order to further investigate the abrupt change that was observed at $N = 82$ within the DFT framework, we traced back the properties of μ in indium directly to those of




Figure 2. A breakdown of the contributions of the polarised core states to the final calculated μ values of the $I^\pi = 9/2^+$ state from the HF DFT calculations, compared to experiment. See text for details.

their polarised cores. To this end, we performed symmetry-restoration HF calculations by filling in the deformed odd-proton holes obtained in the self-consistent DFT states in indium. In this way, we gained access to the corresponding 0^+ , 1^+ , 2^+ , 3^+ , 4^+ , ... states of the polarised tin core. The effect of their contributions to μ in indium is summarised in Figure 2. The coupling to the 0^+ state of the tin core gives results identical to the Schmidt limit. At this point, it is essential to note that without the time-odd mean fields the core is not spin-polarised and therefore its states conserve signature symmetry and odd-angular-momentum components vanish. The effect of the time-odd mean fields on magnetic moments thus proceeds through the coupling to the 1^+ , 3^+ , ..., 9^+ states of the polarised core. As shown in Figure 2, the experimental value in ^{131}In is reproduced by including the 1^+ core state only. Contributions of higher odd- I states were negligible. For the $N < 82$ isotopes, contributions of the 1^+ core states stay approximately constant. Coupling of the unpaired proton to the even- I polarised core states, which are non-existing at the $N = 82$ shell closure, but appear when the neutron shell opens, result in the strong decrease of the magnetic moment when moving away from $N = 82$. Even- I contributions beyond $I^\pi = 6^+$ were negligible. When combined with the 1^+ core polarisation contribution, the experimental values are closely reproduced. As the even- I core excitations would also be blocked at the neutron shell closure of $N = 50$, a measurement of the magnetic moment at $N = 50$ would thus probe its magicity.

Our results highlight a major difference between the HFB and HF treatments, which is related to the effect of pair breaking. As the neutron pairing vanishes at the shell closure, the

calculated magnetic moment of ^{131}In is the same for HF and HFB calculations. For isotopes with $N < 82$, the HF calculations, which include the polarisation induced by the broken neutron pairs, reproduce the steep decrease of the magnetic moment, as soon as two holes occur. On the contrary, the HFB results, which do not entail broken pairs, do not reproduce that decrease. Further theoretical developments, such as a multi-reference version of HFB to include the mixing of deformed two-quasiparticle excited states, are needed in the HFB framework.

Previously, the decreasing μ values of the $1/2^-$ isomeric states with increasing neutron number (see Figure 3) were not explained by nuclear theory. Our new experimental results for $^{129,131}\text{In}$ show a sudden increase, which crosses the extreme single-particle value at $N = 82$. The results are compared with literature values of $^{115-127}\text{In}$ in Table 1. We find that the DFT calculations without pairing correlations also provide a better description than those with pairing correlations (HFB). The introduction of time-odd fields improves the agreement with experiment. However, the pronounced increase at $N = 82$ is not reproduced within this framework, with predictions close to the single-particle limit. Yet, the experimentally observed variation, and crossing of the single-particle limit at $N = 82$ suggests the $1/2^-$ isomers are sensitive to collectivity. Our VS-IMSRG calculations provide some insight here. Overall, these calculations provide a good description of the previously unexplained trend observed for the magnetic moments of the $1/2^-$ states. Analysis of the VS-IMSRG calculations indicates that the $1/2^-$ state contains a significant component resulting from the coupling between the $9/2^+$ single-particle configuration and the 5^- (anti-aligned neutrons in the $h_{11/2}$ and $s_{1/2}$ orbitals) state of the core. This therefore suggests that further refinement of the present DFT calculations would be needed to explicitly include other core configurations e.g. negative-parity states, to accurately describe the $1/2^-$ states.

Summary and Outlook

The indium isotopes have been considered a textbook example for the dominance of single-particle properties in heavy nuclei. Here, we show that their ground-state electromagnetic properties significantly differ at $N = 82$ compared to $N < 82$, despite the single unpaired proton ($\pi g_{9/2}$ proton orbit) dominating the behaviour of this complex many-body system.

Our new experimental results for $^{129,131}\text{In}$ show that upon reaching the neutron-rich magic number $N = 82$, abrupt changes of their electromagnetic properties are observed, and only at the neutron magic number the single-particle structure is almost fully recovered. This challenges our previous understanding of these isotopes, which were assumed to have a single-particle description by invoking the use of empirical effective operators.

We presented two complementary nuclear models to investigate how these seemingly simple structures emerge from the complex interactions among nucleons. Both DFT and *ab initio* calculations provide a good description of the experimental




Figure 3. Nuclear magnetic moments for the $1/2^-$ isomeric states of $^{113-131}\text{In}$ isotopes. Results are compared with *ab initio* and Density Functional Theory calculations. The horizontal dotted line indicates the single-particle value (Schmidt limit). See text for more details

trends. Within the DFT framework, the intricate isotopic dependence of the indium magnetic and electric ground-state moments turns out to require correct treatment of spin and shape polarisation exerted by the proton-hole on the $Z = 50$ magic core. The inclusion of time-odd mean fields was shown to be essential to reproduce the experimental findings.

Indium nuclei can coexist in long-lived excited states with $I^\pi = 1/2^-$. The structure of these states was suggested to be dominated by a single unpaired proton in the $\pi p_{1/2}$ orbit. However, in contrast to the simple trends observed for the ground states, the magnetic moments of the $1/2^-$ states exhibit a large variation with neutron number, previously considered a puzzle in the region⁷. These features and the new experimental observations are well described by our *ab initio* calculations, for which these isotopes in the region of $Z = 50$ are the frontier of such *ab initio* calculations of nuclear moments^{9,10}. Our work indicates a very mixed configuration for the isomeric states, which is not yet captured with the DFT calculations, where negative parity core polarisation needs to be included.

Although our *ab initio* calculations provide a good description of the experimental trends, they fall short in reproducing the magnitude of the $9/2^+$ electromagnetic moments. This may be attributed to the lack of inclusion of many-body currents, which are known to be essential to describe the electromagnetic properties of light nuclei⁵⁴. Future theoretical developments would be needed to include many-body currents and clarify the role that they play in heavy nuclei. On the other hand, a theoretical description of the nuclear quadrupole mo-

ments requires the extensive inclusion of neglected many-body correlations. Thus, our new experimental results at $N = 82$ provide critical data for future developments of both *ab initio* and DFT theory.

Our experimental results for ^{131}In allowed an investigation of the strength of the time-odd mean fields, which until now have been poorly constrained in DFT theories. Such time-odd channels are essential for a correct description of numerous nuclear properties, such as double beta-decay rates³⁴, permanent electric dipole moment measurements^{32,33}, and dark matter searches³⁵. This provides strong motivation to extend experiments to other isotopes possessing single-hole (particle) with respect to suggested nuclear closed shells at extreme proton-to-neutron ratios. In addition, further theoretical developments are needed to investigate the role of meson-exchange currents and additional non-nucleonic degrees of freedom **in nuclear moments**.

References

- Schmidt, T. The electric quadrupole moment of the nucleus. *Nature* **138**, 404, DOI: [10.1038/138404a0](https://doi.org/10.1038/138404a0) (1936).
- Jones, K. L. *et al.* The magic nature of 132 Sn explored through the single-particle states of 133 Sn. *Nature* **465**, 454–457, DOI: [10.1038/nature09048](https://doi.org/10.1038/nature09048) (2010).
- Taniuchi, R. *et al.* 78 Ni revealed as a doubly magic stronghold against nuclear deformation. *Nature* **569**, 53–58, DOI: [10.1038/s41586-019-1155-x](https://doi.org/10.1038/s41586-019-1155-x) (2019). [1912.05978](https://doi.org/10.1038/s41586-019-1155-x).
- Togashi, T., Tsunoda, Y., Otsuka, T., Shimizu, N. & Honma, M. Novel Shape Evolution in Sn Isotopes from Magic Numbers 50 to 82. *Phys. Rev. Lett.* **121**, 062501, DOI: [10.1103/PhysRevLett.121.062501](https://doi.org/10.1103/PhysRevLett.121.062501) (2018). [1806.10432](https://doi.org/10.1103/PhysRevLett.121.062501).
- Jenkins, D. G. Recent advances in nuclear physics through on-line isotope separation. *Nat. Phys.* **10**, 909–913, DOI: [10.1038/nphys3165](https://doi.org/10.1038/nphys3165) (2014).
- Heyde, K. *The Nuclear Shell Model*. Springer Series in Nuclear and Particle Physics (Springer Berlin Heidelberg, Berlin, Heidelberg, 1990).
- Eberz, J. *et al.* Spins, moments and mean square charge radii of 104–127In determined by laser spectroscopy. *Nucl. Phys. A* **464**, 9–28, DOI: [10.1016/0375-9474\(87\)90419-2](https://doi.org/10.1016/0375-9474(87)90419-2) (1987).
- Rosiak, D. *et al.* Enhanced Quadrupole and Octupole Strength in Doubly Magic Sn 132. *Phys. Rev. Lett.* **121**, 252501, DOI: [10.1103/PhysRevLett.121.252501](https://doi.org/10.1103/PhysRevLett.121.252501) (2018).
- Gysbers, P. *et al.* Discrepancy between experimental and theoretical β -decay rates resolved from first principles. *Nat. Phys.* **15**, 428–431, DOI: [10.1038/s41567-019-0450-7](https://doi.org/10.1038/s41567-019-0450-7) (2019).
- Morris, T. D. *et al.* Structure of the Lightest Tin Isotopes. *Phys. Rev. Lett.* **120**, 152503, DOI: [10.1103/PhysRevLett.120.152503](https://doi.org/10.1103/PhysRevLett.120.152503) (2018).
- Schunck, N. (ed.) *Energy Density Functional Methods for Atomic Nuclei*. 2053–2563 (IOP Publishing, 2019).
- Rodríguez, L. V. *et al.* Doubly-magic character of ^{132}Sn studied via electromagnetic moments of ^{133}Sn . *Phys. Rev. C* **102**, 051301, DOI: [10.1103/PhysRevC.102.051301](https://doi.org/10.1103/PhysRevC.102.051301) (2020).
- Hinke, C. B. *et al.* Superaligned Gamow-Teller decay of the doubly magic nucleus ^{100}Sn . *Nature* **486**, 341–345, DOI: [10.1038/nature11116](https://doi.org/10.1038/nature11116) (2012).
- Manea, V. *et al.* First Glimpse of the $N=82$ Shell Closure below $Z=50$ from Masses of Neutron-Rich Cadmium Isotopes and Isomers. *Phys. review letters* **124**, 092502, DOI: [10.1103/PhysRevLett.124.092502](https://doi.org/10.1103/PhysRevLett.124.092502) (2020).
- Neyens, G. Nuclear magnetic and quadrupole moments for nuclear structure research on exotic nuclei. *Reports on Prog. Phys.* **66**, 633–689, DOI: [10.1088/0034-4885/66/4/205](https://doi.org/10.1088/0034-4885/66/4/205) (2003).
- Stroberg, S. R., Bogner, S. K., Hergert, H. & Holt, J. D. Nonempirical Interactions for the Nuclear Shell Model: An Update. *Annu. Rev. Nucl. Part. Sci.* **69**, DOI: [10.1146/annurev-nucl-101917-021120](https://doi.org/10.1146/annurev-nucl-101917-021120) (2019). [1902.06154](https://doi.org/10.1146/annurev-nucl-101917-021120).
- Hebeler, K., Bogner, S. K., Furnstahl, R. J., Nogga, A. & Schwenk, A. Improved nuclear matter calculations from chiral low-momentum interactions. *Phys. Rev. C* **83**, DOI: [10.1103/PhysRevC.83.031301](https://doi.org/10.1103/PhysRevC.83.031301) (2011).
- de Groote, R. *et al.* Precision measurement of the magnetic octupole moment in ^{45}Sc as a test for state-of-the-art atomic- and nuclear-structure theory. *Phys. Lett. B* **827**, 136930, DOI: <https://doi.org/10.1016/j.physletb.2022.136930> (2022).
- Bennaceur, K., Dobaczewski, J., Haverinen, T. & Kortelainen, M. Properties of spherical and deformed nuclei using regularized pseudopotentials in nuclear DFT. *J. Phys. G: Nucl. Part. Phys.* **47**, 23, DOI: [10.1088/1361-6471/ab9493](https://doi.org/10.1088/1361-6471/ab9493) (2020).
- Dobaczewski, J. *et al.* Mean-field description of ground-state properties of drip-line nuclei: Pairing and continuum effects. *Phys. Rev. C - Nucl. Phys.* **53**, 2809–2840, DOI: [10.1103/PhysRevC.53.2809](https://doi.org/10.1103/PhysRevC.53.2809) (1996).
- Dobaczewski, J., Flocard, H. & Treiner, J. Hartree-Fock-Bogolyubov description of nuclei near the neutron-drip line. *Nucl. Physics, Sect. A* **422**, 103–139, DOI: [10.1016/0375-9474\(84\)90433-0](https://doi.org/10.1016/0375-9474(84)90433-0) (1984).
- Sheikh, J. A., Dobaczewski, J., Ring, P., Robledo, L. M. & Yannouleas, C. Symmetry restoration in mean-field approaches. *J. Phys. G: Nucl. Part. Phys.* **48**, 123001, DOI: [10.1088/1361-6471/ac288a](https://doi.org/10.1088/1361-6471/ac288a) (2021).
- Cocolios, T. E. *et al.* High-resolution laser spectroscopy with the Collinear Resonance Ionisation Spectroscopy (CRIS) experiment at CERN-ISOLDE. *Nucl. Instruments Methods Phys. Res. Sect. B: Beam Interactions*

- with *Mater. Atoms* **376**, 284–287, DOI: [10.1016/j.nimb.2015.11.024](https://doi.org/10.1016/j.nimb.2015.11.024) (2016).
24. Sahoo, B. K. *et al.* Analytic response relativistic coupled-cluster theory: the first application to indium isotope shifts. *New J. Phys.* **22**, 012001, DOI: [10.1088/1367-2630/ab66dd](https://doi.org/10.1088/1367-2630/ab66dd) (2020).
 25. Vernon, A. *et al.* Simulation of the relative atomic populations of elements $1 \leq Z \leq 89$ following charge exchange tested with collinear resonance ionization spectroscopy of indium. *Spectrochimica Acta Part B: At. Spectrosc.* **153**, 61–83, DOI: [10.1016/J.SAB.2019.02.001](https://doi.org/10.1016/J.SAB.2019.02.001) (2019).
 26. Garcia Ruiz, R. F. *et al.* High-Precision Multiphoton Ionization of Accelerated Laser-Ablated Species. *Phys. Rev. X* **8**, 041005, DOI: [10.1103/PhysRevX.8.041005](https://doi.org/10.1103/PhysRevX.8.041005) (2018).
 27. Epelbaum, E., Hammer, H.-W. & Meißner, U.-G. Modern theory of nuclear forces. *Rev. Mod. Phys.* **81**, 1773–1825, DOI: [10.1103/RevModPhys.81.1773](https://doi.org/10.1103/RevModPhys.81.1773) (2009). [0811.1338](https://arxiv.org/abs/0811.1338).
 28. Machleidt, R. & Entem, D. R. Chiral effective field theory and nuclear forces. *Phys. Reports* **503**, 1–75, DOI: [10.1016/j.physrep.2011.02.001](https://doi.org/10.1016/j.physrep.2011.02.001) (2011). [1105.2919](https://arxiv.org/abs/1105.2919).
 29. Jiang, W. G. *et al.* Accurate bulk properties of nuclei from $A = 2$ to ∞ from potentials with Δ isobars. *Phys. Rev. C* **102**, 054301, DOI: [10.1103/PhysRevC.102.054301](https://doi.org/10.1103/PhysRevC.102.054301) (2020).
 30. Engel, Y., Brink, D., Goeke, K., Krieger, S. & Vautherin, D. Time-dependent Hartree-Fock theory with Skyrme's interaction. *Nucl. Phys. A* **249**, 215 – 238, DOI: [https://doi.org/10.1016/0375-9474\(75\)90184-0](https://doi.org/10.1016/0375-9474(75)90184-0) (1975).
 31. Perlińska, E., Rohoziński, S. G., Dobaczewski, J. & Nazarewicz, W. Local density approximation for proton-neutron pairing correlations: Formalism. *Phys. Rev. C* **69**, 014316, DOI: [10.1103/PhysRevC.69.014316](https://doi.org/10.1103/PhysRevC.69.014316) (2004).
 32. Dobaczewski, J., Engel, J., Kortelainen, M. & Becker, P. Correlating Schiff moments in the light actinides with octupole moments. *Phys. Rev. Lett.* **121**, 232501, DOI: [10.1103/PhysRevLett.121.232501](https://doi.org/10.1103/PhysRevLett.121.232501) (2018).
 33. Chupp, T. E., Fierlinger, P., Ramsey-Musolf, M. J. & Singh, J. T. Electric dipole moments of atoms, molecules, nuclei, and particles. *Rev. Mod. Phys.* **91**, 015001, DOI: [10.1103/RevModPhys.91.015001](https://doi.org/10.1103/RevModPhys.91.015001) (2019).
 34. Dolinski, M. J., Poon, A. W. & Rodejohann, W. Neutrinoless double-beta decay: Status and prospects. *Annu. Rev. Nucl. Part. Sci.* **69**, 219–251, DOI: [10.1146/annurev-nucl-101918-023407](https://doi.org/10.1146/annurev-nucl-101918-023407) (2019).
 35. Engel, J., Pittel, S. & Vogel, P. Nuclear Physics of Dark Matter Detection. *Int. J. Mod. Phys. E* **1**, 1–37, DOI: [10.1142/S0218301392000023](https://doi.org/10.1142/S0218301392000023) (1992).
 36. Co', G., Donno, V. D., Anguiano, M., Bernard, R. N. & Lallena, A. M. Electric quadrupole and magnetic dipole moments of odd nuclei near the magic ones in a self-consistent approach. *Phys. Rev. C* **92**, 024314, DOI: [10.1103/PhysRevC.92.024314](https://doi.org/10.1103/PhysRevC.92.024314) (2015).
 37. Krane, K. S. & Halliday, D. *Introductory nuclear physics* (Wiley, 1988).
 38. Henderson, J. *et al.* Testing microscopically derived descriptions of nuclear collectivity: Coulomb excitation of ^{22}Mg . *Phys. Lett. B* **782**, 468–473, DOI: [10.1016/j.physletb.2018.05.064](https://doi.org/10.1016/j.physletb.2018.05.064) (2018). [1709.03948](https://arxiv.org/abs/1709.03948).
 39. Schmidt, T. Über die magnetischen Momente der Atomkerne. *Zeitschrift für Physik* **106**, 358–361, DOI: [10.1007/BF01338744](https://doi.org/10.1007/BF01338744) (1937).
 40. Taprogge, J. *et al.* $1p\ 3/2$ Proton-Hole State in ^{132}Sn and the Shell Structure Along $N = 82$. *Phys. Rev. Lett.* **112**, 132501, DOI: [10.1103/PhysRevLett.112.132501](https://doi.org/10.1103/PhysRevLett.112.132501) (2014).
 41. Fogelberg, B. *et al.* Decays of ^{131}In , ^{131}Sn , and the position of the $h\ 11/2$ neutron hole state. *Phys. Rev. C* **70**, 034312, DOI: [10.1103/PhysRevC.70.034312](https://doi.org/10.1103/PhysRevC.70.034312) (2004).
 42. Fogelberg, B. & Blomqvist, J. Single-hole and three-quasiparticle levels in ^{131}Sn observed in the decay of $^{131}\text{g}, m1, m2\text{In}$. *Nucl. Phys. A* **429**, 205–217, DOI: [10.1016/0375-9474\(84\)90205-7](https://doi.org/10.1016/0375-9474(84)90205-7) (1984).
 43. Vaquero, V. *et al.* Fragmentation of Single-Particle Strength around the Doubly Magic Nucleus ^{132}Sn and the Position of the $0f\ 5/2$ Proton-Hole State in ^{131}In . *Phys. Rev. Lett.* **124**, 17, DOI: [10.1103/PhysRevLett.124.022501](https://doi.org/10.1103/PhysRevLett.124.022501) (2020).
 44. Lechner, S. *et al.* Probing the single-particle behavior above ^{132}Sn via electromagnetic moments of $^{133,134}\text{Sb}$ and $n = 82$ isotones. *Phys. Rev. C* **104**, 014302, DOI: [10.1103/PhysRevC.104.014302](https://doi.org/10.1103/PhysRevC.104.014302) (2021).
 45. Weiffenbach, C. V. & Tickle, R. Structure of odd- a indium isotopes determined by the $(d, ^3\text{He})$ reaction. *Phys. Rev. C* **3**, 1668–1678, DOI: [10.1103/PhysRevC.3.1668](https://doi.org/10.1103/PhysRevC.3.1668) (1971).
 46. Kay, B. The SOLARIS spectrometer. In *APS April Meeting Abstracts*, vol. 2019 of *APS Meeting Abstracts*, Q13.002 (2019).
 47. Tang, T. L. *et al.* First exploration of neutron shell structure below lead and beyond $n = 126$. *Phys. Rev. Lett.* **124**, 062502, DOI: [10.1103/PhysRevLett.124.062502](https://doi.org/10.1103/PhysRevLett.124.062502) (2020).
 48. Bender, M., Dobaczewski, J., Engel, J. & Nazarewicz, W. Gamow-Teller strength and the spin-isospin coupling constants of the Skyrme energy functional. *Phys. Rev. C* **65**, 054322, DOI: [10.1103/PhysRevC.65.054322](https://doi.org/10.1103/PhysRevC.65.054322) (2002).
 49. Sassarini, P. L., Dobaczewski, J., Bonnard, J. & Garcia Ruiz, R. F. *arXiv:2111.04675* (2021).
 50. Borzov, I. N., Tolokonnikov, S. V. & Fayans, S. Spin-dependent effective nucleon-nucleon interaction in nuclei. *Sov. J. Nucl. Phys.* **40**, 732 (1984).

51. Wakasa, T., Ichimura, M. & Sakai, H. Unified analysis of spin isospin responses of nuclei. *Phys. Rev. C* **72**, 067303, DOI: [10.1103/PhysRevC.72.067303](https://doi.org/10.1103/PhysRevC.72.067303) (2005).
52. Roca-Maza, X., Colò, G. & Sagawa, H. New skyrme interaction with improved spin-isospin properties. *Phys. Rev. C* **86**, 031306, DOI: [10.1103/PhysRevC.86.031306](https://doi.org/10.1103/PhysRevC.86.031306) (2012).
53. Davesne, D., Pastore, A. & Navarro, J. Linear response theory with finite-range interactions. *Prog. Part. Nucl. Phys.* **120**, 103870, DOI: <https://doi.org/10.1016/j.pnpnp.2021.103870> (2021).
54. Pastore, S., Pieper, S. C., Schiavilla, R. & Wiringa, R. B. Quantum Monte Carlo calculations of electromagnetic moments and transitions in $A \leq 9$ nuclei with meson-exchange currents derived from chiral effective field theory. *Phys. Rev. C* **87**, 035503, DOI: [10.1103/PhysRevC.87.035503](https://doi.org/10.1103/PhysRevC.87.035503) (2013).

Methods

The collinear resonance ionisation setup

The indium isotopes were produced at CERN's on-line isotope separator facility, ISOLDE, by impinging 1.4-GeV protons onto the proton-to-neutron converter of a thick uranium-carbide target¹. The converter suppressed nearby caesium mass contamination². The indium isotopes diffused through the target material and their ionisation was significantly enhanced by the use of the resonance laser ion source, RILIS³. The produced indium ions were then accelerated to 40 keV and mass separated using the ISOLDE high-resolution mass separator before being cooled and bunched using a gas-filled linear Paul trap (ISCOOL)^{4,5}. After a trapping time of up to 10 ms, ion bunches with a temporal width of 2 μ s, were then re-accelerated to 40034(1) eV and deflected into the CRIS beamline^{6,7}. The indium ions were then neutralised with a sodium-filled charge-exchange cell with an efficiency of up to 60% and predicted relative atomic populations of 57% and 37% respectively for the $5p\ ^2P_{3/2}$ metastable state and $5p\ ^2P_{1/2}$ ground state⁸. The remaining ion fraction was removed by electrostatic deflectors, and the neutralized atom bunch was collinearly overlapped with two pulsed lasers, one for excitation and another for non-resonant ionisation. The atoms were resonantly excited using two different UV transitions in separate measurements. The first using 246.8-nm laser light for the $5p\ ^2P_{3/2} \rightarrow 9s\ ^2S_{1/2}$ atomic transition. The second using 246.0-nm laser light for the $5p\ ^2P_{1/2} \rightarrow 8s\ ^2S_{1/2}$ atomic transition, which is more sensitive to nuclear magnetic moments, μ , but alone does not give the nuclear electric quadrupole moment, Q . The resonant laser light was produced by frequency tripling the light from an injection-locked Ti:Sapphire laser system⁹. This laser was seeded using a narrow-band SolsTiS continuous-wave Ti:Sapphire laser, and pumped using a Lee Laser LDP-100MQ Nd:YAG laser, producing pulsed narrow-band 740(738)-nm laser light at 1 kHz repetition rate. This light was then frequency tripled to 246.8(246.0)-nm light by the use of two non-linear crystals¹⁰. About 3 μ J of laser light was used to saturate both transitions. The excited atoms were then ionized by a final non-resonant 1064-nm step, provided by a Litron LPY 601 50-100 PIV Nd:YAG laser at 100 Hz repetition rate. The frequency of the resonant first step was scanned and the resulting ions were deflected onto a detector, producing the hyperfine structure containing spectra from

which the hyperfine parameters were obtained. The wavelengths were measured using a HighFinesse WSU-2 wavemeter, which was drift stabilized by simultaneous measurement of a Toptica DLC DL PRO 780 diode laser locked to the $5s^2S_{1/2} \rightarrow 5p\ ^2P_{3/2}\ F=2 \rightarrow 3$ transition of ^{87}Rb using a saturated absorption spectroscopy unit.

Evaluation of nuclear magnetic and quadrupole moments

The μ values were determined using a reference NMR value of $\mu_{ref} = +5.5408(2)\ \mu_N$ for $^{115}\text{In}^{11}$, and the relation

$$\mu = \mu_{ref} \frac{IA}{I_{ref}A_{ref}} (1 + \Delta), \quad (1)$$

where the differential hyperfine anomaly, Δ , is negligible for these atomic states of indium¹². Here A_{ref} are our experimentally determined values for the $5p^2\ P_{3/2}$, $9s^2\ S_{1/2}$, $5p^2\ P_{1/2}$ and $8s^2\ S_{1/2}$ states of stable ^{115}In , and A are those of short-lived isotopes. The final μ values presented in Table 1 are a weighted average of the μ values from each atomic state, which were self-consistent within 2σ .

The spectroscopic nuclear electric quadrupole moments, Q , were extracted using the relation

$$B_{hf} = eQV_{ZZ}, \quad (2)$$

where a value of $B_{hf}(^2P_{3/2})/Q = 576(4)\ \text{MHz/b}$ was used, obtained from relativistic coupled-cluster atomic calculations¹³. Here V_{ZZ} is the electric field gradient produced by the electrons at the nucleus.

The ratio of A_{hf} factors is depends dominantly on atomic structure and therefore remains constant when with the correct nuclear spin assignment is used for fitting of spectra resulting from hyperfine structure¹⁴. Although reduced χ^2 fitting of the spectra (as shown in Figure 4) resulted in minima when we use $I = 9/2$; the spin assignment from ratio of upper and lower atomic state A_{hf} factors (of the $5p\ ^2P_{3/2} \rightarrow 9s\ ^2S_{1/2}$ transition) could not rule out nuclear spins $7/2$, $11/2$, outside of 1σ uncertainty. The spin of the $I = 9/2$ states are tentatively assigned by experiment and strongly supported by nuclear theory. The spin assignments of the $I = 1/2$ states were confirmed unambiguously from the number of peaks in their spectra.

VS-IMSRG calculations

The VS-IMSRG calculations start from the underlying NN and 3N interactions expressed in the harmonic-oscillator basis. We then construct an effective Hamiltonian designed for a particular valence space where the exact diagonalization is feasible²⁰. The effective Hamiltonian was decoupled from the full A -body Hamiltonian through the application of an approximate unitary transformation derived from the Magnus expansion method²¹. Using the same transformation, the effective $M1$ and $E2$ valence-space operators were then decoupled consistently with the Hamiltonian. During the calculation, all of operators arising in nested commutator expansions were truncated at the two-body level, the IMSRG(2) approximation.

We begin in a spherical harmonic-oscillator basis with frequency of $\hbar\omega = 16\ \text{MeV}$ and $e = 2n + l \leq e_{\max} = 14$ for $1.8/2.0\ (\text{EM})^{22,23}$ (and 12 for N^2LOGO) with an additional cut of $e_1 + e_2 + e_3 \leq E_{3\max} = 16$ imposed due to storage limitations of 3N matrix elements. The valence space was taken as the proton $\{1p_{1/2}, 1p_{3/2}, 0f_{5/2}, 0g_{9/2}\}$ and neutron $\{2s_{1/2}, 1d_{3/2}, 1d_{5/2}, 0g_{7/2}, 0h_{11/2}\}$ single-particle orbits above ^{78}Ni core.

To investigate the effect of valence space on the calculations, the calculations were also performed with the proton $\{1p_{1/2}, 1p_{3/2}, 0f_{5/2}, 0g_{9/2}\}$ and neutron $\{2s_{1/2}, 1d_{3/2}, 0h_{11/2}, 1f_{7/2}\}$ valence




Figure 4. (color online) **Example hyperfine spectra of the ^{129}In and ^{131}In isotopes,** a) measured using the 246.8-nm ($5p\ ^2P_{3/2} \rightarrow 9s\ ^2S_{1/2}$) transition. The $9/2^+$ ground and $1/2^-$ isomer states are indicated. b) Example spectra of the $1/2^-$ isomer structure measured with the 246.0-nm ($5p\ ^2P_{1/2} \rightarrow 8s\ ^2S_{1/2}$) transition.

Table 1. The magnetic hyperfine structure parameters, A_{hf} , measured in this work for the odd-mass $^{113}\text{--}^{131}\text{In}$ isotopes and corresponding extracted magnetic dipole moment values.

A	I	A_{hf} (MHz)				μ^{\ddagger} (μ_N)	$\mu^{\text{Lit.}}$ (μ_N)
		$5p\ ^2P_{3/2}$	$9s\ ^2S_{1/2}$	$5p\ ^2P_{1/2}$	$8s\ ^2S_{1/2}$		
105	$9/2^+$						+5.675(5) ¹⁵
107	$9/2^+$						+5.585(8) ¹⁵
109	$9/2^+$						+5.538(4) ¹⁵
111	$9/2^+$						+5.503(7) ¹⁵
113	$9/2^+$	+241.8(8)	+130(1)	+2276.0(8)	+242.7(8)	+5.5264(19)	+5.5289(2) ¹⁶
113	$1/2^-$	-87(10)	-38(20)	-774(50)	-90(50)	-0.21(1)	-0.21074(2) ¹⁷
115	$9/2^+$	+239.9(5)	+130.3(8)	+2281.9(8)	+243.3(6)	+5.5408(2) [‡]	+5.5408(2) ¹¹
115	$1/2^-$	-96(3)	-48(9)	-903(20)	-66(10)	-0.2405(38)	-0.24398(5) ¹⁸
117	$9/2^+$	+241(4)	+130(5)	+2277(2)	+244(2)	+5.5286(43)	+5.519(4) ¹⁵
117	$1/2^-$	-106(4)	-49(10)	-1028(10)	-113(8)	-0.276(27)	-0.25174(3) ¹⁵
119	$9/2^+$	+240(4)	+130(5)			+5.499(62)	+5.515(1) ¹⁵
119	$1/2^-$	-132(7)	-70(10)			-0.342(12)	-0.319(5) ¹⁵
121	$9/2^+$	+243(4)	133(5)			+5.575(62)	+5.502(5) ¹⁵
121	$1/2^-$	-140(2)	-85(6)			-0.3600(41)	-0.355(4) ¹⁵
123	$9/2^+$	+238(4)	+129(5)			+5.442(61)	+5.491(7) ¹⁵
123	$1/2^{(-)}$	-160(2)	-80(5)			-0.4047(54)	-0.400(4) ¹⁵
125	$9/2^+$	+240.3(6)	+129.9(8)			+5.496(24)	+5.502(9) ¹⁵
125	$1/2^{(-)}$	-176(7)	-90(10)			-0.450(17)	-0.433(4) ¹⁵
127	$9/2^+$	+241.8(7)	+130(1)	+2278.3(6)	+243.8(4)	+5.5321(14)	+5.522(8) ¹⁵
127	$1/2^{(-)}$	-171(3)	-91(10)	-1613(9)	-174(8)	-0.4355(24)	
129	($9/2^+$)	+243.3(8)	+132(1)	+2304.9(9)	+244.8(7)	+5.5961(23)	
129	$1/2^{(-)}$	-156(3)	-80(4)	-1434(2)	-162(10)	-0.3871(6)	
131	($9/2^+$)	+275.9(6)	+149.3(7)			+6.312(14)	
131	$1/2^{(-)}$	-20(7)	-11(4)	-188(20)	-20(2)	-0.051(3)	

[‡] These μ values were determined using a reference NMR value of $\mu = +5.5408(2)\ \mu_N$ ¹¹.

Table 2. The B_{hf} hyperfine structure parameter (from the $^2P_{3/2}$ state) values, measured in this work for the odd-mass $^{113-131}\text{In}$ isotopes and the extracted electric quadrupole moment values.

A	I	B_{hf} (MHz)	Q^\dagger (eb)	$Q^{Lit.}$ (eb)
105	$9/2^+$			$+0.79(5)^{15,19}$
107	$9/2^+$			$+0.77(5)^{15,19}$
109	$9/2^+$			$+0.80(3)^{15,19}$
111	$9/2^+$			$+0.76(2)^{15,19}$
113	$9/2^+$	$+441(15)$	$+0.767(27)$	$+0.759(8)^{15,19}$
115	$9/2^+$	$+454.2(65)$	$+0.789(13)$	$+0.770(8)^{15,19}$
117	$9/2^+$	$+465(13)$	$+0.807(23)$	$+0.788(10)^{15,19}$
119	$9/2^+$	$+462(13)$	$+0.802(23)$	$+0.812(7)^{15,19}$
121	$9/2^+$	$+462(13)$	$+0.803(23)$	$+0.774(10)^{15,19}$
123	$9/2^+$	$+424(13)$	$+0.736(23)$	$+0.720(9)^{15,19}$
125	$9/2^+$	$+382.3(55)$	$+0.664(11)$	$+0.68(3)^{15,19}$
127	$9/2^+$	$+338(16)$	$+0.588(29)$	$+0.56(3)^{15,19}$
129	$(9/2^+)$	$+280.4(73)$	$+0.487(13)$	
131	$(9/2^+)$	$+177.3(57)$	$+0.310(10)$	

† Extracted using a value of $B_{hf}(^2P_{3/2})/Q = +576(4) \text{ MHz/b}^{13}$.

space above the ^{92}Ni core. Values of the $9/2^+$ states varied by $<2\%$ and $1/2^-$ by $<25\%$, due to the increased sensitivity of the $1/2$ states to collectivity. The trends over neutron number in the calculations obtained were unaffected. The final valence-space diagonalization and evaluation of electromagnetic moments were performed with the KSHELL code²⁴. The effect of fully accounting for translation invariance in the $M1$ operator was found to be less than 1%, largely due to the lack of radial dependence of the $M1$ operator.

DFT calculations

We performed the DFT calculations using code HFODD^{25,26} version (3.01m) and a standard Skyrme density functional UNEDF1²⁷. The deformed DFT single-particle wave functions were expanded on the spherical harmonic-oscillator basis up to $N_0 = 16$ quanta. An A -dependent harmonic-oscillator frequency $\hbar\omega$ was fixed according to the prescription defined in Ref.²⁸. For the state $9/2^+$, the proton configuration (with no pairing correlations) was fixed by occupying 49 lowest axially deformed prolate orbitals, that is, the Nilsson orbital $[404]_{7/2}^+$ with the angular-momentum projection on the symmetry axis $\Omega = -9/2$ was left empty. Similarly, for the state $1/2^-$, the hole was kept in the oblate Nilsson orbital $[301]_{1/2}^-$ with $\Omega = -1/2$. It was essential to pick the proton-hole configurations with angular momenta properly aligned along the axial-symmetry axis. Indeed, as discussed in Refs.^{29,30}, the time-odd mean fields and core spin polarisation depend on the relative orientation of the intrinsic angular momentum and shape.

For both proton configurations, the open-shell neutron configurations were constructed either with (HFB results) or without (HF results) pairing correlations included. For all deformed mean-field states $|\Omega\rangle$ obtained in this way, which were eigenstates of the z -component of the angular momentum, $\hat{L}_z|\Omega\rangle = \Omega|\Omega\rangle$, we performed the total angular-momentum restoration³¹ by evaluating the integral

$$|IM\rangle = \frac{1}{2}(2I+1) \int_{\beta=0}^{\pi} d\beta d_{M\Omega}^I(\beta) \exp(-i\beta\hat{L}_y) |\Omega\rangle,$$

where $d_{M\Omega}^I(\beta)$ are the Wigner functions³² and M is the projection of the angular momentum on the laboratory z -axis. This allowed us to

determine the standard spectroscopic magnetic dipole μ and electric quadrupole Q moments³³ as

$$\mu = \sqrt{\frac{4\pi}{3}} \langle II | \hat{M}_{10} | II \rangle \quad \text{and} \quad Q = \sqrt{\frac{16\pi}{5}} \langle II | \hat{Q}_{20} | II \rangle,$$

where \hat{M} and \hat{Q} are the corresponding $M1$ and $E2$ electromagnetic operators, respectively. In addition, for paired configurations the particle-number symmetry was restored in a similar way³¹. Neither effective charges nor effective g -factors were used in the *ab initio* or DFT calculations.

References

- Köster, U. Intense radioactive-ion beams produced with the ISOL method. *The Eur. Phys. J. A* **15**, 255–263, DOI: [10.1140/epja/i2001-10264-2](https://doi.org/10.1140/epja/i2001-10264-2) (2002). [arXiv:1011.1669v3](https://arxiv.org/abs/1011.1669v3).
- Dillmann, I. *et al.* Selective laser ionization of $N \geq 82$ indium isotopes: The new r -process nuclide ^{135}In . *The Eur. Phys. J. A* **13**, 281–284, DOI: [10.1007/s10050-002-8756-7](https://doi.org/10.1007/s10050-002-8756-7) (2002).
- Rothe, S., Marsh, B. A., Mattolat, C., Fedosseev, V. N. & Wendt, K. A complementary laser system for ISOLDE RILIS. *J. Physics: Conf. Ser.* **312**, 052020, DOI: [10.1088/1742-6596/312/5/052020](https://doi.org/10.1088/1742-6596/312/5/052020) (2011).
- Mané, E. *et al.* An ion cooler-buncher for high-sensitivity collinear laser spectroscopy at ISOLDE. *Eur. Phys. J. A* **42**, 503–507, DOI: [10.1140/epja/i2009-10828-0](https://doi.org/10.1140/epja/i2009-10828-0) (2009).
- Fränberg, H. *et al.* Off-line commissioning of the ISOLDE cooler. *Nucl. Inst. Methods Phys. Res. B* **266**, 4502–4504, DOI: [10.1016/j.nimb.2008.05.097](https://doi.org/10.1016/j.nimb.2008.05.097) (2008).
- Vernon, A. R. *et al.* Optimising the collinear resonance ionisation spectroscopy (CRIS) experiment at CERN-ISOLDE. *Nucl. Instruments Methods Phys. Res. Sect. B: Beam Interactions with Mater. Atoms* **463**, 384–389, DOI: <https://doi.org/10.1016/j.nimb.2019.04.049> (2020).
- Cocolios, T. *et al.* High-resolution laser spectroscopy with the Collinear Resonance Ionisation Spectroscopy (CRIS) experiment at CERN-ISOLDE. *Nucl. Instruments Methods Phys. Res.*

- Sect. B: Beam Interactions with Mater. Atoms* **376**, 284–287, DOI: [10.1016/j.nimb.2015.11.024](https://doi.org/10.1016/j.nimb.2015.11.024) (2016).
8. Vernon, A. *et al.* Simulation of the relative atomic populations of elements $1 \leq Z \leq 89$ following charge exchange tested with collinear resonance ionization spectroscopy of indium. *Spectrochimica Acta Part B: At. Spectrosc.* **153**, 61–83, DOI: <https://doi.org/10.1016/j.sab.2019.02.001> (2019).
 9. Sonnenschein, V. *et al.* Characterization of a pulsed injection-locked Ti:sapphire laser and its application to high resolution resonance ionization spectroscopy of copper. *Laser Phys.* **27**, 085701, DOI: [10.1088/1555-6611/aa7834](https://doi.org/10.1088/1555-6611/aa7834) (2017).
 10. Bass, M., Franken, P. A., Hill, A. E., Peters, C. W. & Weinreich, G. Optical Mixing. *Phys. Rev. Lett.* **8**, 18–18, DOI: [10.1103/PhysRevLett.8.18](https://doi.org/10.1103/PhysRevLett.8.18) (1962).
 11. Flynn, C. P. & Seymour, E. F. W. Knight Shift of the Nuclear Magnetic Resonance in Liquid Indium. *Proc. Phys. Soc.* **76**, 301–303, DOI: [10.1088/0370-1328/76/2/415](https://doi.org/10.1088/0370-1328/76/2/415) (1960).
 12. Persson, J. R. Table of hyperfine anomaly in atomic systems. *At. Data Nucl. Data Tables* **99**, 62–68, DOI: [10.1016/j.adt.2012.04.002](https://doi.org/10.1016/j.adt.2012.04.002) (2013). [arXiv:1110.5991v1](https://arxiv.org/abs/1110.5991v1).
 13. Garcia Ruiz, R. F. *et al.* High-Precision Multiphoton Ionization of Accelerated Laser-Ablated Species. *Phys. Rev. X* **8**, 041005, DOI: [10.1103/PhysRevX.8.041005](https://doi.org/10.1103/PhysRevX.8.041005) (2018).
 14. Cheal, B. *et al.* Nuclear spins and moments of Ga isotopes reveal sudden structural changes between N=40 and N=50. *Phys. Rev. Lett.* **104**, 252502, DOI: [10.1103/PhysRevLett.104.252502](https://doi.org/10.1103/PhysRevLett.104.252502) (2010).
 15. Eberz, J. *et al.* Spins, moments and mean square charge radii of $^{104-127}\text{In}$ determined by laser spectroscopy. *Nucl. Phys. A* **464**, 9–28, DOI: [10.1016/0375-9474\(87\)90419-2](https://doi.org/10.1016/0375-9474(87)90419-2) (1987).
 16. Rice, M. & Pound, R. V. Ratio of the Magnetic Moments of $\text{In } 115$ and $\text{In } 113$. *Phys. Rev.* **106**, 953–953, DOI: [10.1103/PhysRev.106.953](https://doi.org/10.1103/PhysRev.106.953) (1957).
 17. Childs, W. J. & Goodman, L. S. Nuclear Spin and Hyperfine Interaction of $\text{In } 113\text{m}$. *Phys. Rev.* **118**, 1578–1581, DOI: [10.1103/PhysRev.118.1578](https://doi.org/10.1103/PhysRev.118.1578) (1960).
 18. Cameron, J. A., King, H. J., Eastwood, H. K. & Summers-Gill, R. G. THE MAGNETIC MOMENT OF INDIUM- ^{115}m . *Can. J. Phys.* **40**, 931–942, DOI: [10.1139/p62-099](https://doi.org/10.1139/p62-099) (1962).
 19. Stone, N. J. Nuclear Magnetic Dipole and Electric Quadrupole Moments: Their Measurement and Tabulation as Accessible Data. *J. Phys. Chem. Ref. Data* **44**, 031215, DOI: [10.1063/1.4917489](https://doi.org/10.1063/1.4917489) (2015).
 20. Stroberg, S. R., Bogner, S. K., Hergert, H. & Holt, J. D. Nonempirical Interactions for the Nuclear Shell Model: An Update. *Annu. Rev. Nucl. Part. Sci.* **69**, DOI: [10.1146/annurev-nucl-101917-021120](https://doi.org/10.1146/annurev-nucl-101917-021120) (2019). [1902.06154](https://arxiv.org/abs/1902.06154).
 21. Morris, T. D., Parzuchowski, N. M. & Bogner, S. K. Magnus expansion and in-medium similarity renormalization group. *Phys. Rev. C* **92**, 34331, DOI: [10.1103/PhysRevC.92.034331](https://doi.org/10.1103/PhysRevC.92.034331) (2015). [1507.06725](https://arxiv.org/abs/1507.06725).
 22. Hebeler, K., Bogner, S. K., Furnstahl, R. J., Nogga, A. & Schwenk, A. Improved nuclear matter calculations from chiral low-momentum interactions. *Phys. Rev. C* **83**, DOI: [10.1103/PhysRevC.83.031301](https://doi.org/10.1103/PhysRevC.83.031301) (2011).
 23. Simonis, J., Stroberg, S. R., Hebeler, K., Holt, J. D. & Schwenk, A. Saturation with chiral interactions and consequences for finite nuclei. *Phys. Rev. C* **96**, DOI: [10.1103/PhysRevC.96.014303](https://doi.org/10.1103/PhysRevC.96.014303) (2017). [1704.02915](https://arxiv.org/abs/1704.02915).
 24. Shimizu, N., Mizusaki, T., Utsuno, Y. & Tsunoda, Y. Thick-restart block Lanczos method for large-scale shell-model calculations. *Comput. Phys. Commun.* DOI: [10.1016/j.cpc.2019.06.011](https://doi.org/10.1016/j.cpc.2019.06.011) (2019).
 25. Schunck, N. *et al.* Solution of the Skyrme-Hartree-Fock-Bogolyubov equations in the Cartesian deformed harmonic-oscillator basis. (VIII) HFODD (v2.73y): A new version of the program. *Comp. Phys. Commun.* **216**, 145 – 174, DOI: <https://doi.org/10.1016/j.cpc.2017.03.007> (2017).
 26. Dobaczewski, J. *et al.* Solution of universal nonrelativistic nuclear DFT equations in the Cartesian deformed harmonic-oscillator basis. (IX) HFODD (v3.06h): a new version of the program. *J. Phys. G: Nucl. Part. Phys.* **48**, 102001 (2021).
 27. Kortelainen, M. *et al.* Nuclear energy density optimization: Large deformations. *Phys. Rev. C* **85**, 024304, DOI: [10.1103/PhysRevC.85.024304](https://doi.org/10.1103/PhysRevC.85.024304) (2012).
 28. Dobaczewski, J. & Dudek, J. Solution of the Skyrme-Hartree-Fock equations in the Cartesian deformed harmonic oscillator basis II. The program HFODD. *Comput. Phys. Comm.* **102**, 183 – 209, DOI: [https://doi.org/10.1016/S0010-4655\(97\)00005-2](https://doi.org/10.1016/S0010-4655(97)00005-2) (1997).
 29. Schunck, N. *et al.* One-quasiparticle states in the nuclear energy density functional theory. *Phys. Rev. C* **81**, 024316, DOI: [10.1103/PhysRevC.81.024316](https://doi.org/10.1103/PhysRevC.81.024316) (2010).
 30. Satuła, W., Bączyk, P., Dobaczewski, J. & Konieczka, M. No-core configuration-interaction model for the isospin- and angular-momentum-projected states. *Phys. Rev. C* **94**, 024306, DOI: [10.1103/PhysRevC.94.024306](https://doi.org/10.1103/PhysRevC.94.024306) (2016).
 31. Sheikh, J. A., Dobaczewski, J., Ring, P., Robledo, L. M. & Yannouleas, C. Symmetry restoration in mean-field approaches. *J. Phys. G: Nucl. Part. Phys.* **48**, 123001, DOI: [10.1088/1361-6471/ac288a](https://doi.org/10.1088/1361-6471/ac288a) (2021).
 32. Varshalovich, D., Moskalev, A. & Khersonskii, V. *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).
 33. Ring, P. & Schuck, P. *The nuclear many-body problem* (Springer-Verlag, Berlin, 1980).

Acknowledgements

This work was supported by ERC Consolidator Grant No.648381 (FNPMLS); STFC grants ST/L005794/1, ST/L005786/1, ST/P004423/1, ST/M006433/1, ST/P003885/1 and Ernest Rutherford Grant No. ST/L002868/1; the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under grant DE-SC0021176; GOA 15/010 from KU Leuven, BriX Research Program No. P7/12; the FWO-Vlaanderen (Belgium); the European Unions Grant Agreement 654002 (ENSAR2); National Key R&D Program of China (Contract No: 2018YFA0404403); the National Natural Science Foundation of China (No:11875073); the Polish National Science Centre under Contract No. 2018/31/B/ST2/02220. TRIUMF receives funding via a contribution through the National Research Council of Canada. The theoretical work was further supported by NSERC and the US Department of Energy (DOE) under contract DE-FG02-97ER41014. The

VS-IMSRG Computations were performed with an allocation of computing resources on Cedar at WestGrid and Compute Canada, and on the Oak Cluster at TRIUMF managed by the University of British Columbia department of Advanced Research Computing (ARC). We would also like to thank the ISOLDE technical group for their support and assistance and the University of Jyväskylä for the use of the injection-locked cavity. We acknowledge the CSC-IT Center for Science Ltd., Finland, for the allocation of computational resources. This project was partly undertaken on the Viking Cluster, which is a high performance compute facility provided by the University of York. We are grateful for computational support from the University of York High Performance Computing service, Viking and the Research Computing team.

Author contributions statement

A.R.V. prepared the manuscript with input from all authors, especially R.F.G.R., J.Bo., J.D., J.D.H., T.M., G.N., K.T.F., T.E.C., R.P.G. and S.R.S. . R.F.G., J.B., C.L.B., M.L.B., T.E.C., K.T.F., W.G., R.P.G., A.K., K.M.L., G.N., S.G.W., A.R.V. and X.F.Y. proposed the experiment(s), A.R.V., C.L.B., M.L.B., T.E.C., K.T.F., G.J.F., G.G., W.G., R.P.G., R.H., A.K., D.L., K.M.L., R.F.G., S.G.W., X.F.Y. and D.Y. conducted the experiment(s), A.R.V., C.L.B., R.F.G. and J.H. analysed the results, J.Bo. and J.D. performed theoretical (DFT) nuclear calculations, J.D.H., T.M., S.R.S. performed theoretical (VS-IMSRG) nuclear calculations. All authors reviewed the manuscript.

Competing interests statement

The authors declare no competing interests.

Data availability statement

Examples of spectra source data for the previously unmeasured ^{129,131}In isotopes, most relevant to this work, are included in this article. The full datasets generated during and/or analysed during the current study are available in the Zenodo repository, <https://doi.org/10.5281/zenodo.6406949>. The code used to analyse the data is also included in the repository.

The data files related to the DFT calculations are available at https://webfiles.york.ac.uk/HFODD/Projects/Magnetic_and_electric_moments_in_Indium/

Code availability statement

The code used to analyse the data is included in the Zenodo repository, <https://doi.org/10.5281/zenodo.6406949>.

The code used to perform the DFT calculations is available at <https://webfiles.york.ac.uk/HFODD/Projects/hf301m/>

Additional information

Correspondence and requests for materials should be addressed to A.R. Vernon or R.F. Garcia Ruiz

Reprints and permissions information is available at <http://www.nature.com/reprints>