

This is a repository copy of *Real-time analysis of endogenous Wnt signalling in 3D mesenchymal stromal cells*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/104840/>

Version: Accepted Version

Article:

Saleh, Fatima, Carstairs, Alice Elizabeth, Etheridge, Sharon Leah et al. (1 more author) (2016) Real-time analysis of endogenous Wnt signalling in 3D mesenchymal stromal cells. Stem Cells International. 7132529.

<https://doi.org/10.1155/2016/7132529>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Real-time analysis of endogenous Wnt signalling in 3D mesenchymal stromal cells

Fatima Saleh^{1,2}, Alice Carstairs¹, S. Leah Etheridge¹, Paul Genever¹

¹Department of Biology, University of York, York, YO10 5DD, UK

²Faculty of Health Sciences, Beirut Arab University, Beirut, P.O. Box 115020, Lebanon

F. Saleh Email: f.saleh@bau.edu.lb

A. Carstairs Email: ac713@york.ac.uk

S.L. Etheridge Email: leah.etheridge@york.ac.uk

P. Genever Email: paul.genever@york.ac.uk

Corresponding Author:

Prof. Paul Genever

Department of Biology,

University of York, York,

YO10 5DD, UK.

Email: paul.genever@york.ac.uk

Abstract

Wnt signalling has been implicated in the regulation of stem cell self-renewal and differentiation, however the majority of *in vitro* studies are carried out using monolayer 2D culture techniques. Here, we used mesenchymal stromal cell (MSC) EGFP-reporter lines responsive to Wnt pathway activation in a 3D spheroid culture system to mimic better the *in vivo* environment. Endogenous Wnt signalling was then investigated under basal conditions and when MSCs were induced to undergo osteogenic and adipogenic differentiation. Interestingly, endogenous Wnt signalling was only active during 3D differentiation whereas 2D cultures showed no EGFP expression throughout an extended differentiation time-course. Furthermore, exogenous Wnt signalling in 3D adipogenic conditions inhibited differentiation compared to unstimulated controls. In addition, suppressing Wnt signalling by Dkk-1 restored and facilitated adipogenic differentiation in MSC spheroids. Our findings indicate that endogenous Wnt signalling is active and can be tracked in 3D MSC cultures where it may act as a molecular switch in adipogenesis. The identification of the signalling pathways that regulate MSCs in a 3D *in vivo*-like environment will advance our understanding of the molecular mechanisms that control MSC fate.

Introduction

Mesenchymal stromal cells (MSCs) are a heterogeneous cell population first isolated from the bone marrow, with other sources of related cell types also present in other tissues, for example in dental pulp and adipose tissue [1–3]. The MSC population contains stem cells able to differentiate into osteoblasts, adipocytes and chondrocytes both *in vitro* and *in vivo* [4], hence the term “mesenchymal stem cell” is often used, although referring to non-clonal undefined cell types. Despite this limitation, MSCs are widely viewed to have a range of therapeutic regenerative applications, particularly in a musculoskeletal setting [5]. By increasing our understanding of the role of signalling pathways in orchestrating the fate of MSCs, we will generate a fundamental basis upon which we can build, to manipulate these cells for novel treatment strategies in a wide range of diseases.

One of the regulatory pathways associated with controlling MSC proliferation, differentiation, apoptosis and migration is the Wnt signalling pathway [6, 7]. Canonical Wnt signalling (Figure 1A) is initiated by Wnt ligands binding to membrane receptors Frizzled (Fzd) and co-receptor low-density lipoprotein receptor-related protein (LRP5/6), activating Dishevelled (Dvl). Dvl then inhibits phosphorylation of β -catenin by glycogen kinase (GSK)-3 β , leading to β -catenin stabilisation and accumulation in the cytosol before its translocation to the nucleus. Nuclear β -catenin, interacts with members of the lymphoid enhancer factor (LEF) and T-cell factor (TCF) transcription factor family, activating target gene expression [6, 7]. At the receptor level, the Wnt-Fzd receptor interaction may be inhibited by proteins such as Dickkopf (Dkk) [8, 9]. Numerous *in vivo* studies have implicated Wnt signalling as being vital in osteoblast development with both disrupting and activating mutations of LRP5 impacting on bone mass [10–12]. It is also evident that all the components for Wnt signalling activation exist in MSCs and can be stimulated [6]. Yet, discrepancies exist between *in vivo* data

using mouse models and differentiation studies performed *in vitro* where in multiple instances, the stimulation of Wnt signalling has been shown to inhibit differentiation [13–17].

To address these issues, we employed a three-dimensional (3D) culture model with real-time fluorescent reporters to investigate canonical Wnt signalling in MSCs, including murine MSCs and a highly characterised immortalised human clonal MSC line, under control and differentiated conditions. Although conventional culture methods are important experimental techniques, cells grown in 2D are not exposed to the same environmental conditions as cells in 3D tissues, which include locally acting signalling gradients [18]. In fact, a study has previously shown that Wnt signalling, both canonical and non-canonical, was enhanced in MSC spheroids (in this case formed using a biomaterial derived from chitin) leading to altered differentiation capacities based on activation of the Wnt pathway. This was not apparent in 2D MSC monolayer culture [19]. Unlike many other 3D culture methods, we used a 3D spheroid system that does not rely on attachment to any surface, such as a scaffold; rather, the cells form self-aggregating spheroids in a viscous medium [20]. Using this 3D culture system in combination with Wnt reporter cell lines, we examined endogenous and exogenous Wnt signalling under undifferentiated and differentiated conditions, providing evidence that Wnt signalling in MSCs is context -dependent.

Methods

Cell Culture in 2D and 3D

The Y201 hTERT-immortalised human clonal MSC line, which we have generated and characterised previously [21], and a mouse MSC line, C3H10T1/2 were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% foetal bovine serum (FBS) and maintained at 37°C in a 5% CO₂/95% air atmosphere. The Y201 cells were used as a model human MSC line, whilst the C3H10T1/2 cells

were used as a model murine MSC line to enable observation to be made in MSC-type cells from different, commonly used species involving Wnt-based analyses. Wnt3a (Biotechne) was supplemented where indicated at varying concentrations as a positive control for Wnt pathway activation. The Wnt inhibitor IWR-1 was obtained from Sigma Aldrich.

For 3D culture of both cell types, 30,000 MSCs were seeded into U-bottomed non-adherent 96-well plates in media containing 0.25% (w/v) methylcellulose, complete media changes were performed every 2-3 days throughout the time-course.

Establishing a Wnt-responsive C3H10T1/2 cell line using a TCF reporter construct with EGFP

To construct the reporter plasmid, the human cytomegalovirus (CMV) promoter from expression vector pEGFP-N1 (Clontech) was removed by AseI and HindIII cleavage, and replaced with a fragment cut from the SuperTopFlash plasmid (M50) (gift from R. Moon, Addgene plasmid # 12456) [22] using KpnI and HindIII. Briefly, pEGFP-N1 and M50 plasmids were linearised with the restriction enzymes AseI and KpnI, respectively, and blunted by DNA Polymerase I Large (Klenow) Fragment (Promega). Linearised plasmids were then purified with a QIAquick PCR Purification Kit (Qiagen) according to the manufacturer's instructions before the plasmids were both cut with HindIII to give rise to a pEGFP-N1 (minus CMV) backbone vector and a TCF binding sites/TK promoter (insert). Both the backbone vector and insert were run on a 1% agarose gel and then gel extracted using the QIAquick Gel Extraction Kit (Qiagen) according to manufacturer's instructions. Prior to ligation with T4 DNA ligase (Promega), the pEGFP-N1 backbone was dephosphorylated using Shrimp Alkaline Phosphatase (SAP) (Promega). The ligated plasmid was then transformed into Hi-Coli-5A chemically competent cells (Advantagen) according to the manufacturer's instructions, plated out on LB/Kanamycin plates and incubated overnight at 37°C. DNA was isolated using the GenElute™ HP Plasmid Purification Kit (Sigma), and the integrity of the SuperTop-EGFP construct checked by double digestion with XhoI and HpaI.

Confluent C3H10T1/2 cells were transfected using Lipofectamine 2000 according to manufacturer's instructions (Invitrogen), and the plasmid linearised prior to transfection using Apal. A positive control with pEGFP-N1 and a mock transfection control were also performed. To select for stable transfectants, 2-3 days after transfection, cells were exposed to 1mg/ml Geneticin (Cabochem). Geneticin-containing medium was changed twice a week and after 2 weeks, cells were cultured in C3H10T1/2 medium containing a reduced concentration of Geneticin (500µg/ml) to maintain stable transfectants. To detect for stable transfection, cells were treated with 5µM BIO, an activator of Wnt signalling, or DMSO as a vehicle control.

Establishing a Wnt pathway responsive Y201 hTERT MSC cell line using a Signal Lenti TCF/LEF Reporter

Y201 hTERT MSCs were transduced using the Signal Lenti TCF/LEF kit (Qiagen) according to manufacturer's instructions. Following transduction, cells were cultured under puromycin selection (10µg/mL) and serially diluted across a 96 well plate (following Corning protocol) to isolate individual clonal Y201-Wnt hTERT MSC lines. These were evaluated for GFP activity in response to Wnt3a activation by flow cytometry and responsive lines were selected.

Wnt signalling in the Y201-Wnt hTERT MSC lines was analysed with a Zeiss Lightsheet Z.1. To allow visualisation of spheroids and cell tracking, hTERT-MSCs were labelled with cell tracker red (Life Technologies) prior to spheroid formation as per manufacturer's instructions. Briefly, the required numbers of cells were aliquoted and centrifuged to remove FBS. Cells were labelled with cell tracker and any excess stain neutralised and removed. Samples were then embedded in a 1% agarose/PBS gel rod using a capillary and tight fitting plunger. After the agarose polymerised the gel was extruded from the capillary into the surrounding chamber media (PBS) for imaging.

Assessment of adipogenesis and osteogenesis

Osteogenic and adipogenic differentiation of Y201 hTERT MSCs and C3H10T1/2 cells were initiated 24 hours after spheroid formation using differentiation cocktails as previously described, namely supplementation with 5mM β -glycerophosphate, 50 μ g/mL L-ascorbic acid phosphate and 10nM dexamethasone or 0.5mM isobutyl-methylxanthine, 1 μ g/ml insulin, 100 μ M indomethacin and 1 μ M dexamethasone respectively [6, 21, 23]. At specific time points, spheroids were embedded in OCT (Tissue Tek) and frozen in liquid nitrogen before sectioning using a Leica CM3050S cryostat (Leica Microsystems).

Adipogenesis was assessed by lipid droplet accumulation using Oil Red O staining as previously described briefly [6, 21, 23], cells were fixed in 4% w/v paraformaldehyde solution before incubation in 60% isopropanol. Lipid droplets were stained in 0.3% Oil Red O (Sigma Aldrich) and washed before imaging using a brightfield microscope.

Osteogenic differentiation was analysed by staining deposited calcium using 40mM Alizarin red solution (pH4.2) after fixing in 4% w/v paraformaldehyde. Plates were allowed to dry fully before imaging.

Immunocytochemistry

To determine and confirm EGFP localisation, undifferentiated, osteogenic and adipogenic spheroids were sectioned before immunostaining with an anti-EGFP, rabbit Alexa Fluor® 488 conjugated antibody (Invitrogen) at 1:500 dilution and counterstained with DAPI. Samples were mounted in Vectashield (Vector laboratories) and examined using LSM 710 confocal microscope (Zeiss).

Analysis of Wnt signalling during adipogenesis

C3H10T1/2 Wnt reporter spheroids were treated with 50% Wnt3a-conditioned medium (CM) or 50% control L-cell-CM and Wnt antagonist Dkk-1 (100ng/ml) (R&D Systems) under adipogenic conditions for 7 days before EGFP expression was detected by confocal microscopy and adipogenic potential

evaluated (Multiphoton Zeiss LSM 510). Wnt3a-CM and L-cell-CM was processed from L-Wnt3a cells, a Wnt3a-overexpressing murine cell line, and the control non-transfected L-cells, respectively, as described previously [6].

Results


Reporter MSCs respond specifically to Wnt pathway activation in a dose-dependent manner

We selected the Y201 hTERT MSCs for generating the Wnt reporter based on their equal propensity for the tri-lineage differentiation, typical of an MSC [21]. These cells were transduced using the Signal Lenti TCF/LEF Reporter (EGFP) Kit where EGFP is expressed under the control of a minimal (m)CMV promoter and repeats of the TCF/LEF transcriptional response element (TRE). Activation of the TCF/LEF TRE, typically seen in Wnt signalling, will therefore result in EGFP expression (Figure S1A). After transduction, 7 individual clonal lines were isolated by serial dilution and expanded, these clones were tested for their ability to respond to Wnt3a stimulation by flow cytometry (Figure S1B) and analysed by fluorescent microscopy (Figure S1C). Figure 1B demonstrates the responsiveness of the clonal line (A5) selected for further study, minimal expression can be detected when cells are unstimulated, but upon exposure to Wnt3a, a dose-dependent increase in fluorescence was observed. EGFP fluorescence was then reduced when cells were stimulated with Wnt3a in the presence of IWR-1, a Wnt signalling inhibitor. The ability to respond to Wnt3a was also detectable by flow cytometry, which is more sensitive to low levels of fluorescence. Here, fluorescence was plotted on two green channels to identify cellular autofluorescence, which will be equal in both channels so resulting in events plotted in a straight line. Where EGFP expression is present due to Wnt3a stimulation, events deflect away from the autofluorescence mid-line and is captured in a gate. Analyses in this way typically showed that unstimulated cells had a fluorescent


population of less than 5% (Figure 1C), after cells have been gated to remove dead cells and debris. Upon stimulation with 600ng/mL Wnt3a, 80% of Y201-Wnt hTERT MSCs were fluorescent (Figure 1C). Therefore clone A5 of the Y201 MSC-EGFP reporter cells were used throughout for analyses of Wnt effects in a human MSC model.

Figure 1

A



B



C




FIGURE 1: Stimulation using Wnt3a results in observable fluorescence in **human Y201 MSC-EGFP** reporter lines. (A) Overview of activated canonical Wnt signalling pathway. (B) Fluorescence microscopy of **human Y201 MSC-EGFP reporter cells** treated with varying concentrations of Wnt3a with or without a Wnt pathway inhibitor, IWR-1. Scale bar = 200µm. (C) Flow cytometry scatter plot histograms of Wnt3a treated **Y201 Wnt EGFP reporters** versus untreated, after removal of dead cells and debris by gating. Shift in population away from autofluorescence indicates positive EGFP expression (lower right quadrant).

Wnt signalling is not activated in Y201-Wnt hTERT MSCs during osteogenesis or adipogenesis in 2D culture

Wnt signalling activity is typically implicated in regulating stem cell proliferation and differentiation, particularly in MSCs and skeletal lineages. We therefore sought to determine the status of endogenous Wnt signalling during both osteogenesis and adipogenesis, as an inverse relationship is thought to exist in their lineage specification. Interestingly, we were unable to detect Wnt signalling activation in either differentiation route (Figure 2A and 2C). Over a 20 day time-course of osteogenic and adipogenic induction, no EGFP-positive cells could be detected by flow cytometry at any stage and were even significantly below the low EGFP levels in basal undifferentiated conditions at the same time points. Histological staining demonstrated that the Y201-Wnt hTERT MSCs were successfully differentiating and depositing calcium under osteogenic conditions (Figure 2B) or producing lipid droplets under adipogenic conditions (Figure 2D). It would therefore appear that differentiation of MSCs using typical 2D culture systems does not require activation of endogenous canonical Wnt signalling and the lack of Wnt activation during osteogenesis *in vitro* does not replicate *in vivo* observations.

Figure 2




FIGURE 2: Endogenous Wnt signalling is not activated during 2D MSC differentiation. (A) Percentage of positively expressing **human Y201 MSC-EGFP reporter cells** during osteogenic differentiation as measured by flow cytometry, $p < 0.0001$ versus basal control **with representative flow cytometry plot**. (B) Alizarin red staining for calcium deposition during osteogenic differentiation. Scale bar = 200 μ m. (C) Percentage of positively expressing **human Y201 MSC-EGFP cells** during adipogenic differentiation as measured by flow cytometry, **** $p < 0.0001$ versus basal control **with representative flow cytometry plot**. (D) Oil red O staining for lipid droplet formation during adipogenic differentiation. Scale bar = 200 μ m.

Endogenous Wnt signalling pathway is activated in 3D MSC cultures

2D culture systems do not accurately represent the complexity of an *in vivo* system, therefore to determine if a 3D culture system would better replicate *in vivo* Wnt pathway activation we tested the Y201-Wnt hTERT MSCs in a self-aggregating spheroid culture system. Results showed that after 1 day of spheroid formation there was low level, sporadic EGFP expression indicative of endogenous Wnt signalling in individual cells which increased by day 3 (Figure 3A). Y201-Wnt hTERT MSCs spheroids were able to respond robustly to Wnt3a stimulation which resulted in the majority of cells in the spheroid expressing high levels of EGFP. Therefore this 3D culture system is suitable for examining Wnt pathway activation.

To enable species-specific comparison with previous *in vivo* studies using mouse models, we used a murine MSC line, CH310T1/2, which also has tri-lineage differentiation potential. These cells behave similarly to the human Y201-Wnt hTERT MSC cell line in that they are responsive to Wnt in a dose-dependent manner and no Wnt activation could be detected in 2D differentiation studies (results not shown). In basal conditions, sporadic EGFP expression was detected in CH310T1/2 spheroids, which increased over time in culture, **similar to the pattern observed in human MSC spheroids, but with a much more widespread distribution, appearing throughout the spheroid** (Figure 3B, control, compare to Figure 3A). Active Wnt signalling was also observed in spheroids during early

osteogenesis, however TCF-dependent EGFP expression in adipogenic spheroids remained low (Figure 3B). To determine more precisely the location of EGFP-positive cells in 3D structures and confirm EGFP expression, spheroids were cryosectioned and immunostained with an anti-EGFP antibody. After 7 days of 3D culture, an EGFP-positive pool was predominantly located in the spheroid centre with EGFP-negative cells at the periphery in undifferentiated and osteogenic sections. Again, no EGFP expression was detected in sections of adipogenic spheroids; **it should be noted that adipogenic spheroids increase in size over time due to the accumulation of lipid droplets** (Figure 3C).

Finally, we determined the effects of exogenous Wnt stimulation on 3D cultures. In undifferentiated control samples endogenous EGFP expression was displayed over the whole spheroid, an effect repressed by the Wnt antagonist, Dkk-1, and retrieved by treating with Wnt3a-CM (**Figure S2**). However, in agreement with previous results, EGFP expression could not be detected during adipogenic differentiation over a time-course of 7 days (Figure 3D). Supplementing the media with Wnt3a-CM induced sporadic EGFP expression in individual cells within the adipogenic spheroid and inhibited adipogenic differentiation as shown by a lack of Oil red O staining. In contrast, EGFP expression was absent following treatment with the Wnt inhibitor, Dkk-1, which promoted lipid accumulation and increased spheroid size, seemingly accelerating adipogenic differentiation compared to L-cell-CM controls (Figure 3D).

Figure 3




FIGURE 3: Wnt pathway activation is observed reproducibly in 3D MSC culture and during osteogenic differentiation but not during adipogenic differentiation. (A) Lightsheet microscopy of cell tracker red stained **human Y201 Wnt-EGFP MSCs** showing sporadic activation of Wnt signalling increasing with time in culture, with Wnt3A-treated controls (right panels). Scale bar = 100Fμm. (B) Confocal microscopy of **murine C3H10T1/2 MSC** 3D cultures showing Wnt signalling is active in basal conditions and during osteogenic differentiation but not adipogenic differentiation. Scale bar = 100μm. (C) Immunofluorescence of sectioned spheroids stained for EGFP demonstrates location of EGFP fluorescence in 3D culture. Scale bar = 100μm. **Adipogenic differentiation causes spheroids to enlarge in size; a lower magnification image of the whole spheroid is included (top right, merge panel).** (D) Adipogenic C3H10T1/2 MSC spheroids show inverse relationship between adipogenesis and Wnt signalling by Oil Red O staining.

Discussion

Previous studies have identified Wnt signalling as a major developmental signalling pathway, particularly critical in stem cell maintenance and differentiation. However, Wnt function and the onset of its activation is not always clear using typical *in vitro* culture methods, which we have sought to address in MSCs by analysing endogenous and exogenous Wnt signalling in a 3D context.

Human tissues are complex three-dimensional multi-cellular structures. Recently, *in vitro* techniques have been developed that allow the growth and manipulation of cells in 3D, including the use of cell aggregates such as those examined in this study. The imaging of these aggregates is not always trivial and often dependent on the laser penetration. Previous studies, including ours, have used a multiphoton confocal microscope granting deeper penetration into a sample with less damage as longer wavelengths can be used to excite fluorophores [24–26]. However, imaging in this capacity may still only grant a limited picture of an entire spheroid. New advances in Lightsheet microscopy have employed the use of suspending spheroids in agarose solution allowing the spheroid to be rotated and images taken from multiple angles [27–28]. The combination of reporter cell lines, 3D spheroid culture and advanced imaging techniques allow for the analysis of signalling at a single cell resolution, and offer greater insight into the mechanisms controlling tissue function.

First, we developed stable reporter cell lines in the human Y201 hTERT MSCs and a commonly used murine cell line, C3H10T1/2, both capable of multi lineage differentiation *in vitro* when given appropriate stimuli [21, 24–25]. These cell lines were responsive to exogenous Wnt stimulation **with almost undetectable expression** of GFP during 2D culture, yet no Wnt signalling activation could be detected during 2D differentiation towards either the osteogenic or adipogenic lineages. These observations suggest strongly that active Wnt signalling is not a biological requirement using these *in vitro* MSC systems, which contradicts most *in vivo* evidence. It was hypothesised that by growing MSCs in 3D cultures, the *in vivo* environment is more closely recapitulated in very simplistic terms,

which could influence Wnt signalling activity in MSCs compared to conventional 2D methods. Under osteogenic conditions, Wnt signalling was active in a manner similar to undifferentiated spheroids. However, under adipogenic conditions, Wnt signalling was suppressed.

Interestingly, in both undifferentiated and differentiated 3D cultures, Wnt signalling activity appeared to be much more prominent in the centre of the spheroid. Here it would be expected that cells would be subjected to different signalling inputs compared to cells at the surface. Some of these inputs will be altered concentration gradients such as nutrients, signalling molecules, oxygen, mechanical forces and cytoskeletal tensions [31]. All these factors play a major role in driving cell-contact mediated signalling including the Wnt pathway. This is particularly important considering the locally-acting nature of Wnt molecules, which has recently been demonstrated using a 3D in vitro culture of intestinal stem cells [32].

To examine these observations further, adipogenic spheroids were treated with medium conditioned from Wnt3a overexpressing murine cells or a Wnt antagonist. Wnt3a activated EGFP expression but only in sporadic, isolated cells under adipogenic conditions, however staining for adipocytes using Oil red O showed that activating Wnt signalling inhibited adipogenesis in 3D cultures. Moreover, suppressing Wnt signalling using Dkk-1 enhanced adipogenic differentiation in C3H10T1/2 spheroids. Taken together, our results show that Wnt signalling is inactive during and inhibitory to adipogenesis, supporting previous observations. For example, it has been found Wnt3a decreases the expression of adipogenic transcription factors peroxisome proliferator-activated receptor gamma (PPAR γ) and fatty acid binding protein 4 (FABP4) in C3H10T1/2 cells, inhibiting adipogenesis [33]. Our data also suggest that Wnt activation during adipogenesis results in a weak TCF-mediated transcriptional response (observed as low EGFP expression), though this may be sufficiently inhibitory to further adipogenic responses. Alternatively, Wnt3a may be acting through non-canonical mechanisms to regulate adipogenesis as previously described [34].

Conclusions

We have generated two EGFP-reporter MSC lines for the real-time study of Wnt signalling. Using these tools we have demonstrated that Wnt signalling is activated in basal and osteogenic conditions only when examined in 3D culture and not in conventional 2D conditions. In contrast, no Wnt-dependent EGFP expression was detected in adipogenic differentiating spheroids and exogenous stimulation of Wnt signalling had a negative effect on adipogenesis. Our study shows the importance of considering MSC signalling mechanisms in a 3D context to avoid potential discrepancies and apparently contradictory observations.

References

- [1] M. F. Pittenger, A. M. Mackay, S. C. Beck, R. K. Jaiswal, R. Douglas, J. D. Mosca, M. A. Moorman, D. W. Simonetti, S. Craig, and D. R. Marshak, 'Multilineage potential of adult human mesenchymal stem cells', *Science*, vol. 284, no. 5411, pp. 143–147, Apr. 1999.
- [2] S. Gronthos, M. Mankani, J. Brahimi, P. G. Robey, and S. Shi, 'Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo', *Proc. Natl. Acad. Sci. U. S. A.*, vol. 97, no. 25, pp. 13625–13630, Dec. 2000.
- [3] P. A. Zuk, M. Zhu, P. Ashjian, D. A. De Ugarte, J. I. Huang, H. Mizuno, Z. C. Alfonso, J. K. Fraser, P. Benhaim, and M. H. Hedrick, 'Human Adipose Tissue Is a Source of Multipotent Stem Cells', *Mol. Biol. Cell*, vol. 13, no. 12, pp. 4279–4295, Dec. 2002.
- [4] M. Dominici, K. Le Blanc, I. Mueller, I. Slaper-Cortenbach, F. Marini, D. Krause, R. Deans, A. Keating, D. Prockop, and E. Horwitz, 'Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement', *Cytotherapy*, vol. 8, no. 4, pp. 315–317, 2006.
- [5] A. Carstairs and P. Genever, 'Stem cell treatment for musculoskeletal disease', *Curr. Opin. Pharmacol.*, vol. 16, pp. 1–6, Jun. 2014.
- [6] S. L. Etheridge, G. J. Spencer, D. J. Heath, and P. G. Genever, 'Expression Profiling and Functional Analysis of Wnt Signaling Mechanisms in Mesenchymal Stem Cells', *STEM CELLS*, vol. 22, no. 5, pp. 849–860, Sep. 2004.

- [7] T. Reya and H. Clevers, 'Wnt signalling in stem cells and cancer', *Nature*, vol. 434, no. 7035, pp. 843–850, Apr. 2005.
- [8] A. Glinka, W. Wu, H. Delius, A. P. Monaghan, C. Blumenstock, and C. Niehrs, 'Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction', *Nature*, vol. 391, no. 6665, pp. 357–362, Jan. 1998.
- [9] P. Fedi, A. Bafico, A. Nieto Soria, W. H. Burgess, T. Miki, D. P. Bottaro, M. H. Kraus, and S. A. Aaronson, 'Isolation and biochemical characterization of the human Dkk-1 homologue, a novel inhibitor of mammalian Wnt signaling', *J. Biol. Chem.*, vol. 274, no. 27, pp. 19465–19472, Jul. 1999.
- [10] Y. Cui, P. J. Niziolek, B. T. MacDonald, C. R. Zylstra, N. Alenina, D. R. Robinson, Z. Zhong, S. Matthes, C. M. Jacobsen, R. A. Conlon, R. Brommage, Q. Liu, F. Msee, D. R. Powell, Q. Yang, B. Zambrowicz, H. Gerrits, J. A. Gossen, X. He, M. Bader, B. O. Williams, M. L. Warman, and A. G. Robling, 'Lrp5 functions in bone to regulate bone mass', *Nat. Med.*, vol. 17, no. 6, pp. 684–691, Jun. 2011.
- [11] P. J. Niziolek, T. Farmer, Y. Cui, C. H. Turner, M. L. Warman, and A. G. Robling, 'High-bone-mass-producing mutations in the Wnt signaling pathway result in distinct skeletal phenotypes', *Bone*, vol. 49, no. 5, pp. 1010–1019, Nov. 2011.
- [12] R. Brommage, J. Liu, G. M. Hansen, L. L. Kirkpatrick, D. G. Potter, A. T. Sands, B. Zambrowicz, D. R. Powell, and P. Vogel, 'High-throughput screening of mouse gene knockouts identifies established and novel skeletal phenotypes', *Bone Res.*, vol. 2, p. 14034, Oct. 2014.
- [13] J. A. Rudnicki and A. M. Brown, 'Inhibition of chondrogenesis by Wnt gene expression in vivo and in vitro', *Dev. Biol.*, vol. 185, no. 1, pp. 104–118, May 1997.
- [14] S. E. Ross, N. Hemati, K. A. Longo, C. N. Bennett, P. C. Lucas, R. L. Erickson, and O. A. MacDougald, 'Inhibition of adipogenesis by Wnt signaling', *Science*, vol. 289, no. 5481, pp. 950–953, Aug. 2000.
- [15] G. M. Boland, G. Perkins, D. J. Hall, and R. S. Tuan, 'Wnt 3a promotes proliferation and suppresses osteogenic differentiation of adult human mesenchymal stem cells', *J. Cell. Biochem.*, vol. 93, no. 6, pp. 1210–1230, Dec. 2004.
- [16] J. De Boer, H. J. Wang, and C. Van Blitterswijk, 'Effects of Wnt signaling on proliferation and differentiation of human mesenchymal stem cells', *Tissue Eng.*, vol. 10, no. 3–4, pp. 393–401, Apr. 2004.
- [17] J. de Boer, R. Siddappa, C. Gaspar, A. van Apeldoorn, R. Fodde, and C. van Blitterswijk, 'Wnt signaling inhibits osteogenic differentiation of human mesenchymal stem cells', *Bone*, vol. 34, no. 5, pp. 818–826, May 2004.
- [18] F. A. Saleh and P. G. Genever, 'Turning round: multipotent stromal cells, a three-dimensional revolution?', *Cytotherapy*, vol. 13, no. 8, pp. 903–912, Sep. 2011.

- [19] S. Hsu and G.-S. Huang, 'Substrate-dependent Wnt signaling in MSC differentiation within biomaterial-derived 3D spheroids', *Biomaterials*, vol. 34, no. 20, pp. 4725–4738, Jul. 2013.
- [20] F. A. Saleh, J. E. Frith, J. A. Lee, and P. G. Genever, 'Three-dimensional in vitro culture techniques for mesenchymal stem cells', *Methods Mol. Biol. Clifton NJ*, vol. 916, pp. 31–45, 2012.
- [21] S. James, J. Fox, F. Afsari, J. Lee, S. Clough, C. Knight, J. Ashmore, P. Ashton, O. Preham, M. Hoogduijn, R. D. A. R. Ponzoni, Y. Hancock, M. Coles, and P. Genever, 'Multiparameter Analysis of Human Bone Marrow Stromal Cells Identifies Distinct Immunomodulatory and Differentiation-Competent Subtypes', *Stem Cell Rep.*, vol. 4, no. 6, pp. 1004–1015, Sep. 2015.
- [22] M. T. Veeman, D. C. Slusarski, A. Kaykas, S. H. Louie, and R. T. Moon, 'Zebrafish prickles, a modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements', *Curr. Biol. CB*, vol. 13, no. 8, pp. 680–685, Apr. 2003.
- [23] M. J. Hoogduijn, E. Gorjup, and P. G. Genever, 'Comparative Characterization of Hair Follicle Dermal Stem Cells and Bone Marrow Mesenchymal Stem Cells', *Stem Cells Dev.*, vol. 15, no. 1, pp. 49–60, Feb. 2006.
- [24] P. C. de Sampaio, D. Auslaender, D. Krubasik, A. V. Failla, J. N. Skepper, G. Murphy, and W. R. English, 'A Heterogeneous In Vitro Three Dimensional Model of Tumour-Stroma Interactions Regulating Sprouting Angiogenesis', *PLOS ONE*, vol. 7, no. 2, p. e30753, Feb. 2012.
- [25] E. E. Hoover and J. A. Squier, 'Advances in multiphoton microscopy technology', *Nat. Photonics*, vol. 7, no. 2, pp. 93–101, Feb. 2013.
- [26] J. Laurent, C. Frongia, M. Cazales, O. Mondesert, B. Ducommun, and V. Lobjois, 'Multicellular tumor spheroid models to explore cell cycle checkpoints in 3D', *BMC Cancer*, vol. 13, p. 73, 2013.
- [27] C. Lorenzo, C. Frongia, R. Jorand, J. Fehrenbach, P. Weiss, A. Maandhui, G. Gay, B. Ducommun, and V. Lobjois, 'Live cell division dynamics monitoring in 3D large spheroid tumor models using light sheet microscopy', *Cell Div.*, vol. 6, p. 22, 2011.
- [28] F. Pampaloni, N. Ansari, and E. H. K. Stelzer, 'High-resolution deep imaging of live cellular spheroids with light-sheet-based fluorescence microscopy', *Cell Tissue Res.*, vol. 352, no. 1, pp. 161–177, Feb. 2013.
- [29] T. Katagiri, A. Yamaguchi, T. Ikeda, S. Yoshiki, J. M. Wozney, V. Rosen, E. A. Wang, H. Tanaka, S. Omura, and T. Suda, 'The non-osteogenic mouse pluripotent cell line, C3H10T1/2, is induced to differentiate into osteoblastic cells by recombinant human bone morphogenetic protein-2', *Biochem. Biophys. Res. Commun.*, vol. 172, no. 1, pp. 295–299, Oct. 1990.
- [30] S. M. Taylor and P. A. Jones, 'Multiple new phenotypes induced in 10T1/2 and 3T3 cells treated with 5-azacytidine', *Cell*, vol. 17, no. 4, pp. 771–779, Aug. 1979.
- [31] L. G. Griffith and M. A. Swartz, 'Capturing complex 3D tissue physiology in vitro', *Nat. Rev. Mol. Cell Biol.*, vol. 7, no. 3, pp. 211–224, Mar. 2006.

- [32] H. F. Farin, I. Jordens, M. H. Mosa, O. Basak, J. Korving, D. V. F. Tauriello, K. de Punder, S. Angers, P. J. Peters, M. M. Maurice, and H. Clevers, 'Visualization of a short-range Wnt gradient in the intestinal stem-cell niche', *Nature*, vol. 530, no. 7590, pp. 340–343, Feb. 2016.
- [33] G. Rawadi, B. Vayssière, F. Dunn, R. Baron, and S. Roman-Roman, 'BMP-2 Controls Alkaline Phosphatase Expression and Osteoblast Mineralization by a Wnt Autocrine Loop', *J. Bone Miner. Res.*, vol. 18, no. 10, pp. 1842–1853, Oct. 2003.
- [34] W. Qiu, L. Chen, and M. Kassem, 'Activation of non-canonical Wnt/JNK pathway by Wnt3a is associated with differentiation fate determination of human bone marrow stromal (mesenchymal) stem cells', *Biochem. Biophys. Res. Commun.*, vol. 413, no. 1, pp. 98–104, Sep. 2011.

Declaration

The authors declare that there is no conflict of interest regarding the publication of this paper.

Acknowledgements

We would like to thank Randall Moon, University of Washington, Seattle, WA for the kind gift of the M50 (8x) SuperTopFlash plasmid. We would also like to thank the Biotechnology and Biological Sciences Research Council, Smith & Nephew, National Centre for the Replacement, Refinement & Reduction of Animals in Research (NC/K001671/1) and the Arthritis Research UK Tissue Engineering Centre (19429) for funding this work.

Supplementary Material

Supplementary Figure 1: Generation and isolation of clonal Wnt responsive human Y201 MSC-EGFP reporter lines.

Supplementary Figure 2: Stimulation of Wnt signalling in murine C3H10T1/2 MSC 3D cultures by exogenous application of conditioned media.