UNIVERSITY of York

This is a repository copy of A Direct Demonstration of Functional Differences between Subdivisions of Human V5/MT+.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/108903/</u>

Version: Accepted Version

Article:

Strong, Samantha L, Silson, Edward H, Gouws, André D et al. (2 more authors) (2017) A Direct Demonstration of Functional Differences between Subdivisions of Human V5/MT+. Cerebral Cortex. pp. 1-10. ISSN 1460-2199

https://doi.org/10.1093/cercor/bhw362

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

A Direct Demonstration of Functional Differences between Subdivisions of Human V5/MT+.

RunningTitle: Functional differences between subdivisions of hV5/MT+

Samantha L. Strong^{a,b}, Edward H. Silson^{b,c}, André D. Gouws^b, Antony B. Morland^{b,d} & Declan J. McKeefry^a

^aSchool of Optometry and Vision Science, University of Bradford, Bradford, West Yorkshire, BD7 1DP, UK.

^bYork Neuroimaging Centre, Department of Psychology, University of York, York, YO10 5DD, UK.

- ^cLaboratory of Brain and Cognition, National Institute of Mental Health, Bethesda, USA
- ^dCentre for Neuroscience, Hull-York Medical School, University of York, York, YO10 5DD, UK.

Corresponding Author: Declan McKeefry Email: d.mckeefry@bradford.ac.uk Address: School of Optometry and Vision Science, University of Bradford, Bradford, West Yorkshire, BD7 1DP, UK. Tel: +44(0)1274 234648.

Abstract

Two subdivisions of human V5/MT+; one located posteriorly (MT/TO-1), the other more anteriorly (MST/TO-2), were identified in human participants using functional magnetic resonance imaging (fMRI) on the basis of their representations of the ipsi- versus contra-lateral visual field. These subdivisions were then targeted for disruption by the application of repetitive transcranial magnetic stimulation (rTMS). rTMS was delivered to cortical areas whilst participants performed direction discrimination tasks involving three different types of moving stimuli defined by the translational, radial or rotational motion of dot patterns. For translational motion, performance was significantly reduced relative to baseline when rTMS was applied to both MT/TO-1 and MST/TO-2. For radial motion there was a differential effect between MT/TO-1 and MST/TO-2, with only disruption of the latter area affecting performance. rTMS failed to reveal a complete dissociation between MT/TO-1 and MST/TO-2 in terms of their contribution to the perception of rotational motion. On the basis of these results MT/TO-1 and MST/TO-2 appear to be functionally distinct subdivisions of hMT/V5+. Whilst both areas appear to be implicated in the processing of translational motion, only the anterior region (MST/TO-2) makes a causal contribution to the perception of radial motion.

Key Words: transcranial magnetic stimulation, fMRI, psychophysics, V5/MT+.

Introduction

Moving visual stimuli elicit neural activity across an extensive network of areas in the human cerebral cortex (Dupont et al., 1994; Watson et al., 1993; Tootell et al., 1995; McKeefry et al., 1997; Smith et al., 1998; Culham et al., 2001). Within this network, human V5/MT+ (hV5/MT+) has come to be regarded as the cortical area most closely associated with the perception of visual motion (Zeki et al., 1991; Watson et al., 1993; Dumoulin et al., 2000). However, as its name implies, hV5/MT+ is not a single area but instead forms a complex, containing multiple areas that have been differentiated on the basis of differences in visual field representation and the receptive field (RF) sizes of constituent neurons (Dukelow et al., 2001; Huk et al., 2002; Amano et al., 2009). In this respect, hV5/MT+ mirrors its homolog in the monkey brain which also comprises multiple visual areas. Importantly, these subdivisions in monkey V5/MT+ contain neurons which respond selectively to different types of moving stimuli (Desimone and Ungerleider, 1986; Komatsu and Wurtz, 1988; Tanaka et al., 1993; Nelissen et al., 2006; Kolster et al., 2009; Albright, 1984; 2012). One subdivision, MT, contains directionally selective neurons which respond to many different types of motion (Lagae et al., 1994). Another subdivision, MST (Ungerleider and Desimone, 1986), has been further differentiated into a dorsal region (MSTd) where neurons respond preferentially to radial flow field stimuli, and a ventrolateral region (MSTv or MSTI) which contains neurons that are more responsive to planar motion and are important in the generation of pursuit eye movements and object tracking (Saito et al., 1986; Mikami et al., 1986; Komatsu and Wurtz, 1988; Duffy and Wurtz, 1991a; b; Tanaka et al., 1993, Eifuku and Wurtz, 1998; Duffy, 1998).

Neuroimaging studies have managed to parcellate hV5/MT+ into at least two subdivisions (Dukelow et al., 2001; Huk et al., 2002; Amano et al., 2009) and the existence of additional areas seems highly likely (see: Kolster et al., 2010). The two subdivisions most consistently identified form posterior and anterior regions within hV5/MT+ and have been differentiated on various grounds including; their respective representations of the ipsi- and contra-lateral visual field, retinotopy and population receptive field properties (Dukelow et al., 2001; Huk et al., 2002; Amano et al. 2009). Despite the different criteria employed by these studies, they all propose that the posterior subdivision of the hV5/MT+ complex is homologous with macaque area MT, whilst the more anterior region corresponds to MST. Differences do arise, however, in the nomenclature used across these studies to name the

constituent components of hV5/MT+. Certain studies have adhered to the use of terms that reflect the functional differences first described in the monkey brain and refer to human MT and MST (Dukelow et al., 2001; Huk et al., 2002; Kolster et al., 2010). Others have employed the terms TO-1 and TO-2 to refer, respectively, to the posterior and anterior subdivisions which have been differentiated on the basis of retinotopic and population receptive field properties (Amano et al., 2009). In this study we have adopted the hybrid terms MT/TO-1 for the posterior and MST/TO-2 for the anterior sub-divisions, to reflect their potential differentiation on the basis of both functional and retinotopic grounds. However, currently it is not entirely clear the extent to which these subdivisions of hV5/MT+ are coextensive or to what degree they correspond to the functional properties displayed by monkey MT and MST (dorsal or ventral). fMRI studies have demonstrated the existence of functional differences between constituent areas of the hV5/MT+ complex (Morrone et al., 2000; Smith et al., 2006; Kourtzi et al., 2002; Kolster et al., 2010). One observed difference, consistent with the functional specialisations reported for monkey MT and MST, is that the anterior subdivision of hV5/MT+, MST/TO2, is more responsive to radial motion or optic flow stimuli and appears to be more specialised for encoding the global flow properties of complex motion stimuli, compared to its posterior counterpart MT/TO-1 (Smith et al., 2006). Along similar lines, Wall et al., (2008) reported that human MST/TO-2, unlike MT/TO-1, exhibits adaptation to optic flow stimuli further emphasising the sensitivity of the former to more complex optic flow stimuli.

The purpose of this study was to investigate the existence of functional differences between the subdivisions of hV5/MT+ and demonstrate the extent to which the perception of different kinds of motion stimuli is critically dependent on neural activity within these subdivisions. Whilst fMRI experiments can identify cortical areas which exhibit the appropriate response properties that correlate with a given perceptual function, it is necessary to use interventional techniques in order to demonstrate causality. One such technique is transcranial magnetic stimulation (TMS). TMS can be used to induce transient and localised disruption of human cortical function (Pascual-Leone et al., 2000; Walsh and Cowey, 2000). Motion perception has proven to be very amenable to study using TMS (Beckers and Homberg, 1992; Hotson et al., 1994; Beckers and Zeki, 1995; Anand et al., 1998; Walsh et al., 1998; Cowey et al., 2006; Laycock et al., 2007; McKeefry et al., 2008) and in this study we deployed TMS in order to selectively disrupt neural activity in areas MT/TO-1 and MST/TO-2,

having first identified these regions of interest (ROIs) using previously employed fMRI localisers. We then measured the effects of targeted disruption to MT/TO-1 and MST/TO-2 whilst human observers performed directional judgement tasks for different kinds of motion stimuli (translational, radial and rotational). Our aim was to provide evidence of a direct causal relationship between the neural activity within areas MT/TO-1 and MST/TO-2 and the perception of different kinds of motion stimuli.

Materials and Methods

Participants

Nine participants (age range 21-46 years; mean age 29.3; six male) took part in this study, all of whom had normal or corrected-to-normal vision at time of testing and had no history of neurological or psychiatric disorders. Experiments were conducted in accordance with the Declaration of Helsinki and were approved by both York Neuroimaging Centre Ethics Committee and the University of Bradford Ethics Committee.

MRI Imaging and Analysis

Functional T2* magnetic resonance (MR) images were acquired using a GE 3-Tesla Sigma Excite HDX MRI scanner. Gradient recalled echo pulse sequences were used to measure blood oxygenation level-dependent (BOLD) signal as a function of time (TR=3000ms, TE=29ms, FOV=192cm, 128x128 matrix, 39 contiguous slices, $1.5 \times 1.5 \times 1.5 \text{ mm}^3$, interleaved slice order with no gap). A 16-channel phased-array half-head coil positioned at the occipital pole of the subject was used to measure MR signal focused on the visual cortex. A high-resolution T1-weighted three-dimensional (3D) anatomical data set was used for co-registration of functional and structural data. This was acquired using an eight-channel phased-array full-head coil (TR=7.8ms, TE=3ms, TI=450ms, FOV=290 x 290 x 276, 256 x 256 x 176 matrix, flip angle= 20°, $1.13 \times 1.13 \times 1.0 \text{ mm}^3$).

The data obtained from these functional scans were analysed using BrainVoyager QX software (Version 3.0, Brain Innovation). Pre-processing of this data included spatial smoothing (3mm Gaussian kernel, full width at half maximum), 3D motion correction, slice scan timing correction, and high pass (GLM-Fourier) temporal filtering (0.01 Hz). Multiple linear regression was then applied to the

data allowing contrasts to be made between moving-static conditions within each subject across multiple runs. Hemodynamic responses were corrected appropriately for neuro-vascular lag (Boynton et al., 1996).

Identification and Localisation of Target and Control Sites

Two subdivisions of hV5/MT+, designated MT/TO-1 and MST/TO-2, were identified using techniques similar to those described previously (Dukelow et al., 2001; Huk et al., 2002). In a block-design paradigm subjects viewed 15s periods of moving and static dots, interspersed with blank intervals. The moving dots were restricted to a 15° (diameter) circular aperture the centre of which was horizontally displaced by 15° from the central fixation point in either the left or right visual field depending on the trial. The aperture contained 300 white dots (each ~0.2° in diameter) presented on a black background and moved at a speed of 7°/s radially inwards or outwards with the direction alternating every second. The static dots were restricted to the same circular aperture as the moving dots. Each trial contained five repeats of the cycle and throughout the trial subjects fixated on a central fixation point (see Figure 1a). The more anteriorly located MST/TO-2 was localised in each hemisphere by identifying activations in the hemisphere ipsilateral to stimulation of either the right or left visual field (see Figure 1b). MT/TO-1 was then identified by subtracting the anterior MST/TO-2 activity from the whole hV5/MT+ complex activation found for contralateral presentations. By contrasting moving with static activity from the MST/TO-2 localiser, ipsilateral increases in BOLD signal were found in 15/16 hemispheres. Of these 15 hemispheres, stimuli presented in the left visual field produced a significant cluster of activation in anterior left hV5/MT+ (left MST/TO-2), and stimuli in the right visual field produced activation in right anterior hV5/MT+ (right MST/TO-2). This is consistent with the findings of previous literature that has characterised the large receptive field sizes of these anterior regions (e.g. Amano et al, 2009).

(figure 1 here)

Although some local spread of the TMS magnetic field occurs across tissue adjacent to the targeted site, previous research has shown that the differential effects of TMS are measurable in target sites

with centroids as little as 10mm apart in human cortex (cf. Pitcher et al., 2009; Silson et al., 2013). Following this, 10mm was used as the minimum criteria for distance between target points in each subject. Target points for each of our sites of interest were created by overlaying the functional data onto a 3D structural scan and creating target points for both MT/TO-1 and MST/TO-2 based on their respective centres-of-mass. The Euclidean distance (d) between these target points was then computed (see Figure 2). In the right hemisphere, it was found that MT/TO-1 and MST/TO-2 were at least 10mm apart in 8/9 hemispheres. One subject (S5) did not meet the minimum criteria for inter-target distance and so was removed from the subset of subjects that were carried forward to take part in the TMS experiment.

(figure 2 here)

Retinotopic mapping techniques (Sereno et al., 1995; DeYeo et al., 1996; Engel et al., 1997; Wandell et al., 2007) using a 90° anti-clockwise rotating wedge (contrast reversal rate 6Hz), and an expanding annulus (≤15° radius), both lasting 36s per cycle, were used to identify the control site (LO-1) in each subject. Consistent with previous data (Larsson and Heeger, 2006; Silson et al., 2013), LO-1 was found adjacent to V3d representing the contralateral lower visual field posteriorly, and the contralateral upper visual field anteriorly (see Figure 3). Cortical area LO-1 was chosen as a control site because it lies in close proximity to areas MT/TO-1 and MST/TO-2, but in contrast to these areas LO-1 has no known role in the processing of visual motion, appearing instead to be involved in processing orientation information related to the recognition of objects (Larsson and Heeger, 2006; Silson et al., 2013). The use of this control site should determine whether there are any effects of proximity to TMS on performance. It should also allow us to confirm that any effects found from applying TMS to the target ROIs are not simply due to the general effect of applying TMS to the extra-striate visual cortex.

(figure 3 here)

We did attempt to differentiate MT/TO-1 and MST/TO-2 using retinotopic criteria but we found that whilst MT/TO1 did seem to possess retinotopy, the maps obtained from MT/TO2 were less reliable. This is consistent with the results of Huk et al., (2002) and presumably is a reflection of the increased

receptive field sizes for MST/TO2 that are less likely to be shown to be retinotopic by the spatial and temporal parameters of the standard rotating wedge and ring stimuli employed here (see: Kolster et al., 2010 for a discussion on this point).

Co-Registration of fMRI and TMS Target sites

The target points for repetitive TMS delivery were chosen as the centre-of-mass co-ordinates within the areas identified as MT/TO-1, MST/TO-2 and LO-1 following the fMRI experiments described above. All of the target ROIs selected for TMS were located in the right cerebral hemisphere and the mean Talairach co-ordinates for each of these sites are given in Table 1. Note that the target sites for areas in this study were identified primarily on the basis of the ipsi- versus contra-lateral responses to motion stimuli. The control site, LO-1, was identified on the basis of its previously described retinotopy (Larsson and Heeger, 2006; Silson et al., 2013). Table 1 also includes Talairach co-ordinates for MT/TO-1 and MST/TO-2 which in previous studies have identified on both retinotopic and functional grounds (Dukelow et al., 2001; Kolster et al. 2010). Despite the different criteria employed there appears to be a close degree of correspondence across the studies in terms of the location of the centres of these ROIs.

Table 1 here

Following identification of the three target points in 3D space, co-registration between each subject's head and their structural scans was achieved using a 3D ultrasound digitizer CMS30P (Zebris) in conjunction with BrainVoyager QX (McKeefry et al., 2008). This method creates a local spatial co-ordinate system which is able to link the spatial positions of ultrasound transmitters on the subject and the coil with pre-specified fiducials on the 3D representations.

Stimuli and Psychophysical Procedures

All motion stimuli were displayed on a high-resolution CRT monitor with a refresh rate of 75Hz (Mitsubishi DiamondPro 2070SB). Stimuli were generated using Psychophysics Toolbox Version 3

(Brainard, 1997; Pelli, 1997; Kleiner et al., 2007) in 32-Bit MATLAB (Version 7.6.0. Natick, Massachusetts: The MathWorks Inc., 2008). Dot stimuli were restricted to a 10° circular aperture containing 300 white dots on a black background. Each dot subtended 0.2° of visual angle (dot density ~3.82/deg²) and all dots moved at a speed of 7°/s regardless of direction. The centre of this aperture was horizontally displaced by 15° to the left of the fixation point (see Figure 4). Three kinds of motion stimulus were used; 1) translational motion – where the dots moved either up or down, 2) radial motion – where the dots moved outwards or inwards from the centre of the aperture and 3) rotational motion – where the dots rotated in a clock-wise or counter-clockwise about the centre of the aperture.

(Figure 4 here)

In each type of motion stimulus a low, but detectable, percentage of dots moved in a coherent direction (signal dots) whilst the rest moved randomly (noise dots). Participants were required to identify the coherent direction of the signal dots using a two alternative-forced-choice (2AFC) paradigm for three different conditions: translational coherent dots (up or down), radial coherent dots (inward or outward), and rotational coherent dots (clockwise or anticlockwise). Subjects were instructed to record their decision regarding the direction of the signal dots using an appropriate button on the keyboard as quickly and as accurately as possible. Our decision to employ the 3 different motion stimuli tasks was motivated by earlier single-unit monkey neurophysiology (e.g. Duffy and Wurtz, 1991a; b) and human neuroimaging studies (e.g. Smith et al., 2006). These studies have shown that different selectivities (particularly for radial motion stimuli) exist across the subdivisions of human and monkey V5/MT+ which we wished to exploit using similar stimulus types. For each participant, and separately for each motion stimulus type, preliminary psychophysical measurements were used to establish the proportion of dot coherence required to result in correct motion discrimination 75% of the time. The psychophysical data were then fitted by a two-parameter logistic function of the form:

 $P(x) = [1 + \exp\{\delta(\theta - x)\}]^{-1}$ [equation 1]

Where: *x* represents the stimulus value (coherence), P(x) is the response probability at *x*, δ is the slope parameter, and θ represents the threshold parameter corresponding to the stimulus level at which response probability is 75%. These subject- and stimulus-specific coherence values (see Table 2) were used in the main TMS experiments where subjects performed ~100 trials for each condition in each task. Trials were removed if the subject took longer than 3s to respond. Overall <3% of trials were removed from the complete dataset.

(Table 2 here)

TMS Protocol

During the task, participants viewed a centrally placed fixation cross with their right eye (left eye occluded) from a distance of 57cm. The centre of a 10° (diameter) stimulus was placed 15° horizontally relative to the fixation cross in the left visual field (LVF). This was done in order to minimise any involvement of ipsi-lateral V5/MT+ in the performance of the motion discrimination tasks. In these experiments TMS was delivered to the target sites in the right hemisphere, leaving their counterparts in the left hemisphere functioning normally. Amano et al (2009) have demonstrated that the receptive fields of hV5/MT+ neurones can extend well beyond the vertical meridian into the ipsilateral (in this case the left) visual field. Our stimulus placement was therefore an attempt to minimise contributions from the ipsi-lateral non-stimulated motion area. Similar reasoning lies behind the choice of stimulus size (10° diameter) for the TMS experiments, in that larger stimulus sizes extending towards the midline would also allow the involvement of neurons with large receptive fields from the undisrupted ipsi-lateral hV5/MT+. The choice of stimulus size is at first glance at odds with results from monkey single-unit neurophysiology, where studies have shown that MSTd neurons give weak responses to relatively small stimuli (Komatsu and Wurtz, 1988). However, more recent neuroimaging studies in human have clearly shown that activation of human MST/TO-2 (and its differentiation from MT/TO-1) can be achieved for stimuli of sizes 8°x 8° and greater (Becker et al., 2008).

In the combined TMS and psychophysical experiments the onset of the motion stimulus was synchronous with onset of a train of five biphasic (equal relative amplitude) repetitive TMS (rTMS)

pulses (Figure 4). Previous results had demonstrated that this temporal configuration was the most effective at inducing effects in hV5/MT+ (McKeefry et al., 2008). These pulses were applied to the participant's scalp using a figure-of-eight coil (50 mm diameter) connected to a Magstim Super Rapid2 stimulator (Magstim, Wales, UK). The rTMS trains were applied at a frequency of 25Hz, at a level of 70% of the maximum output. Participants undertook two blocks of the motion task for each TMS site and condition. Only one condition was tested in each session and the order of presentation of conditions was counter-balanced across participants.

Statistical Analysis

Statistical analysis of the results was performed using the SPSS software package (IBM). Repeatedmeasures ANOVAs were calculated across all conditions (baseline, MT/TO-1, MST/TO-2, LO-1 control) for each of the three tasks individually (translational, radial, rotational). When a significant main effect was present, pair-wise comparisons were applied to the data sets (Bonferroni corrected for multiple comparisons). The assumption of normal distribution was confirmed with Mauchly's Test of Sphericity. If this assumption was met (i.e. sphericity is non-significant) then the ANOVA was calculated assuming sphericity, however if the assumption was violated, the degrees of freedom (dF) would be corrected to allow appropriate interpretation of the F value of the ANOVA. These dF corrections included the Greenhouse-Geisser correction when sphericity was less than 0.75, and Huynh-Feldt correction when sphericity exceeded 0.75.

Results

Percent correct (pCorrect) is the main dependent variable measured within this experiment. This variable quantifies the subject- and condition-specific variance in performance around a 75% threshold as a function of the task performed and TMS stimulation condition. Applying repetitive TMS to MT/TO-1 and MST/TO-2 during motion coherence direction discrimination tasks appears to produce effects that are task-specific (see Figure 5). These effects were estimated using repeated measures ANOVAs and all pair-wise comparisons were Bonferroni corrected. Significant main effects

of experimental condition on performance were found for all three motion direction discrimination tasks: translational (F(3,21) = 30.35, p<0.001), radial (F(3,21) = 13.40, p<0.001), and rotational (F(3,21) = 8.34, p = 0.001).

For translational motion, pair-wise comparisons showed that application of TMS to both MT/TO-1 and MST/TO-2 produced significant reductions in ability of subjects to determine the direction of motion of the dots relative to both baseline and control conditions (MT/TO-1 versus baseline, p = 0.003; MT/TO-1 versus control, p = 0.007; MST/TO-2 versus baseline, p = 0.001; MST/TO-2 versus control (LO1), p = 0.002). No other pair-wise comparisons were found to be significant (p = 1.00 in all cases). Clearly, for the discrimination of the translational motion direction, the consequences of disruption by TMS are the same for the two areas, indicating that neural activity in both MT/TO-1 and MST/TO-2 is essential for the perception of such stimuli.

In contrast, for radial motion there was a significant differential effect on perception when TMS was applied to MT/TO-1 and MST/TO-2. When MT/TO-1 was targeted, there was no significant effect on performance (p = 1.00), whereas application of TMS to MST/TO-2 resulted in a decrease in subjects' ability to perceive radial motion relative to baseline and the control site (MST/TO-2 versus baseline, p = 0.005; MST/TO-2 versus control, p = 0.005). Importantly, there is a significant dissociation between the effects of TMS on the perception of radial motion between MT/TO-1 and MST/TO-2 (MST/TO-2 versus MT/TO-1, p = 0.007) demonstrating that neural activity within area MST/TO-2, but not MT/TO-1, is required for performance of the radial motion discrimination task.

For the rotational motion task, application of TMS to MST/TO-2 significantly reduced performance relative to baseline and control (MST/TO-2 versus baseline, p = 0.017; MST/TO-2 versus control, p = 0.035). Application of TMS to MT/TO-1 produced no significant effects (significant (MT/TO-1 versus baseline, p = 0.289). However, whilst there are clear deficits relative to the baseline and control conditions when TMS is applied to MST/TO-2, comparisons with performance when TMS is applied to MT/TO-1 fall short of showing complete dissociation between the two areas (MT/TO-1 versus MST/TO-2, p = 0.687). No other comparisons were found to be significant (baseline versus control, p = 0.371; MT/TO-1 versus control, p = 1.00).

Response times were recorded for every trial in all three tasks. A significant main effect of response time was found for translational motion (F(3,21) = 4.24, p = 0.017), however pairwise comparisons failed to identify any significant differences between conditions (baseline versus MT/TO-1, p = 0.189; baseline versus MST/TO-2, p = 0.336; baseline versus control, p = 0.307; MT/TO-1 versus control, p = 0.607; all other comparisons, p = 1.00). No significant main effects of response time were found for either radial (F(3,21) = 1.88, p = 0.165), or rotational motion (F(3,21) = 0.99, p = 0.416), showing that response time did not differ significantly across any of the TMS conditions in either of those tasks (Figure 6). However it is noteworthy that longer reaction times occurred under conditions where discrimination deficits were found – the opposite of a speed-accuracy trade off.

(figure 5 here)

Across all tasks and all motion conditions, there were no significant differences between baseline performance and performance during TMS of the control site. This similarity supports the decision to use this site (LO-1) as a suitable visual cortex control and we can therefore conclude that any experimental effects found are not a result of confounding variables associated with general application of TMS as this would also affect the performance associated with control site. No significant effects were found for response times across radial and rotational motion, confirming that the response times will not have had any confounding effects on the performance of the subject.

(figure 6 here)

Discussion

This study has demonstrated that when TMS is used to disrupt neural activity within two major subdivisions of hV5/MT+, MT/TO-1 and MST/TO-2, dissociable effects on the perception of different kinds of moving stimuli can be induced in human observers. The application of TMS to either MT/TO-1 or MST/TO-2 impairs significantly the perception of translational motion, but only disruption of MST/TO-2 induces significant deficits in the perception of radial flow patterns. Previous neuroimaging experiments have successfully managed to distinguish between these subdivisions of hMT/V5+ on the basis of retinotopy and receptive field properties (Dukelow et al., 2001; Huk et al., 2002; Amano et al., 2009). In addition, differences in the response selectivities of MT/TO-1 and MST/TO-2 to different kinds of motion stimuli have also been demonstrated (Morrone et al., 2000; Smith et al., 2006; Wall et al., 2008; Kolster et al. 2010; Pitzalis et al. 2013). This study provides further evidence of functional differences between the constituent sub-divisions of hV5/MT+ by establishing the existence of causal dependencies between neural activity within these sub-divisions and the perception of different kinds of moving stimuli.

Our results demonstrate that the anterior subdivision of hV5/MT+, area MST/TO-2, appears integral to the perception of radial flow patterns. In this respect, human MST/TO-2 appears functionally similar to monkey MSTd; the neurons of which exhibit a similar response selectively (Saito et al., 1986; Tanaka and Saito, 1989; Tanaka et al., 1989; Duffy and Wurtz, 1991a; b; Lagae et al., 1994). However, this is directly at odds with previous proposals of homology between human MST/TO-2 and monkey MSTv made on the basis of similarities between their RF characteristics (Amano et al., 2009). The justification for our proposal, that MST/TO-2 is more closely allied to MSTd rather than MSTv, is based on the functional similarities between MST/TO-2 and MSTd, coupled with its functional differences with MSTv, where neurons are unresponsive to radial flow stimuli and more important for the generation of smooth pursuit eye movements (Eifuku and Wurtz, 1998). These conflicting findings realting to the RF and functional properties highlight the difficulties in establishing clear homologies between the subdivisions of human and monkey V5/MT+. This is further complicated by the fact that other studies of hV5/MT+ suggest there may be at least four separate representations of the visual field (MT, pMSTv, pFST & pV4t (p = putative)) within hV5/MT+ (Kolster et al., 2010). The Talairach coordinates for MT and pMSTv as defined by Kolster et al. (2010), agree very closely with our locations

of MT/TO-1 and MST/TO-2, respectively (see Table 1). The use of the term pMSTv by Kolster et al., (2010) clearly suggests homology with monkey MSTv and, in view of the coinciding cortical locations, also with our MST/TO-2. However, Kolster et al (2010) define pMSTv purely in terms of retinotopic criteria. Whilst their data highlight that pMSTv is responsive to motion and less so to shape, no data are presented to assess whether it adheres to the functional criteria (e.g. lack of selectivity to radial flow stimuli) that have previously been used to differentiate MSTv and MSTd in monkeys (Desimone and Ungerleider, 1986; Komatsu and Wurtz, 1988; Tanaka et al., 1993). Therefore, a link between MST/TO-2 and MSTd cannot be conclusively ruled out until we have a clearer understanding of the functional properties of human pMSTv.

It has been argued that the four-component composition of hV5/MT+ revealed by Kolster et al. (2010) is more consistent with the structure of monkey V5/MT+ and, moreover, constitutes a key organisational feature of V5/MT+ across all primates (Nelissen et al., 2006; Kolster et al., 2010). Similar to the monkey brain, the multiple visual field maps present in human V5/MT+ have been linked to the analysis of different aspects of motion processing. They are hypothesised as forming the basis of separate processing pathways, emanating from V5/MT, which are involved in the analysis of different kinds of moving stimuli (Xiao et al., 1997; Kolster et al., 2009; Kourtzi et al., 2002). In the present study we were unable to find evidence of separate ventral retinotopic maps corresponding to FST and V4t. However, absence of evidence does not constitute evidence of absence and it may be that our retinotopic mapping protocols are simply not suited to revealing all of the individual maps that may be present in hV5/MT+. For example, the constituent sub-divisions of hV5/MT+ are thought to share a foveal confluence (see: Kolster et al., 2010), it is likely that the size of wedge employed for this experiment may have been too large to accurately distinguish between phase reversals in foveal cortex (see: Schira et al., 2009). It may very well prove to be the case that the subdivision of V5/MT+ into four separate areas, rather than just two, is the common organisation feature across all primate brains.

Our results are consistent with previous fMRI studies that have examined functional differences between the subdivisions of hV5/MT+ defined by the same criteria (Dukelow et al., 2001; Huk et al., 2002; Amano et al., 2009). The data show that human MST/TO-2 is differentially responsive to the

basic components of optic flow (Pitzalis et al., 2013; Smith et al., 2006; Wall et al., 2008). However, the precise pattern of stimulus selectively reported for human MST/TO-2 is not consistent across all studies. Smith and colleagues (Smith et al., 2006; Wall et al., 2008) found human MST/TO-2 to be responsive to radial and rotational flow patterns whilst Pitzalis et al (2013) showed the same area to be most responsive to radial motion, but less so to rotational stimuli. The reasons for these discrepancies are not clear but the results described by Pitzalis et al (2013) appear to be more in line with the pattern of deficits induced by TMS in this study where we have failed to reveal a complete dissociation between the contribution of MT/TO-1 and MST/TO-2 to the perception of rotational motion. This finding is consistent with neuropsychological case studies (Beardsley and Vaina, 2005) which point to the existence of separate cortical loci for the neural activity that supports the perception of radial and rotational motion.

Radial motion stimuli are considered as having particular significance in that the expansion/contraction from/to a central point provides an important cue for the guidance of self-motion and for ecologically important visual tasks within visual environments (Gibson, 1950; Warren and Hannon, 1988; Warren and Rushton, 2009). Evidence from the monkey brain suggests that neurons in area MSTd play a key role in the analysis of this information (Tanaka and Saito, 1989; Duffy and Wurtz, 1991a; 1995) and that electrical stimulation of this area can bias directional judgements of self-motion made by monkeys (Britten and Van Wezel, 1998). Reports that regions within hV5/MT+ exhibit similar response selectivity for radial motion stimuli has led to the proposal that area MST/TO-2 plays a similar role in the encoding of self-motion in the human brain (Morrone et al., 2000; Pitzalis et al., 2013). In addition to MST/TO-2, radial flow patterns have also been shown to elicit responses in other brain areas such as V3A, V3B, V6 and the intra-parietal sulcus (Morrone et al., 2000; Smith et al., 2006; Wall and Smith, 2008; Cardin et al., 2012). MST/TO-2 possibly forms just one component area within a hierarchical processing network comprising multiple cortical areas within which increasingly more complex analyses allow the extraction of behaviourally relevant information from radial flow stimuli (Wall and Smith, 2008).

TMS can alter the signal:noise ratio of neural activity within a cortical region in various ways which will result in functional impairments. Current TMS research has yet to agree on exactly how the effects of

TMS are mediated within the cortex. Some studies posit that TMS induces neural noise (Ruzzoli et al., 2010), others that it reduces neural signal (Harris et al., 2008), while another school of thought proposes that TMS may be both reducing signal and increasing noise at the same time (Allen et al., 2007). Regardless of how TMS disrupts cortical function we demonstrate that measurable and specific functional deficits are induced by the delivery of TMS to particular target sites. We are able to demonstrate that there is a causal dependence between different aspects of motion perception and neural activity within separate sub-divisions of hV5/MT+. Crucially, though the functional deficits induced by the application of TMS to areas MT/TO-1 and MST/TO-2 do not constitute a double dissociation. Such a finding would have carried the implication that the analysis of translational and radial motion occurs independently within MT/TO-1 and MST/TO-2. Instead, disruption to both MT/TO-1 and MST/TO-2 induces deficits in the perception of translational motion, while the perception of radial motion is only affected by disruption of MST/TO-2. Rather than independent processing, this suggests a more serial form of processing where information is passed on from MT/TO-1 and subsequently subjected to more complex analysis from which sensitivity to radial flow emerges only at the level of MST/TO-2. Neurons in monkey MT have smaller receptive fields that those found in MST which as a result are able to integrate motion signals over much larger spatial extents (e.g. Saito et al., 1986). Mechanisms supporting this kind of transformation have been described in monkey MST (Yu et al., 2010) where sensitivity to optic flow has been found to be based upon local selectivity to translational motion within sub-regions of the large RFs of MST neurons. This localised response is then combined across the whole RF to generate global sensitivity to optic flow (Yu et al., 2010). Similar local versus global RF sensitivities may explain the functional deficits found in human MST/TO-2 in this study.

In summary, this study has provided direct evidence of causal links between neural activity in different subdivisions of hV5/MT+ and the perception of different kinds of motion stimuli. The results provide further confirmation of what has long been suspected; namely, that hV5/MT+ comprises multiple visual areas that are separable, not only on retinotopic, but also on functional bases. In this respect, motion processing in the human brain may be organised along similar lines to that found in the monkey and is based on a hierarchy of retinotopically distinct visual areas, the neurons of which possess increasing RF size, response selectivity and processing complexity (Orban, 2008). Areas

MT/TO-1 and MST/TO-2 in the human brain, like MT and MSTd in the monkey, may provide the origins of different cortical processing networks that are involved in the analysis of planar motion and radial flow, respectively.

Acknowledgements

This work was funded by a BBSRC award (BB/N003012/1). We are grateful to Jim Gilchrist, Sam Lawrence and Richard Vernon for their assistance in data analysis. We acknowledge the Faculty of Life Sciences, University of Bradford for the PhD studentship award to SS.

References

Albright TD. 1984. Direction and orientation selectivity of neurons in visual area MT of the macaque. J Neurophysiol. 52:1106-1130.

Albright TD. 2012. On the perception of probable things: Neural substrates of associative memory, imagery, and perception. Neuron. 74(2):227-245.

Allen EA, Pasley BN, Duong T, Freeman RD. 2007. Transcranial magnetic stimulation elicits coupled neural and hemodynamic consequences. Science. 317:1918-1921.

Amano K, Wandell BA, Dumoulin SO. 2009. Visual field maps, population receptive field sizes, and visual field coverage in the human MT+ complex. J Neurophysiol. 102:2704-2718.

Anand S, Olson JD, Hotson JR. 1998. Tracing the timing of human analysis of motion and chromatic signals from occipital to tempero-parieto-occipital cortex: a transcranial magnetic stimulation study. Vis Res. 38:2619-2627.

Beardsley SA, Vaina LM. 2005. How can a patient blind to radial motion discriminate shifts in the center-of-motion?. J Comp Neurosci. 18:55-66.

Becker HG, Erb M, Haarmeier T. 2008. Differential dependency on motion coherence in sub-regions of the human MT+ complex. Eur J Neurosci. 28:1674-1685.

Beckers G, Homberg V. 1992. Cerebral visual motion blindness: transitory akinetopsia induced by transcranial magnetic stimulation of human area V5. Proc Roy Soc (B). 249:173-178.

Beckers G, Zeki S. 1995. The consequences of inactivating areas V1 and V5 on visual motion perception. Brain. 118:49-60.

Brainard DH. 1997. The psychophysics toolbox. Spat Vis. 10:433-436.

Britten KH, Van Wezel RJA (1998) Electrical microstimulation of cortical area MST biases heading perception in monkeys. Nat Neurosci. 1:59-63.

Boynton GM, Engel SA, Glover GH, Heeger, DJ. 1996. Linear systems analysis of functional magnetic resonance imaging in human V1. J Neurosci. 16:4207–4221.

Cardin V, Hemsworth L, Smith AT. 2012. Adaptation to heading direction dissociates the roles of human MST and V6 in the processing of optic flow. J Neurophysiol. 108:794-801.

Cowey A, Campana G, Walsh V, Vaina LM. 2006. The role of human extra-striate visual areas V5/MT and V2/V3 in the perception of the direction of global motion: a transcranial magnetic stimulation study. Exp Brain Res. 171:558-562.

Culham J, He S, Dukelow S, Verstraten FAJ. 2001. Visual motion and the human brain: what has neuroimaging told us? Acta Psychologica. 107:69-94.

Desimone R, Ungerleider LG. 1986. Multiple visual areas in the caudal superior temporal sulcus of the macaque. J Comp Neurol. 248:164–189.

DeYoe EA, Carman GJ, Bandettini P, Glickman S, Wieser J, Cox R, Miller D, Neitz J. 1996 Mapping striate and extra-striate visual areas in human cerebral cortex. Proc Natl Acad Sci.[USA] 93:2382–2386.

Duffy CJ, Wurtz RH. 1991a. Sensitivity of MST neurons to optic flow stimuli. I. A continuum of response selectivity to large-field stimuli. J Neurophysiol. 65:1329–1345.

Duffy CJ, Wurtz RH. 1991b. Sensitivity of MST neurons to optic flow stimuli. II. Mechanisms of response selectivity revealed by small-field stimuli. J Neurophysiol. 65:1346-1359.

Duffy CJ, Wurtz RH. 1995. Response of monkey MST neurons to optic flow stimuli with shifted centers of motion. J Neurosci. 15:5192-5208.

Duffy CJ. 1998. MST neurons respond to optic flow and translational movement. J Neurophysiol. 80:1816-1827.

Dukelow SP, DeSouza JFX, Culham JC, van den Berg AV, Menon RS, Vilis T. 2001. Distinguishing subregions of the human MT+ complex using visual fields and pursuit eye movements. J Neurophysiol. 86:1991–2000.

Dumoulin SO, Bittar RG, Kabani NJ, Baker CL, LeGoualher G, Pike GB, Evans AC. 2000. A new anatomical landmark for reliable identification of human area V5/MT: a quantitative analysis of sulcal patterning. Cereb Cortex. 10:454–463.

Dupont P, Orban GA, De Bruyn B, Verbruggen A, Mortelmans L. 1994. Many areas in the human brain respond to visual motion. J Neurophysiol. 72:1420–1424.

Eifuku S, Wurtz RH. 1998. Response to motion in extra-striate area MST1: Center-surround interactions. J Neurophysiol. 80:282-296.

Engel SA, Glover GH, Wandell BA. 1997. Retinotopic organization in human visual cortex and the spatial precision of functional MRI. Cereb Cortex. 7:181-192.

Gibson JJ (1950) The perception of the Visual World. Boston (MA): Houghton Mifflin.

Harris JA, Clifford CW, Miniussi C. 2008. The functional effect of transcranial magnetic stimulation: signal suppression or neural noise generation? J Cog Neurosci. 20:734-740.

Hotson J, Braun D, Herzberg W, Boman D. 1994. Transcranial magnetic stimulation of extra-striate cortex degrades human motion direction discrimination. Vis Res. 34:2115-2123.

Huk AC, Dougherty RF, Heeger DJ. 2002. Retinotopy and functional subdivision of human areas MT and MST. J Neurosci. 22:7195–7205.

Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C. 2007. What's new in Psychtoolbox-3. Percept. 36:1.

Kolster H, Mandeville JB, Arsenault JT, Ekstrom LB, Wald LL, Vanduffel W. 2009. Visual field map clusters in macaque extra-striate visual cortex. J Neurosci. 29:7031-7039.

Kolster H, Peeters R, Orban GA. 2010. The retinotopic organization of the human middle temporal area MT/V5 and its cortical neighbors. J Neurosci. 30:9801-20.

Komatsu H, Wurtz RH. 1988. Relation of cortical areas MT and MST to pursuit eye movements. I. Localization and visual properties of neurons. J Neurophysiol. 60:580-603.

Kourtzi Z, Bulthoff HH, Erb M, Grodd W. 2002. Object-selective responses in the human motion area MT/MST. Nat Neurosci. 5:17–18.

Larsson J, Heeger DJ. 2006. Two retinotopic visual areas in human lateral occipital cortex. J Neurosci. 26:13128-13142.

Lagae L, Maes H, Raiguel S, Xiao D-K, Orban GA. 1994. Responses of macaque STS neurons to optic flow components: a comparison of areas MT and MST. J Neurophysiol. 71:1597–1626.

Laycock R, Crewther DP, Fitzgerald PB, Crewther SG. 2007. Evidence for fast signals and later processing in human V1/V2 and V5/MT+: a TMS study of motion perception. J Neurophysiol. 98:1253-1262.

McKeefry DJ, Watson JDG, Frackowiak RSJ, Fong K, Zeki S. 1997. The activity in Human areas V1, V2, V3 and V5 during the perception of coherent and incoherent motion. Neuroimage. 5:1-12.

McKeefry DJ, Burton MP, Vakrou C, Barrett BT, Morland AB. 2008. Induced deficits in speed perception by transcranial magnetic stimulation of human cortical areas V5/MT+ and V3A. J Neurosci. 28:6848-57.

Mikami A, Newsome WT, Wurtz RH. 1986. Motion selectivity in macaque visual cortex. I. Mechanisms of direction and speed selectivity in extrastriate area MT. J Neurophysiol. 55:1308-1327.

Morrone MC, Tosetti M, Montanaro D, Fiorentini A, Cioni G, Burr DC. 2000. A cortical area that responds specifically to optic flow, revealed by fMRI. Nat Neurosci. 3:1322–1328.

Nelissen K, Vanduffel W, Orban GA. 2006. Charting the lower superior temporal region, a new motion-sensitive region in monkey superior temporal sulcus. J Neurosci. 26:5929-47.

Orban GA (2008) Higher order visual processing in macaque extra-striate cortex. Physiolo Rev. 88:59–89.

Pascual-Leone A, Walsh V, Rothwell J. 2000. Transcranial magnetic stimulation in cognitive neuroscience – virtual lesion, chronometry and functional connectivity. Curr Opin Neurobiol. 10:232-237.

Pelli DG. 1997. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. Spat Vis. 10:437-442.

Pitcher D, Charles L, Devlin JT, Walsh V, Duchaine B. 2009. Triple dissociation of faces, bodies, and objects in extrastriate cortex. Curr Biol. 19:319-324.

Pitzalis S, Sdoia S, Bultrini A, Committeri G, Di Russo F, Fattori P, Galletti C, Galati G. 2013. Selectivity to translational egomotion in human brain motion areas. PLoS ONE. 8(4): e60241.doi 10.1371.

Ruzzoli M, Marzi CA, Miniussi C. 2010. The neural mechanisms of the effects of transcranial magnetic stimulation on perception. J Neurophysiol. 103:2982-2989.

Sack AT, Kadosh RC, Schuhmann T, Moerel M, Walsh V, Goebel R. 2009. Optimizing functional accuracy of TMS in cognitive studies: a comparison of methods. J Cog Neurosci. 21:207-221.

Saito H, Yukie M, Tanaka K, Hikosaka K, Fukada Y, Iwai E. 1986. Integration of direction signals of image motion in the superior temporal sulcus of the macaque monkey. J Neurosci. 6:145-157.

Schira MM, Tyler CW, Breakspear M, Spehar B. 2009. The foveal confluence in human visual cortex. J Neurosci. 29:9050-9058.

Sereno MI, Dale AM, Reppas JB, Kwong KK, Belliveau JW, Brady TJ, Rosen BR, Tootell RB. 1995. Borders of multiple human visual areas in humans revealed by functional MRI. Science. 268:889-893.

Silson EH, McKeefry DJ, Rodgers J, Gouws AD, Hymers M, Morland AB. 2013. Specialized and independent processing of orientation and shape in visual field maps LO1 and LO2. Nat Neurosci. 16:267-269.

Smith AT, Greenlee MW, Singh KD, Kraemer FM, Hennig J. 1998. The processing of first- and second-order motion in human visual cortex assessed by functional magnetic resonance imaging (fMRI). J Neurosci. 18:3816–3830.

Smith AT, Wall MB, Williams AL, Singh KD. 2006. Sensitivity to optic flow in human cortical areas MT and MST. Eur J Neurosci. 23:561–569.

Tanaka K, Fukada Y, Saito H. 1989. Underlying mechanisms of the response specificity of the expansion/contraction and rotation cells in the dorsal part of the medial superior temporal area of the macaque monkey. J Neurophysiol. 62:642–656.

Tanaka K, Saito H. 1989. Analysis of motion of the visual field by direction, expansion/contraction, and rotation cells clustered in the dorsal part of the medial superior temporal area of the macaque monkey. J Neurophysiol. 62:626–641.

Tanaka K, Sugita Y, Moriya M, Saito H. 1993. Analysis of object motion in the ventral part of the medial superior temporal area of the macaque visual cortex. J Neurophysiol. 69:128-142.

Tootell RBH, Reppas JB, Kwong KK, Malach R, Born IT, Brady TJ, Rosen BR, Belliveau JW. 1995. Functional analysis of human MT and related visual cortical areas using magnetic resonance imaging. J Neurosci. 15:3215-3230.

Ungerleider LG, Desimone R. 1986. Cortical connections of visual area MT in the macaque. J Comp Neurol. 248:190–222.

Wall MB, Lingnau A, Ashida H, Smith AT. 2008. Selective visual responses to expansion and rotation in the human MT complex revealed by functional magnetic resonance imaging adaptation. European J Neurosci. 27:2747-2757.

Walsh V, Ellison A, Battelli L, Cowey A. 1998. Task-specific impairments and enhancements induced by magnetic stimulation of human visual area V5. Proc Roy Soc (B). 265:537-543.

Walsh V, Cowey A. 2000. Transcranial magnetic stimulation and cognitive neuroscience. Nat Rev Neurosci. 1:73–79.

Wandell BA, Dumoulin SO, Brewer AA. 2007. Visual field maps in human cortex. Neuron. 56:366-383.

Warren WH, Hannon DJ. 1988. Direction of self-motion is perceived from optical flow. Nature. 336:162–168.

Warren PA, Rushton SK. 2009. Optic flow processing for the assessment of object movement during ego movement. Curr Biol. 19:1555-1560.

Watson JD, Myers R, Frackowiak RS, Hajnal JV, Woods RP, Mazziotta JC, Shipp S, Zeki S. 1993. Area V5 of the human brain: evidence from a combined study using positron emission tomography and magnetic resonance imaging. Cereb Cortex. 3:79–94.

Xiao DK, Marcar VL, Raiguel SE, Orban GA. 1997. Selectivity of macaque MT/V5 neurons for surface orientation in depth specified by motion. Eur J Neurosci. 9:956–964.

Yu CP, Page WK, Gaborski R, Duffy CJ. 2010. Receptive field dynamics underlying MST neuronal optic flow selectivity. J Neurophysiol. 103:2794-2807.

Zeki S, Watson JDG, Lueck CJ, Friston KJ, Kennard C, Frackowiak RSJ. 1991. A direct demonstration of functional specialization in the human visual cortex. J Neurosci. 11:641–649.

Figure Legends

Figure 1. Stimulus specification and identification of MT/TO-1 and MST/TO-2. a) (Top Row) Example stimuli showing dots presented in either left or right visual field. (Bottom row) Axial fMRI data from one representative subject (S3) showing BOLD signal (p<0.001) generated by moving vs static functional localisers presented in both left and right visual field (averaged across four runs). Anterior MST/TO-2 (white dotted line) can be seen relative to whole V5/MT+ complex in both hemispheres and is identified as the anterior portion of the complex activated by ipsilateral stimuli. b) Magnified view of posterior occipital lobe in the same subject when viewing dots in right visual field demonstrating the identification of MT/TO-1. Here, left MT/TO-1 (yellow dotted line) is shown as the subtraction of the MST/TO-2 ipsilateral activation (white dotted line) from the whole V5/MT+ complex activated by contralateral stimulation. c) Areas MST/TO-2 and MT/TO-1 shown as increases in BOLD signal superimposed onto 3D inflated surfaces of the left cerebral hemisphere of subject S3. The black circle highlights the magnified area of the two images, with the left image showing ipsilateral activation of MST/TO-2 (white dotted line) produced when dots were viewed in the left visual field. Similarly the right image shows identification of MT/TO-1 (yellow dotted line) when MST/TO-2 is subtracted from the full contralateral activation of V5/MT+ produced by viewing dots in the right visual field.

Figure 2. Bar chart showing Euclidean distance (mm) between MT/TO-1 and MST/TO-2 in the right hemisphere (RH) for each subject. The black dashed line denotes the 10mm separation criterion.

Figure 3. Location of the LO-1 control stimulation site. In subject S3 a portion of the lateral surface of the posterior occipital lobe is highlighted. This region is magnified in the lower part of the diagram and shown as an inflated 3D mesh with a pseudo colour representation of the visual field.

Figure 4. Psychophysical procedure for direction discrimination task using radial motion as the example stimulus. In the repetitive TMS trials, pulse trains were delivered synchronously with onset of stimuli and persisted for an equivalent duration (200ms). Following stimulus offset, the subjects reported the perceived motion direction relevant to the task (up/down for translational motion, in/out for radial motion, clockwise/anti-clockwise for rotational motion) by an appropriate key press.

Figure 5. Bar charts showing average proportion correct for Translational (a), Radial (b), and Rotational (c) motion tasks (a low value suggests the task was found to be more difficult). Single asterisks (*) highlight significant differences at 0.05 level; double asterisks (**) highlight significant differences at 0.01 level. Error bars show S.E.M.

Figure 6. Bar chart showing average response time (RT) for Translational (a), Radial (b), and Rotational (c) motion tasks. The data show no significant differences at 0.05 level. Error bars show S.E.M.

Table 1. Table comparing average Talairach co-ordinates for MT/TO-1 and MST/TO-2 from the current study with co-ordinates from Dukelow et al. (2001) and Kolster et al. (2010).

Right Hemisphere	MT/TO-1			MST/TO-2		
	X	У	Ζ	X	У	Ζ
This study	42 ± 2.7	-76 ± 3.0	-3 ± 7.4	43 ± 4.1	-69± 6.7	0 ± 8.8
Dukelow et al,. 2000	44 ± 3	-64 ± 7	5 ± 4	45 ± 3	-60 ± 5	5 ± 4
Kolster et al., 2010	46	-78	6	44	-70	5
Left Hemisphere	MT/TO-1			MST/TO-2		
	X	У	Ζ	X	У	Z
This study	-46 ± 2.3	-79 ± 2.7	-1 ± 8.9	-47 ± 5.7	-71 ± 5.3	-2 ± 9.1
Kolster et al., 2010	-48	-75	8	-45	-67	6

Subject	Translational (%)	Radial (%)	Rotational (%)
S1	22.6	13.6	10.9
S2	27.4	24.4	13
S3	16.0	10.1	7.9
S4	20.9	13.0	6.6
S6	26.3	15.4	10.2
S7	39.8	20.2	4.4
S8	29.4	24.4	8.7
S9	35.1	23.7	5.0
Average	27.2 ± 7.7	18.1 ± 5.8	8.3 ± 3.0

Table 2. Table showing individual 75% thresholds for all three types of moving dot pattern for each subject. Table also demonstrates average values ± standard deviation.



Figure 2.







Figure 4.



Figure 5.



