

This is a repository copy of *Gambling problems among patients in primary care:a cross-sectional study of general practices*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/118855/>

Version: Accepted Version

Article:

Cowlshaw, Sean, Gale, Lone, Gregory, Alison et al. (2 more authors) (2017) Gambling problems among patients in primary care:a cross-sectional study of general practices. The British journal of general practice : the journal of the Royal College of General Practitioners. e274-e279. ISSN: 1478-5242

<https://doi.org/10.3399/bjgp17X689905>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 Running head: GAMBLING PROBLEMS IN PRIMARY CARE

2

3

4

5

6

7

8

9 Gambling problems in primary care: A cross-sectional study of general practices

10

11 Cowlshaw, S^{1,2}

12 Gale, L¹

13 Gregory, A¹

14 McCambridge, J³

15 Kessler, D¹

16

17 ¹ Centre for Academic Primary Care, School of Social and Community Medicine, University
18 of Bristol, United Kingdom.

19 ² Centre for Gambling Research, College of Arts and Social Sciences, the Australian National
20 University (ANU), Australia.

21 ³ Department of Health Sciences, University of York, United Kingdom.

22

23 Author for correspondence: Sean Cowlshaw (sean.cowlshaw@bristol.ac.uk).

24

25 Word count: 2,492

Abstract

Background: Primary care is an important context for addressing health-related behaviours, and may provide a setting for identification of gambling problems. **Aim:** To indicate the extent of gambling problems among patients attending general practices, and explore settings or patient groups that experience heightened vulnerability. **Design and Setting:** Cross-sectional study of patients attending 11 general practices in southwest England. **Method:** Adult patients ($n = 1,058$) were recruited from waiting rooms of practices that were sampled on the basis of population characteristics. Patients completed anonymous questionnaires comprising measures of mental health problems (e.g., depression) and addictive behaviours (e.g., risky alcohol use). The Problem Gambling Severity Index (PGSI) measured gambling problems, along with a single-item measure of problems among family members. Estimates of extent and variability according to practice and patient characteristics were produced.

Results: There were 0.9% of all patients exhibiting problem gambling (PGSI 5+), and 4.3% reporting problems that were low to moderate in severity (PGSI 1-4). Around 7% of patients reported gambling problems among family members. Further analyses indicated that rates of any gambling problems (PGSI 1+) were higher among males and young adults, and more tentatively, within a student healthcare setting. They were also elevated among patients exhibiting drug use, risky alcohol use and depression. **Conclusion:** There is need for improved understanding of the burden of, and responses to, gambling problems in general practices, and new strategies to increase identification in order to facilitate improved care and early intervention.

KEY WORDS: gambling, general practice, cross-sectional study, England

48 **How this fits in:** Gambling problems are emerging concerns for public health in the UK, and
49 primary care is a potential context for identifying patients who would benefit from early
50 intervention or specialist services, as such patients may already attend for related reasons.
51 However, there are no data on gambling problems in UK general practices, and this study
52 assessed the extent of these issues, and sought to identify patient groups that may be
53 particularly vulnerable. It suggests that gambling problems are important clinical issues for
54 primary care attenders, with around 1 in 20 patients reporting pasting year problems, which
55 were mostly of low to moderate severity. The findings highlight need for increased
56 acknowledgement and capacities to respond to gambling problems in general practices (e.g.,
57 through training and support for GPs in order to identify patients and help facilitate access to
58 specialist services).

59 Participation in gambling is increasing in the UK, with surveys indicating that around
60 59% of British adults reported gambling activities (excluding National Lottery) in 2010,
61 which was an increase of 7% from 2007 [1]. These trends have occurred in the context of
62 developments in gambling technologies (e.g., electronic gambling machines, online
63 gambling) and increased exposure (for example, gambling-related advertisements grew by
64 almost 500% between 2007 and 2012) [2], and larger numbers of people experiencing
65 problems with gambling [1]. These problems encompass a spectrum of difficulties that are
66 defined mainly by gambling-related harms (e.g., financial crises, relationship breakdown) [3],
67 and can sometimes reach levels of severity that warrant diagnoses of pathological gambling
68 or gambling disorder (in the ICD-10 [4] and DSM-5 [5], respectively). Prevalence studies
69 indicate that around 7% of men (2% of women) experience at least some problems with
70 gambling annually in the UK, with higher levels among young adults (e.g., 17% of males
71 aged 16-24 reported at least some problems in 2012) [6]. There is also a socio-economic
72 gradient of risk, whereby elevated risk of gambling problems is associated with low income
73 and high deprivation [7].

74 Gambling problems cluster with other health-related behaviours [8], and are
75 associated with anxiety disorders and psychosomatic complaints, and high rates of suicidal
76 ideation and attempts [9]. These problems are also associated with overuse of health-care
77 services, with problem gamblers being twice as likely to consult their GP for mental health
78 concerns, five times as likely to be hospital inpatients, and eight times as likely to access
79 psychological counselling, when compared to people with no such problems [9]. However,
80 help-seeking for gambling is infrequent and usually crisis-driven [10], and thus occurs only
81 after experiencing severe gambling-related harms. Accordingly, there is a strong need for
82 initiatives to increase help-seeking and early intervention, and these include new means of
83 identification and response within generalist healthcare settings.

Primary care is an established context for addressing health-related behaviours (e.g., alcohol misuse) [11], and may be an important setting for identification of problematic gambling [12]. High use of services [9] suggests overrepresentation of gambling problems in primary care, and particularly within practices that serve vulnerable populations. This is supported by U.S. data suggesting rates of gambling disorders ranging from 6% [13] to 15% [14] among primary care attenders (relative to estimates from population-based studies that range from 0.2-1.0%) [15], and higher levels within low income populations [14]. It is already recommended that UK GPs screen high risk groups (e.g., those reporting financial problems), and refer cases for specialist treatment [12]. This is notwithstanding the lack of any evaluation of gambling problems in UK general practices, whereby the prevalence of conditions remains unknown. In this context, the aims of this project were to:

- 1) Provide data on the extent of gambling problems among patients attending general practices in England;
- 2) Explore variability according to practice and patient characteristics, and thus indicate clinical settings or patient groups that experience heightened vulnerability.

Methods

Participants and procedure

The target population comprised patients attending general practices in the Bristol region of southwest England. Eleven practices were purposively sampled according to population deprivation and patient characteristics, as follows: (1) deprivation levels were quantified using data from the Office for National Statistics, which indicated four practices from deprived areas (top 30% for deprivation in England), two practices in areas of low deprivation (bottom 30%), and three practices in a moderate band (middle 40% for deprivation); (2) one practice provided care to young adults in a student health service, and

one practice provided services to a homeless population. The latter were targeted to assess risk according to key population sub-groups.

Patients aged over 18 years and attending practices for any reason were eligible, but were excluded if they were unable to understand English, required immediate medical attention, or were unable to give consent. Patients were approached by a researcher in waiting rooms before appointments, and were provided with information about the study. Those who provided consent were given anonymous questionnaires. These were self-completed and returned in the waiting room or using pre-paid envelopes, and yielded $n = 1,058$ questionnaires. Across practices sampled according to deprivation, the patient numbers ranged from $n = 58$ to $n = 122$. There was $n = 17$ and $n = 163$ participants recruited from the practice for homeless patients and the student health service, respectively. Socio-demographic characteristics are shown in Table 1.

TABLE 1

Measures

Brief measures identified mental health concerns and addictive behaviours. These included the 2-item Whooley scale for depression [16], and the GAD-2 [17] scale for anxiety, which are recommended in primary care [18]. Risky alcohol use was measured using the three consumption items from the Alcohol Use Disorders Identification Test (AUDIT-C) [19, 20]. Non-prescription and recreational drug use was assessed using a Single-Item Screening Question (SISQ) for unhealthy drug use [21]. The format of this item, which required numeric indications of number of times (in the past year) using an illegal drug or prescription medication for non-medical reasons, was modified and comprised a binary response (*yes* or *no*) indicating any past year usage.

Gambling frequency was assessed using items derived from the British Gambling Prevalence Surveys [1], and asked about purchases of lottery or instant win / scratch tickets,

play on bingo, casino table games, slot machines and other electronic gambling machines, games of skill against other individuals, or betting money on sporting events. These items used past year timeframes (0 = *Never*, 6 = *4+ times a week*), along with an item about any other gambling. Patients reporting gambling were then asked to complete the *Problem Gambling Severity Index (PGSI)* [22], which consists of 9-items scored on 4-point response scales (0 = *Never*, 3 = *Almost always*) that relate to past year experiences. The study used a criterion of PGSI 5+ for problem gambling (which has been shown to yield greatest classification accuracy relative to clinician ratings involving detailed case conceptualisations) [23], with scores of PGSI 1-4 indicating low to moderate severity problems (given all such respondents were demonstrating at least some signs of problematic gambling). There was a single-item about whether family members or close relatives had ever had problems with gambling, which was adapted from epidemiological surveys [24] and had a binary response format.

Data analyses

Data-file preparation was conducted using SPSS Version 21, while analyses were conducted using Program R. These comprised descriptive analyses of rates of gambling problems and other mental health concerns and addictive behaviours. Exploratory analyses of variability according to practice characteristics were conducted, followed by evaluations of associations with patient-level characteristics. These comprised Pearson χ^2 -tests and logistic regression models that explored significant effects. The latter specified gambling problems as endogenous variables, and with patient characteristics treated as exogenous. These were evaluated in separate models, which thus estimated bivariate associations through Odds Ratios (ORs) and 95% Confidence Intervals (CIs).

Results

Preliminary analyses indicated modest levels of missing data ranging from around 5% (depression) to 13% (alcohol) across most measures, and were managed through pairwise deletion. However, there were higher levels for the PGSI, with around 45% of eligible participants (i.e., reporting gambling in the past year) having missing data across items. Exploratory analyses indicated around 90% of these patients that reported gambling on lottery or with instant win tickets only, and suggested that missing data were attributable mainly to such patients failing to define these activities as gambling. Missing data were addressed using zero-fill techniques, and thus assumed no gambling problems.

Table 2 indicates frequencies of gambling problems and mental health problems or addictive behaviours. There were around 1% of patients demonstrating problem gambling (PGSI 5+), and 4% exhibiting problems that were low to moderate in severity (PGSI 1-4). Thus, a total of 5.2% of patients (95% CI = 4.0% to 6.8%) exhibited at least some gambling problems across a spectrum of severity. There were 7.2% of patients reporting gambling problems among family members, and this included eight patients reporting problems with their own gambling (PGSI 1+). Levels were lower than rates of other mental health problems and addictive behaviours.

TABLE 2

Subsequent analyses explored variability in gambling problems (PGSI 1+) according to practice characteristics. Given small numbers of practices in this study, the results (see Table 2) are highly exploratory. However, they yielded trends ($p < .10$) suggesting elevated rates in the student health service, when compared to practices characterised by low (OR = 2.57, 95% CI = 0.99 to 7.47) and moderate deprivation (OR = 2.12, 95% CI = 0.95 to 4.73). Modest elevations were observed for highly deprived practices but were not significantly different when compared to practices characterised by low (OR = 1.91, 95% CI = 0.81 to

5.25) or moderate deprivation (OR = 1.58, 95% CI = 0.95 to 4.73). These deprived practices included the clinic for homeless patients, which were too few for statistical comparison ($n = 17$), but exhibited extremely high rates of gambling problems (29.4%).

Bivariate associations involving any gambling problems (PGSI 1+) and patient characteristics are shown in Table 3. These indicated significant associations with gender, age and relationship status. Logistic regression illustrated higher rates among: males (compared to females: OR = 2.55, 95% CI = 1.44 to 4.55), patients aged 18-24 years (compared to 35-44 year olds: OR = 2.43, 95% CI = 1.21 to 5.06), and patients who were single / never married (compared to married or cohabitating: OR = 2.35, 95% CI = 1.32 to 4.29). Patients screening positive for depression demonstrated a 2-fold increase in rates of gambling problems (OR = 2.08, 95% CI = 1.15 to 3.94), while risky alcohol use was associated with a near 3-fold increase (OR = 2.78, 95% CI = 1.60 to 4.89). Drug use was associated with a 5-fold increase in gambling problems (OR = 5.03, 95% CI = 2.78 to 8.99).

Discussion

Summary

The results indicated around 5% of patients reporting problems with gambling across a spectrum of severity, including approximately 1% who were problem gamblers (PGSI 5+), and 4% reporting problems that were low to moderate in severity (PGSI 1-4). There were around 7% reporting gambling problems among family members or close relatives, and were also likely to encounter gambling-related harms [25]. These rates were lower than other mental health concerns (e.g., depression: 56%) and addictive behaviours (e.g., risky alcohol use: 32%) that have stronger traditions of recognition in primary care. Notwithstanding, the study indicated groups and perhaps clinical contexts that were characterised by heightened vulnerability. There were high rates among males and young adults (the extent of any gambling problems among males aged 18-24 was 25.4%, 95% CI = 15.6% to 38.2%), and

208 more tentatively, within the student healthcare setting. Gambling problems were elevated
209 among patients demonstrating drug use, alcohol risk and depression.

210 *Strengths and limitations*

211 The study involved purposive sampling of practices, and recruitment of a sample that
212 was a reasonable approximation of patients encountered regularly in primary care. However,
213 the number of practices was small and participants were not randomly sampled, while data on
214 response rates were not recorded. Findings may be affected by refusals to participate and
215 missing data, which was high for the gambling problem measure. This comprised the PGSI
216 [22], which does not assess the full breadth of gambling-related harms [26]. To reduce
217 burden, the study used a single-item measure of gambling problems among family members,
218 while clinical characteristics were measured using brief screens that possess moderate
219 specificity [20, 27], and do not correspond to severe mental health concerns and addiction
220 problems.

221 *Comparison with existing literature*

222 Estimates of the extent of gambling problems were lower than those in prior research
223 from the U.S. [13, 14], and are similar to levels in population-based studies in the UK [1].
224 Notwithstanding, the present findings highlight that gambling problems are important clinical
225 issues for primary care attenders, that are strongly linked with poor mental health [9] and
226 have major impacts that extend beyond the individual [28]. There is evidence that people
227 with gambling problems can benefit from therapeutic interventions, including intensive and
228 brief interventions [29, 30], and alongside minimal interventions for ‘concerned significant
229 others’ [31]. These provide the basic components of an intervention framework that aligns
230 with models of care for alcohol misuse, and comprises multiple tiers of intervention [32].
231 These address a spectrum of severity (e.g., simple advice or brief interventions for hazardous

or harmful drinking, intensive therapies for dependence), as well as support needs of families, and have bases in identification strategies that are situated within primary care [11].

Implications for research and practice

The study indicates around 1 in 20 patients that report some degree of gambling problem in routine primary care, and highlights need for improved acknowledgement and capacities to respond to these issues. It supports the recommendation that GPs and clinical staff should be vigilant for gambling problems [12], and particularly among young males and patients who are depressed or using alcohol and drugs. At a minimum, there should be training and support for clinical staff in identification and pathways to care. However, in the absence of visible signs of gambling problems that are low to moderate in severity, it seems unlikely that such strategies (which exclude questioning in the absence of visible risk factors) will identify many individuals who would benefit from early intervention. As such, it may also be that selective screening [33] of high risk groups (e.g., depressed and/or young males), or within particular contexts (e.g., university clinics), are potentially appropriate.

There is need for further evidence that indicates the burden of gambling problems in primary care at a national level, and particularly illustrating co-occurrence and impacts on other presenting problems. The development of strategies to identify gambling problems is associated with particular research needs, including studies which demonstrate that initiatives can yield improved access to interventions, and also that patients in primary care, who are not seeking help for gambling, will benefit from interventions. Finally, these identification strategies can only be justified if adequate services are available to deliver interventions. It appears that such requirements are lacking in the UK, where intervention research for gambling is virtually non-existent, while treatment services are grossly inadequate [34]. Such inadequacies are notwithstanding the best efforts of service providers (which mainly comprise voluntary sector organisations), and can be attributed to an unusual situation in the

257 UK whereby research and treatment are commissioned almost exclusively by gambling
258 industry affiliated bodies. Given that between 15-40% of most gambling revenues (depending
259 on type of activity) [35] is derived from people reporting problems with gambling in the UK,
260 there are conflicts of interest between public health and economic policy goals (whereby even
261 small reductions in numbers of people gambling heavily implies far larger reductions in
262 economic yield) [36]. Because of the vested interests of addiction industries [36, 37],
263 evidence and interventions that are supported through independent funding are needed.
264 Gambling should be formally recognised as a health-related issue in the UK, and included
265 within the remits of mainstream commissioning bodies that are responsible for public health
266 and service provision.
267

Additional Information

Funding: This project was funded by the NIHR School for Primary Care Research (SPCR).

Ethical approval: Ethical approval for the study was granted by the NHS Health Research

Authority (HRA), IRAS project ID: 192004, REC reference: 16/WA/0055.

Competing interests: The authors have no competing interests to declare.

Acknowledgements: We would like to thank Graham England and Sarah Flourentzou from

the Bristol Addiction Recovery Agency (ARA) for supporting the project, and the

management teams of general practices that hosted the study. The project was supported by

an advisory committee comprising Jody Clarke (Bristol City Council), Graham England

(Bristol ARA), Graham Box (patient/public involvement representative) and Gail Thornton

(patient/public involvement representative). Finally, we are extremely grateful to the patients

who also participated in the study.

285

References

- 286 1. Wardle H, Griffiths MD, Orford J, Moody A, Volberg R. Gambling in Britain: A time
287 of change? Health implications from the British Gambling Prevalence Survey 2010.
288 Int J Ment Health Addict. 2012;10:273-7.
- 289 2. Ofcom. *Trends in advertising activity – Gambling*. Independent Regulator and
290 Competition Authority for the UK Communications Industries (Ofcom). 2013.
291 Available from: [http://stakeholders.ofcom.org.uk/binaries/research/tv-](http://stakeholders.ofcom.org.uk/binaries/research/tv-research/Trends_in_Ad_Activity_Gambling.pdf)
292 [research/Trends_in_Ad_Activity_Gambling.pdf](http://stakeholders.ofcom.org.uk/binaries/research/tv-research/Trends_in_Ad_Activity_Gambling.pdf)
- 293 3. Delfabbro P. Problem and pathological gambling: A conceptual review. J Gambl Bus
294 Econ. 2013;7:35-53.
- 295 4. World Health Organization. The ICD-10 classification of mental and behavioural
296 disorders: Clinical descriptions and diagnostic guidelines. Geneva: Author; 1992.
- 297 5. American Psychiatric Association. The Diagnostic and Statistical Manual of Mental
298 Disorders. 5th ed. Washington: Author; 2013.
- 299 6. Wardle H, Seabury C. Gambling behaviour. In: Craig R, Mindell J, editors. Health
300 Survey for England 2012: Health, Social Care and Lifestyles. London: Health and
301 Social Care Information Centre; 2013.
- 302 7. Orford J, Wardle H, Griffiths M, Sproston K, Erens B. The role of social factors in
303 gambling: Evidence from the 2007 British Gambling Prevalence Survey. Community,
304 Work & Family. 2010;13:257-71.
- 305 8. Goodyear-Smith F, Arroll B, Kerse N, Sullivan S, Coupe N, Tse S, et al. Primary care
306 patients reporting concerns about their gambling frequently have other co-occurring
307 lifestyle and mental health issues. BMC Fam Pract. 2006;7:25.
- 308 9. Cowlishaw S, Kessler D. Problem gambling in the UK: Implications for health,
309 psychosocial adjustment and health care utilization. Eur Addict Res. 2016;22:90-98.
- 310 10. Evans L, Delfabbro PH. Motivators for change and barriers to help-seeking in
311 Australian problem gamblers. J Gambl Stud. 2005;21:133-55.
- 312 11. O'Flynn N. Harmful drinking and alcohol dependence: Advice from recent NICE
313 guidelines. Br J Gen Pract. 2011;61:754-56.
- 314 12. George S, Gerada C. Problem gamblers in primary care: Can GPs do more? Br J Gen
315 Pract. 2011;61:248-49.
- 316 13. Pasternak AV, Fleming MF. Prevalence of gambling disorders in a primary care
317 setting. Arch Fam Med. 1999;8:515-20.
- 318 14. Morasco BJ, Vom Eigen KA, Petry NM. Severity of gambling is associated with
319 physical and emotional health in urban primary care patients. Gen Hosp Psychiatry.
320 2006;28:94-100.
- 321 15. Petry NM, Blanco C. National gambling experiences in the United States: Will history
322 repeat itself? Addiction. 2013;108:1032-37.
- 323 16. Whooley MA, Avins AL, Miranda J, Browner WS. Case-finding instruments for
324 depression. J Gen Intern Med. 1997;12:439-45.
- 325 17. Kroenke K, Spitzer RL, Williams JB, Monahan PO, Lowe B. Anxiety disorders in
326 primary care: Prevalence, impairment, comorbidity, and detection. Ann Intern Med.
327 2007;146:317-25.
- 328 18. Kendrick T, Pilling S. Common mental health disorders—identification and pathways
329 to care: NICE clinical guideline. Br J Gen Pract. 2012;62:47-49.
- 330 19. Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR.
331 AUDIT-C as a brief screen for alcohol misuse in primary care. Alcohol Clin Exp Res.
332 2007;31:1208-17.

- 333 20. Rumpf HJ, Hapke U, Meyer C, John U. Screening for alcohol use disorders and at-
334 risk drinking in the general population: Psychometric performance of three
335 questionnaires. *Alcohol Alcohol*. 2002;37:261-68.
- 336 21. McNeely J, Cleland CM, Strauss SM, Palamar JJ, Rotrosen J, Saitz R. Validation of
337 Self-Administered Single-Item Screening Questions (SISQs) for unhealthy alcohol
338 and drug use in primary care patients. *J Gen Intern Med*. 2015;30:1757-64.
- 339 22. Ferris J, Wynne H. The Canadian Problem Gambling Index. Ottawa: Canadian Centre
340 on Substance Abuse; 2001.
- 341 23. Williams RJ, Volberg RA. The classification accuracy of four problem gambling
342 assessment instruments in population research. *International Gambling Studies*.
343 2014;14:15-28.
- 344 24. Salonen AH, Castren S, Alho H, Lahti T. Concerned significant others of people with
345 gambling problems in Finland: A cross-sectional population study. *BMC Public*
346 *Health*. 2014;14:398.
- 347 25. Li E, Browne M, Rawat V, Langham E, Rockloff M. Breaking bad: Comparing
348 gambling harms among gamblers and affected others. *J Gambl Stud*. 2016. Epub 2016
349 July 21. DOI:10.1007/s10899-016-9632-8.
- 350 26. Langham E, Thorne H, Browne M, Donaldson P, Rose J, Rockloff M. Understanding
351 gambling related harm: A proposed definition, conceptual framework, and taxonomy
352 of harms. *BMC Public Health*. 2016;16:80.
- 353 27. Bosanquet K, Bailey D, Gilbody S, Harden M, Manea L, Nutbrown S, McMillan D.
354 Diagnostic accuracy of the Whooley questions for the identification of depression: A
355 diagnostic meta-analysis. *BMJ Open*. 2015;5:e008913.
- 356 28. Cowlishaw S, Suomi A, Rodgers B. Implications of gambling problems for family
357 and interpersonal adjustment: Results from the Quinte Longitudinal Study. *Addiction*.
358 2016;111:1628-36.
- 359 29. Cowlishaw S, Merkouris S, Dowling N, Anderson C, Jackson A, Thomas S.
360 Psychological therapies for pathological and problem gambling. *Cochrane Db Syst*
361 *Rev*. 2012;11. doi:10.1002/14651858.CD008937.pub2.
- 362 30. Yakovenko I, Hodgins DC. Latest developments in treatment for disordered
363 gambling: Review and critical evaluation of outcome studies. *Current Addiction*
364 *Reports*. 2016;3:299-306.
- 365 31. Hodgins DC, Toneatto T, Makarchuk K, Skinner W, Vincent S. Minimal treatment
366 approaches for concerned significant others of problem gamblers: A randomized
367 controlled trial. *J Gambl Stud*. 2007;23:215-30.
- 368 32. Department of Health. Models of care for alcohol misusers (MoCAM). London; NHS
369 National Treatment Agency for Substance Misuse: 2006.
- 370 33. O'Doherty LJ, Taft A, Hegarty K, Ramsay J, Davidson LL, Feder G. Screening
371 women for intimate partner violence in healthcare settings: Abridged Cochrane
372 systematic review and meta-analysis. *BMJ*. 2014;348:g2913.
- 373 34. George S, Copello A. Treatment provision for Britain's problem gamblers: Present
374 gaps and future opportunities. *Advances in Psychiatric Treatment*. 2011;17:318-22.
- 375 35. Orford J, Wardle H, Griffiths M. What proportion of gambling is problem gambling?
376 Estimates from the 2010 British Gambling Prevalence Survey. *International Gambling*
377 *Studies*. 2013;13:4-18.
- 378 36. Adams PJ. Moral jeopardy: Risks of accepting money from tobacco, alcohol and
379 gambling industries. Cambridge; Cambridge University Press: 2016.
- 380 37. Babor T, Hall W, Humphreys K, Miller P, Petry N, West R. Who is responsible for
381 the public's health? The role of the alcohol industry in the WHO global strategy to
382 reduce the harmful use of alcohol. *Addiction*. 2013;108:2045-47.

Table 1. Sample socio-demographic characteristics ($n = 1,058$).

	<i>n</i>	%
Gender (female)	636	64.7%
Age		
18-24	211	20.7%
25-34	154	15.1%
35-44	137	13.4%
45-64	284	27.8%
65+	235	23.0%
Relationship status		
Single (never married)	341	33.3%
Married / living with partner	526	51.4%
DSW / other	156	15.2%
Education		
Secondary school or less	270	27.0%
Post-secondary school education	627	62.6%
Postgraduate education	60	6.0%
Other	44	4.4%
Employment		
Employed	398	39.33%
Unemployed	126	12.45%
Retired	226	22.33%
Student	166	16.40%
Other	96	9.49%
Ethnicity (white)	889	87.67%

NB: Due to small amounts of missing data, patient numbers across categories may not aggregate to 100%.

Table 2. Estimates of the extent of mental health problems and addictive behaviours, including gambling problems.

		<i>n</i>	%	95% CI	
				LB	UB
Gambling					
	PGSI 5+	10	0.9%	0.5%	1.8%
	PGSI 1-4	45	4.3%	3.2%	5.7%
	Problems in the family	73	7.2%	5.7%	9.0%
Mental health / addictive behaviours					
	Depression (Whooley 1+)	561	55.8%	52.7%	58.9%
	Anxiety (GAD-2 2+)	262	27.0%	24.3%	30.0%
	Alcohol (AUDIT-C 5+)	307	32.4%	29.4%	35.5%
	Drug use (SISQ)	140	14.3%	12.2%	16.7%
PGSI 1+ across practice characteristics					
	High deprivation ($k = 4$, $n = 380$)	23	6.1%	4.0%	9.1%
	Moderate deprivation ($k = 3$, $n = 331$)	13	3.9%	2.2%	6.8%
	Low deprivation ($k = 2$, $n = 184$)	6	3.3%	1.3%	7.3%
	Student health service ($k = 1$, $n = 163$)	13	8.0%	4.5%	13.5%

NB: Whooley = Whooley depression scale, GAD-2 = 2-item GAD scale for anxiety, AUDIT-C = 3-item consumption scale from the AUDIT, SISQ = single-item screening question for unhealthy drug use.

Table 3. Analyses of associations with any gambling problems (PGSI 1+) and patient-level socio-demographic and clinical characteristics.

		<i>n</i>	%	χ^2	<i>p</i>
Socio-demographic characteristics					
Gender	Male	29	8.4%	10.0	0.002
	Female	22	3.5%		
Age	18-24	22	10.4%	18.2	0.001
	23-44	8	5.2%		
	35-44	4	2.9%		
	45-64	13	4.6%		
	65+	5	2.1%		
Relationship status	Single, never married	29	8.5%	11.9	0.003
	Married/cohabitating	20	3.8%		
	DSW/other	4	2.6%		
Education	Secondary school or less	9	3.3%	2.6	0.272
	Post-secondary school education	37	5.9%		
	Postgraduate/other	6	5.8%		
Employment	Employed	21	5.3%	7.7	0.052
	Unemployed	11	8.7%		
	Student	11	6.6%		
	Retired/other	9	2.8%		
Ethnicity	White	45	5.1%	0.0	0.969
	Non-white	7	5.6%		
Clinical characteristics					
Depression	Whooley (1+)	38	6.8%	5.1	0.024
	Whooley (0)	15	3.4%		
Anxiety	GAD-2 (3+)	19	7.3%	2.3	0.127
	GAD-2 (<3)	32	4.5%		
Alcohol	AUDIT-C (5+)	30	9.8%	12.9	0.000
	AUDIT -C (<5)	24	3.7%		
Drug use	SISQ Yes	22	15.7%	32.8	0.000
	SISQ No	30	3.6%		

NB: Positive endorsement of either item from the Whooley was used to indicate possible depression. Scores of 3+ on the GAD-2 were used to indicate potential anxiety. Scores of 5+ on the AUDIT-C were used to indicate high risk (including hazardous and harmful) drinking.