UNIVERSITY of York

This is a repository copy of *Isocyanurate Formation During Oxazolidinone Synthesis from Epoxides and Isocyanates Catalysed by a Chromium(Salphen) Complex.*

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/id/eprint/118975/</u>

Version: Accepted Version

Article:

Wu, Xiao orcid.org/0000-0002-3089-9200, Mason, Jess and North, Michael orcid.org/0000-0002-6668-5503 (2017) Isocyanurate Formation During Oxazolidinone Synthesis from Epoxides and Isocyanates Catalysed by a Chromium(Salphen) Complex. Chemistry : A European Journal. pp. 12937-12943. ISSN: 0947-6539

https://doi.org/10.1002/chem.201702948

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Isocyanurate Formation During Oxazolidinone Synthesis from Epoxides and Isocyanates Catalysed by a Chromium(Salphen) Complex

Xiao Wu,^[a] Jess Mason^[a] and Michael North*^[a]

Abstract: Chromium(salphen) complex **10** is found to be a catalyst for the preparation of oxazolidinones from epoxides and isocyanates. Using the optimal reaction conditions (1.5 mol% of chromium(salphen) complex **10** at 80 °C in toluene for 4 hours), six epoxides were reacted with five isocyanates, providing **15** oxazolidinones in up to 90% yield. With electron-deficient isocyanates, cyclotrimerisation of the isocyanate to the corresponding isocyanurates is a competing reaction, showing the importance of matching catalyst activity to that of the substrates.

Introduction

Epoxides **1** are useful building blocks in organic synthesis and a number of five-membered heterocycles can be prepared when they are treated with heterocumulenes. The reaction with carbon dioxide to furnish either cyclic- or poly-carbonates **2** and **3** has been extensively studied.^{1,2,3} Other areas of interest include the formation of either cyclic dithiocarbonates **4** or trithiocarbonates **5**,⁴ and oxazolidinones **6** and **7**⁵ (Scheme 1).

chemistry,⁶ as well as being valuable synthetic intermediates.⁷ The reaction of epoxides with isocyanates is one of the most useful and efficient methods for the synthesis of oxazolidinones (Scheme 1). A number of catalyst systems have been reported for this reaction, including ammonium salts,⁸ lanthanide salts,⁹ lithium halides,¹⁰ magnesium halides,¹¹ tetraphenylantimony iodide¹² and trialkyltin halides.¹³

Recently, our group has reported a number of highly efficient metal-based catalysts **8–10** for the formation of cyclic carbonates from epoxides and carbon dioxide (Figure 1).^{14,15,16} Compounds **8** and **9** also exhibit excellent catalytic activity for the preparation of oxazolidinones.^{5a,b} Thus, 5 mol% of bimetallic aluminium(salen) complex **8** in the absence of a co-catalyst was found to be an active catalytic system for the synthesis of a range of mono- and di-substituted oxazolidinones. In the presence of tetrabutylammonium bromide (5 mol%), aluminium heterosorpionate complex **9** (5 mol%) was also shown to be an effective catalyst for the preparation of oxazolidinones. Six epoxides were reacted with six aromatic isocyanates to give 25 oxazolidinones, showing a broad substrate scope.



Scheme 1. Synthesis of heterocycles from epoxides and heterocumulenes.

Oxazolidinones are an important class of heterocycles and have been found to be useful compounds in medicinal

 [a] Dr. X. Wu, Miss J. Mason, Professor M. North Green Chemistry Centre of Excellence, Department of Chemistry University of York
 Heslington, York, YO10 5DD
 E-mail: Michael.north@york.ac.uk

Supporting information for this article is given via a link at the end of the document.



Figure 1. Catalysts for the synthesis of cyclic carbonates and oxazolidinones.

More recently, our group has reported the synthesis of cyclic carbonates from epoxides and carbon dioxide using chromium(salphen) complex **10**.¹⁶ In this case, only 1.5 mol% of complex **10** was necessary to catalyse the synthesis of a range of cyclic carbonates, either from monosubstituted epoxides at room temperature and one bar carbon dioxide pressure, or from disubstituted epoxides at elevated temperature and pressure. In view of the high catalytic activity shown by complex **10**, we decided to investigate its use in the related reaction between epoxides and isocyanates. Herein, we report the use of chromium(salphen) complex **10** as an efficient catalyst for the formation of oxazolidinones from epoxides and isocyanates.

Results and Discussion

Screening studies were carried out using styrene oxide 1a and phenyl isocyanate 11a in a 1:1 ratio under various reaction conditions (Scheme 2) and the results are shown in Table 1. Initial experiments were undertaken using 1.5 mol% of both chromium(salphen) complex 10 and tetrabutylammonium bromide (TBAB) as a co-catalyst at 80 °C for 24 hour under solvent-free conditions (Table 1, entry 1). However, the reaction mixture solidified after 30 minutes. This suggested that a solvent was necessary to increase the conversion to the oxazolidinone products. Toluene was selected based on literature precedent^{5a,b} and the reaction mixture was heated to 80 °C. After 24 hours, 75% conversion was obtained (Table 1, entry 2). Control experiments proved to be very informative. In the absence of TBAB, chromium(salphen) complex 10 was able to catalyse the reaction itself and a 90% conversion was achieved, whereas in the absence of complex 10, only 7% conversion was observed (Table 1, entries 3 and 4). Increasing the catalyst loading to 2.5 mol% was not beneficial while lowering the catalyst loading to 0.5 mol% resulted in low conversion (Table 1, entries 5 and 6). To further optimise the reaction conditions, a sample was taken after each hour and it was found that the reaction reached maximum conversion after 4 hours. It was therefore decided that 1.5 mol% of catalyst 10 and heating in toluene at 80 °C for 4 hours were the optimal reaction conditions for the synthesis of oxazolidinones from epoxides and isocyanates.



Scheme 2. Reaction of styrene oxide 1a with phenylisocyanate 11a

Table	1.	Optimisation	of	the	synthesis	of	oxazolidinones	6a	and	7a	using
comple	ex -	10 and TBAB.									

					27	
Entry	Complex 10 (mol%)	TBAB (mol%)	t (h)	Toluene (mL)	Conversion ^[a] (%)	6a:7b ^[a]
1	1.5	1.5	24	0	70	50:50
2	1.5	1.5	24	0.5	75	66:34
3	1.5	0	24	0.5	90	35:65
4	0	1.5	24	0.5	7	75:25
5	2.5	0	24	0.5	88	50:50
6	0.5	0	24	0.5	28	50:50
7	1.5	0	4	0.5	90	35:65

[a] Conversion of epoxide **1a** into **6a** and **7a** and ratio of **6a:7a** were determined by ¹H NMR spectroscopy of the unpurified reaction mixture.

Having determined the optimal reaction conditions for the synthesis of oxazolidinones 6a and 7a from styrene oxide 1a and phenyl isocyanate 11a, a series of epoxides was used together with phenyl isocyanate 11a to give oxazolidinones 6b-g and 7b-g (Scheme 3). The results of this study are shown in Table 2. Functionalised aliphatic epoxides (3phenoxypropropylene oxide 1b and epichlorohydrin 1c) and unfunctionalised aliphatic epoxide 1d were all excellent substrates, giving high conversions and excellent regioselectivity to the 3,5-isomers 6b-d (Table 2, entries 2-4). In contrast, glycidol 1e was found to be a poor substrate, giving a complex mixture from which no product could be isolated (Table 2, entry 5). When aromatic epoxides 1f and 1g, containing a halogen at the para-position were used as substrates, good conversions were obtained though with no regioselectivity (Table 2, entries 6 and 7).



1,6,7. a: $R^1 = Ph$; b: $R^1 = CH_2OPh$; c: $R^1 = CH_2CI$; d: $R^1 = C_8H_{17}$; e: $R^1 = CH_2OH$; f: $R^1 = 4$ -CIC₆H₄; g: $R^1 = 4$ -BrC₆H₄

Scheme 3. Reaction of epoxide	s 1a-g with phenylisocyanate 11a.
-------------------------------	-----------------------------------

able 2. Synthesis of oxazolidinone 6a-g and 7a-g using epoxide 1a-g. ^[a]					
Entry	Epoxide	Conversion ^[b] (%)	Yield ^[c] (%)		
	1a R^1 = Ph	90 (35:65)	6a,7a (65)		
2	1b $R^1 = CH_2OPh$	80 (1:0)	6b (62)		
3	1c $R^1 = CH_2CI$	100 (1:0)	6c (90)		
Ļ	1d $R^1 = C_8 H_{17}$	100 (92:8)	6d (59)		
5	1e R^1 = CH ₂ OH	0	-		
6	$\mathbf{1f} \operatorname{R}^1 = 4\operatorname{-}\operatorname{ClC}_6\operatorname{H}_4$	75 (1:1)	6f , 7f (59)		
7	$\mathbf{1g} \ R^1 = 4\text{-}BrC_6H_4$	70 (1:1)	6g, 7g (62)		

[a] Reactions were carried out using 1.5 mol% chromium(salphen) complex **10** in toluene at 80 °C for 4 hours. [b] Determined by ¹H NMR spectroscopy of the unpurified reaction mixture and figure in brackets is the ratio of **6:7**. [c] Yield of isolated product after purification by column chromatography.

To further expand the substrate scope, styrene oxide **1a** and epichlorohydrin **1c** were selected as representative aromatic and aliphatic epoxides respectively for the synthesis of oxazolidinones from various isocyanates (Scheme 4). Five substituted aromatic isocyanates **11b-f** and two aliphatic isocyanates **11g** and **11h** were reacted with these epoxides and

the results are summarised in Table 3. It was found that aromatic isocyanates 11b,c gave good to excellent conversions after 4 hours. A 1:1 ratio of 3,4- and 3,5- regioisomers was observed when styrene oxide 1a was used, while with epichlorohydrin 1c, complete regioselectivity for the 3,5-isomer was achieved (Table 3, entries 1-5 and 8-12). In comparison, no conversion was obtained when aliphatic isocyanates 11g,h were used (entries 6,7 and 13,14). With aromatic isocyanates that have halogen groups at the para-position, chromium(salphen) complex 10 was found to catalyse the formation of both oxazolidinones and isocyanurates (perhydro-1,3,5-triazine-2,4,6triones) 12 (Figure 2) (Table 3, entries 3-5). Isocyanurates are known to enhance the physical properties of polyurethanes and coating materials,¹⁷ typically by increasing their flame retardation and filming characteristics, and commercial products containing polymeric isocyanurates have increased chemical and thermal resistance.18



6,7. h,o: $R^2 = 4$ -MeOC₆H₄; i,p: $R^2 = 4$ -MeC₆H₄; j,q: $R^2 = 4$ -FC₆H₄; k,r: $R^2 = 4$ -ClC₆H₄; l,s: $R^2 = 4$ -BrC₆H₄; m,t: $R^2 = CH_2Ph$; n,u: $R^2 = Et$;

11. a: $R^2 = Ph$; b: $R^2 = 4$ -OMeC₆H₄; c: $R^2 = 4$ -MeC₆H₄; d: $R^2 = 4$ -FC₆H₄; e: $R^2 = 4$ -ClC₆H₄; f: $R^2 = 4$ -BrC₆H₄; g: $R^2 = CH_2Ph$; h: $R^2 = Et$

Scheme 4. Reaction of epoxides 1a,c with isocyanate 11a-h.



Figure 2. Structure of isocyanurates 12 and urea 13.

Isocyanurates have previously been prepared by the cyclotrimerisation of isocyanates catalysed by species including organotin¹⁹ and organozinc compounds,²⁰ copper and nickel halides²¹ and palladium,²² magnesium²³ and iron complexes.⁰ The formation of isocyanurates **12a-c** was much less important when the competing reaction was between epichlorohydrin **1c** and isocyanates than when styrene oxide **1a** was used. However, since the separation of oxazolidinones **6q-s** from

isocyanurates **12a-c** proved difficult, the isolated yields of these oxazolidinones were less than satisfactory (Table 3, entries 10-12). The formation of isocyanurates as by-products from the reaction between epoxides and isocyanates has not previously been reported. A control experiment using 4-fluorophenyl isocyanate **11d** in the presence of chromium complex **10** under the standard reaction conditions but in the absence of an epoxide also gave isocyanurate **12a**, indicating that chromium complex **10** is a strong Lewis acid and capable of catalysing the cyclotrimerisation of isocyanates. In the absence of chromium complex **10**, only the starting material 4-fluorophenyl isocyanate was recovered. When isocyanates that have an electron-donating substituent at the *para*-position such as **11b** were used, the corresponding urea **13** was isolated rather than the isocyanurate.

Table 3. Synthesis of oxazolidinones 6,7h-u using complex 10. ^[a]						
	Conv ^[b]	1.1				

Entry	Epoxide	Isocyanate	(%)	Yield ^[c] (%)
1	1a R ¹ = Ph	11b $R^2 = 4$ -MeOC ₆ H ₄	60 (1:1)	6h, 7h (56)
2	1a R ¹ = Ph	11c $R^2 = 4$ -MeC ₆ H ₄	90 (1:1)	6i , 7i (62)
3	1a R ¹ = Ph	11d $R^2 = 4-FC_6H_4$	30 (1:1)	7j (12) ^[d]
4	1a R ¹ = Ph	11e $R^2 = 4 - CIC_6H_4$	22 (1:1)	7k (10) ^[e]
5	1a R ¹ = Ph	11f $R^2 = 4$ -BrC ₆ H ₄	23 (1:1)	7I (10) ^[f]
6	1a R ¹ = Ph	11g $R^2 = CH_2Ph$	0	
7	1a R ¹ = Ph	11h R ² = Et	0	
8	$1c R^1 = CH_2CI$	11b $R^2 = 4$ -MeOC ₆ H ₄	98 (1:0)	60 (80)
9	$\textbf{1c} \ \textbf{R}^1 = \textbf{C}\textbf{H}_2\textbf{C}\textbf{I}$	11c $R^2 = 4$ -MeC ₆ H ₄	96 (1:0)	6p (83)
10	$\textbf{1c} \ \textbf{R}^1 = \textbf{C}\textbf{H}_2\textbf{C}\textbf{I}$	11d $R^2 = 4-FC_6H_4$	90 (1:0)	6q (21) ^[g]
11	$\textbf{1c} \ \textbf{R}^1 = \textbf{C}\textbf{H}_2\textbf{C}\textbf{I}$	11e $R^2 = 4 - CIC_6H_4$	85 (1:0)	6r (42)
12	$\textbf{1c} \ \textbf{R}^1 = \textbf{C}\textbf{H}_2\textbf{C}\textbf{I}$	11f $R^2 = 4$ -BrC ₆ H ₄	90 (1:0)	6s (36)
13	$\textbf{1c} \ \textbf{R}^1 = \textbf{C}\textbf{H}_2\textbf{C}\textbf{I}$	11g $R^2 = CH_2Ph$	0	
14	1c R ¹ = CH ₂ Cl	11h $R^2 = Et$	0	

[a] Reactions were carried out using 1.5 mol% of chromium(salphen) complex 10 in toluene at 80 °C for 4 hours. [b] Conversion of epoxides to oxazolidinones 6,7 determined by ¹H NMR spectroscopy of the unpurified reaction mixture and figure in brackets is the ratio of 6:7. [c] Yield of isolated product after purification by column chromatography. [d] Mixture of oxazolidinone 7j and isocyanurate 12a in 40:60 ratio. [e] Mixture of oxazolidinone 7l and isocyanurates 12c in 50:50 ratio. [g] Mixture of oxazolidinone 6g and isocyanurate 12a in 95:5 ratio.

The chemo- and regio-selectivities observed when using complex **10** as a catalyst can be explained by the catalytic cycles shown in Scheme 5. Complex **10** can act both as a Lewis

acid and as a source of a good nucleophile (chloride). Substrates 1 and 11 possess three sites which are susceptible to nucleophilic attack: both ends of the epoxide and the central carbon atom of the isocyanate. They also possess two sites for Lewis acid activation: the epoxide oxygen and the isocyanate oxygen. For oxazolidinone formation, the chromium(salphen) species first acts as a Lewis acid, activating the epoxide towards ring-opening by chloride anion. The regioselectivites observed with epoxides 1a-g are consistent with ring-opening of the activated epoxide normally occurring at the less-hindered, unsubstituted position by an S_N2 type mechanism, eventually giving the 3,5-isomer of the oxazolidinone. However, when R¹ is an aromatic group capable of stabilising a benzylic carbenium ion, then ring-opening of the epoxide at the more-hindered, substituted position ultimately leading to the 3,4-isomer of the oxazolidinone becomes competitive and little or no regioselectivity is observed. The chromium coordinated alkoxide then reacts with the isocvanate to form a carbamate and this ring closes to give oxazolidinone 6 or 7 with regeneration of catalyst 10. For isocyanurate formation, the isocyanate oxygen atom coordinates competitively to the metal centre, followed by the nucleophilic attack of the chloride anion to give a chromium complexed carbomovl chloride complex A. Following this, the nucleophilic nitrogen can then react further with two molecules of isocyanates before ring-closing with elimination of chloride to form isocyanurates 12 and regenerate catalyst 10.

Conclusions

Chromium(salphen) complex 10 was found to catalyse the formation of oxazolidinones from epoxides and isocyanates in the absence of a co-catalyst. Compared to previously reported aluminium(salen) complexes, chromium(salphen) complex 10 appears to be more Lewis acidic. This allows the catalyst loading to be reduced to 1.5 mol% and the reaction time to be reduced to 4 hours. However, the enhanced Lewis acidity of complex 10 causes the reaction to take a different pathway with electron-deficient isocyanates, leading to the formation of isocyanurates 12 by a pathway that involves Lewis acid activation and subsequent trimerisation of the isocyanate rather than Lewis acid activation of the epoxide. These results illustrate that although a strongly Lewis acidic catalyst is desirable to activate the epoxide for the formation of oxazolidinones from epoxides and isocyanates, the optimal catalyst for the process is substrate dependent and requires the electronic properties of both substrates and catalyst to be matched.

Experimental Section

Catalyst **10** was prepared as previously reported.¹⁶ All other compounds were commercially available and used as supplied. ¹H NMR and ¹³C NMR were recorded in CDCl₃ at 25 °C on a JEOL 400 spectrometer operating at 400 and 100 MHz, respectively. Melting points were determined using a Stuart

SMP3 apparatus. Infrared spectra were recorded on a Bruker Vertex 70 instrument.



Scheme 5. Proposed catalytic cycles for the synthesis of oxazolidinones and isocyanurates induced by chromium(salphen) complex 10.

General procedure for oxazolidinone synthesis

Epoxide **1a-g** (0.83 mmol) and isocyanate **11a-h** (0.83 mmol) were added to a solution of catalyst 10 (0.013 mmol) in toluene (0.5 mL). The resulting mixture was stirred at 80 °C for 4 hours. After being allowed to cool to room temperature, toluene was removed under pressure to give the crude oxazolidinone products. The conversion of epoxide to oxazolidinone was determined by ¹H NMR spectroscopy of the crude mixture. The products were purified by flash chromatography to give compound **6/7 a-s**.

3,5-Diphenyloxazolidin-2-one (**6a**): Obtained as a yellow solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (5:1), then hexane–EtOAc (3:1). (85 mg, 43%). M.p. 80–82 °C (lit. 79–82 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.56 (2H, d, *J* 8.0 Hz, ArH), 7.46–7.37 (7H, m, ArH), 7.15 (1H, t, *J* 8.0 Hz, ArH), 5.65 (1H, d, *J* 8.0, CHO), 4.39 (1H, t, *J* 8.0 Hz, CH₂N), 3.98 (1H, t, *J* 8.0 Hz, CH₂N); ¹³C (100

3,4-Diphenyloxazolidin-2-one (**7a**): Obtained as a yellow solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (5:1), then hexane–EtOAc (3:1). (44 mg, 22%). M.p. 80–82 °C (lit. 79–82 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.28 (9H, m, ArH), 7.07 (1H, t, *J* 8.0 Hz, ArH), 7.15 (1H, t, *J* 7.4 Hz, ArH), 5.40 (1H, dd, *J* 8.0, 6.0 Hz, CHN), 4.79 (1H, t, *J* 8.0 Hz, CH₂O), 4.21 (1H, dd, *J* 8.0, 6.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 155.9, 138.2, 137.0, 129.4, 128.9, 128.8, 126.2, 120.8, 69.8, 60.7. Mass spec. (ESI): calcd. m/z 240.1019 [C₁₅H₁₄NO₂]⁺; found: 240.1024. calcd. m/z 262.0838 [C₁₅H₁₃NO₂+Na]⁺; found: 262.0842. IR (neat, cm⁻¹): 1741.

3-Phenyl-5-phenoxymethyloxazolidin-2-one (**6b**): Obtained as a yellow solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (3:1), then hexane–EtOAc (2:1) (218 mg, 93%). M.p. 138–141 °C (lit. 139–140 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.58 (2H, d, *J* 8.0 Hz, ArH), 7.40 (2H, t, *J* 8.0 Hz, ArH), 7.30 (2H, t, *J* 8.0 Hz, ArH), 7.16 (1H, t, *J* 8.0 Hz, ArH), 7.00 (1H, t, *J* 8.0 Hz, ArH), 6.91 (2H, d, *J* 8.0 Hz, ArH), 5.09–4.96 (1H, m, CHO), 4.24–4.19 (3H, m, CH₂O, CH₂N), 4.09 (1H, dd, *J* 8.0, 4.0 Hz, CH₂N); ¹³C (100 MHz, CDCl₃) δ 158.0, 154.4, 138.1, 129.6, 129.1, 124.2, 121.7, 118.3, 114.6, 70.3, 67.8, 47.4. Mass spec. (ESI): calcd. m/z 292.0944 [C₁₆H₁₅NO₃+Na]⁺; found: 292.0939. IR (neat, cm⁻¹): 1732.

3-Phenyl-5-chloromethyloxazolidin-2-one (6c): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of hexane–EtOAc (2:1) (158 mg, 90%). M.p. 108–110 °C (lit. 101–103 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.56 (2H, m, ArH), 7.40 (2H, t, *J* 8.0 Hz, ArH), 7.17 (1H, t, *J* 8.0 Hz, ArH), 4.91–4.85 (1H, m, CHO), 4.18 (1H, t, *J* 8.0, Hz, CH₂N), 3.97 (1H, dd, *J* 8.0, 4.0 Hz, CH₂N), 3.81 (1H, dd, J 12.0, 4.0 Hz, CH₂Cl); ¹³C (100 MHz, CDCl₃) δ 153.9, 137.7, 129.2, 124.4, 118.3, 70.8, 48.2, 44.5. Mass spec. (ESI): calcd. m/z 234.0292 [C₁₀H₁₀CINO₂+Na]⁺; found: 234.0296. IR (neat, cm⁻¹): 1728.

3-Phenyl-5-octyloxazolidin-2-one (**6d**): Obtained as a white solid after purification by flash chromatography using a solvent system of hexane–EtOAc (9:1) (135 mg, 59%). M.p. 71–73 °C (lit. 70–71 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.54 (2H, d, *J* 8.0 Hz, ArH), 7.37 (2H, t *J* 8.0 Hz, ArH), 7.13 (1H, t, *J* 8.0 Hz, ArH), 4.66–4.61 (1H, m, CHO), 4.08 (1H, t, *J* 8.0 Hz, CH₂N), 3.66 (1H, dd, *J* 8.0, 7.5 Hz, CH₂N), 1.91–1.68 (2H, m, CH₂), 1.42–1.22 (12H, m, 6×CH₂), 0.90 (3H, t, J 8.0 Hz, CH₃); ¹³C (100 MHz, CDCl₃) δ 155.0, 138.4, 129.0, 123.9, 118.1, 73.1, 50.5, 35.0, 31.8, 29.4, 29.2, 29.1, 24.5, 22.6, 14.1. Mass spec. (ESI): Calcd. m/z 298.1777 [C₁₇H₂₅NO₂+Na]⁺; found:298.1767. IR (neat, cm⁻¹): 1717.

3-Phenyl-5-(4-chlorophenyl)-oxazolidin-2-one (**6f**): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of CH₂Cl₂-hexane (4:1) (68 mg, 30%). M.p. 124–125 °C (lit. 124–126 °C).^[5b] ¹H NMR (400 MHz, CDCl₃) δ 7.55 (2H, d, *J* 8.0 Hz, ArH), 7.43–7.36 (6H, m, ArH), 7.16 (1H, t, *J* 8.0 Hz, ArH), 5.63 (1H, t, *J* 8.0 Hz, CHO), 4.39 (1H, t, *J* 8.0 Hz, CH₂N), 3.93 (1H, dd, *J* 8.0, 7.5 Hz, CH₂N); ¹³C (100 MHz, CDCl₃) δ 154.4, 137.9, 136.6, 135.0, 129.3, 129.1, 127.1, 124.3, 118.3, 73.3, 52.6. Mass spec. (ESI): Calcd. m/z 296.0449 [C₁₅H₁₂CINO₂+Na]⁺; found: 296.0449. IR (neat, cm⁻¹): 1733.

3-Phenyl-4-(4-chlorophenyl)-oxazolidin-2-one (**7f**): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of CH₂Cl₂-hexane (4:1) (66 mg, 29%). M.p. 140–143 °C (lit. 140–142 °C).^[5b] ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.24 (8H, m, ArH), 7.11–7.07 (1H, m, ArH), 5.38 (1H, dd, *J* 12.0, 8.0 Hz, CHN), 4.78 (1H, t, *J* 12.0 Hz, CH₂O), 4.17 (1H, dd, *J* 8.0, 4.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 155.7, 139.4, 136.7, 134.9, 129.6, 129.0, 127.6, 124.9, 120.9, 69.6, 60.1. Mass spec. (ESI): calcd. m/z 274.0629 [C₁₅H₁₃CINO₂]⁺; found: 274.0633. Calcd. m/z 296.0449 [C₁₅H₁₂CINO₂+Na]⁺; found: 296.0448. IR (neat, cm⁻¹): 1733.

3-Phenyl-5-(4-bromophenyl)-oxazolidin-2-one (**6g**): Obtained as a white solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (7:1), then hexane–EtOAc (4:1), then hexane–EtOAc (3:1), (77 mg, 30%). M.p 132–134 °C (lit. 132–134 °C).^[5b] ¹H NMR (400 MHz, CDCl₃) δ 7.58–7.53 (4H, m, ArH), 7.41–7.37 (2H,m, ArH), 7.32–7.29 (2H, m, ArH), 7.18–7.14 (1H, m, ArH), 5.61 (1H, t, *J* 8.0 Hz, CHO), 4.39 (1H, t, *J* 8.0 Hz, CH₂N), 3.92 (1H, dd, *J* 8.0, 7.5 Hz, CH₂N); ¹³C (100 MHz, CDCl₃) δ 154.3, 137.8, 137.1, 132.2, 129.2, 127.3, 124.4 123.2, 118.3, 73.3, 52.5. Mass spec. (ESI): calcd. m/z 339.9944 [C₁₅H₁₂BrNO₂+Na]⁺; found: 339.9940. IR (neat, cm⁻¹): 1726.

3-Phenyl-4-(4-bromophenyl)-oxazolidin-2-one (**7g**): Obtained as a yellow solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (7:1), then hexane–EtOAc(4:1), then hexane–EtOAc (3:1), (82 mg, 32%). M.p. 157–158 °C (lit. 156–158 °C).^[5b] ¹H NMR (400 MHz, CDCl₃) δ 7.49 (2H, d, *J* 8.0 Hz, ArH), 7.37–7.25 (4H, m, ArH), 7.19 (2H, d, *J* 8.0, ArH), 7.10 (1H, m, ArH), 5.37 (1H, dd, *J* 8.0, 6.0 Hz, CHN), 4.78 (1H, t, *J* 8.0 Hz, CH₂O), 4.17 (1H, dd, *J* 8.0, 6.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 155.7, 137.2, 132.6, 129.2, 129.0, 127.9 125.0, 120.8, 116.5, 69.5, 60.1. Mass spec. (ESI): calcd. m/z 339.9944 [C₁₅H₁₂BrNO₂+Na]⁺; found: 339.9943. IR (neat, cm⁻¹): 1728.

3-(4-Methoxyphenyl)-5-phenyloxazolidin-2-one (6h): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of hexane:EtOAc (5:1) (66 mg, 28%). M.p. 100–102 °C (lit. 105–107 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.46–7.38 (7H, m, ArH), 6.93–6.91 (2H, m, ArH), 5.63 (1H, t, *J* 8.0 Hz, CHO), 4.35 (1H, t, *J* 8.0 Hz, CH₂N), 3.94 (1H, dd, *J* 12.0, 8.0 Hz, CH₂N), 3.80 (3H, s, OCH₃); ¹³C (100 MHz, CDCl₃) δ 156.4, 155.0, 138.2, 131.3, 129.0, 128.9, 125.6, 120.3, 114.3, 74.0, 55.5, 53.3. Mass spec. (ESI): calcd. m/z 292.0944 [C₁₆H₁₅CINO₂+Na]⁺; found: 292.0933. IR (neat, cm⁻¹): 1732.

 ^{13}C (100 MHz, CDCl₃) δ 156.8, 156.4, 138.1, 129.8, 129.2, 128.8, 126.5, 123.3, 114.1, 69.7, 61.3, 55.3. Mass spec. (ESI): calcd. m/z 2792.0944 [C16H15CINO2+Na]^+; found: 292.0934. IR (neat, cm^{-1}): 1739.

3-(4-Methylphenyl)-5-phenyloxazolidin-2-one (**6i**): Obtained as a white solid after purification by flash chromatography using a solvent system of hexane–EtOAc (9:1) (68 mg, 32%). M.p. 96–98 °C (lit. 98–100 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.43–7.41 (7H, m, ArH), 7.18 (2H, d, *J* 8.0 Hz, ArH), 5.62 (1H, t, *J* 8.0 Hz, CHO), 4.35 (1H, t, *J* 8.0 Hz, CH₂N), 3.94 (1H, dd, J 8.0, 7.0 Hz, CH₂N), 2.32 (3H, s, CH₃); ¹³C (100 MHz, CDCl₃) δ 153.9, 141.7, 129.5, 129.3, 128.4, 128.3, 126.6, 121.3, 110.0, 74.0, 52.6, 20.8. Mass spec. (ESI): calcd. m/z 276.0995 [C₁₆H₁₅NO₂+Na]⁺; found: 279.0991. IR (neat, cm⁻¹): 1735.

3-(4-Methylphenyl)-4-phenyloxazolidin-2-one (**7i**): Obtained as a white solid after purification by flash chromatography using a solvent system of hexane–EtOAc (9:1) (63 mg, 30%). M.p. 106–108 °C (lit. 105–107 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.36–7.20 (7H, m, ArH), 7.03 (2H, d, *J* 8.0 Hz, ArH), 5.35 (1H, dd, *J* 8.0, 6.0 Hz, CHN), 4.76 (1H, t, *J* 8.0 Hz, CH₂O), 4.19 (1H, dd, *J* 8.0, 4.0 Hz, CH₂O), 2.23 (3H, s, CH₃); ¹³C (100 MHz, CDCl₃) δ 156.2, 138.3, 134.5, 134.3, 129.5, 129.3, 128.8, 126.3, 121.1, 69.8, 60.8, 20.8. Mass spec. (ESI): calcd. m/z 276.0995 [C₁₆H₁₅NO₂+Na]⁺; found: 279.1005. IR (neat, cm⁻¹): 1739.

3-(4-Fluorophenyl)-4-phenyloxazolidin-2-one (**7j**): Obtained as a white solid in a mixture with **12a** after purification by flash chromatography using a solvent system of first hexane–EtOAc (9:1), then hexane– EtOAc (5:1) (28 mg, 12%). ¹H NMR (400 MHz, CDCl₃) δ 7.4–7.25 (7H, m, ArH), 6.95–6.93 (2H, m, ArH), 5.33 (1H, dd, *J* 12.0, 8.0 Hz, CHN), 4.79 (1H, t, *J* 8.0 Hz, CH₂O), 4.22 (1H, dd, *J* 8.0, 6.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 161.5, 156.1, 137.8, 133.0, 129.2, 129.0, 126.4, 123.0 (d, *J* 8 Hz), 115.7 (d, *J* 23 Hz), 69.8, 61.1; ¹⁹F NMR (376 MHz, CDCl₃): δ 117.1 (s). Mass spec. (ESI): calcd. m/z 280.0744 [C₁₅H₁₂FNO₂+Na]⁺; found: 280.0716.

3-(4-Chlorophenyl)-4-phenyloxazolidin-2-one (**7k**): Obtained as a pale yellow solid in a mixture with **12b** after purification by flash chromatography using a solvent system of hexane–EtOAc (9:1) (22 mg, 10%). ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.29 (7H, m, ArH), 7.21 (2H, d, *J* 8.0 Hz, ArH), 5.36 (1H, dd, *J* 12.0, 8.0 Hz, CHN), 4.79 (1H, t, *J* 8.0 Hz, CH₂O), 4.21 (1H, dd, *J* 8.0, 6.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 155.8, 137.7, 135.6, 129.7, 129.6, 128.9, 126.2, 121.9, 69.8, 60.6. Mass spec. (ESI): calcd. m/z 296.0449 [C₁₅H₁₂CINO₂+Na]⁺; found: 296.0453.

3-(4-Bromophenyl)-4-phenyloxazolidin-2-one (**7I**): Obtained as a yellow solid in a mixture with **12c** after purification by flash chromatography using a solvent system of hexane–EtOAc (9:1) (26 mg, 10%). M.p. 132–135 °C (lit. 134–137 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.37 (9H, m, ArH), 5.35 (1H, dd, *J* 8.0, 6.0 Hz, CHN), 4.79 (1H, t, *J* 8.0 Hz, CH₂O), 4.21 (1H, dd, *J* 8.0, 6.0 Hz, CH₂O); ¹³C (100 MHz, CDCl₃) δ 156.2, 132.7, 131.8, 130.0, 129.5, 128.5, 126.0, 123.7, 122.2, 67.2, 61.8. Mass spec. (ESI): calcd. m/z 318.0124 [C₁₅H₁₂BrNO₂+H]⁺; found: 318.0130.

3-(4-Methoxyphenyl)-5-chloromethyloxazolidin-2-one (60): Obtained as a yellow solid after purification by flash chromatography using a solvent system of hexane–EtOAc (3:1) (160 mg, 80%). M.p. 106–108 °C (lit. 105–106 °C).^[5a] ¹H NMR $\begin{array}{l} (400 \text{ MHz, CDCl}_3) \ \delta \ 7.43 \ (2H, \ d, \ J \ 8.0 \ Hz, \ ArH), \ 6.92 \ (2H, \ d, \ J \ 8.0 \ Hz, \ ArH), \ 4.89-4.83 \ (1H, \ m, \ CHO), \ 4.17-4.10 \ (2H, \ m, \ CH_2N), \ 3.79 \ (3H, \ s, \ OCH_3), \ 3.78-3.72 \ (2H, \ m, \ CH_2Cl); \ ^{13}C \ (100 \ MHz, \ CDCl_3) \ \delta \ 156.6, \ 154.2, \ 130.9, \ 120.4, \ 114.3, \ 70.8, \ 55.5, \ 48.7, \ 44.6. \ Mass \ spec. \ (ESI): \ calcd. \ m/z \ 264.0398 \ [C_{11}H_{12}CINO_3+Na]^+; \ found: \ 264.0395. \ IR \ (neat, \ cm^{-1}): \ 1728. \end{array}$

3-(4-Methylphenyl)-5-chloromethyloxazolidin-2-one (**6p**): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of hexane–EtOAc (3:1) (170 mg, 83%). M.p. 104–106 °C (lit. 104–107 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.42 (2H, d, *J* 8.0 Hz), 7.19 (2H, d, *J* 8.0 Hz), 4.89–4.82 (1H, m, CHO), 4.15 (1H, t, *J* 8.0 Hz, CH₂N), 3.92 (1H, dd, *J* 12.0, 8.0 Hz, CH₂N), 3.89 (1H, dd, *J* 12.0, 4.0 Hz, CH₂Cl), 3.73 (1H, dd, *J* 12.0, 8.0 Hz, CH₂Cl), 2.32 (3H, s, CH₃); ¹³C (100 MHz, CDCl₃) δ 154.0, 135.2, 134.1, 129.6, 118.4, 70.8, 48.3, 44.5, 20.7. Mass spec. (ESI): calcd. m/z 226.0629 [C₁₁H₁₃CINO₂+H]⁺; found: 226.0630. IR (neat, cm⁻¹): 1732.

3-(4-Fluorophenyl)-5-chloromethyloxazolidin-2-one (6q): Obtained as a pale yellow solid after purification by flash chromatography using a solvent system of hexane–EtOAc (5:1) (125 mg, 67%). ¹H NMR (400 MHz, CDCl₃) δ 7.51 (2H, dd, *J* 8.0, 4.0 Hz, ArH), 7.09 (2H, t, *J* 8.0 Hz, ArH), 4.91–4.85 (1H, m, CHO), 4.16 (1H, t, *J* 8.0 Hz, CH₂N), 3.96 (1H, dd, *J* 8.0, 4.0 Hz, CH₂N), 3.81 (1H, dd, *J* 12.0, 4.0 Hz, CH₂Cl), 3.75 (1H, dd, *J* 12.0, 8.0 Hz, CH₂Cl); ¹³C (100 MHz, CDCl₃) δ 159.5 (d, J 244.5 Hz), 154.0, 133.9 (d, J 3.2 Hz), 120.2 (d, J 8.1 Hz), 115.9 (d, J 22.6 Hz), 70.8, 48.4, 44.5. ¹⁹F NMR(376 MHz, CDCl₃): δ 117.8 (s). Mass spec. (ESI): calcd. m/z 252.0198 [C₁₀H₉CIFNO₂+Na]⁺; found: 252.0195.

3-(4-Chlorophenyl)-5-chloromethyloxazolidin-2-one (**6r**): Obtained as an orange solid after purification by flash chromatography using a solvent system of hexane-EtOAc (1:1) (100 mg, 42%). M.p. 129-132 °C (lit. 130-133 °C).^{[5a] 1}H NMR (400 MHz, CDCl₃) δ 7.49 (2H, d, J 8.0 Hz, ArH), 7.34 (2H, d, J 8.0 Hz, ArH), 4.91-4.85 (1H, m, CHO), 4.14 (1H, t, J 8.0 Hz, CH₂N), 3.93 (1H, dd, J 8.0, 4.0 Hz, CH₂N), 3.79 (1H, dd, J 12.0, 4.0 Hz, CH₂Cl), 3.75 (1H, dd, *J* 12.0, 8.0 Hz, CH₂Cl); ¹³C (100 MHz, CDCl₃) & 153.7, 136.3, 129.6, 129.1, 119.4, 70.8, 48.0, 267.9903 44.5. Mass spec. (ESI⁺): calcd.m/z [C₁₀H₉Cl₂NO₂+Na]⁺; found: 267.9905. IR (neat, cm⁻¹): 1741.

3-(4-Bromophenyl)-5-chloromethyloxazolidin-2-one (6s): Obtained as a yellow solid after purification by flash chromatography using a solvent system of first hexane–EtOAc (3:1), then hexane–EtOAc (2:1) (92 mg, 36%). M.p. 126–128 °C (lit. 125–128 °C).^[5a] ¹H NMR (400 MHz, CDCl₃) δ 7.51–7.44 (4H, m, ArH), 4.92–4.86 (1H, m, CHO), 4.15 (1H, t, *J* 8.0 Hz, CH₂N), 3.94 (1H, dd, *J* 8.0, 4.0 Hz, CH₂N), 3.80 (1H, dd, *J* 12.0, 4.0 Hz, CH₂Cl), 3.75 (1H, dd, *J* 12.0, 8.0 Hz, CH₂Cl); ¹³C (100 MHz, CDCl₃) δ 153.7, 132.7, 132.1, 119.7, 117.3, 70.8, 48.0, 44.4. Mass spec. (ESI+): calcd. m/z 311.9397 [C₁₀H₉BrCINO₂+Na]⁺; found: 311.9389. IR (neat, cm⁻¹): 1732.

Keywords: oxazolidinone • isocyanurate • chromium • epoxide • catalysis

 For recent reviews of cyclic carbonate syntheses see: a) C. Martín, G. Fiorani, A. W. Kleij, ACS Catal. 2015, 5, 1353; b) A. C. Kathalikkattil, R. Babu, J. Tharun, R. Roshan, D.-W. Park, *Catal. Surv. Asia* 2015, *19*, 223; c) B. Yu, L.-N. He, *ChemSusChem* 2015, *8*, 52; d) M. Cokoja, M. E. Wilhelm, M. H. Anthofer, W. A. Herrmann, F. E. Kühn, *ChemSusChem* 2015, *8*, 2436; e) B.-H. Xu, J.-Q. Wang, J. Sun, Y. Huang, J.-P. Zhang, X.-P. Zhang, S.-J. Zhang, *Green Chem*. 2015, *17*, 108; f) V. D'Elia, J. D. A. Pelletier, J.-M. Basset, *ChemCatChem* 2015, *7*, 1906.

- [2] For recent reviews of polycarbonate syntheses see: a) S. Klaus, M. W. Lehenmeier, C. E. Anderson, B. Rieger, *Coord. Chem. Rev.* 2011, *255*, 1460; b) M. R. Kember, A. Buchard, C. K. Williams, *Chem. Commun.* 2011, *47*, 141; c) X.-B. Lu, W.-M. Ren, G.-P. Wu, *Acc. Chem. Res.* 2012, *45*, 1721; d) D. J. Darensbourg, S. J. Wilson, *Green Chem.* 2012, *14*, 2665; e) N. Ikpo, J. C. Flogeras, F. M. Kerton, *Dalton Trans.* 2013, *42*, 8998; f) D. J. Darensbourg, A. D. Yeung, *Polym. Chem.* 2014, *5*, 3949; g) M. Taherimehr, P. P. Pescarmona, *J. Appl. Polym. Sci.* 2014, *131*, 41141; h) Y. Qin, X. Sheng, S. Liu, G. Ren, X. Wang, F. J. Wang, J. *CO2 Util.* 2015, *11*, 3; i) S. Paul, Y. Zhu, C. Romain, R. Brooks, P. K. Saini, C. K. Williams, *Chem. Commun.* 2015, *51*, 6459.
- For recent reviews covering cyclic and polycarbonate syntheses see: a)
 R. Martín, A. W. Kleij, *ChemSusChem* 2011, *4*, 1259; b) X.-B. Lu, D. J.
 Darensbourg, *Chem. Soc. Rev.* 2012, *41*, 1462; c) W. N. R. Wan
 Isahak, Z. A. Che Ramli, M. W. Mohamed Hisham, M. A. Yarmo, *Renew. Sust. Energy Rev.* 2015, *47*, 93.
- a) J. A. Durden, H. A. Stansbury, W. H. Catlette, J. Am. Chem. Soc. [4] 1960, 82, 3082; b) C. G. Overberger, A. Drucker, J. Org. Chem. 1964, 29, 360; c) S. Hayashi, M. Fukurawa, Y. Fujino, T. Nakao, K. Nagato, Chem. Pharm. Bull. 1971, 19, 1594; d) G. E. McCasland, A. B. Zanlungo, L. J. Durham, J. Org. Chem. 1974, 39, 1462; e) Y. Taguchi, K. Yanagiya, I. Shibuya, Y. Suhara, Bull. Chem. Soc. Jpn. 1988, 61, 921; f) Y. Taguchi, M. Yasumoto, I. Shibuya, Y Suhara, Bull. Chem. Soc. Jpn. 1989, 62, 474; j) N. Kihara, Y. Nakawaki, T. Endo, J. Org. Chem. 1995, 60, 473; h) R. Maggi, C. Malmassari, C. Oro, R. Pela, G. Sartori, L. Soldi, Synthesis 2008, 53; i) I. Yavari, M. Ghazanfarpour-Darjani, Z. Hossaini, M. Sabbaghan, N. Hosseini, Synlett 2008, 889; j) A. Z. Halimehjani, F. Ebrahimi, N. Azizi, M. R. Saidi, J. Heterocycl. Chem. 2009, 46, 347; k) W. Clegg, R. W. Harrington, M. North, P. Villuendas, J. Org. Chem. 2010, 75, 6201; I) M. North, P. Villuendas, Synlett 2010, 623; m) J. Cao, M. Yu, H. Li, L. Wang, X. Zhu, G. Wang, Y. Shi, C. Cao, Res. Chem. Intermed. 2015, 41, 5323; n) C. Beattie, M. North, ChemCatChem 2014, 6, 1252.
- [5] a) T. Baronsky, C. Beattie, R. W. Harrington, R. Irfan, M. North, J. G. Osende, C. Young, ACS Catal. 2013, *3*, 790; b) J. A. Castro-Osma, A. Earlam, A. Lara-Sánchez, A. Otero, M. North, *ChemCatChem* 2016, *8*, 2100; c) C. Beattie, M. North, *RSC Adv.* 2014, *4*, 31345.
- [6] a) G. S. K. K. Reddy, A. Ali, M. N. L. Nalam, S. G. Anjum, H. Cao, R. S. Nathans, C. A. Schiffer, T. M. Rana, *J. Med. Chem.* 2007, *50*, 4316; b) N. Fresno, M. Macías-González, A. Torres-Zaguirre, M. Romero-Cuevas, P. Sanz-Camacho, J. Elguero, F. J. Pavón, F. Rodríguez de Fonseca, P. Goya, R. Pérez-Fernández, *J. Med. Chem.* 2015, *58*, 6639.
- a) W. R. Roush, R. A. James, *Aust. J. Chem.* 2002, *55*, 141; b) J. A.
 Birrell, E. N. Jacobsen, *Org. Lett.* 2013, *15*, 2895; c) V.Laserna, W. Guo,
 A. W. Kleij, *Adv. Synth. Catal.* 2015, *357*, 2849.
- [8] G. P. Speranza and W. J. Peppel, J. Org. Chem. 1958, 23, 1922.
- a) C. Qian, D. Zhu, Synlett 1994, 129; b) H.-Y. Wu, J. -C. Ding, Y.-K.
 Liu, J. Indian Chem. Soc. 2003, 80, 36; c) M. T. Barros, A. M. F. Phillips, Tetrahedron: Asymmetry 2010, 21, 2746.
- [10] a) L. Aroua, A. Baklouti, *Synth. Commun.* 2007, *37*, 1935; b) J. E. Herweh, W. J. Kauffman, *Tetrahedron Lett.* 1971, *12*, 809; c) J. E. Herweh, *J. Heterocycl. Chem.* 1968, *5*, 687; d) J. E. Herweh, T. A. Foglia, D. Swern, *J. Org. Chem.* 1968, *33*, 4029; e) H. Siegel, H. Wittmann, *Monatsh. Chem.* 1982, *113*, 1005.
- [11] X. Zhang, W. Chen, C. Zhao, C. Li, X. Wu, W. Z. Chen, Synth. Commun. 2010, 40, 3654.
- [12] a) A. Baba, M. Fujiwara, H. Matsuda, *Tetrahedron Lett.* **1986**, *27*, 77; b)
 M. Fujiwara, A. Baba, H. Matsuda, *J. Heterocycl. Chem.* **1988**, *25*,

1351; c) M. Fujiwara, A. Baba, H. Matsuda, *Bull. Chem. Soc. Jpn.* **1990**, *63*, 1069.

- [13] a) I. Shibata, A. Baba, H. Iwasaki, H. Matsuda, J. Org. Chem. 1986, 51, 2177; b) M. Fujiwara, A. Baba, Y. Tomohisa, H. Matsuda, Chem. Lett. 1986, 1963; c) A. Baba, K. Seki, H. Matsuda, J. Heterocycl. Chem. 1990, 27, 1925; d) K. Yano, N. Amishiro, A. Baba, H. Matsuda, Bull. Chem. Soc. Jpn. 1991, 64, 2661.
- [14] For a review of compound **8** see: M. North, *ARKIVOC* **2012**, (part (i)), 610.
- [15] A. Otero, A. Lara-Sánchez, J. Fernández-Baeza, C. Alonso-Moreno, J. Tejeda, J. A. Castro-Osma, I. Márquez-Segovia, L. F. Sánchez-Barba, A. M. Rodríguez, M. V. Gómez, *Chem. Eur. J.* 2010, *16*, 8615.
- [16] J. A. Castro-Osma, M. North, X. Wu, Chem. Eur. J. 2016, 22, 2100.
- [17] H. A. Duong, M. J. Cross, J. Louie, *Org. Lett.* **2004**, *6*, 4679.
- [18] a) Z. Wirpsza in *Polyurethanes: Chemistry, Technology and Application*, Ellis Horwood, London, UK, **1993**; b) A. K. Zitinkina, N. A. Sibanova, O. G. Tarakanov, *Russ. Chem. Rev.* **1985**, *54*, 1866; c) T. Nawata, J. E. Kresta, , K. C. J. Frisch, *Cell. Plast.* **1975**, 267; d) L. Nicholas, G. R. J. Gmitter, *Cell. Plast.* **1965**, 85.
- [19] a) A. J. Bloodworth, A. G. Davies, *J. Chem. Soc.* **1965**, 6858; b) S. R.
 Foley, G. P. A. Yap, D. S. Richesom, *Organometallics* **1999**, *18*, 4700.
- [20] Z. Guo, X. Wei, H. Tong, D. Liu, J. Organomet. Chem. 2015, 776, 136.
- [21] J. F. Villa, H. B. Powell, Synth. React. Inorg. Met.-Org. Chem. 1976, 6, 59.
- [22] a) S. G. Lee, K. Y. Choi, Y. J. Kim, S. Park, S. W. Lee, *Dalton Trans.*,
 2015, *44*, 6537; b) F. Paul, S. Moulin, O. Piechaczyk, P. Le Floch, J. A. Osborn, *J. Am. Chem. Soc.* **2007**, *129*, 7294.
- [23] H. R. Sharpe, A. M. Geer, H. E. L. Williams, T. J. Blundell, W. Lewis, A. J. Blake, D. L. Kays, *Chem Commun.* 2017, *53*, 937.

Entry for the Table of Contents (Please choose one layout)

FULL PAPER



A tale of two reactions. A chromimium(salphen) complex catalyses the formation of oxazolidinones from epoxides and isocyanates. However, with particularly electron-deficient isocyanates, the complex catalyses cyclotrimerisation of the isocyanate to an isocyanurate instead.

Xiao Wu, Jess Mason and Michael North*

Page No. – Page No.

Isocyanurate Formation During Oxazolidinone Synthesis from Epoxides and Isocyanates Catalysed by a Chromium(Salphen) Complex