



UNIVERSITY OF LEEDS

This is a repository copy of *Anticommutative extension of the Adler map*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/100480/>

Version: Accepted Version

---

**Article:**

Konstantinou-Rizos, S and Mikhailov, AV (2016) Anticommutative extension of the Adler map. *Journal of Physics A: Mathematical and Theoretical*, 49 (30). 30LT03. pp. 1-7. ISSN 1751-8113

<https://doi.org/10.1088/1751-8113/49/30/30LT03>

---

© 2016 IOP Publishing Ltd. This is an author-created, un-copyedited version of an article accepted for publication/published in *Journal of Physics A: Mathematical and Theoretical*. IOP Publishing Ltd is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at <https://dx.doi.org/10.1088/1751-8113/49/30/30LT03>

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Anticommutative extension of the Adler map

S. Konstantinou-Rizos<sup>\*1,2</sup> and A.V. Mikhailov<sup>†3</sup>

<sup>1</sup>Institute of Mathematical Physics and Seismodynamics, Chechen State University, Russia

<sup>2</sup>Faculty of Mathematics and Computer Technology, Chechen State University, Russia

<sup>3</sup>Department of Applied Mathematics, University of Leeds, UK

May 31, 2016

## Abstract

We construct a noncommutative (Grassmann) extension of the well-known Adler Yang-Baxter map. It satisfies the Yang-Baxter equation, it is reversible and birational. Our extension preserves all the properties of the original map except the involutivity.

## 1 Introduction

The study of the set-theoretical solutions of the Yang-Baxter equation, which was formally proposed by Drinfeld in [4], gained a more algebraic flavour in [3]. The term “Yang-Baxter maps” for such solutions was proposed by Veselov in [13]. Those Yang-Baxter maps which possess Lax representation [12] are of particular interest, since they are associated to integrable mappings [13, 14], and they also possess a natural connection with integrable partial differential equations through Darboux transformations [9]. On the other hand, noncommutative extensions of integrable systems have been constantly of interest over the last few decades. Recently, in [7], Grassmann extensions of Darboux transformations were constructed for a class of nonlinear Schrödinger equations, together with their associated super (Grassmann-extended) differential-difference and difference-difference systems. Following the way proposed in [7], super differential-difference and difference-difference systems were constructed for a noncommutative Darboux transformation for the supersymmetric KdV equation in [15], and a year later for a generalised super KdV system in [16]. Moreover, the study of the extension of the theory of Yang-Baxter maps in noncommutative (Grassmann) setting was initiated in [6], where the first examples of Grassmann extended Yang-Baxter maps were constructed.

In this paper, we construct a noncommutative (Grassmann) extension of the well-celebrated Adler map [1], which, over the last couple of decades, has attracted the interest of many authors in the area of integrable systems. This is due to the fact that, not only it is associated with several concepts of integrability, such as Darboux & Bäcklund transformations, integrable lattice equations and the theory of Yang-Baxter maps, but also it possesses a quite simple form; thus, it constitutes a convenient map in terms of applications. However, the Adler map lacks the property of being non-involutive, which means

---

\*skonstantin84@gmail.com, skonstantin@chesu.ru

†A.V.Mikhailov@leeds.ac.uk

that it is a trivial map in terms of its dynamics. The noncommutative extension of the Adler map, which we present in our paper, has equally elegant form and preserves all the properties of the original map; namely, it satisfies the Yang-Baxter equation, it is reversible and birational. Nonetheless, it has the extra significant property of being non-involutive.

The paper is organised as follows. In the next section, we present the basic properties of Grassmann algebras. For more information on Grassmann analysis one can consult [2]. Moreover, we explain some properties of the Grassmann extended Yang-Baxter maps possessing Lax representation. In section 3, using some recent results on Grassmann extensions of Darboux transformations in the case of a generalised super KdV system [16], we construct a noncommutative extension of the Adler map. Additionally, we show that, in general, noncommutative extensions of involutive Yang-Baxter maps do not preserve their property of involutivity in their noncommutative version. Finally, in section 4, we close with some concluding remarks.

## 2 Preliminaries

### 2.1 Grassmann algebra

Let  $G$  be a  $\mathbb{Z}_2$ -graded algebra over  $\mathbb{C}$  with a unity element  $e$ . As a linear space  $G$  is a direct sum  $G = G_0 \oplus G_1 \pmod{2}$ , such that  $G_i G_j \subseteq G_{i+j}$ . Those elements of  $G$  that belong either to  $G_0$  or to  $G_1$  are called *homogeneous*, the ones in  $G_1$  are called *odd* (or *fermionic*), while those in  $G_0$  are called *even* (or *bosonic*) and  $e \in G_0$ .

By definition, the parity  $|a|$  of an even homogeneous element  $a$  is 0, and it is 1 for odd homogeneous elements. The parity of the product  $|ab|$  of two homogeneous elements is a sum of their parities:  $|ab| = |a| + |b|$ . Grassmann commutativity means that  $ba = (-1)^{|a||b|}ab$  for any homogeneous elements  $a$  and  $b$ . In particular,  $\alpha^2 = 0$ , for all odd elements  $\alpha \in G_1$ , and even elements commute with all the elements of  $G$ .

Lastly, for a square matrix,  $M$ , of the following block-form

$$M = \begin{pmatrix} P & \Pi \\ \Lambda & L \end{pmatrix},$$

where  $P$  and  $L$  are matrices with even entries, while  $\Pi$  and  $\Lambda$  possess only odd entries (the block matrices are not necessarily square matrices), we can define the *supertrace* –and denote it by  $\text{str}(M)$ – to be the following quantity

$$\text{str}(M) = \text{tr}(P) - \text{tr}(L),$$

where  $\text{tr}(\cdot)$  is the usual trace of a matrix. We shall use the supertrace later on to generate invariants of the Grassmann extended Adler map.

### 2.2 Yang-Baxter map and its Lax representation in the Grassmann case

Let  $V_G = \{(a, \alpha) \mid a \in G_0, \alpha \in G_1\}$ . The definitions of Yang-Baxter maps in the Grassmann and commutative cases formally coincide; the only difference is the set of the objects of the maps. In particular, we say that a map  $S \in \text{End}(V_G \times V_G)$

$$((x, \chi), (y, \psi)) \xrightarrow{S} ((u, \xi), (v, \eta)), \quad (1)$$

is a *Grassmann extended Yang-Baxter map* if it satisfies the *Yang-Baxter equation*:

$$S^{12} \circ S^{13} \circ S^{23} = S^{23} \circ S^{13} \circ S^{12}.$$

Here  $S^{ij} \in \text{End}(V_G \times V_G \times V_G)$ ,  $i, j = 1, 2, 3$ ,  $i \neq j$ , are defined by the following relations

$$S^{12} = S \times id, \quad S^{23} = id \times S \quad \text{and} \quad S^{13} = \pi^{12} S^{23} \pi^{12},$$

where  $\pi^{12}$  is the involution defined by  $\pi^{12}((x, \chi), (y, \psi), (z, \zeta)) = ((y, \psi), (x, \chi), (z, \zeta))$ . Moreover, map (1) is called *reversible* if  $S^{21} \circ S^{11} = id$ , where  $S^{21}$  follows from the definition of  $S^{12}$  if we change  $(u, \xi) \leftrightarrow (v, \eta)$  and  $(x, \chi) \leftrightarrow (y, \psi)$ .

Furthermore, we shall be using the term *Grassmann extended parametric Yang-Baxter map* if two parameters  $a, b \in G_0$  are involved in the definition of (1), namely we have a map

$$S_{a,b} : ((x, \chi), (y, \psi)) \mapsto ((u, \xi), (v, \eta)), \quad (2)$$

satisfying the *parametric Yang-Baxter equation*

$$S_{a,b}^{12} \circ S_{a,c}^{13} \circ S_{b,c}^{23} = S_{b,c}^{23} \circ S_{a,c}^{13} \circ S_{a,b}^{12}. \quad (3)$$

Similar to [12], we can consider a problem of refactorisation

$$L_a(u, \xi; \lambda) L_b(v, \eta; \lambda) = L_b(y, \psi; \lambda) L_a(x, \chi; \lambda) \quad (4)$$

where  $L_a(x, \chi; \lambda)$  is a matrix whose entries are Grassmann-valued,  $x, a \in G_0$ ,  $\chi \in G_1$  and  $\lambda$  is a complex variable (a *spectral parameter*). The problem can be formulated as following: for a given set  $\{a, b, x, y \in G_0, \psi, \chi \in G_1\}$  find such  $\{u, v \in G_0, \xi, \eta \in G_1\}$  that equation (4) is satisfied identically in  $\lambda$ . If this problem of refactorisation has a unique solution, then it defines a map (2) which satisfies the YB equation (3) and it is reversible. The proof of the latter statement is exactly the same as in the commutative case [14].

The invariants of a Grassmann-valued Yang-Baxter map can be found using the supertrace. Applying the supertrace to equation (4), we obtain that  $\text{str}(L_b(y, \psi) L_a(x, \chi))$  is a generating function of the invariants associated Yang-Baxter map  $S_{a,b}$ .

### 3 Grassmann extension of the Adler map

A generalised super KdV system

$$\begin{aligned} u_t &= u_{xxx} - 6uu_x + 6\xi_{xx}\eta + 6\eta_{xx}\xi, \\ \xi_t &= 4\xi_{xxx} - 6u\xi_x - 3u_x\xi, \\ \eta_t &= 4\eta_{xxx} - 6u\eta_x - 3u_x\eta, \end{aligned} \quad (5)$$

where  $u = u(x, t)$  is an even variable, whereas  $\xi = \xi(x, t)$  and  $\eta = \eta(x, t)$  are odd, was presented and studied in [8]. The Darboux-Bäcklund transformations and corresponding noncommutative discrete integrable systems associated with (5) were recently constructed in [16]. A Lax matrix-operator associated with (5) is given by the following

$$\mathcal{L} = D_x - \lambda \begin{pmatrix} 0 & 0 & 0 \\ e & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & e & 0 \\ u & 0 & \xi \\ \eta & 0 & 0 \end{pmatrix}, \quad (6)$$

which is the spatial part of the Lax pair for (5). We omit the temporal part of the Lax pair, since we shall not be using it in this text.

A Grassmann extended Darboux transformation associated to the Lax operator (6) is represented by the following matrix (see [16])

$$W = \lambda \begin{pmatrix} 0 & 0 & 0 \\ e & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} \frac{v_1}{2} & e & 0 \\ \frac{v_1^2}{4} - p_1 + \xi\eta_{[1]} & \frac{v_1}{2} & \xi \\ \eta_{[1]} & 0 & e \end{pmatrix}, \quad (7)$$

where  $v_1$  is an even variable,  $p_1 \in \mathbb{C}$  and  $\xi, \eta_{[1]}$  are odd. In what follows, we use matrix (7) to construct a Grassmann extended Yang-Baxter map.

### 3.1 Noncommutative Adler map

According to the Grassmann extended Darboux transformation in (7), changing  $(v_1/2, \xi, \eta_{[1]}; p_1) \rightarrow (x, \chi_1, \chi_2; a)$ , we consider the following matrix

$$M_a(x, \boldsymbol{\chi}) = \lambda M_1 + M_0 = \lambda \begin{pmatrix} 0 & 0 & 0 \\ e & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} x & e & 0 \\ x^2 - a + \chi_1\chi_2 & x & \chi_1 \\ \chi_2 & 0 & e \end{pmatrix}, \quad (8)$$

where  $\boldsymbol{\chi} = (\chi_1, \chi_2)$ , and we substitute it into the Lax equation. Then, we have the following.

**Proposition 3.1.1.** *(Noncommutative Adler map) Let  $x + y$  be an invertible element of the Grassmann algebra. Then, the matrix refactorization problem*

$$M_a(u, \boldsymbol{\xi})M_b(v, \boldsymbol{\eta}) = M_b(y, \boldsymbol{\psi})M_a(x, \boldsymbol{\chi}), \quad (9)$$

where  $M_a = M_a(x, \boldsymbol{\chi})$  is given by (8) is equivalent to a map

$$((x, \boldsymbol{\chi}), (y, \boldsymbol{\psi})) \xrightarrow{S_{a,b}} ((u, \boldsymbol{\xi}), (v, \boldsymbol{\eta})), \quad (10)$$

given by the following

$$x \mapsto u = y + \frac{a - b}{x + y - \chi_1\psi_2}; \quad (11a)$$

$$\chi_1 \mapsto \xi_1 = \psi_1 - \frac{a - b}{x + y}\chi_1; \quad (11b)$$

$$\chi_2 \mapsto \xi_2 = \psi_2; \quad (11c)$$

$$y \mapsto v = x + \frac{b - a}{x + y - \chi_1\psi_2}; \quad (11d)$$

$$\psi_1 \mapsto \eta_1 = \chi_1; \quad (11e)$$

$$\psi_2 \mapsto \eta_2 = \chi_2 + \frac{a - b}{x + y}\psi_2, \quad (11f)$$

which satisfies the Yang-Baxter equation (3), is a parametric, reversible, and birational map with invariants

$$I_1 = x + y, \quad I_0 = \chi_1\chi_2 + \psi_1\psi_2. \quad (12)$$

*Proof.* Equation (9) implies that  $\xi_2 = \psi_2$  and  $\eta_1 = \chi_1$ , and, using the latter, the following system of equations

$$u + v = x + y, \quad (13a)$$

$$uv + v^2 - b + \chi_1\eta_2 = yx + x^2 - a + \chi_1\chi_2, \quad (13b)$$

$$u^2 - a + \xi_1\psi_2 + uv = y^2 - b + \psi_1\psi_2 + yx, \quad (13c)$$

$$u\chi_1 + \xi_1 = y\chi_1 + \psi_1, \quad \psi_2v + \eta_2 = \psi_2x + \chi_2, \quad (13d)$$

$$(u^2 - a + \xi_1\psi_2)v + u(v^2 - b + \chi_1\eta_2) + \xi_1\eta_2 = (y^2 - b + \psi_1\psi_2)x + y(x^2 - a + \chi_1\chi_2) + \psi_1\chi_2, \quad (13e)$$

for variables  $u, v, \xi_1$  and  $\eta_2$ . Now, from equations (13b) and (13c) we obtain

$$v(x + y) - b + \chi_1\eta_2 = x(y + x) - a + \chi_1\chi_2, \quad (14a)$$

$$u(x + y) - a + \xi_1\psi_2 = y(x + y) - b + \psi_1\psi_2, \quad (14b)$$

where we have made use of (13a). Moreover, expressing  $\xi_1$  and  $\eta_2$  in terms of  $u$  and  $v$ , using equations (13d), and substituting to (14) we find that  $u$  and  $v$  are given by (11a) and (11d), respectively. Using the latter, from (13d) follows that

$$\xi_1 = \psi_1 - \frac{a - b}{x + y - \chi_1\psi_2}\chi_1, \quad \eta_2 = \chi_2 + \frac{a - b}{x + y - \chi_1\psi_2}\psi_2.$$

In the above expressions, we multiply both the numerators and the denominators with the conjugate expression of the latter, namely  $x + y + \chi_1\psi_2$ , and we use the fact that  $\chi_1^2 = \psi_2^2 = 0$ . Then, it follows that  $\xi_1$  and  $\eta_2$  are given by (11b) and (11f), respectively. Lastly, equation (13e) is satisfied in view of the rest ((13a)- (13d)).

The supertrace of the quantity  $M_b(y, \boldsymbol{\psi})M_a(x, \boldsymbol{\chi})$  reads:

$$\text{str}(M_b(y, \boldsymbol{\psi})M_a(x, \boldsymbol{\chi})) = 2\lambda + (x + y)^2 + \chi_1\chi_2 + \psi_1\psi_2 - a - b - e.$$

Thus,  $I = (x + y)^2 + \chi_1\chi_2 + \psi_1\psi_2 = I_1^2 + I_0$  is invariant of the map (10)-(11), where  $I_1$  and  $I_0$  are given by (12). However, it can be readily verified that  $I_1$  and  $I_0$  are invariants themselves.

From the uniqueness of the refactorisation problem (9) it follows that the map (10)-(11) satisfies the Yang-Baxter equation and it is reversible. The birationality of the map follows from the fact that (9) admits the symmetry  $(u, \boldsymbol{\xi}, v, \boldsymbol{\eta}, a, b) \leftrightarrow (y, \boldsymbol{\psi}, x, \boldsymbol{\chi}, b, a)$ . □

Setting all odd variables in (11) equal to zero and considering even variables and parameters  $a, b$  to be  $\mathbb{C}$ -valued (the bosonic limit) we obtain the standard Adler map (15)

$$(x, y) \xrightarrow{Y_{a,b}} \left( y + \frac{a - b}{x + y}, x + \frac{b - a}{x + y} \right). \quad (15)$$

The Adler map (15) occurs from the 3-D consistent discrete potential KdV equation [10, 11]. Its Lax representation [12, 14] occurs from the bosonic limit of (9); that is, it is given by

$$L_a(u)L_b(v) = L_b(y)L_a(x),$$

where  $L_a = L_a(x)$  is given by  $M_a(x, \boldsymbol{\chi})$  in (8) if one sets the odd variables equal to zero, namely

$$L_a(x) = \begin{pmatrix} x & e \\ x^2 + a - \lambda & x \end{pmatrix}.$$

It can be readily verified that the Adler map (15) is involutive; yet, for its Grassmann extension (10)-(11) we have the following.

**Proposition 3.1.2.** *The map  $S_{a,b} \in \text{End}(V_G \times V_G)$  given by (11) is non-involutive.*

*Proof.* If we denote

$$(x, \chi_1, \chi_2, y, \psi_1, \psi_2) \xrightarrow{S_{a,b}} (x', \chi'_1, \chi'_2, y', \psi'_1, \psi'_2) \xrightarrow{S_{a,b}} (x'', \chi''_1, \chi''_2, y'', \psi''_1, \psi''_2),$$

then it follows from (11c) and (11f) that

$$\chi''_2 = \psi'_2 = \chi_2 + \frac{a-b}{x+y} \psi_2 \neq \chi_2$$

which implies that

$$S_{a,b} \circ S_{a,b} \neq id,$$

and this completes the proof. □

## 4 Conclusions

We contributed to the direction initiated in [6], by constructing a noncommutative (Grassmann) extension of the famous Adler map, using some recent results on noncommutative Darboux transformations for a generalised super KdV system [16]. We showed that, in contrast to the original map, the Grassmann extended Adler map is not-involutive. This is an unexpected and quite interesting phenomenon, since involutive maps possess trivial dynamics.

In [16], Darboux transformations for the system (5) are constructed in three different cases. The first case corresponds to the Darboux matrix (7) which we used to construct the noncommutative Adler map (10)-(11). The second case of Darboux transformation, in the same paper [16], is associated with the following matrix

$$W_2 = \lambda \begin{pmatrix} 0 & 0 & 0 \\ e & 0 & 0 \\ 0 & 0 & e \end{pmatrix} + \begin{pmatrix} \frac{v_1}{2} & e & -\xi_{[1]} \\ \frac{v_1^2}{4} - p_1 & \frac{v_1}{2} & -\frac{v_1}{2} \xi_{[1]} \\ -\frac{v_1}{2} \eta & -\eta & -p_1 - \xi_{[1]} \eta \end{pmatrix}. \quad (16)$$

Changing in the above  $(v_1/2, -\eta, -\xi_{[1]}; p_1) \rightarrow (x, \chi_1, \chi_2; a)$ , we obtain a Lax operator

$$N_a(x, \boldsymbol{\chi}) = \lambda \begin{pmatrix} 0 & 0 & 0 \\ e & 0 & 0 \\ 0 & 0 & e \end{pmatrix} + \begin{pmatrix} x & e & \chi_2 \\ x^2 - a & x & x\chi_2 \\ x\chi_1 & \chi_1 & \chi_1\chi_2 - a \end{pmatrix},$$

which we used for the construction of a Yang-Baxter map (similar to Proposition 3.1.1). Surprisingly the map obtained in this way proved to be exactly the same as (10)-(11) and, moreover, the differential-difference equations corresponding to (7) and (16) obtained in [16] coincide after a relabeling of variables. This suggests that the Darboux transformations (7) and (16) are equivalent in a certain sense, namely they should be related via a transformation, although not necessarily a trivial one.

## Acknowledgements

A.V.M. acknowledges support from the Leverhulme Trust.

## References

- [1] ADLER V 1993 Recuttings of polygons *Funktsional. Anal. i Prilozhen.* **27** 79–82
- [2] BEREZIN F. 1987 Intoduction to superanalysis *Reidel Publishing Co. Dordrecht* **9**
- [3] BUCHSTABER V 1998 The Yang-Baxter transformation *Russ. Math. Surveys* **53:6** 1343-1345
- [4] DRINFELD V 1992 On some unsolved problems in quantum group theory *Lecture Notes in Math.* **1510** 1–8
- [5] GONCHARENKO, V.M. AND VESELOV, A.P. 2004 Yang-Baxter maps and matrix solitons *New trends in integrability and partial solvability* Springer191–197
- [6] GRAHOVSKI G.G., KONSTANTINOU-RIZOS S., MIKHAILOV A.V. 2016 Grassmann extensions of Yang-Baxter maps *Symmetry, Integrability and Geometry: Methods and Applications J. Phys. A* **49** 145202
- [7] GRAHOVSKI G AND MIKHAILOV A 2013 Integrable discretisations for a class of nonlinear Scrodinger equations on Grassmann algebras *Phys. Lett. A* **377** 3254–3259
- [8] HOLOD P.I. AND PAKULIAK S.Z. 1989 On the superextension of the Kadomtsev-Petviashvili equation and finite-gap solutions of Korteweg-de Vries superequations *Problems of Modern Quantum Field Theory* Springer 107–116
- [9] KONSTANTINOU-RIZOS S AND MIKHAILOV A 2013 Darboux transformations, finite reduction groups and related Yang-Baxter maps *J. Phys. A* **46** 425201
- [10] NIJHOFF F AND CAPEL H 1995 The discrete Korteweg-de Vries equation *Acta Appl. Math.* **39** 133–158
- [11] PAPAGEORGIOU V, NIJHOFF F, AND CAPEL H 1990 Integrable mappings and nonlinear integrable lattice equations *Phys. Lett. A* **147** 106–114
- [12] SURIS Y AND VESELOV A 2003 Lax matrices for Yang-Baxter maps *J. Nonlinear Math. Phys.* **10** 223–230
- [13] VESELOV A 2003 Yang-Baxter maps and integrable dynamics *Phys. Lett. A* **314** no. 3 214–221
- [14] VESELOV A 2007 Yang-Baxter maps: dynamical point of view *J. Math. Soc. Japan* **17** 145–167
- [15] XUE LING-LING, LEVI D. AND LIU Q.P. 2013 Supersymmetric KdV equation: Darboux transformation and discrete systems *J. Phys. A: Math. Theor. (FTC)* **46**502001
- [16] XUE LING-LING AND LIU Q.P. 2014 Bäcklund–Darboux transformations and discretizations of super KdV equation *Symmetry, Integrability and Geometry: Methods and Applications* **10**