

This is a repository copy of Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/100858/

Version: Accepted Version

Article:

Eftychiou, C, Barmby, DS, Wilson, SJ et al. (7 more authors) (2016) Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience. Catheterization and Cardiovascular Interventions, 88 (4). pp. 546-553. ISSN 1522-1946

https://doi.org/10.1002/ccd.26587

© 2016 Wiley Periodicals, Inc. This is the peer reviewed version of the following article: Eftychiou, C., Barmby, D. S., Wilson, S. J., Ubaid, S., Markwick, A. J., Makri, L., Blaxill, J. M., Spratt, J. C., Gunning, M. and Greenwood, J. P. (2016), Cardiovascular Outcomes Following Rotational Atherectomy: A UK Multicentre Experience. Cathet. Cardiovasc. Intervent., 88: 546–553. doi: 10.1002/ccd.26587, which has been published in final form at https://doi.org/10.1002/ccd.26587. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Table 1. Baseline characteristics of patients

DEMOGRAPHICS	
Median age (years)	72 (37-94)
Male	347 (68.3%)
BMI (kg/m ²)	27.3 (16.0-45.0)
PROCEDURAL URGENCY	
Elective	337 (65.4%)
Acute coronary syndrome	178 (34.6%)
RISK FACTORS	
Smokers (current and ex-smokers)	283 (55.7%)
DM	148 (28.7%)
Hypertension	382 (73.9%)
Dyslipidaemia	237 (65.1%)
HISTORY	
Prior MI	234 (45.4%)
Prior PCI	133 (25.7%)
Heart Failure	43 (9.8%)
Prior CABG	63 (12.2%)
Stroke	70 (13.6%)
Chronic Kidney Disease	82 (16.2%)
PVD	111 (21.5%)
Permanent Pacemaker	7 (1.4%)
Haemoglobin (g/dL)	12.9 (6.2-17.0)
Creatinine (µmol/L)	94.0 (46-1462)

Data shown as N (%) or median (range).

BMI=Body Mass Index; CABG=Coronary artery bypass surgery; DM=Diabetes Mellitus; MI=Myocardial Infarction; PCI=Percutaneous Coronary Intervention; PVD=Peripheral Vascular Disease.