



This is a repository copy of *"Stress Control" as a Large Group Psychoeducational Intervention at Step 2 of IAPT Services: Acceptability of the Approach and Moderators of Effectiveness.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/102717/>

Version: Accepted Version

---

**Article:**

Burns, P., Kellett, S. and Donohoe, G. (2016) "Stress Control" as a Large Group Psychoeducational Intervention at Step 2 of IAPT Services: Acceptability of the Approach and Moderators of Effectiveness. *Behavioural and Cognitive Psychotherapy* , 44 (4). pp. 431-443. ISSN 1352-4658

<https://doi.org/10.1017/S1352465815000491>

---

This article has been published in a revised form in *Behavioural and Cognitive Psychotherapy* [<https://doi.org/10.1017/S1352465815000491>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © British Association for Behavioural and Cognitive Psychotherapies.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Abstract

**Background:** ‘Stress Control’ (SC) has been adopted as a core intervention in step 2 of Improving Access to Psychological Therapies (IAPT) services, but contemporary evidence of effectiveness has lagged behind service uptake. **Aims:** To investigate the acceptability and effectiveness of SC and to explore moderators of outcome. **Method:** Analysis of acceptability (via attendance rates) and effectiveness (via IAPT minimum dataset). **Results:** SC was well tolerated with 73.3% of all patients and 75.4% of ‘clinical cases’ attending three or more sessions. Of the 546 ‘clinical cases’ attending SC and not in receipt of other interventions, 37% moved to recovery. Attendance improved outcome as for those patients attending all SC sessions, the recovery rate rose to 59.2%. **Conclusion:** SC appears a well-tolerated and effective intervention that enables large numbers to gain access to treatment in an organisationally efficient manner. Attendance appears important in facilitating SC outcomes.

**Keywords:** stress control; psychoeducation; PWP; IAPT stepped care

The landscape of psychological services in the UK has been transformed via the introduction of the Improving Access to Psychological Therapies (IAPT) programme. IAPT was introduced as a response to the Depression Report (Layard et al. 2006) highlighting the scarcity of availability of evidence-based psychological therapies for common mental health problems. A frequent criticism from patients of mental services has been the lack of accessibility to such evidence based psychological interventions (Turpin, Richards, Hope, & Duffy, 2008). The core philosophy of IAPT is the delivery of treatments consistent with the National Institute for Health and Clinical Excellence (NICE) guidelines for depression and anxiety (Clark, 2011). Nascent IAPT organisational models were evaluated via demonstration sites in 2006 (Clark et al., 2009; Parry et al., 2011) and then rolled out nationally in England in 2008 (CSIP Choice & Access Team, 2008). NICE recommends the provision of stepped-care service delivery models for the treatment of mild-moderate depression and anxiety disorders (excluding PTSD and social anxiety disorder). Reviews comparing stepped care with usual or enhanced usual care favour stepped care (Firth, Barkham & Kellett, 2014).

SC was developed to provide a clinically effective and organisationally efficient approach to treating common mental health problems (White, 2008). The SC approach is defined by its 'low contact-high volume' psychoeducational group-based approach. This is in contrast to the 'high contact-low volume' traditional one to one therapies (Brown et al. 2006). Psychoeducation is amongst the most effective of the range of evidenced-based practices across mental health disorders (Lukens & McFarlane, 2004). In IAPT services, psychoeducational interventions are delivered by Psychological Well-Being Practitioners (PWPs) at step 2 of the stepped care service delivery model (CSIP, 2008). The role of the PWP is that of a 'coach' as opposed to therapist (Turpin, 2010). In one-to-one low intensity work there have been three estimates thus far of the size of the PWP therapist effect. These range from 1 (Ali, Littleworth, McMillan, Delgadillo, Miranda, Croudace & Gilbody, 2014) to 7-9 % (Green, Barkham, Kellett & Saxon, 2014; Firth, Barkham, Kellett & Saxon, 2015).

## SC; effectiveness and moderators

The initial development of SC stimulated a broad range of evidence in terms of satisfaction, acceptability, clinical effectiveness/efficacy, organisational efficiency and durability of effect. SC users report high satisfaction rates (Houghton & Saxon, 2007; Kellett et al. 2004), with 96% highly recommending the treatment to others (White, 1995). Kellett, Clarke and Matthews (2007a) reported a dropout rate of 31%. White, Keenan and Brooks (1995) tested the efficacy of SC in a controlled trial. Post-intervention, SC showed highly significant changes compared to wait-list. Kellett et al. (2007b) benchmarked SC outcomes against individual CBT and individual psychodynamic-interpersonal psychotherapy to find few differences. Attendees show significant and reliable changes over the course of SC, with a 50% reduction in anxiety and depression (Wood et al, 2005; Joice & Mercer, 2010). Kellett et al (2007b) found that applying practice-based selection criteria improved outcomes. Kellett et al. (2007b) stated that SC was organisationally efficient due to both the high patient:facilitator ratios and also the low rates (20%) requiring further input. Gains are maintained in both the short (White et al., 1995; White & Keenan-Ross, 1997; Kellett et al., 2007b; Van Deale, 2013) and long-term (White, 1998).

Since this initial work, research regarding SC has atrophied - this has occurred despite SC being adopted as a common psychoeducational intervention within IAPT. A schism has occurred between the popularity of the SC approach and the standard of the contemporary evidence. The present research is novel in being the first to report SC outcomes from an IAPT service and also consider factors which moderate outcome. The aims were to (1) assess SC acceptability and effectiveness and (2) understand the moderating role of deprivation, presenting problem, dual delivery of interventions and problem severity.

## **Method**

### **Design and Context**

A pre-post design examined the effectiveness and acceptability of SC as an intervention for patients presenting with common mental health problems at step 2 of a city-wide IAPT service in the North of England.

## **Participants**

N = 2814 patients (1813 females) attended SC. The total number of patients referred to the service during this period was N = 42,968. Ages ranged from 16-88 years, with a mean age of 44.27 years (SD=13.85). Of the 2814 participants, 1062 were considered to be ‘clinical cases’ at the start of SC, meaning that they scored above clinical cut-off on the PHQ or the GAD (or both). To be considered as having received adequate dose of SC, patients need to have attended 3 or more sessions and this categorically defined attendance. All analyses of effectiveness were based upon the sample of N=801 ‘clinical cases’ (see Measures section) who attended SC (i.e. 3+ sessions). A number of these patients also received additional help within the IAPT service during SC. Participants who received other interventions were therefore considered part of a ‘SC+’ research sample (N=388), versus a SC only sample (N=413). Attendees scoring above clinical cut-offs on both PHQ-9 and GAD-7 (see measures section), were coded as comorbid anxiety and depression. If a patient scored above clinical cut-off on GAD-7 and not the PHQ-9, they were considered to have an anxiety disorder (and visa versa for the PHQ-9 and depression). Figure 1 details the various research samples.

Insert figure 1 here please

## **Measures and Outcomes**

The Patient Health Questionnaire (PHQ-9; Kroenke & Spitzer, 2002; clinical caseness score = 10) and the Generalised Anxiety Disorder-7 (GAD-7; Spitzer, Kroenke, & Williams, 2006; clinical caseness score = 8) are valid and reliable case identifier and outcome measures of depression and anxiety. The criteria for clinical change occurring during SC was GAD-7 final score  $\leq 7$  and PHQ-9  $\leq 9$ , as is used to define moving to recovery rates in IAPT (Gyani, Shafran Layard & Clark, 2009). Reliable change calculations (Evans et al. 2014) were employed to investigate whether reliable improvements/deteriorations occurred. A change of 6 points (PHQ-9) and 4 points

SC; effectiveness and moderators

(GAD-7) in either direction represented a reliable change (increase equals deterioration and decrease equals improvement). Deprivation was measured using the Index of Multiple Deprivation 2010 (IMD, Department for Communities and Local Government, 2011a, 2011b). The IMD is an aggregation of deprivation indices (income, employment, health and disability, education, skills/training, barriers to housing and services, crime and living environment). Postcodes were used to establish IMD rank; a higher rank (0-100) indicates an area with higher proportion of people living in deprivation.

### **Intervention**

Patients attended SC through two routes (1) referred to IAPT from GPs and screened by PWP who offered SC as an intervention option within the suite of low intensity treatments or (2) via self-referral through gaining knowledge of SC through the service website, posters, leaflets or word of mouth. All participants were required to book on to SC prior to attending. The specific nature of the other interventions received was not recorded for SC+ participants, but was at step 2 was cCBT ('Beating the Blues' and 'FearFighter'), one to one PWP work or healthy living workshops. Patients that were also stepped up to step 3 interventions received CBT, counselling, group behavioural activation or couples therapy. It was not possible to determine whether extra therapeutic interventions from outside of the service (e.g. private therapy) also occurred. SC is intended as a stand-alone intervention and so patients were discouraged from accessing other IAPT interventions simultaneously.

SC was delivered using the White (2005) treatment model, which superseded the White and Keenan (1990) approach. The White (2005) approach entails providing psychoeducative low intensity cognitive behaviourally informed self-help for patients across the anxiety disorders, with a management of depressed mood component. Sessions were didactic and patients were informed that they could simply attend, listen and complete the exercises. Patients can attend SC with carers/friends/family should this facilitate engagement (White, 2000). SC was delivered in

## SC; effectiveness and moderators

community settings and often outside of normal office hours, in order to enable uptake and reduce stigma (White, 2000). Thirty-eight groups ran between October 2009-April 2014. Group size ranged from 23-106, with a mean size of  $N=74$ . Each SC group was facilitated by two PWP's; each session lasted for 2 hours, half an hour of which was devoted to a comfort-break, entailing a total treatment time of 9 hours. SC ran weekly over six sessions containing the following elements: week 1, introduction to psychoeducation and the cognitive behavioural model; week 2, management of physiology; week 3, management of mental events; week 4, management of behaviour; week 5, management of panic attacks and sleep and week 6, self-care. At the end of each session, material for the next session was distributed containing homework exercises. At the final session, relapse prevention materials were distributed. Participants were not followed-up if they missed sessions and were not reviewed on completion.

## Results

Out of a total sample of  $N = 2814$  patients, 2062 (73.3%) attended SC (i.e. 3+ sessions). In terms of total patients referred to the IAPT service (see method), SC saw 6.55% of referrals. Figure 1 contains a summary of the research samples and associated attendance rates and Table 1 describes the demographics and deprivation ranks. Patients who attended  $<3$  SC sessions were typically younger than those who attended full SC ( $t(2812) = 5.694, p<.001, d = 0.24$ ) and also lived in areas of greater deprivation ( $t(2798) = 4.295, p<.001, d = 0.19$ ). In terms of those patients that met caseness criteria prior to intervention,  $N = 801$  (75.4%) attended more than three SC sessions.

Insert table 1 here please

Table 2 reports the group outcomes and the individual outcome rates for the SC and SC+ samples. There was no association between purity of intervention and whether or not patients

SC; effectiveness and moderators

moved to recovery. Patients that received SC+ lived in areas of greater deprivation ( $t(781.16) = 1.975, p < .05, d = 0.14$ ). In order to evidence the effectiveness of SC as an intervention in its own right, the subsequent analysis excluded the SC+ sample. Of the  $N=413$  SC only patients, 194 (47.1%) moved to recovery. Table 3 reports the recovery rate by session attendance analysis. When patients attended all SC sessions, the recovery rate was 59.2%, with a significant association between number of sessions attended and movement to recovery ( $\chi^2(3) = 44.537, p < .001$ ). The recovery ratio increased proportionally with attendance; the odds in favour of recovery were 9.06 times higher if all sessions were attended. There was no significant difference in GAD scores at pre-intervention between those who attended  $<3$  sessions and those who attended full SC ( $t(109.042) = 0.71, ns$ ). However, patients at assessment who then went onto attend  $<3$  sessions had significantly higher PHQ scores ( $t(222) = 2.839, p < .01, d = 0.42$ ) than those who attended full SC. Patients who attended less SC lived in areas of greater deprivation than who attended full SC ( $t(222) = 2.175, p < .05, d = 0.32$ ).

Insert tables 2 and 3 here please

Table 4 reports recovery rates and reliable change by presentation. Patients with either depression or anxiety were more likely to move to recovery than those with co-morbidity ( $\chi^2(2) = 10.901, p < .01$ ). Depression presentations were 2.5 times and anxiety presentations 1.89 times more likely to move to recovery. Of the 387 patients who met caseness on the GAD-7 before SC (anxiety and comorbid samples), 228 (58.9%) reliably improved.  $N=11$  (2.8%) reliably deteriorated (anxiety). Of the  $N=302$  meeting depression caseness criteria (depression and comorbid samples), 137 (45.4%) reliably improved.  $N= 6$  (2%) reliably deteriorated (depression). Figure 2 displays a scatter plot showing that presentation severity was significantly correlated with change in distress score following SC ( $r(412) = 0.298, p < .001$ ).



Insert table 4 and figure 2 here please

Table 5 reports SC outcomes by severity. ‘Severely depressed’ patients prior to intervention showed a significantly greater reduction in depression, than those categorised with ‘mild to moderate depression’ ( $t(64.963) = 4.621, p < .001, d = 1.09$ ). Recovery rates were higher for patients in the ‘mild to moderate depression’ category; 55.6% moved to recovery in comparison to 26.3% in the ‘severe depression’ cluster ( $\chi^2(1) = 15.922, p < .001$ ). A similar pattern was also apparent for anxiety outcomes. ‘Severely anxious’ patients showed significantly greater improvement than those presenting with mild anxiety ( $t(248.88) = 7.235, p < .001, d = 1.23$ ). Recovery rates were higher for those with mild anxiety: 60.5% moved to recovery, whereas 32.7% of the severe anxiety cluster recovered ( $\chi^2(1) = 20.504, p < .001$ ). A biserial correlation found that deprivation was significantly related to moving to recovery ( $r_b = .142; p < .005$ ). Patients who did not move to recovery were more deprived; 2% of variance in recovery status was accounted for by deprivation ( $r_b^2 = .02$ ).

Insert table 5 here please

## Discussion

This study has provided contemporary IAPT evidence of the uptake and effectiveness of SC and investigated the role of moderating factors. SC was delivered an intervention to nearly 7% of total referrals to the service, indicating the prominence of the intervention and the plurality of other service provision. SC was well tolerated in terms of attendance; more than 70% attended at least three SC sessions, with attendance rates higher for those with pre-intervention clinically significant distress. Rates were higher than extant attendance evidence (e.g. Kellett et al. 2007a). Those patients that dropped out of SC before attending at least three sessions lived in areas of greater deprivation. SC appears comparatively clinically equivalent to the other IAPT interventions (Gyani,

Shafran, Layard & Clark, 2013) and produced higher recovery rates than the Green et al. (2014) and Firth et al. (2015) analyses of one-to-one PWP work. This may be due to the rapid and overt normalising effect of attending a large group (Kellett et al. 2007b). When SC was delivered as the sole intervention then recovery rates were higher than for those who also received a supplementary intervention (i.e. the SC+ research sample). This should not be construed as an interference effect, as those who received extra intervention were found to have higher levels of distress pre-intervention, in addition to living in areas of higher deprivation.

Analysis of the impact of attendance on outcome showed a clear pattern, as recovery rates were higher when patients attended more sessions. For example, 59.2% of participants who attended all SC sessions moved to recovery, whereas only 13.4% of those who attended three sessions did so. Recovery rates were similar to extant SC evidence, with 47.1% of those who attended at least three sessions moving to recovery. Patients who presented with a single mental health concern (i.e. the depression-only or anxiety-only research samples) had enhanced recovery rates. There was a higher proportion of reliable change for anxiety as opposed to depression. This maybe because SC contains greater anxiety management, as opposed to mood management, component (Kellett et al. 2007a).

The study highlights the importance of attendance in relation to generating positive outcomes, as chance of recovery increased with number of sessions attended. Strategies to maintain engagement with patients at risk of dropping out of SC need to be developed and evaluated. A trial could compare attendance for SC groups that have an attendance intervention embedded within them to TAU rates. Strategies for increasing attendance might be the antibiotic metaphor of ‘finishing the course of treatment’ and informing patients that chance of recovery more than doubles when they fully attend. Future research is also required to discover the reasons why patients dropout and studies employing qualitative methods would be at a premium. Similarly, the reasons why patients receive more than one intervention also need investigating. It is possible that screening PWPs felt overwhelmed when highly symptomatic and deprived patients attended and

SC; effectiveness and moderators

therefore attempted to ‘rescue’ the patient through offering multiplying provision (Stean, 2014).

The findings related to IMD rank suggest a relationship between living in areas of higher deprivation and both lower attendance and poorer outcomes. This suggests that the socio-economic context impinges on outcomes and that a ‘perfect storm’ can be created of deprivation being associated with poor attendance and then associated poorer outcomes. Methods to engage people from such areas are again vitally important to develop and evaluate.

The separate analyses for depression and anxiety severity at assessment showed a similar pattern: for both measures, the moving to recovery rates were higher for patients reporting milder symptom distress. SC was designed for people with mild to moderate common mental health problems and Kellett et al. (2004) showed that selection of less severe cases improved outcomes. However, SC in this evaluation was delivered to patients across the spectrum of presentation severities. Across both outcome measures, the average reduction in scores was around double in the severe presentation group, when compared with the mild to moderate group. This finding suggests that SC may provide a pragmatic approach to meeting the needs of patients experiencing a range of distress. Solely focusing on moving to recovery rates might suggest that SC is not effective for people with more severe presentations, and therefore IAPT services need to consistently factor in reliable change calculations to supplement moving to recovery rates. The consistency of the intervention could also be called into question. Although SC delivery was consistent with the SC package and all groups were facilitated by two PWPs, the intervention was facilitated by different PWPs with varying levels of experience. This could also be interpreted as evidence that SC can be facilitated effectively by a variety of staff. There is a need to develop a competency measure for delivery of psychoeducation. The lack of follow-up data in the current study is a weakness, particularly as contemporary evidence concerning durability of SC effects is required.

In conclusion, SC appears to be a well-tolerated and effective intervention for patients presenting to IAPT services and treated at step 2 with a large-group psychoeducational approach. SC can be delivered to groups of up to 150 services users by two PWPs, at a total time investment

## SC; effectiveness and moderators

of 24 hours clinical contact time. This further endorses SC as an organisationally efficient intervention (Kellett et al. 2007b). Attendance appeared important regarding outcome and people who dropped out tended to live in areas of higher deprivation. IAPT services need to adopt and evaluate 'in reach' strategies to such communities. This would ensure that living in an area of deprivation does not also mean that the chances of benefitting from an evidenced based psychological intervention are also suppressed.

## References

- Ali, S., Littlewood, E., McMillan, D., Delgado, J., Miranda, A., Croudace, T., & Gilbody, S.** (2014). Heterogeneity in patient-reported outcomes following low-intensity mental health interventions: A multilevel analysis. *PloS ONE*, 9, e99658.
- Brown, J., Elliott, S. A., & Butler, C.** (2006). Can large-scale self-referral psychoeducational stress workshops help improve the psychological health of the population? *Behavioural and Cognitive Psychotherapy*, 34, 165–177.
- Clark, D. M.** (2011). Implementing NICE guidelines for the psychological treatment of depression and anxiety disorders: The IAPT experience. *International Review of Psychiatry*, 23, 375–384.
- Clark, D. M., Layard, R., Smithies, R., Richards, D. A., Suckling, R. & Wright, B.** (2009). Improving access to psychological therapy: Initial evaluation of two UK demonstration sites. *Behaviour Research and Therapy*, 47, 910-920.
- CSIP Choice and Access Team** (2008). *Improving Access to Psychological Therapies (IAPT) Commissioning Toolkit*. London, UK: Department of Health.
- Department of Communities and Local Government** (2011a). *English Indices of Deprivation 2010*. Retrieved from:  
[www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/6320/1870718.pdf](http://www.gov.uk/government/uploads/system/uploads/attachment_data/file/6320/1870718.pdf)
- Department of Communities and Local Government** (2011b). *English IMD 2010 Data*. Retrieved from: [www.communities.gov.uk/publications/corporate/statistics/indices2010](http://www.communities.gov.uk/publications/corporate/statistics/indices2010)
- Firth, N. Barkham, M. & Kellett, S.** (2014). The clinical effectiveness of stepped care systems for depression in working age adults: A systematic review. *Journal of Affective Disorders* (online)

SC; effectiveness and moderators

- Firth, N., Barkham, M., Kellett, S., & Saxon, D. (2015).** Therapist effects and moderators of effectiveness and efficiency in Psychological Wellbeing Practitioners: A multilevel modelling analysis. *Behaviour Research and Therapy*, 69, 54-62.
- Green, H, Barkham, M., Kellett, S., & Saxon, D. (2014).** Therapist effects and IAPT psychological wellbeing practitioners (PWP): A multilevel modelling and mixed methods analysis. *Behaviour Research and Therapy*, 63, 43-54.
- Gyani, A., Shafran, R., Layard, R., & Clark, D. M. (2013).** Enhancing recovery rates: Lessons from year one of IAPT. *Behaviour Research and Therapy*, 51, 597-606.
- Houghton, S., & Saxon, D. (2007).** An evaluation of large group CBT psycho-education for anxiety disorders delivered in routine practice. *Patient Education and Counselling*, 68, 107-110.
- Joice, A., & Mercer, S. (2010).** An evaluation of the impact of a large group psycho-educational programme (Stress Control) on patient outcome: does empathy make a difference? *The Cognitive Behaviour Therapist*, 3, 1-17.
- Kellett, S., Newman, D. W., Matthews, L., & Swift, A. (2004).** Increasing the effectiveness of large group format CBT via the application of practise-based evidence. *Behavioural and Cognitive Psychotherapy*, 32, 231-234.
- Kellett, S., Clarke, S., & Matthews, L. (2007a).** Session impact and outcome in group psycho-educative cognitive behavioural therapy. *Behavioural and Cognitive Psychotherapy*, 35, 335-342.
- Kellett, S., Clarke, S., & Matthews, L. (2007b).** Delivering group psychoeducational CBT in Primary Care: Comparing outcomes with individual CBT and individual psychodynamic-interpersonal psychotherapy. *British Journal of Clinical Psychology*, 46, 211-222.
- Kroenke, K., & Spitzer, R. L. (2002).** The PHQ-9: A New Depression Diagnostic and Severity Measure. *Psychiatric Annals*, 32, 509–515.

SC; effectiveness and moderators

- Layard, R., Bell, S., Clark, D., Knapp, M., Meacher, M., Priebe, S., Thornicroft, G., Turnbull, A. & Wright, B.** (2006). The depression report: A new deal for depression and anxiety disorders. Centre for Economic Performance's Mental Health Policy Group.
- Lukens, E. P., & McFarlane, W. R.** (2004). Psychoeducation as evidence-based practice: Considerations for practice, research and policy. *Brief Treatment and Crisis Intervention*, 4, 205-225.
- Parry, G., Barkham, M., Brazier, J., Dent-Brown, K., Hardy, G., Kendrick, T., Rick, J., Chambers, E., Chan, T., Connell, J., Hutten, R., de Lusignan, S., Mukuria, C., Saxon, D., Bower, P. & Lovell, K.** (2011). An evaluation of a new service model: Improving Access to Psychological Therapies demonstration sites 2006-2009. Final report. NIHR Service Delivery and Organisation programme.
- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B.** (2006). A Brief Measure for Assessing Generalized Anxiety Disorder. *Archives of Internal Medicine*, 166, 1092-1097.
- Turpin, G.** (2010): IAPT Good Practice Guide to using Self-help Materials. NMH DU/IAPT, 1-40.
- Turpin, G., Richards, D., Hope, R., & Duffy, R.** (2008). Improving access to psychological therapies in: A national initiative to ensure the delivery of high quality evidence-based therapies, *Papeles del Psicologo*, 29, 271-280.
- Van Deale, T.** (2013). A psychoeducational approach to stress management: an implementation and effectiveness study (unpublished Ph.D. dissertation). University of Leuven, Belgium.  
Retrieved from  
[http://www.kuleuven.be/lucas/Docs/Nieuwsflitsen/47\\_LUCAS\\_nieuwsbrief/Doctoral\\_dissertation\\_TomVanDaele.pdf](http://www.kuleuven.be/lucas/Docs/Nieuwsflitsen/47_LUCAS_nieuwsbrief/Doctoral_dissertation_TomVanDaele.pdf).
- White, J.** (1995). An analysis of components in large group didactic therapy – 'Stress Control'. *Clinical Psychology Forum*, 76, 11-13.
- White, J.** (1998). "Stress Control" large group therapy for generalised anxiety disorder: two year follow-up. *Behavioural and Cognitive Psychotherapy*, 26, 237-245.

SC; effectiveness and moderators

**White, J.** (2000). Clinical psychology in primary care. *Primary Care Psychiatry*, 6, 127-136.

**White, J.** (2008). CBT and the challenge of primary care: Developing effective, efficient, equitable, acceptable and accessible services for common mental health problems. *Journal of pUBLIC Mental Health*, 7, 32-41.

**White, J., & Keenan, M.** (1990). 'Stress Control': A pilot study of large group therapy for generalized anxiety disorder, *Behavioural Psychotherapy*, 18, 143-146.

**White, J., Keenan, M., & Brooks, N.** (1995). 'Stress Control': A controlled comparative investigation of large group therapy for generalized anxiety disorder. *Behavioural Psychotherapy*, 20, 97-114.

**White, J., & Keenan-Ross, M.** (1997). Stress Control large group didactic therapy for anxiety: an approach for managed care systems. *The Behavior Therapist*, 20, 192-196.

**Wood, S., Kitchiner, N. J., & Bisson, J. I.** (2005). Experiencing and implementing an adult educational approach to treating anxiety disorder. *Journal of Psychiatric and Mental Health Nursing*, 12, 95-99.






Figure 1. Defining the patient population and access/uptake of Stress Control.

SC; effectiveness and moderators

	N	Mean Age (SD)	Mean Deprivation Rank (SD)
<b>SC</b>	1698	44.20 (14.16)	23.88 (17.30)
Non-attenders (<3)	467	41.24 (14.01)	27.04 (18.28)
Attenders (>2)	1231	45.32 (14.07)	22.68 (16.77)
<b>SC+</b>	1116	44.38 (13.37)	26.08 (18.38)
Non-attenders (<3)	285	42.77 (14.27)	27.43 (18.35)
Attenders (>2)	831	44.93 (13.01)	25.62 (18.38)
<b>Whole sample</b>	2814	44.27 (13.85)	24.76 (17.77)
Non-attenders (<3)	752	41.82 (14.12)	27.18 (18.29)
Attenders (>2)	2062	45.16 (13.65)	23.87 (17.49)

Table 1; age and deprivation ranks for whole sample and subgroups

SC; effectiveness and moderators

	Group Outcomes						Individual Outcomes					
	N	Pre-SC Mean (SD)	Post-SC Mean (SD)	Pre-post change Mean (SD)	95% CI range	t	d	Positive reliable change	Positive clinically significant change (i.e. moving to recovery)	Reliable and clinically significant positive change	Stasis	Negative reliable change (i.e. deterioration)
<b>Stress Control (SC)</b>												
<i>Whole 'clinical case' sample</i>												
PHQ-9	414	15.50 (4.47)	11.58 (6.31)	3.92 (5.32)	3.41-4.43	15.006*	0.88	143 (34.5%)	165 (39.9%)	129 (31.2%)	264 (63.8%)	7 (1.7%)
GAD-7	512	13.88 (3.83)	9.90 (5.70)	3.98 (5.00)	3.54-4.41	18.008*	1.04	240 (46.9%)	204 (39.8%)	180 (35.2%)	260 (50.8%)	12 (2.3%)
<i>Attended &gt;2 sessions</i>												
PHQ-9	302	15.29 (4.46)	10.10 (6.03)	5.20 (5.45)	4.58-5.81	16.578*	1.17	137 (45.4%)	131 (43.4%)	124 (41.1%)	158 (52.5%)	6 (2%)
GAD-7	387	13.66 (3.79)	8.60 (5.43)	5.06 (5.14)	4.54-5.57	19.365*	1.34	228 (58.9%)	194 (50.1%)	173 (44.7%)	148 (38.2%)	11 (2.8%)
<b>Stress Control Plus (SC+)</b>												
<i>Whole 'clinical case' sample</i>												
PHQ-9	422	15.91 (4.46)	11.68 (6.39)	4.23 (5.23)	3.73-4.73	16.621*	0.95	160 (37.9%)	169 (40.0%)	123 (29.1%)	254 (60.2%)	8 (1.9%)
GAD-7	490	14.31 (3.91)	10.21 (5.72)	4.09 (5.10)	3.64-4.55	17.775*	1.05	249 (50.8%)	187 (38.2%)	172 (35.1%)	227 (46.3%)	14 (2.9%)
<i>Attended &gt;2 sessions</i>												
PHQ-9	311	15.59 (4.29)	10.38 (5.89)	5.22 (5.18)	4.64-5.80	17.774*	1.22	144 (46.3%)	151 (48.6%)	112 (36%)	164 (52.7%)	3 (1%)
GAD-7	368	14.16 (3.87)	9.34 (5.47)	4.82 (5.19)	4.29-5.35	17.794*	1.25	217 (58.6%)	163 (44.1%)	149 (40.3%)	143 (38.6%)	10 (2.7%)

P < .001\*

Table 2; group and individual outcomes rates for the SC and the SC+ research samples

SC; effectiveness and moderators

Moving to Recovery	Number of SC sessions attended				Total
	3	4	5	6	
<b>SC</b>					
Yes	9 (13.4%)	31 (39.7%)	61 (55.5%)	93 (59.2%)	194
No	58	47	49	64	218
<b>SC+</b>					
Yes	20 (27.4%)	28 (40%)	44 (41.1%)	72 (52.9%)	164
No	53	42	63	64	222

Table 3; recovery rates by session attendance

SC; effectiveness and moderators

	Group Outcomes						Individual Outcomes					
	N	Pre-SC Mean (SD)	Post-SC Mean (SD)	Pre-post change Mean (SD)	95% CI range	t	d	Positive reliable change	Positive clinically significant change (i.e. moving to recovery)	Reliable and clinically significant positive change	Stasis	Negative reliable change (i.e. deterior ation)
<b>Anxiety-only</b>												
GAD-7	110	11.46 (3.04)	6.80 (4.34)	4.66 (4.98)	3.72-5.61	9.816*	1.53	65 (59.1%)	63 (57.3%)	54 (49.1%)	40 (36.4%)	5 (4.5%)
<b>Depression- only</b>												
PHQ-9	25	13.20 (3.15)	8.20 (5.36)	5.00 (4.87)	2.99-7.01	5.130*	1.59	13 (52%)	16 (64%)	13 (52%)	12 (48%)	0 (0%)
<b>Comorbid</b>												
GAD-7	277	14.53 (3.70)	9.32 (5.65)	5.21 (5.20)	4.60-5.83	16.692*	1.41	163 (58.8%)	115 (41.5%)	106 (38.3%)	108 (39%)	6 (2.2%)
PHQ-9	277	15.48 (4.51)	10.27 (6.07)	5.21 (5.50)	4.56-5.86	15.768*	1.16	124 (44.8%)	115 (41.5%)	97 (35%)	147 (53.1%)	6 (2.2%)

P < .001\*

Table 4; recovery and reliable change rates by clinical presentation




Figure 2; Scatter plot of relationships between pre-intervention distress and amount of change pre-post SC

SC; effectiveness and moderators

	Group Outcomes						Individual Outcomes					
	N	Pre-SC Mean (SD)	Post-SC Mean (SD)	Pre-post change Mean (SD)	95% CI range	t	d	Positive reliable change	Positive clinically significant change (i.e. moving to recovery)	Reliable and clinically significant positive change	Stasis	Negative reliable change (i.e. deterior ation)
<b>GAD-7</b>												
<b>Severity</b>												
Mild	114	8.39 (1.34)	5.74 (3.49)	2.66 (3.29)	2.05-3.27	8.638*	1.99	51 (44.7%)	69 (60.5%)	44 (38.6%)	58 (50.9%)	5 (4.4%)
Severe	153	17.61 (2.01)	10.89 (6.14)	6.72 (5.81)	5.79-7.67	14.313*	3.34	101 (66%)	50 (32.7%)	50 (32.7%)	51 (33.3%)	1 (0.7%)
<b>PHQ-9</b>												
<b>Severity</b>												
Mild- moderate	250	10.17 (2.54)	6.98 (4.05)	3.18 (4.19)	2.66-3.71	12.029*	1.25	80 (32%)	139 (55.6%)	76 (30.4%)	163 (65.2%)	7 (2.8%)
Severe	57	22.75 (2.11)	15.02 (7.32)	7.74 (7.17)	5.84-9.64	8.152*	3.67	27 (47.4%)	15 (26.3%)	15 (26.3%)	30 (52.6%)	0 (0%)

P < .001\*

Table 5; recovery and reliable change rates by initial presentation severity