



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/103909/>

Version: Accepted Version

Article:

Khurram, S.A., Warner, C., Henry, A.M. et al. (2016) Accuracy of clinical coding for procedures in oral and maxillofacial surgery. *British Journal of Oral and Maxillofacial Surgery*. ISSN: 0266-4356

<https://doi.org/10.1016/j.bjoms.2016.05.031>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Accuracy of Clinical Coding for Oral and Maxillofacial Surgery Procedures

Syed Ali Khurram,¹ Claire Warner,² Alastair M Henry,² Anand Kumar,² Ricardo I Mohammed-Ali.²

¹ Unit of Oral and Maxillofacial Pathology, School of Clinical Dentistry, 19 Claremont Crescent, Sheffield, S10 2TA, UK.

² Department of Oral and Maxillofacial Surgery, Charles Clifford Dental Hospital, Wellesley Road, Sheffield Teaching Hospitals NHS Foundation Trust, S10 2SZ, UK.

Syed Ali Khurram- Academic Clinical Lecturer, Unit of Oral and Maxillofacial Pathology (address as above).² Email: s.a.khurram@sheffield.ac.uk

Claire Warner- Dental foundation trainee, Oral and Maxillofacial Surgery (address as above).¹ Email- claire.warner4@nhs.net

Alastair M Henry- Senior House Officer, Oral and Maxillofacial Surgery (address as above).¹ Email: amhenry84@mac.com

Anand Kumar- Specialist Registrar, Oral and Maxillofacial Surgery (address as above).¹ Email: anandkumar@doctors.org.uk

Ricardo I Mohammed-Ali- Consultant, Oral and Maxillofacial Surgery (address as above).¹ Email: ricardomohammed-ali@nhs.net

Corresponding author:

Dr. Syed Ali Khurram. BDS, MSc., PhD., MFDS-RCS, FHEA

Academic Clinical Lecturer, Unit of Oral and Maxillofacial Pathology,

School of Clinical Dentistry, 19 Claremont Crescent, Sheffield, S10 2TA, UK

Email: s.a.khurram@sheffield.ac.uk, Tel: 44 114 2717951, Fax: 44 114 2717894

Conflict of interest: None.

Funding: None received.

Keywords: clinical coding, oral and maxillofacial, payment by results, OPCS, HRG.

Word count- 1684

Abstract

Clinical coding has important financial implications, and discrepancies in the assigned codes can directly affect the funding of a department and hospital. Over the last few years, numerous oversights have been noticed in the coding of oral and maxillofacial (OMF) procedures. To establish the accuracy and completeness of coding, we retrospectively analysed the records of patients during two time periods: March to May 2009 (324 patients), and January to March 2014 (200 patients). Two investigators independently collected and analysed the data to ensure accuracy and remove bias. A large proportion of operations were not assigned all the relevant codes, and only 32% - 33% were correct in both cycles. To our knowledge, this is the first reported audit of clinical coding in OMFS, and it highlights serious shortcomings that have substantial financial implications. Better input by the surgical team and improved communication between the surgical and coding departments will improve accuracy.

Introduction

Since the introduction of Payment by Results by the UK Department of Health in April 2002 healthcare providers are paid for the procedures they do instead of being provided a lump sum (1). For this to happen, treatments and care must be translated into codes. These form the basis of the Healthcare Resource Group (HRG) code, which is a nationally agreed tariff based on the Office of Population Censuses and Survey (OPCS) classification 2 and the International Statistical Classification of Diseases (ICD)(3). The OPCS system classifies surgical activity, and the ICD relates to diagnoses and coexisting conditions. Accuracy of codes is therefore essential to generate the correct payment. Inaccuracies in clinical coding in the specialties of plastic surgery, urology, and ear, nose, and throat (ENT) have been reported to have an impact on the HRG code and subsequently on the income received (4–7). Although factors such as complex anatomy, wide range of procedures, and the diverse range of diseases can potentially cause inaccuracies in clinical coding, there are only two reported studies in literature that have explored this in oral and maxillofacial surgery (OMFS) procedures (8, 9).

The aim of the audit was to determine the accuracy of the OPCS codes generated from OMFS procedures in our department, as anecdotal evidence had suggested inaccuracies that resulted in errors in payments to the department.

Material and methods

A retrospective review of all operations carried out under general anaesthetic at two time points in an OMFS department at a tertiary referral hospital was undertaken. Approval was obtained from the local Clinical Effectiveness Unit. Two cycles of operations were analysed: in the first (March to May 2009) 324 operations were assessed, and in the second (January to March 2014) 200 procedures were analysed. Operations included the full spectrum of OMFS subspecialties. Digital operative notes were obtained from the hospital's Informatics Department; these were originally recorded on the Operating Room Management Information System (ORMIS) during the procedures. Any identifiable data was removed to ensure anonymity, and the hospital number replaced with an audit number to identify the cases. OMFS dental core trainees (formerly senior house officers) systematically analysed all the operative notes under the supervision of an OMFS specialist trainee (StR) and a consultant. Codes generated by professional hospital coders using OPCS version 4.2 were cross-checked with the operative notes to determine accuracy and completeness (10). A standard of 100% accuracy and 100% completeness of generated codes were set as the gold standard. Corrected or missing codes were recorded and grouped together. To improve accuracy after the first cycle, information about the most common coding oversight was disseminated to the coders and clinicians. This took a much longer time than anticipated because of the large numbers and initial difficulties with data collection and analysis. Numerous discussions between the management and coding department resulted in the short-term placement of a dedicated procedural coder on the OMFS inpatient ward and in the dental hospital to gain a better insight into the procedures and to improve dialogue between clinicians and coders. The second cycle, which took place two years after dissemination of the original data, was consistent in terms of the time interval studied, data collection, and analysis.

Results

In the first cycle, 67.6% (219/324) of the recorded codes were correct. The second review showed no improvement in accuracy, with 66.5% (133/200) being correct (Figure 1).




Figure 1. Percentage accuracy of clinical coding in the two cycles of the audit (cycle 1: n=320; cycle 2: n=200) (black = correct; grey = incorrect).

There was a considerable improvement in the coding of bilateral procedures: in the first cycle, 31 of 96 were correct (32.3%) and in the second, 61 of 77 were correct (79.2%) (Figure 2).




Figure 2. Percentage of correctly-coded bilateral procedures (cycle 1: n=320; cycle 2: n=200) (black = correct; grey = incorrect).

Inaccuracies appeared to be distributed across most OMFS operations (Fig. 3). The coding of dentoalveolar procedures improved, but there was no improvement in coding of trauma and oncology operations, and accuracy was considerably reduced for temporomandibular joint (TMJ) and orthognathic procedures.




Figure 3. Percentage accuracy of clinical coding across different OMFS procedures (cycle 1: n=320; cycle 2: n=200) (black = correct; grey = incorrect).

Discussion

Coding errors are routinely seen in surgical practice, and reports of inaccuracies range from 22% to 47.7%.^{4–6,8} The assignment of incorrect HRG codes affects the income received by the healthcare provider (9) and even small errors can have huge financial repercussions (Table 1). This is supported by the findings from both cycles of our audit.

Table 1- Comparison of distribution of correct clinical codes across different OMFS procedures between the two audit cycles.

	Cycle 1 %	Cycle 2 %
TMJ	55.6	0.0
Dentoalveolar	56.4	72.1
Orthognathic	93.8	42.9
Trauma	79.1	64.4
Salivary	100.0	100.0
Oncology	68.5	63.3

The audit also highlighted numerous oversights and short-comings in the OPCS and HRG systems, which form the main framework of rules and regulations for coders. The tariff set for the removal of one tooth is the same as that for a full dental clearance under general anaesthetic, which may require the removal of multiple teeth and is likely to take more time and resources. Discussions with coders showed that, unlike operations in other specialties, bilateral dental or dentoalveolar procedures, such as removal of impacted molars, are considered as single procedures and do not generate a higher tariff (11). These inconsistencies between specialties may contribute to errors in the assignment of HRG codes and result in the wrong amount being paid to the healthcare provider. ICD codes modify the tariffs set, and tariffs are higher for patients with coexisting medical conditions (Table 2).

Table 2- Comparison of clinical code related tariffs between elective and emergency OMFS procedures.

Procedure	Tariff (day case) £	Tariff (emergency) £
Simple extraction	476	993
Surgical removal of tooth	612	1469
Surgical Excision of lesion- face	546	922
Laser excision- face	1408	1802
Dental Clearance (single tooth extraction, upper, lower or full clearance)	612	1469
Suture (face)	546	929
Suture (Mouth)	612	1469

The audit did not look specifically at the recording of coexisting conditions, but revenue can potentially be lost in this area. Accurate coding also depends on the recording of detailed

histories for each admission or treatment, and written medical notes need to be clear and legible to allow coders to extract the relevant information. ORMIS allows clinicians and theatre staff to add coexisting conditions and relevant codes while writing operative notes. However, informal discussions with colleagues suggested that they were unaware of this, and those who were aware, assumed that other staff members would have filled in the information. A better understanding of the importance of clinical coding and the income it generates is needed, and this could be achieved through publications in peer-reviewed journals and locally arranged audits, or through continued professional development. The OPCS classification (formulated in 1987) requires procedures to be divided into multiple codes, which complicate the system and open it up to inaccuracies. Further mistakes occur because some newer procedures do not have precise codes and staff has to choose those that best describe them. Errors could also have an impact on performance league tables, as differences in clinical practice could be attributed to coding errors rather than clinical performance (6). The recording of bilateral procedures in the second cycle, which had improved by 2.5 times, suggested that the recommendations implemented after the first cycle had raised the awareness of coders. However, accuracy did not improve for most of the others, and was considerably reduced for TMJ and orthognathic procedures in the second cycle. This suggests that coders would benefit from a better understanding of the complex anatomy of the head and neck, and from the help of clinicians for more complex procedures. The lack of improvement and reduction in accuracy suggests that a single dissemination of findings is not enough, and continual dialogue between the two departments is essential. Evidence suggests that clerical workers are better at coding than clinical staff, and that partnerships between clinicians and coders improves accuracy (4,12). The appointment of a dedicated OMFS procedural coder for a short time on the OMFS ward after the first review aimed to provide coders a better insight into OMFS procedures and facilitate better communication with clinicians. However, the lack of improvement inaccuracy indicates that it was unsuccessful, possibly because the placement involved only one coder and was for a short time. A longer placement, or frequent sessions on the OMFS ward, theatres, or clinics might enable coders to learn more about the procedures. If more were encouraged to do this, information could be disseminated throughout the department. Clinicians have a critical role in ensuring that all medical records are meticulously recorded and legibly written. A collective effort by clinicians, nurses, and theatre staff to correctly describe procedures and record coexisting conditions will give the coders enough information to generate the correct HRG code for payment. Most importantly, we must ensure that the Health and Social Care

Information Centre (HSCIC) is made aware of the shortcomings in the existing OPCS classification and revenue codes, and the ambiguity concerning bilateral procedures in OMFS must be removed so that all operations are coded appropriately.

Conflict of Interest

The authors declare no conflict of interest.

Funding

None received.

References

1. Secretary of State for Health. *Delivering the NHS Plan: next steps on investment, next steps on reform*. April 2002.
2. Office of Population Censuses and Surveys. *Classification of Surgical Operations, 4th Revision*. London: HMSO; 1990.
3. World Health Organization. *International Statistical Classification of Diseases, 10th Revision*. Geneva: WHO; 1992.
4. Colville RJI, Laing JHE, Murison MSC. Coding plastic surgery operations: an audit of performance using OPCS-4. *Br J Plast Surg* 2000;53:420–422.
5. James NK, Reid CD. Plastic surgery audit codes: are the results reproducible? *Br J Plast Surg* 1991; 44: 62–4.
6. Ballaro A, Oliver S, Emberton M. Do we do what they say we do? *BJU Int* 2000;85: 389-391.
7. Mirza S, Arshad F, Watson A. Clinical coding in ear, nose and throat (ENT) operations. *BJHCM* 2011;17(12):541–544.
8. Mitra I, Malik T, Homer JJ et al. Audit of clinical coding of major head and neck operations. *Ann R Coll Surg Engl* 2009;91(3):245-248.
9. Moar, K. Impact of coding errors on departmental income: an audit of coding of microvascular free tissue transfer cases using OPCS-4 in UK. *Br J Oral Maxillofac Surg* 2012;50(1):85.

10. Health & Social Care Information Centre. *OPCS-4 eVersion*.
<http://systems.hscic.gov.uk/data/clinicalcoding/codingstandards/opcs4/eversion> (accessed 3 August 2015).
11. Health & Social Care Information Centre. *Details on mappings between primary classification codes and HRGs and the logics required to generate HRGs*.
<http://www.hscic.gov.uk/article/6103/HRG4-201516-Local-Payment-Grouper> (accessed 12 September 2015)
12. Price E. Professional Practice and Innovation: The Coding Masterpiece - a Framework for the Formal Pathways and Processes of Health Classification. *HIMJ* 2011;40(1):14-20.