



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/109475/>

Version: Accepted Version

Article:

Bulmer, Maria, Boehnke, Jan Rasmus and Lewis, Gary J. (2017) Predicting moral sentiment towards physician-assisted suicide: The role of religion, conservatism, authoritarianism, and Big Five personality. *Personality and Individual Differences*. 244–251. ISSN: 0191-8869

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

**Predicting moral sentiment towards physician-assisted suicide:
The role of religion, conservatism, authoritarianism, and Big Five
personality**

Maria Bulmer^{a,*}, Jan R. Böhnke^b, Gary J. Lewis^{c,*}

^aDepartment of Psychology, University of York, Heslington, YO10 3WF, United Kingdom

^bHull York Medical School and Department of Health Sciences, University of York, Heslington, YO10 5DD, United Kingdom

^cDepartment of Psychology, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, United Kingdom

*Corresponding authors.

E-mail addresses: bulmermaria@gmail.com (M. Bulmer)

gary.lewis@rhul.ac.uk (G.J. Lewis).

!!Do not circulate without permission of the authors!!

This is a pre-print version of the paper published as:

Bulmer, M., Böhnke, J. R., & Lewis, G. J. (2017). Predicting moral sentiment towards physician-assisted suicide: The role of religion, conservatism, authoritarianism, and Big Five personality. *Personality and Individual Differences*, 105, 244–251. <https://doi.org/10.1016/j.paid.2016.09.034>

Abstract

The issue of physician-assisted suicide is a highly contentious social issue and thus there is importance in understanding the factors that predict attitudes in this domain. In the current study we sought to examine individual differences in moral sentiment towards physician-assisted suicide with a particular focus on religion/religiosity, political ideology, authoritarianism, and Big Five personality traits, all of which were identified in an extensive review of previous studies as potentially relevant predictors. Based on N=1598 respondents from the Baylor Religion Survey (US) our results indicated an independent role for each of the predictors: being a Protestant or a Catholic (vs. no religion), higher levels of religiosity, higher levels of conservatism (vs. liberalism), and higher levels of authoritarianism uniquely predicted lower levels of support for physician-assisted suicide. Moreover, higher levels of extraversion independently predicted greater support for physician-assisted suicide. These results confirm a set of previously described predictors in an independent data set and extend prior research by showing that they independently predict moral sentiment towards physician-assisted suicide when modelled jointly. In summary, moral sentiment towards physician-assisted suicide reflects individual differences in a broad range of social and psychological factors.

Key words: physician-assisted suicide; personality; religiosity; political ideology; authoritarianism.

1. Introduction

The issue of physician-assisted suicide is one of the most contentious contemporary social debates with considerable variation in public opinion on this matter (Cohen, Van Landeghem, Carpentier, & Deliens, 2014; Emanuel, 2002). Examining the demographic, social, and psychological factors that predict such attitudes is thus of importance in order to better understand the etiology of views on this important social issue. Previous research has highlighted that education, religious denomination and religiosity, and political attitudes, among other factors, are predictive of attitudes towards physician-assisted suicide and euthanasia in general (e.g. Baume, O'Malley & Bauman, 1995; Burdette, Hill & Moulton, 2005; Sørbye, Sørbye, & Sørbye, 1995; Verbakel & Jaspers, 2010). However, this work has often been restricted to modest sample sizes (i.e. $n < 200$; Anderson & Caddell, 1993; Ho & Penney, 1992; Kemmelmeier, Wiczorkowska, Erb & Burnstein, 2002). Moreover, little work to date has comprehensively examined whether these established predictors reflect independent effects, a question of some interest given the close links between constructs such as religiosity, political conservatism, and authoritarianism (Ludeke, Johnson, & Bouchard, 2013; Saucier, 2000).

To address these issues, we used a survey sample of adults from the United States to answer the following questions: 1) are religiosity, political conservatism, and authoritarianism independently associated with moral sentiment towards physician-assisted suicide?; 2) do the Big Five personality traits provide incremental prediction for moral sentiment towards physician-assisted suicide? Next we provide a brief overview of work in the field to date.

1.1. Predicting Sentiment Towards Physician-Assisted Suicide: A Brief Overview

Although our focus in the current study specifically centers on moral sentiment towards physician-assisted suicide, many studies have used the terms active euthanasia (i.e. acting intentionally to end a person's life: Ho, 1998) and physician-assisted suicide/euthanasia (i.e. providing a patient with the knowledge or means necessary to end life: Canadian Medical Association; 2007) interchangeably (Emanuel, Daniels, Fairclough & Clarridge, 1996; Kemmelmeier, et al., 2002) and participants tend not to distinguish between these types (Ho, 1998). As such, our review of previous research includes findings concerning both forms.

A number of studies have identified predictors of attitudes towards physician-assisted suicide/euthanasia (see Table 1 for a more detailed overview). For example, several studies have reported that those with higher levels of education are more likely to be in favor of physician-assisted euthanasia (Cohen et al., 2006; Holden, 1993; Ward, 1980; Verbakel & Jaspers, 2010). Similarly, a broad body of research has overwhelmingly shown that both religious denomination and levels of religiosity predict attitudes toward euthanasia. Perhaps unsurprisingly given the condemnation of euthanasia by most organized religions (Larue, 1996), atheists are more likely to hold favorable opinions of euthanasia than Protestants and Catholics (Baume et al., 1995; Burdette et al., 2005; Cohen et al., 2006; Verbakel & Jaspers, 2010). Differences are also apparent across religious denominations with Protestants being more accepting of physician-assisted suicide than Catholics in the United States (Anderson & Caddell, 1993; Verbakel & Jaspers, 2010), Australia (Baume et al., 1995), and in much of Europe (Cohen et al., 2006; Verbakel & Jaspers, 2010). Of note, however, Cohen et al. (2006) found widely

varying attitudes toward euthanasia throughout European countries with religiosity and religious group as main predictors, which points to the importance of cultural and/or societal influences. More broadly, whereas religious denomination predicts attitudes towards physician-assisted suicide, level of religious commitment is also of relevance. For instance, a study using General Social Survey (1977-1991) data to examine the attitudes of the elderly found attendance at church services (religious denomination was not detailed) to be associated with lower levels of support for euthanasia (Leinbach, 1993), suggesting that it is not only denominational affiliation that influences attitudes towards euthanasia but also religious commitment (also see Anderson & Caddell, 1993 and Burdette et al., 2005).

Although religiosity and religious denomination are robustly associated with attitudes towards euthanasia, this effect has been noted to be accounted for by conservatism (Ho & Penny, 1992); however, other studies report independent effects of religion and political ideology (e.g. Burdette et al., 2005). Moreover, while further studies have confirmed negative links between conservatism and attitudes towards euthanasia (e.g. Burdette et al., 2005; Sørbye, Sørbye, & Sørbye, 1995), in some studies this effect has been accounted for by level of education (Ward, 1980). Finally, related work has highlighted that authoritarianism – the tendency to value traditions and social hierarchy (Altemeyer, 1981) – may also be associated with lower levels of support for euthanasia. In a sample of German university students those who self-reported higher in authoritarianism were less supportive of euthanasia (Kimmelmeier et al., 2002)¹. Of note,

¹ Note, the study by Kimmelmeier et al. (2002) was primarily concerned with the links between individualism and support for euthanasia (with the authors finding a robust positive association): the Baylor Religion Survey does not provide an individualism variable for our secondary analyses and so we do not discuss this observation further here.

however, the same study reported a null effect in a Polish sample of university students indicating that this link requires further examination. And Verbakel & Jaspers (2010), using World and European Values Survey data from 33 countries, reported that those who value autonomy more highly were more likely to be in support of euthanasia.

Relatively few studies have examined personality traits as predictors of attitudes toward euthanasia. However, of the research in this domain to date, support for euthanasia has been negatively associated with conscientiousness (Aghababaei & Wasserman, 2013) and agreeableness (Aghababaei, Wasserman & Hatami, 2014; Wasserman, Aghababaei & Nannini, 2016), and positively associated with openness (Aghababaei et al., 2014; Wasserman et al., 2016).

!!Do not circulate without permission of the authors!!

Table 1. Overview of previous work assessing multiple psychosocial predictors of attitudes towards euthanasia

| Authors | Sample | Measures | Core Findings |
|--|---|---|--|
| Aghababaei & Wasserman (2013) | <p>Participants: 284</p> <p>Demographics: 40% male, 60% female (age M=20.8, SD=2.9). All participants Muslim.</p> <p>Country: Iran</p> | <p>Definition of PAS/euthanasia: Attitude Toward Euthanasia Scale (ATE), includes active/passive, voluntary/involuntary PAS</p> <p>Variables: HEXACO Personality Inventory; Ashton & Lee, 2009), motivations toward religion (intrinsic/extrinsic/ extrinsic social), interest in religion, life satisfaction</p> | <ul style="list-style-type: none"> • Males more supportive of PAS than females • Life satisfaction (-), interest in religion (-), intrinsic and extrinsic motivations for religion (-), honesty-humility (-), conscientiousness (-) correlated with acceptance of euthanasia <p>Regression:</p> <ul style="list-style-type: none"> • Intrinsic (-) and extrinsic motivations for religion (-), interest in religion (-) significant predictors when personality, life satisfaction, age, and gender controlled for |
| Aghababaei, Hatami & Rostami (2011) | <p>Participants:233</p> <p>Demographics: 49.3% male, 50.2% female (age M=23.18)</p> <p>Country: Iran</p> | <p>Definition of PAS/euthanasia: Active and passive euthanasia examined separately using Euthanasia Attitude Scale (Tordella & Neutens, 1979)</p> <p>Variables: Big Five personality traits, motivations toward religion (intrinsic/ external social/ external individual), trolley dilemma</p> | <p>Regression:</p> <ul style="list-style-type: none"> • Internal religious orientation (-) associated with attitudes toward active euthanasia • Internal (-) and external religious orientation (-) predict combined euthanasia attitudes • Individual external religious orientation (-) predicted attitudes toward passive euthanasia |

**Aghababaei,
Wasserman &
Hatami (2014)**

Participants: 165
Demographics: 64.8% male,
35.2% female (age M=23.3,
SD=3.4). All participants
Muslim.
Country: Iran

**Definition of
PAS/euthanasia:** Euthanasia
Attitude Scale (Tordella &
Neutens, 1979), omitting “I
have faith in the medical
system to implement
euthanasia properly”
Variables: HEXACO
Personality Inventory
(examining honesty-humility,
emotionality, extraversion,
agreeableness,
conscientiousness, openness;
Ashton & Lee, 2009),
curiosity/exploration,
religiosity

- Openness (+), agreeableness (-), honesty-humility (-), extraversion (-) correlated with positive attitudes toward euthanasia
- Stepwise regression:**
- Honesty-humility, extraversion, agreeableness no longer significant when controlling for the above, religiosity, and openness
 - Openness (+) predictor of attitudes toward euthanasia

**Anderson &
Caddell (1993)**

Participants: 63 health care
(oncology) professionals
including nurses (63.5%),
pharmacists (20.6%), social
service workers (9.5%), and
others (6.3%)
Demographics: 12.7% male,
87.3% female (age M=38.43,
SD=9.26); Protestants (65%),
Catholics (22.2%), and others
(12.7%)
Country: Midwest, USA

**Definition of
PAS/euthanasia:** “Active
euthanasia”, demonstrated
through vignettes given to
participants
Variables: Religious
denomination, religiosity,
previous experience in
withholding treatment, years
in medical profession, age,
gender, marital status

- Catholics less accepting of PAS than Protestants
- Multivariate regression:**
- Religiosity (-) predicts attitudes toward PAS
 - Religious denomination not significantly related to attitudes on PAS

| | | | |
|--|---|--|--|
| Baume, O'Malley & Bauman (1995) | <p>Participants: 1,238 doctors Demographics: Catholics (19.4%), Anglicans (18.6%), non-theists (29.2%) and others; gender/age not reported Country: New South Wales, Australia</p> | <p>Definition of PAS/euthanasia: "Active voluntary euthanasia" and "Physician-assisted suicide" Variables: Religious denomination</p> | <ul style="list-style-type: none"> • Non-theists more accepting of PAS than theists • Protestants more accepting of PAS than Catholics <p>Logistic regression:</p> <ul style="list-style-type: none"> • Catholics, Protestants less accepting of PAS than non-theists |
| Burdette, Hill & Moulton (2005) | <p>Participants: 1,111 Demographics: 57% female, 43% male (age M=45); mainly white (80%); average of 13 years in education; 27% conservative religious groups, 17% no religion Country: USA</p> | <p>Definition of PAS/euthanasia: "Physician-assisted suicide" Variables: Religious denomination, religiosity, age, sex, education, region, political orientation, race, previous contact with terminal illness, support of palliative care</p> | <p>Regression:</p> <ul style="list-style-type: none"> • With all variables controlled for, race (non-whites less supportive than whites; mediated through church attendance), political conservatism (-), denomination (Conservative Protestants less supportive than non-religious), and religiosity (-) predict PAS attitudes • Religiosity accounts for effects of moderate Protestantism and Catholicism |
| Cohen et al. (2006) | <p>Participants: 41,125 Demographics: 47.5% female, 52.5% male; ages range from 18-29 (23.6%), 30-39 (19.8%), 40-49 (18.9%), 50-59 (14.7%), 60-69 (12.9%), and 70+ (9.5%) Country: 33 European countries</p> | <p>Definition of PAS/euthanasia: "Euthanasia (terminating the life of the incurably sick)" Variables: Religious denomination, self-determination, religiosity, country, age, sex, marital status, education level, social class, agricultural class</p> | <ul style="list-style-type: none"> • Acceptance of PAS varied between countries • Men more accepting than women • Education (+), age (-) correlated with acceptance of PAS • Effect of religious denomination different in different countries <p>Multivariate analysis:</p> <ul style="list-style-type: none"> • Religiosity partially explained effect of age, country, education, class |

**Danyliv & O’Neill
(2015)**

Participants: 8099, consisting of 6 different groups measured in 1983, 1984, 1989, 1994, 2005, and 2012, respectively

Demographics: Across all years: no religion (36.6%), Catholic (10%), Church of England (34.1%), other (19.3%); age/gender not reported

Country: Britain

Definition of

PAS/euthanasia: “Suppose a person has a painful incurable disease. Do you think that doctors should be allowed by law to end the patient’s life, if the patient requests it?”

(Considered active voluntary)

Variables: Religious denomination, religiosity, age, sex, household income, marital status, satisfaction with health care system, autonomy

Multivariate logistic regression

- Increase in support for PAS over time
- Religiosity strongest predictor across all years, negatively predicts support of PAS
- Catholics less supportive of PAS than the non-religious

!!!Do not circulate without permission of the author!!!

**Emmanuel,
Daniels, Fairclough
& Clarridge (1996)**

Participants: 703
Demographics: 355
oncologists (age M=48.3; 87%
male; mainly white (87.8%);
29.5% Protestant, 22.1%
Catholic, 33.7% Jewish), 155
oncology patients (age M=52.5;
39.4% male; mainly white
(94.1%); 19% Protestant, 52.9%
Catholic, 18.3% Jewish), 193
members of public (age
M=54.5; 40.4% male, mainly
white (85.5%); 23.4%
Protestant, 65.5% Catholic,
7.6% Jewish)
Country: USA

**Definition of
PAS/euthanasia:**
Description active voluntary
PAS
Variables: age, sex,
ethnicity, marital status,
religious denomination,
importance of religion,
religiosity, income,
education, employment,
health, possession of advance
care directive, participation in
decisions of end of life
Oncology patients/public:
depression, pain, physical
functioning
Patients only: support group,
self-perceived chance of cure,
disease status
Oncologists: hospital
admission in past year

Multivariate Logistic Regression:

- Religious denomination (Catholics least supportive), age (-) predicted PAS attitude
- Non-religious and higher income participants more likely to have taken steps toward euthanasia
- High religiosity predicted less consideration of euthanasia

Ho (1998)

Participants: 420
Demographics: 38.3% male,
61.7% female; aged 17 to 60
(M=31); 63% employed
Country: Australia

**Definition of
PAS/euthanasia:**
Considered active, passive,
voluntary, involuntary
euthanasia separately and in
combination
Variables: gender, age,
education, employment
status, occupation

- Active and passive euthanasia considered similarly
- Strong distinction between voluntary and involuntary euthanasia

**Ho & Penney
(1992)**

Participants: 168
Demographics: Men (40.4%), women (59.5%), aged 16 to 61 (M=29); 50% enrolled in or finished tertiary education
Country: Australia

Definition of PAS/euthanasia: Passive euthanasia and active euthanasia, examined individually
Variables: Religiosity, conservatism, abortion sex, age, education level, employment, type of employment, income

- No gender difference for PAS; weak correlations between age, education, SES, income and attitudes toward PAS
 - Religiosity (-) and conservatism (-) correlated with attitudes toward PAS and abortion
- Multiple regression:**
- With conservatism controlled, religiosity no longer predicts PAS
 - Conservatism predicted attitudes toward active and passive PAS

Holden (1993)

Participants: 922 ,
Demographics: 785 right-to-die group members (38.9% male, 61.1% female, age M=64.9), and 161 pro-life group members (34.2% male, 65.8% female, age M=41). Right-to-die group had higher proportion of Whites, Jews, non-theists, white-collar-workers, and was more educated, non-Christian, and older.
Country: California, USA

Definition of PAS/euthanasia: Not specified beyond “approval/disapproval of a terminally ill person’s right to euthanasia”
Variables: religious denomination, political stance and philosophy, sex, age, race, income, marital status, education, occupation, belief in afterlife, abortion

- Being Christian (as opposed to non-Christian) (-) associated with support for PAS
- Pro-life group: PAS attitude more strongly influenced by religious upbringing than death-proximate experiences
- Right-to-die group: PAS attitude more strongly influenced by death-proximate experiences than religious upbringing

**Kemmelmeier et al.
(2002)**

Participants:
Study 1, 100
Study 2, 102;
Study 3, 72;
Study 4, 1158
Demographics:
Study 1: 56% male, 44% female (age M=22.5, SD=2.2)
Study 2: 21.6% male, 78.4%

Definition of PAS/euthanasia: *Study 1* Euthanasia: “Help of a physician in ending the life of terminally ill person”; participants used euthanasia and PAS interchangeably
Study 2 Examined both PAS (active, voluntary), and

- Regression analyses:**
- Study 1:*
- Horizontal collectivism (+) predicts PAS attitudes
- Study 2:*
- Horizontal individualism (+) and authoritarianism (-) predict PAS attitudes
- Study 3:*
- Individualistic priming led to more positive PAS

female (age M=24.1, SD=7.5)

Study 3: 44.4% male, 55.6%

female (age M= 19.4, SD=1)

Study 4: 43.6% male, 56.4%

female

Country: Study 1: Poland,

Study 2: Germany, Studies 3, 4:

USA

involuntary euthanasia

Study 3 PAS

Study 4 Active voluntary
euthanasia/PAS

Variables:

Study 1: PAS attitude

importance, individualism,
authoritarianism

Study 2: PAS attitude

importance, individualism,
authoritarianism

Study 3: Collectivist or

individualistic self-
manipulation, PAS attitude
importance

Study 4: State individualism
(previously measured by
state, not measured for each
participant)

attitudes

Study 4:

• Individualism by state (+) correlates with PAS
attitudes

!!Do not circulate without permission of the authors!!

Leinbach (1993)

Participants: 3,980 (9 cohorts across 15 years)

Demographics: Aged 45 to 85

Country: USA

Definition of

PAS/ethanasia: “When a person has a disease that cannot be cured, do you think doctors should be allowed by law to end the patient’s life by some painless means if the patient and his family request it?”

Variables: Religious attendance, region, income, age, race, political party, socializing, employment, religious conviction, political views (27 total included for regression)

Definition of

PAS/ethanasia: “Turning off the life-sustaining machines for someone who is in a coma and will never recover consciousness” (passive involuntary), “Ending the life of someone who is severely ill and disoriented and is expected to get worse, as in Alzheimer’s Disease” (unspecific), “Ending the life of a child who is severely retarded and deformed and who will have to endure considerable pain and be institutionalized for all of his/her life” (unspecific)

• Age did not affect PAS attitude as cohort became older

Multiple Classification Analysis:

• Religious attendance, strength of religious conviction, race, region accounted for most variance in PAS attitude

Lester, Hadley & Lucas (1990)

Participants: 223

Demographics: 48% male, 52% female (age M=20, SD=1.5)

Country: Not given. Authors work in USA

Definition of

PAS/ethanasia: “Turning off the life-sustaining machines for someone who is in a coma and will never recover consciousness” (passive involuntary), “Ending the life of someone who is severely ill and disoriented and is expected to get worse, as in Alzheimer’s Disease” (unspecific), “Ending the life of a child who is severely retarded and deformed and who will have to endure considerable pain and be institutionalized for all of his/her life” (unspecific)

• Viewing euthanasia as moral (-) associated with lying

Factor analysis:

• Factor 1 defined by: viewing suicide, refusal of medical treatment, abortion, and euthanasia as moral; (+) associated with psychoticism
• Factor 2 defined by: viewing war, execution, and cannibalism as moral; (-) associated with neuroticism, lying, and irrationality

| | | | |
|---|---|--|--|
| Sorbye, Sorbye & Sorbye (1995) | <p>Participants: 289 nursing students</p> <p>Demographics: 16% male, 84% female (age M=25.4, SD=5.11)</p> <p>Country: Norway</p> | <p>Variables: Psychoticism, extraversion, neuroticism, lying, irrationality, sex, age; attitudes toward war, executions, cannibalism, suicide, refusal of treatment, abortion, euthanasia</p> <p>Definition of PAS/euthanasia: Active voluntary euthanasia</p> <p>Variables: Strength of religious belief, political conservatism, perception of life as meaningful, 3 vignettes about people of varying levels of illness wanting to die</p> | <p>Regression:</p> <ul style="list-style-type: none"> • Religious belief (-), political conservatism (-), life as meaningful (-) predict PAS attitudes |
| Verbakel & Jaspers (2010) | <p>Participants: 37,393</p> <p>Demographics: Aged 18 to 75</p> <p>Country: 31 European countries, USA, Canada</p> | <p>Definition of PAS/euthanasia: “Euthanasia” unspecified</p> <p>Variables: Religiosity, religious denomination, “slippery slope” (control over one’s life, age, employment, marital status, dependent children), social activity, widowhood, autonomy (education, attachment to personal autonomy), sex, country-level variables (permissiveness toward euthanasia, religiosity, health system, autonomy value,</p> | <ul style="list-style-type: none"> • Protestants have more favourable attitudes toward PAS than Catholics, who have more favourable attitudes than Muslims • Age (-) control over one’s life (+) social activity (+), education (+), autonomy (+), religiosity (-) predict PAS attitudes • Country-level religiosity (-), denomination (Catholic less permissive than Protestant), health case (+), autonomy (+; although insignificant with all other country-level variables controlled), suicide (+) predict PAS attitudes |

| | | | |
|--|--|--|--|
| <p>Ward (1980)</p> | <p>Participants: 1,530 Demographics: 45.3% male, 54.7% female (age M=44.7); 87.5% white Country: USA</p> | <p>health, suicide rate)</p> <p>Definition of PAS/euthanasia: “When a person has a disease that cannot be cured, do you think doctors should be allowed by law to end that patient’s life by some painless means if the patient and his family request it?”</p> <p>Variables: religiosity, political conservatism, death penalty, abortion, attitude toward suicide, age, sex, race, education, income, health, satisfaction</p> | <p>Regression:</p> <ul style="list-style-type: none"> • Age (-), education (+), religiosity (-), religion (highest acceptance in non-religious > Jews > Catholics > Protestants) predicted PAS attitudes • Males and whites more positive PAS attitudes • Those accepting PAS also accepted abortion and capital punishment • Correlation between political conservatism and PAS attitude accounted for by education level |
| <p>Wasserman, Aghababaei & Nannini (2016)</p> | <p>Participants: 165 Iranians, 156 Americans Demographics: Iran: 64.8% male, 35.2% female; USA: 38.5% male, 61.5% female. Americans were significantly older than Iranians Country: Iran, USA</p> | <p>Definition of PAS/euthanasia: Euthanasia Attitude Scale (Tordella & Neutens, 1979), omitting “I have faith in the medical system to implement euthanasia properly”</p> <p>Variables: HEXACO Personality Inventory, spirituality</p> | <ul style="list-style-type: none"> • Americans significantly more supportive of euthanasia than Iranians • Honesty-humility (-), agreeableness (-), openness (+), spirituality (-) correlated with euthanasia attitudes <p>Regression:</p> <ul style="list-style-type: none"> • With all variables controlled for, openness (+) and spirituality (-) predict euthanasia attitudes in both Iranians and Americans • Groups differed when analysed on ethical and practical consideration subscales |

Notes. PAS = physician-assisted suicide; (-) = negative association/correlation, (+) = positive association/correlation; M = mean; NB this table only includes studies that assessed multiple psychosocial predictors of physician-assisted suicide/euthanasia in order to most closely relate to the approach taken in the current study. As such, we do not include studies that, for example, only assessed religiosity.

1.2. The Current Study

While previous work has provided important foundations for understanding individual differences in attitudes towards physician-assisted euthanasia, at least two important questions remain unanswered. Firstly, while religious denomination and religiosity are robustly associated with attitudes towards physician-assisted euthanasia, it is currently unclear whether these associations reflect independent effects, or whether related constructs, such as authoritarianism and political ideology more accurately define the link. This issue of interest because the link between religious denomination and being opposed to physician-assisted suicide may be a reflection of adherence to doctrinal teachings (e.g. Christian leaders broadly condemn physician-assisted suicide), or attributable to psychological characteristics associated with religiosity – e.g. rigidity to change, traditionalism, authoritarianism (Altemeyer & Hunsberger, 1992). And these perspectives are of course not mutually exclusive. Secondly, limited work to date has addressed broader psychological links to physician-assisted suicide, such as basic dimensions of personality. To this end we sought to also examine how Big Five personality traits predict moral sentiment towards physician-assisted euthanasia.

2. Methods

2.1. Participants

We used data collected from the Baylor Religion Survey, Wave II (2007), administered by the Gallup Organization. In the first phase of data collection, Gallup contacted by telephone 1000 households using a random digit telephone sample. Of these, 624 agreed to be sent questionnaires by mail, 456 of which were completed and

returned. In a second phase, Gallup sent by mail 1836 additional questionnaires to pre-selected households in the national Random Digit Dialing database. Of these, 1192 responded, for a final sample of 1648.

Participants were aged between 18 and 96 (mean=50.95, SD=16.42). The sample consisted of 775 males (47%) and 873 females (53%) living across the United States in both rural and urban areas, and of varying socio-economic classes. Participants completed a self-administered 16-page booklet addressing a variety of issues.

2.2. Measures

2.2.1. Moral sentiment toward physician-assisted suicide

Moral sentiment towards physician-assisted suicide was assessed with the following question: *How do you feel about the morality of the following? Physician-assisted suicide*. Possible responses ranged from 1 (Always wrong) to 4 (Not wrong at all).

2.2.2. Religion

Religious denomination was measured with a question asking participants to select their religious tradition from a list of seven options. For the purpose of this study these responses were then condensed into Protestant, Catholic, Other, and None.

Religiosity was assessed with the question: *How religious do you consider yourself to be?* Possible responses ranged from 1 (Not at all religious) to 4 (Very religious).

2.2.3. Authoritarianism

Authoritarianism was measured with the following three items: *Obedience and respect are the most important things kids should learn; we must crack down on troublemakers to save our moral standards and keep law and order; people should be*

made to show respect for America's traditions. Responses were made on a 5-point Likert scale, from 1 (Strongly disagree) to 5 (Strongly agree). A score for each participant was constructed as the mean response across the three questions. Cronbach's alpha was .79.

2.2.4. Political ideology

Participants' political sentiment was measured with the question: How would you describe yourself politically? Possible responses ranged from 1 (Extremely conservative) to 7 (Extremely liberal), with the midway point (4) being Moderate.

2.2.5. Personality

Big Five personality traits – Extraversion, Agreeableness, Conscientiousness, Emotional stability/Neuroticism, and Openness to experiences – were assessed using the Ten Item Personality Measure (TIPI; Gosling, Rentfrow & Swann, 2003). Participants were asked: *Here are a number of personality traits which may or may not apply to you. Please indicate the extent to which you agree or disagree with each trait. I see myself as...*[adjective]. The adjectives were as follows: extroverted, quiet (measuring extraversion), dependable, disorganized (measuring conscientiousness) open to new experiences, uncreative (measuring openness to experiences), anxious, calm (measuring emotional stability/neuroticism) and critical, sympathetic (measuring agreeableness). Participants answered on a 5-point Likert scale from 1 (Strongly disagree) to 5 (Strongly agree). Item scores were reversed where relevant. A score for each participant for each of the Big Five traits was constructed as the mean response across the relevant two items measured from 1 (Strongly disagree) to 5 (Strongly agree). The Spearman-Brown reliability statistic ranged from .17 (openness) to .62 (extraversion).

2.2.6. Demographics

Demographic information was collected with questions about age, sex, education (“What is the highest level of education you have completed?” 1 = 8th grade or less; 7 = postgraduate work/degree), and race (White; Black or African American; American Indian or Alaska Native; Asian; Native Hawaiian or other Pacific Islander; Other: separate yes/no questions for each race).

3. Results

Descriptive statistics are presented in Table 2. In summary, the sample was largely white, with over half of participants reporting as Protestant, and almost all having at least a high school diploma. Moral sentiment towards physician-assisted suicide was fairly evenly spread, as were political orientation and religiosity.

!!Do not circulate without permission of the authors!!

Table 2. Descriptive statistics of study variables

| | N | Valid Percent |
|---|------------------|---------------|
| Moral Sentiment Towards Physician-assisted Suicide | | |
| Always wrong | 589 | 35.7% |
| Almost always wrong | 240 | 14.6% |
| Only wrong sometimes | 367 | 22.3% |
| Not wrong at all | 402 | 24.4% |
| Missing | 50 | 3.0% |
| Sex | | |
| Male | 735 | 44.6% |
| Female | 913 | 55.4% |
| Education | | |
| 8 th grade or less | 16 ^a | 1.0% |
| 9 th -12 th grade no diploma | 92 ^a | 5.6% |
| High school graduate | 369 ^b | 22.4% |
| Some college | 392 ^b | 23.8% |
| Trade/technical/vocational training | 123 ^b | 7.5% |
| College graduate | 316 | 19.2% |
| Postgraduate work/degree | 305 | 18.5% |
| Missing | 35 | 2.1% |
| Race^c | | |
| White | 1432 | 86.9% |
| Black or African American | 106 | 6.4% |
| American Indian or Alaska Native | 66 ^d | 4% |
| Asian | 13 ^d | 0.8% |
| Native Hawaiian or other Pacific Islander | 7 ^d | 0.4% |
| Other | 54 ^d | 3.3% |
| Not a single one specified | 42 ^d | 2.5% |
| Religion | | |
| Catholic | 384 | 23.3% |
| Protestant | 900 | 54.6% |
| Other | 139 | 8.4% |
| None | 175 | 10.6% |
| Missing | 50 | 3.0% |
| How religious do you consider yourself to be | | |
| Not at all religious | 173 | 10.5% |
| Not too religious | 233 | 14.1% |
| Somewhat religious | 676 | 41% |
| Very religious | 520 | 31.6% |
| Don't know | 15 | 0.9% |
| Missing | 31 | 1.9% |
| Political Liberalism | | |
| Extremely Conservative | 84 | 5.1% |
| Conservative | 433 | 26.3% |
| Leaning conservative | 159 | 9.6% |
| Moderate | 470 | 28.5% |
| Leaning liberal | 152 | 9.2% |
| Liberal | 231 | 14.0% |
| Extremely liberal | 66 | 4.0% |

| | | |
|----------------------------------|-------|-------|
| Missing | 53 | 3.2% |
| | M | SD |
| Authoritarianism (missing = 34) | 3.72 | .97 |
| Age (missing = 0) | 50.95 | 16.12 |
| Personality Traits | | |
| Extraversion (missing = 59) | 2.93 | 1.04 |
| Agreeableness (missing = 54) | 2.50 | 0.74 |
| Conscientiousness (missing = 42) | 4.04 | 0.73 |
| Neuroticism (missing = 53) | 2.66 | 0.90 |
| Openness (missing = 43) | 3.90 | 0.73 |

NOTE. ^aThese categories were combined for regression since too few respondents were in the 8th grade or less category; ^bThese categories were merged since they are not clearly ordered in terms of increase in education; ^cRespondents could choose more than one; ^dThese categories were combined since too few respondents were in them individually dummy code: “Race Other”.

!!Do not circulate without permission of the publishers!!

Ordinal logistic regression with survey weights provided by the survey team was used to examine the role of our key variables as predictors of physician-assisted suicide. Analyses were run in Stata 14 (Stata Corp, 2015) and used the SPost commands (Long & Freese, 2014). Dummy variables were created for sex (male=1) using female as the reference category, and race (White, Black, American Indian, Asian, Native Hawaiian) using White as the reference category. Dummy variables were also created for religious denomination (Catholic, Protestant, other, and no religion) with 'no religion' as the reference category. For Education, we merged 8th grade with 9-12th grade because of the low numbers of 8th graders (n=16) in the data set. We also merged the categories High School Graduate, Some College, and Trade/technical/vocational training, since they are not clearly ordered in terms of increases in education level and reflect broadly equivalent levels of achievement.

The model revealed a number of significant effects. Support for physician-assisted suicide was positively predicted by age, level of education, being White (compared to being Black), having no religious denomination (compared to being Protestant or Catholic), higher levels of political liberalism, lower levels of religiosity, and higher levels of extraversion (see Table 3).

Table 3. Results of weighted ordinal logistic regression analyses predicting Moral Sentiment Towards Physician-assisted Suicide; results presented in odds-ratios

| | Observed Data ^a | CI95% | Imputed Data ^b | CI95% |
|-----------------------------|----------------------------|--------------|---------------------------|--------------|
| Age | 1.01 ^{**} | 1.00 - 1.02 | 1.01 ^{**} | 1.00 - 1.02 |
| Education | 1.18 [*] | 1.01 - 1.38 | 1.20 [*] | 1.03 - 1.40 |
| Sex | 0.83 | 0.64 - 1.07 | 0.84 | 0.66 - 1.07 |
| Black ^c | 0.21 ^{***} | 0.09 - 0.50 | 0.23 ^{***} | 0.11 - 0.46 |
| Race Other ^c | 1.12 | 0.73 - 1.70 | 1.09 | 0.74 - 1.59 |
| Protestant ^d | 0.45 ^{**} | 0.28 - 0.73 | 0.56 ^{**} | 0.36 - 0.86 |
| Catholic ^d | 0.46 ^{**} | 0.28 - 0.78 | 0.60 [*] | 0.37 - 0.97 |
| Religion Other ^d | 0.91 | 0.48 - 1.73 | 1.02 | 0.57 - 1.86 |
| Religiosity | 0.45 ^{***} | 0.38 - 0.54 | 0.48 ^{***} | 0.40 - 0.57 |
| Political Liberalism | 1.37 ^{***} | 1.25 - 1.50 | 1.35 ^{***} | 1.24 - 1.47 |
| Authoritarianism | 0.87 | 0.76 - 1.01 | 0.83 [*] | 0.73 - 0.96 |
| Extraversion | 1.19 ^{**} | 1.06 - 1.34 | 1.17 ^{**} | 1.05 - 1.31 |
| Agreeableness | 0.98 | 0.82 - 1.17 | 1.01 | 0.85 - 1.20 |
| Conscientiousness | 1.06 | 0.89 - 1.26 | 1.06 | 0.90 - 1.25 |
| Neuroticism | 1.15 | 1.00 - 1.34 | 1.21 | 0.97 - 1.29 |
| Openness | 1.05 | 0.87 - 1.27 | 1.05 | 0.88 - 1.26 |
| Threshold 1 | 0.31 | 0.06 - 1.55 | -1.05 | -2.62 - 0.53 |
| Threshold 2 | 0.79 | 0.16 - 3.92 | -0.18 | -1.75 - 1.39 |
| Threshold 3 | 3.00 | 0.60 - 15.00 | 1.13 | -0.44 - 2.70 |
| Observations | 1,427 | | 1,598 | |

Note. *** p<0.001, ** p<0.01, * p<0.05; ^aPseudo-R² = .16; ^bBased on 40 multiple imputation chained equation runs based on all independent variables; ^cWhite as reference category; ^dNo religion as reference.

As a sensitivity analysis we used multiple imputation by chained equations (Azur, Stuart, Frangakis, & Leaf, 2011) to impute missing values on the independent variables (40 imputations, 100 burn-in iterations, overall 4000 iterations). The imputed data results are presented in Table 3. The results from this subsidiary analysis remained largely unchanged from those in our principal analysis, with two exceptions. First, the odds ratio for Catholic (compared to No Religion) was notably different – changing from 0.46 to 0.60, and with wider confidence intervals – although still in the same direction and still significant. Second, authoritarianism was now formally significant, with higher values of authoritarianism predicting lower levels of support for physician-assisted suicide.

The ordinal logistic regression model assumes that the link function between each predictor and each category of the dependent variable has the same shape. This can be examined with the Brant test (Brant, 1990; Williams, 2006), which assesses whether binary logistic regressions result in the same set of regression coefficients, independent of how the dependent variable has been dichotomised (i.e., 1 vs. 2+3+4; 1+2 vs. 3+4; 1+2+3 vs. 4). The test indicated potential violations for four of the sixteen variables. For education level ($\chi^2_{df=2} = 10.10, p = .006$) the relationship with moral sentiment towards physician-assisted suicide decreased in strength from $b = .39$ to $.06$; for religiosity ($\chi^2_{df=2} = 21.21, p < .001$) the relationship decreased from $b = -1.11$ to $-.62$; for political liberalism ($\chi^2_{df=2} = 16.52, p < .001$) the relationship decreased from $b = .42$ to $.18$; and for openness ($\chi^2_{df=2} = 10.41, p = .006$) the relationship with moral sentiment towards physician assisted suicide increased from $b = -.01$ to $.22$. In summary, then, education level, religiosity, and political liberalism were predictors of moral sentiment towards physician-assisted suicide; however, the magnitude of these predictions was less

pronounced among those holding higher levels of support for physician-assisted suicide. And the reverse was true for openness, here showing greater predictive power among those holding lower levels of support for physician-assisted suicide.

4. Discussion

A range of studies have examined individual differences in attitudes towards physician-assisted suicide, highlighting a number of predictors, including education level (Verbakel & Jaspers, 2010), religious denomination and religiosity (Cohen et al., 2006; Verbakel & Jaspers, 2010), authoritarianism (Kemmelmeier et al., 2002), and political ideology (Ho & Penny, 1992). Little work to date, though, has sought to examine the independent effects of such predictors. This is an important task because of the often moderate-to-large associations observed for variables such as religiosity, political ideology, and authoritarianism (e.g. Ludeke et al., 2013; Saucier, 2000). In addition, we sought to examine whether Big Five personality traits provided incremental prediction.

We observed a number of independent predictors of support for physician-assisted suicide: specifically, age (older respondents were more supportive), higher levels of education, being White (compared to being Black), having no religious denomination (compared to being Protestant or Catholic), higher levels of political liberalism, lower levels of religiosity, and higher levels of extraversion. Authoritarianism was not a significant predictor in our initial analysis, but in our sensitivity analyses (using multiple imputation to handle missing values) we observed that lower levels of authoritarianism predicted support for physician-assisted suicide.

These results broadly conform to findings of previous studies (e.g. Kemmelmeier et al., 2002; Leinbach, 1993; Sørbye et al., 1995; Verbakel & Jaspers, 2010), although provide the additional information that the reported effects represent independent associations (see more discussion on this point below), as well as showing that personality – notably, trait extraversion – holds incremental prediction. It should be noted, however, that our finding of a positive association between extraversion and physician-assisted suicide sits in contrast to work by Aghababaei and colleagues (Aghababaei & Wasserman, 2013; Aghababaei et al., 2014; Wasserman et al., 2016) who reported negative links with agreeableness and conscientiousness, and positive links with openness. These contrasting findings might reflect differences between the US and Iran (where the majority of the prior personality/euthanasia research was conducted), or measurement instrument (TIPI vs. HEXACO), and so further research is recommended. More generally, these observations highlight that moral sentiment towards physician-assisted suicide reflect a large number of underpinning factors, some of which provide moderate prediction (e.g. religious denomination) whereas other factors are more modest in their levels of prediction (e.g. authoritarianism, extraversion). These results, then, highlight that physician-assisted suicide is a complex social issue with many underlying determinants.

A number of these findings are of particular interest. Prior to our study, while it was apparent that both religion and authoritarianism were associated with moral sentiment towards physician-assisted suicide, it was unclear whether these associations represented independent effects. As noted earlier, such a relationship may be a reflection of adherence to doctrinal teachings, or because of psychological characteristics that are

associated with religiosity – e.g. rigidity to change, traditionalism, authoritarianism (Altemeyer & Hunsberger, 1992) – driving attitudes towards physician-assisted suicide. Our findings are consistent with both accounts, although the link with religious denomination was most pronounced and these results might be taken as evidence for the role of religious identity driving attitudes concerning physician-assisted suicide rather than rigidity to social norms per se. In addition, the results of the Brant test illustrate that some predictors may matter more for differentiating between those who are less supportive of physician-assisted suicide (i.e. Education, Religiousness, Political Liberalism), while others may only matter for differentiating between those showing greater support for physician-assisted suicide (i.e. Openness to Experience). To our knowledge such non-linear relationships have not yet been explored and thus may represent a promising avenue for future research.

Moral sentiment towards physician-assisted suicide and its determinants matter in several contexts. For example, patients' moral sentiments towards physician-assisted suicide are more favorable and homogenous once they are facing severe illness or death, which has been interpreted as a call for legislative/ societal action (Hendry et al., 2013). Attitudes of doctors (Cohen, Van Wesemael, Smets, Bilsen, & Deliens, 2012; Emanuel, Onwuteaka-Philipsen, Urwin, & Cohen, 2016) and the general population (as discussed above) are far more varied and see this as a more contentious issue. The determinants of moral sentiment towards physician-assisted suicide can thus help to clarify the underlying issues at least within a cultural context and help building a framework for discussion and consensus finding on this topic.

4.1. Strengths and Limitations

A clear strength of the study is the use of a large survey sample, which improves on the quality of a number of related studies currently in the field (Anderson & Caddell, 1993; Ho & Penney, 1992; Kemmelmeier et al., 2002). It further allowed us to control for a number of factors that are known to be relevant correlates of attitudes towards physician-assisted suicide. In addition, the use of an imputation procedure as a sensitivity analysis further reduced bias introduced by selective non-response.

A number of limitations require mention. Firstly, our single-item measure of moral sentiment towards physician assisted suicide. The term, although previously accepted as interpreted similarly to active euthanasia (Baume et al., 1995), does not differentiate between active and passive euthanasia, leaving the potential for open interpretation by participants. In addition, the observation of non-linear prediction of moral sentiment towards physician-assisted suicide may reflect methodological artifacts such as response-styles (Wetzel, Böhnke, Brown, 2016) that are more prevalent in single-item measures. Future work, then, is recommended to use more sophisticated assessment of attitudes regarding physician-assisted suicide. Secondly, this study used archival data and was unable to determine the selection of questions. As such, we were unable to include some broader variables that previous studies have found to be relevant, such as individualism (Kemmelmeier et al., 2002). In addition, the abbreviated version of our measures for authoritarianism (3 items) and Big Five traits (2 items per dimension) were not ideal (see reliabilities reported in method section). The challenge of balancing large-scale data collection with psychometrically sound instruments is well-known especially for personality research (Gosling et al., 2003; Rammstedt, & Beierlein, 2014). It is

important to note, however, that scales with just a small number of items, particularly when attempting to assess a broad construct space, such as is the case with Big Five personality traits, will typically produce conventionally unacceptable internal reliability estimates (Rammstedt, & Beierlein, 2014). With this in mind, some authors have recommended using alternative metrics for validating short-form instruments, such as test-retest reliability and convergent validity (Rammstedt, & Beierlein, 2014; Ziegler, Kemper, & Krueger, 2014). Of note, the TIPI has shown acceptable performance in both of these domains (Gosling et al., 2003; Rammstedt & John, 2007) indicating the utility of this instrument. Nonetheless, such brief instruments should only be used when time constraints force the choice between a short-form personality assessment versus no personality assessment (Rammstedt, & Beierlein, 2014). Therefore, future work is recommended to use longer-form measures or adaptive assessments (Makransky, Mortensen, & Glas, 2013) in order to more accurately assess personality traits and their links to attitudes concerning physician-assisted suicide. Fourthly, while the significant predictors were largely robust across the full range of the dependent variable, we observed that this was not the case for education, religiosity, political liberalism, and openness. These variables were less able to differentiate respondents at the top end (at the bottom end for openness) of our dependent variable. Finally, while this was a large survey sample and the use of the survey weights should adjust for over-/under-sampling from the US population, our results are limited in their ability to be generalized outside the United States as there are wide differences in euthanasia attitudes across European countries, depending on factors such as religious belief and national traditions (Cohen et

al., 2006). Moreover, this data was collected in 2007 and attitudes toward euthanasia change over time (Danyliv & O'Neill, 2015).

4.2. Conclusions

In conclusion, this study built upon previously identified predictors of attitudes toward physician-assisted suicide by controlling for other, often linked, predictors and determined that education, race, religious denomination, strength of religiosity, political orientation, and authoritarianism were all independent predictors of these attitudes. In addition, we found that extraversion provided incremental prediction for attitudes towards physician-assisted suicide.

5. Declaration of Conflicting Interest

The authors declare that they do not have any conflicts of interest.

!!Do not circulate without permission of the authors!!

References

- Aghababaei, N., Hatami, J., & Rostami, R. (2011). The role of individual characteristics and judgment pattern in attitude toward euthanasia. *Iranian Journal of Critical Care Nursing, 4*, 23-32.
- Aghababaei, N., & Wasserman, J. A. (2013). Attitude toward euthanasia scale: Psychometric properties and relations with religious orientation, personality, and life satisfaction. *American Journal of Hospice and Palliative Medicine, 30*, 781-785.
- Aghababaei, N., Wasserman, J. A., & Hatami, J. (2014). Personality factors and attitudes toward euthanasia in Iran: Implications for end-of-life research and practice. *Death Studies, 38*, 91-99.
- Altemeyer, R. A. (1981). *Right-wing authoritarianism*. Winnipeg: University of Manitoba Press.
- Altemeyer, B., & Hunsberger, B. (1992). Authoritarianism, religious fundamentalism, quest, and prejudice. *International Journal for the Psychology of Religion, 2*, 113-133.
- Anderson, J. G., & Caddell, D. P. (1993). Attitudes of medical professionals toward euthanasia. *Social Science & Medicine, 37*, 105-114.
- Azur, M. J., Stuart, E. A., Frangakis, C., & Leaf, P. J. (2011). Multiple imputation by chained equations: what is it and how does it work?. *International Journal of Methods in Psychiatric Research, 20*, 40-49.
- Baume, P., O'Malley, E., & Bauman, A. (1995). Professed religious affiliation and the practice of euthanasia. *Journal of Medical Ethics, 21*, 49-54.

- Baylor University. (2007). *The Baylor religion survey, wave II*. Waco, TX: Baylor Institute for Studies of Religion.
- Brant, R. (1990). Assessing proportionality in the proportional odds model for ordinal logistic regression. *Biometrics*, *46*, 1171-1178.
- Burdette, A. M., Hill, T. D., & Moulton, B. E. (2005). Religion and attitudes toward physician-assisted suicide and terminal palliative care. *Journal for the Scientific Study of Religion*, *44*, 79-93.
- Canadian Medical Association. (2007). *Euthanasia and assisted suicide* (update 2007). Ottawa, ON
- Cohen, J., Marcoux, I., Bilsen, J., Deboosere, P., van der Wal, G., & Deliens, L. (2006). European public acceptance of euthanasia: socio-demographic and cultural factors associated with the acceptance of euthanasia in 33 European countries. *Social Science & Medicine*, *63*, 743-756.
- Cohen, J., Van Landeghem, P., Carpentier, N., & Deliens, L. (2014). Public acceptance of euthanasia in Europe: a survey study in 47 countries. *International Journal of Public Health*, *59*, 143-156.
- Cohen, J., Van Wesemael, Y., Smets, T., Bilsen, J., & Deliens, L. (2012). Cultural differences affecting euthanasia practice in Belgium: One law but different attitudes and practices in Flanders and Wallonia. *Social Science and Medicine*, *75*, 845-853.
- Danyliv, A., & O'Neill, C. (2015). Attitudes towards legalising physician provided euthanasia in Britain: The role of religion over time. *Social Science & Medicine*, *128*, 52-56.

- Emanuel, E. J. (2002). Euthanasia and physician-assisted suicide: a review of the empirical data from the United States. *Archives of Internal Medicine*, *162*, 142-152.
- Emanuel, E. J., Daniels, E. R., Fairclough, D. L., & Clarridge, B. R. (1996). Euthanasia and physician-assisted suicide: attitudes and experiences of oncology patients, oncologists, and the public. *The Lancet*, *347*, 1805-1810.
- Emanuel, E. J., Onwuteaka-Philipsen, B. D., Urwin, J. W., & Cohen, J. (2016). Attitudes and practices of euthanasia and physician-assisted suicide in the United States, Canada, and Europe. *Journal of the American Medical Association*, *316*, 79-90.
- Gosling, S. D., Rentfrow, P. J., & Swann, W. B., Jr. (2003). A very brief measure of the Big Five personality domains. *Journal of Research in Personality*, *37*, 504-528.
- Hendry, M., Pasterfield, D., Lewis, R., Carter, B., Hodgson, D., & Wilkinson, C. (2013). Why do we want the right to die? A systematic review of the international literature on the views of patients, carers and the public on assisted dying. *Palliative Medicine*, *27*, 13-26.
- Ho, R. (1998). Assessing attitudes toward euthanasia: An analysis of the subcategorical approach to right to die issues. *Personality and Individual Differences*, *25*, 719-734.
- Ho, R., & Penney, R. K. (1992). Euthanasia and abortion: Personality correlates for the decision to terminate life. *Journal of Social Psychology*, *132*, 77-86.
- Holden, J. (1993). Demographics, attitudes, and afterlife beliefs of right-to-life and right-to-die organization members. *Journal of Social Psychology*, *133*, 521-527.
- Kemmelmeier, M., Wieczorkowska, G., Erb, H. P. & Burnstein, E. (2002). Individualism, authoritarianism, and attitudes toward assisted death: Cross-cultural, cross-regional, and experimental evidence. *Journal of Applied Social Psychology*, *32*, 60-85.

- Larue, G. A. (1996). *Playing God: fifty religions' views on your right to die (Vol. 9)*. Moyer Bell Ltd.
- Leinbach, R. M. (1993). Euthanasia attitudes of older persons a cohort analysis. *Research on Aging, 15*, 433-448.
- Lester, D., Hadley, R. A., & Lucas, W. A. (1990). Personality and a pro-death attitude. *Personality and Individual Differences, 11*, 1183-1185.
- Long, J.S. & Freese, J. (2014). *Regression models for categorical dependent variables using Stata* (3rd ed.). College Station, TX: Stata Press.
- Ludeke, S., Johnson, W., & Bouchard, T. J. (2013). Obedience to traditional authority: A heritable factor underlying authoritarianism, conservatism and religiousness. *Personality and Individual Differences, 55*, 375-380.
- Makransky, G., Mortensen, E. L., & Glas, C. A. (2013). Improving personality facet scores with multidimensional computer adaptive testing: an illustration with the NEO PI-R. *Assessment, 20*, 3-13.
- Rammstedt, B., & Beierlein, C. (2014). Can't we make it any shorter? The limits of personality assessment and ways to overcome them. *Journal of Individual Differences, 35*, 212-220.
- Rammstedt, B., & John, O. P. (2007). Measuring personality in one minute or less: A 10-item short version of the Big Five Inventory in English and German. *Journal of Research in Personality, 41*, 203-212.
- Saucier, G. (2000). Isms and the structure of social attitudes. *Journal of Personality and Social Psychology, 78*, 366-385.

- Sørbye, L. W., Sørbye, S., & Sørbye, S. W. (1995). Nursing students' attitudes towards assisted suicide and euthanasia. *Scandinavian journal of caring sciences*, 9, 119-122.
- StataCorp (2015). *Stata Statistical Software: Release 14*. College Station, TX: StataCorp LP.
- Verbakel, E., & Jaspers, E. (2010). A comparative study on permissiveness toward euthanasia religiosity, slippery slope, autonomy, and death with dignity. *Public Opinion Quarterly*, 74(1), 109-139.
- Ward, R. A. (1980). Age and acceptance of euthanasia. *Journal of Gerontology*, 35, 421-431.
- Wasserman, J. A., Aghababaei, N., & Nannini, D. (2016). Culture, personality, and attitudes toward euthanasia: A comparative study of university students in Iran and the United States. *OMEGA-Journal of Death and Dying*, 72, 247-270.
- Wetzel, E., Böhnke, J.R., Brown, A. (2016). Response biases. In F.T.L. Leong, D. Bartram, F.M. Cheung, K.F. Geisinger, & D. Ilescu (Eds.), *The ITC International Handbook of Testing and Assessment* (p. 349-363). New York: Oxford University Press.
- Williams, R (2006). Generalized ordered logit/partial proportional odds models for ordinal dependent variables. *The Stata Journal*, 6, 58-82.
- Ziegler, M., Kemper, C. J., & Krueger, P. (2014). Short scales - Five misunderstandings and ways to overcome them. *Journal of Individual Differences*, 35, 185-189.