

This is a repository copy of Transcription factors of the alternative NF-κB pathway are required for germinal center B-cell development.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/111317/

Version: Supplemental Material

Article:

De Silva, NS, Anderson, MM, Carette, A et al. (4 more authors) (2016) Transcription factors of the alternative NF-κB pathway are required for germinal center B-cell development. Proceedings of the National Academy of Sciences, 113 (32). pp. 9063-9068. ISSN 0027-8424

https://doi.org/10.1073/pnas.1602728113

© 2016, National Academy of Sciences. This is an author produced version of a paper published in Proceedings of the National Academy of Sciences. Uploaded in accordance with the publisher's self-archiving policy. In order to comply with the publisher requirements the University does not require the author to sign a non-exclusive licence for this paper.

Reuse


Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/



BLIMP1


DAPI


p100/p52

Ε DLBCL MM *CL CL C* $NF-\kappa B$ IRF4 p100 p52 NF-κB RELB c-REL RELA p50 IRF4 β-actin

В


GC light zone





p100/p52 DAPI

p100/p52 CD20



p100/p52

c-REL