White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Leeds.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/112175/

Version: Accepted Version

Article:
Williscroft, K, Grasby, SE, Beauchamp, B et al. (2017) Extensive Lower Cretaceous (Albian) methane seepage on Ellef Ringnes Island, Canadian High Arctic. Geological Society of America Bulletin, 129 (7-8). pp. 788-805. ISSN: 0016-7606

https://doi.org/10.1130/B31601.1

© 2017 Geological Society of America. This is an author produced version of a paper published in Geological Society of America Bulletin. Uploaded in accordance with the publisher's self-archiving policy.

Reuse
Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Extensive Lower Cretaceous (Albian) methane seepage on Ellef Ringnes Island, Canadian High Arctic

Krista Williscroft[1], Stephen E. Grasby[1,2*], Benoit Beauchamp[1], Crispin T.S. Little[3], Keith Dewing,[1,2], Daniel Birgel[4], Terry Poulton[2], Krzysztof Hryniewicz[5]

[1]Department of Geoscience, University of Calgary, 2500 University DR. NW, Calgary, AB, T2N 1N4, Canada

[2]Geological Survey of Canada, Natural Resources Canada 3303 33rd Street NW, Calgary, AB, T2L 2A7, Canada

[3] School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK

[4] Institute of Geology, University Hamburg, 20146 Hamburg, Germany

[5] Institute of Paleobiology, Polish Academy of Sciences, ul. Twarda 51/55, 00-818 Warszawa, Poland

*Corresponding author: steve.grasby@canada.ca

ABSTRACT

During field mapping of Ellef Ringnes Island, Canadian Arctic Archipelago, 139 isolated Lower Cretaceous methane seep deposits were found from 75 field sites. Stable isotopes of the carbonates have values of $\delta^{13}C$= -47‰ to -35‰ and $\delta^{18}O$= -4.0‰ to +0.7‰. Isoprenoids in organics from one of the seeps are significantly depleted in ^{13}C, with the most negative $\delta^{13}C$ of = -118 ‰ and -113 ‰ for PMI and phytane/crocetane, respectively. These values indicate an origin through methane oxidation, consistent with biomarkers that are characteristic for anaerobic methanotrophic archaea within the seep deposits, accompanied by terminally-branched fatty acids sourced by sulphate-reducing bacteria, showing similar ^{13}C values (-92‰). The seep deposits contain a moderate diversity macrofaunal assemblage containing ammonites, bivalves, gastropods, scaphopods, 'vestimentiferan' worm tubes and brachiopods. The assemblage is dominated numerically by species that probably had chemosymbionts. The seep deposits formed in the subsurface with strong redox zones, in an otherwise normal marine setting, characterised by oxic waters at high paleolatitudes.

While geographically widespread, over an area of ~10,000 km^2, seep deposits on Ellef Ringnes Island occur in a narrow stratigraphic horizon, suggesting a large release of biogenic methane occurred over a brief period of time. This gas release was coincident with a transition from a cold to warm climate during the latest Early Albian, and we hypothetize that this may relate to gas hydrate release.

1.0 INTRODUCTION

Methane seepage into modern marine environment was first recognized at the foot of the Florida Escarpment (Paull et al., 1984). Since then, the seepage of methane-enriched fluids from the seabed into the water column has been found to be a common feature along continental margins, forming in many cases distinctive seafloor feature such as pockmarks and mud volcanoes, and ranging from isolated occurrences to high density fields on the sea floor (e.g. Judd and Hovland, 2007; Kulm et al., 1986; Suess, 2014). At seep sites, a major proportion of the methane is consumed in sub-surface sediments by sulfate-dependent anaerobic oxidation of methane (AOM) biogeochemical process (Hoehler et al., 1994), mediated by anaerobic methane oxidizing archaea (ANMEs) and sulfate-reducing bacteria (SRB) (Boetius et al., 2000; Hinrichs et al., 1999; Milucka et al., 2012; Orphan et al., 2001). AOM produces an excess of dissolved inorganic carbon (DIC), promoting the rapid precipitation of authigenic carbonates depleted in ^{13}C as a by-product (Paull et al., 1984; Ritger et al., 1987). These carbonates have complex cement fabrics (e.g. Aloisi et al., 2000; Haas et al., 2010; Naehr et al., 2007; Ritger et al., 1987), and a wide variety of morphologies (e.g. nodules, tubular/tabular concretions, cemented breccias and pavements) and sizes (e.g. Campbell, 2006). The seep carbonates are often exhumed onto the seafloor by sediment erosion and can then act as hard substrates for attached epifaunal animals (e.g. serpulid tubeworms). Active methane seeps support diverse and high-biomass communities of macrofauna, which are dominated by animals having symbiotic relationships with chemotrophic bacteria (principally methanotrophs and thiotrophs). These taxa include bivalves (e.g. solemyid, vesicomyid, lucinid and thyasirid clams, and bathymodiolin mussels) and siboglinid (vestimentiferan) tubeworms (e.g. Dubilier et al., 2008; Levin, 2005; Paull et al., 1984; Sibuet and Olu, 1998).

One of the first identifications of methane seeps in the geological record was from Lower Cretaceous strata in the Sverdrup Basin (Fig. 1) (Beauchamp et al., 1989). In that study, four methane seep deposits were described, two each on Ellef Ringnes and Prince Patrick islands in the remote Canadian High Arctic, and the first linkage with modern cold methane seeps was made. Subsequent detailed petrographic studies of samples from these deposits (Beauchamp and Savard, 1992; Savard et al., 1996) identified a suite of early phase carbonate cements (calcite microspar, fibrous botryoidal and splayed calcite and anhedral yellow calcite) with very negative $\delta^{13}C$ isotope values derived from methane and, on Ellef Ringnes Island only, late phase carbonate cements (saddle dolomite, bladed calcite and coarse anhedral calcite) with less negative $\delta^{13}C$ values formed during burial diagenesis. Savard et al. (1996) showed that original aragonite is present in the early phase cements from the seep deposits from both islands; this constituted the first recognition of aragonite precipitation at ancient methane seeps deposited under 'calcite sea' conditions. Beauchamp et al. (1989) and Beauchamp and Savard (1992) recorded a biota of molluscs (abundant bivalves, smaller numbers of ammonites and gastropods), worm tubes (larger diameter serpulids and smaller diameter possible serpulids), foraminifera, fish teeth and wood fragments from all four seep deposits, and, in addition, terebratulid brachiopods (later described as *Modestella jeletzkyi* (Sandy, 1990)) and coiled spirorbid tubes in the southern deposit of Prince Patrick Island.

Since recognition of the Cretaceous Sverdrup Basin examples, many ancient fossil-rich seep deposits have been recognised in the rock record throughout the world, from as old as the Devonian, or even Silurian, with similar paragenetic cement phases and stable isotopic compositions (see Campbell, 2006 and references therein). In addition, organic biomarkers for AOM have been discovered in seep deposits back to the Carboniferous (Birgel et al., 2008a).

Like modern examples, ancient seep deposits have a wide variety of morphologies, but lens shapes at various scales (metres to 10s metres in diameter and centimetres to metres high) are common (Campbell, 2006). These are often exposed on the land surface as indurated mound-shaped structures, due to the erosion of enclosing fine grained sediments. Ancient seep carbonates can occur as isolated deposits, but are usually found with other examples in the same sedimentary sequence, sometimes at high density. For example, in Late Cretaceous sediments of the Western Interior Seaway, USA, up to 13 'Tepee Buttes' seep deposits crop out over 25 km^2 area (Cochran et al., 2015), 15 Late Jurassic to Early Cretaceous seep deposits crop out along 10 km of exposure in Spitsbergen (Hammer et al. 2011), and numerous Paleocene seep deposits are exposed over 5 km in the Panoche Hills, California (Schwartz et al., 2003).

The original studies of the Sverdrup Basin seep deposits (Beauchamp et al., 1989; Beauchamp and Savard, 1992; Savard et al., 1996). were based on only a few hand samples collected as part of a regional mapping expedition in the mid 1970s (Nassichuk and Roy, 1975). While this allowed recognition of methane seeps in the geologic record, the overall context of the sites was poorly constrained. For three decades no additional field work had been conducted on Ellef Ringnes Island given the remoteness, and the seep deposits there were thought to be isolated occurrences. However, a series of expeditions by the Geological Survey of Canada between 2009 and 2011 led to the discovery of 137 new methane seep deposits across the island, a remarkable density that rivals any other occurrence of seepage in the geologic record (Campbell, 2006). Here we give an integrated morphological, geochemical and paleontological description of these newly discovered seep deposits, analyse their geographic and stratigraphic context, and constrain the style, duration and distribution of seafloor seepage in the Sverdrup Basin in context of Lower Cretaceous climate change.

2.0 GEOLOGIC SETTING

Ellef Ringnes Island is part of the Canadian High Arctic, Nunavut, Canada (Fig. 1). It comprises Jurassic through Paleogene sediments of the Sverdrup Basin, a major Carboniferous to Paleocene depocentre (Embry and Beauchamp, 2008). The basin is a ~1000 km long, north-easterly trending depression filled with up to 13 km of marine and non-marine sediments, as well as Cretaceous (predominantly Aptian) basaltic flows and mafic dykes and sills (Balkwill, 1978) (Fig. 2). The Sverdrup Basin originated during Carboniferous-Early Permian rifting (Balkwill, 1978). Deepening and enlargement of the basin resulted in marine transgression and the deposition of over 400 m of evaporites of the Upper Carboniferous Otto Fiord Formation (Nassichuk and Davies, 1980; Thorsteinsson, 1974). Later lithostatic loading, related to the high clastic sediment flux in the Triassic, initiated diapirism of Otto Fiord Formation evaporites in the central part of the Sverdrup Basin (Gould and DeMille, 1964; Macauley et al., 2000; Thorsteinsson, 1974). Stratigraphic evidence shows that diapirs were still mobile during the Cretaceous (Dewing et al., 2016), during the time when seep deposits described here were formed.

Subsidence and sedimentation increased in the basin during renewed rifting and extension in the Upper Jurassic to Lower Cretaceous (Embry and Beauchamp, 2008). Sediment supply rates greatly increased and normal faults were active. Thick, coarse-grained fluvial sediments (Isachsen Formation) and offshore muds and silts (Christopher Formation) were deposited from the Hauterivian to the Late Albian. Marginal uplift in the mid-Aptian occurred in the northeast Sverdrup Basin, where common volcanic units (sills and basalt flows) were emplaced between 130 and 90 Ma (Embry and Beauchamp, 2008; Saumur et al., 2016).

Methane seep deposits in this study are found within the Upper Aptian to Upper Albian Christopher Formation that is composed of medium-grey to black silty mudstone and fine siltstone, and is divided into the lower Invincible Point and upper MacDougall Point members. The two members are separated by a resistant reddish-brown volcanogenic sandstone and hyalotuff (marker bed in Fig. 3) (Embry, 1985), that was dated at 105.4 +/-0.22 Ma (Evenchick et al., 2015). During Christopher Formation deposition, the Sverdrup Basin was located at ~72° N, relatively close to its current latitude of ~78° N (Wynne et al., 1988),

The Sverdrup Basin rock units on Ellef Ringnes Island are characterized by a general southward dip, which exposes Lower Jurassic rocks in the northwest regions of the island and uppermost Cretaceous rocks in the southeast (Evenchick and Embry, 2012a; Evenchick and Embry, 2012b; Stott, 1969). Large scale, northwest-trending open folds are present over most of the island and deform Cretaceous and older sediments (Evenchick and Embry, 2012a; Evenchick and Embry, 2012b) (Fig. 2). Salt diapirism caused localized deformation and tilting of overlying sediments (Boutelier et al., 2011; Dewing et al., 2016). Steep faults associated with salt diapirs are radially positioned in relation to individual diapirs, displaying strikes 30° to 60° relative to the diapir boundary (Evenchick and Embry, 2012a; Evenchick and Embry, 2012b), and offsetting Cretaceous strata. Diapirs exposed at the surface on Ellef Ringnes Island include the Contour, Dumbells, Hakkon, Helicopter, Hoodoo, Isachsen, and Malloch domes (Fig. 2).

3.0 METHODS

3.1 Field Sampling

As part of the Geological Survey of Canada's Geo-Mapping for Energy and Minerals (GEM) Program, a geological mapping campaign to Ellef Ringnes Island was undertaken from 2009 to

2011, which included a focussed search for methane seep deposits in addition to those previously discovered by Beauchamp et al. (1989). The seep deposits were readily identified in the field as they form resistive, positive relief mounds of carbonate, projecting from the otherwise mudstone-dominated strata that forms a rolling landscape of Arctic tundra (Figs. 4a,b). The two seep sites originally reported by Beauchamp et al. (1989) were visited and sampled as well. All seep sites were located with a high-yield GPS receiver relative to NAD-83 (Fig. 2; Table 1). Some seep sites constituted a single carbonate body, while others had several (up to 8) individual deposits, and are listed in Table 1 as a single site. We defined an individual carbonate mound with defined edges as a 'seep deposit' and a 'seep site' as a location with one or more seep deposits in close proximity (< 5 m). The size of the seep deposits was measured as length and width parallel to bedding and height perpendicular to bedding. Multiple hand samples ranging in size from a few cm up to 30 cm in diameter were collected from different areas of 80 seep deposits. Macrofossils were collected from 35 deposits, with numbers ranging from three to 80 specimens from each site (Table 1).

3.2 Petrography

A subset of hand samples collected in the field were prepared for 30 μm thick uncovered and unpolished thin sections. Observation of these thin sections was conducted with a Nikon Eclipse E600W Polarizing Microscope at the Geological Survey of Canada in Calgary, to identify sediment type and carbonate cements.

178 **3.3 Geochemistry**

179 One hundred and eleven samples from the Ellef Ringnes seep deposits were analyzed for $\delta^{13}C$
180 and $\delta^{18}O$ values of distinct carbonate phases (micrite, yellow calcite, boytroidal calcite, and
181 blocky calcite), carbonate infilling articulated fossils ('fossil fill'), calcified wood, and carbonate
182 cemented sandstones underlying the seep deposits (Table 2). Powdered samples were obtained
183 by drilling cut rock faces, and then analysed at the Isotope Science Laboratory at the University
184 of Calgary (ISL-UofC). Approximately 2 mg of powdered material was reacted with anhydrous
185 phosphoric acid in a Y tube reaction vessel at 25 °C. The evolved CO_2 was cryogenically
186 distilled from the reaction vessel into a 6 mm Pyrex tube and flame sealed. The CO_2 gas was
187 then inlet to the ion source of a VG 903, stable isotope ratio mass spectrometer and analyzed for
188 $^{13}C/^{12}C$ and $^{18}O/^{16}O$ ratios. Selected internal standards ($\delta^{13}C$ values of -40.31 and -2.51 ‰)
189 were run at the beginning and end of the sample set and were used to normalize the data as well
190 as to correct for any instrument drift. The results are expressed as conventional δ-values relative
191 to the Vienna Peedee Belemnite standard (V-PDB). The precision and accuracy of the analysis
192 was 0.2‰ for both $\delta^{13}C_{VPDB}$ and $\delta^{18}O_{VPDB}$.

193 Carbon isotopes of organic matter were measured in four seep deposit samples from site
194 10KRW001. To characterise background host shale chemistry, an additional, 38 samples of
195 Christopher Formation mudstone were collected at approximately every 10 m in section
196 11KRW0045, from 25 m above the zone of seep deposits (which is 34 m thick; see section 4.1)
197 to the base of the formation, 400 m below the seep zone (Table 3). The mudstone and seep
198 carbonate samples were washed with hydrochloric acid, and rinsed with hot distilled water to
199 remove any carbonate before determination of $\delta^{13}C$ of organic carbon using the same methods as
200 above.

Molybdenum (Mo) concentrations in mudstones are frequently used as an indicator of paleo-redox conditions. Under anoxic/euxinic conditions molybdate ions are converted to oxythiomolybdate ions, which react with, and are sequestered by sulphide minerals or organic matter, becoming enriched in sediments (Emerson and Huested, 1991; Voegelin et al., 2009), relative to values for marine carbonates of 0.4 ppm and Post Archean Average Shale (PAAS) concentration of 1.0 ppm (Taylor and McLennan, 1985; Turekian and Wedepohl, 1961). We made elemental determinations for Mo concentration on the Christopher Formation shale samples detailed above (Table 3) and 62 seep carbonate samples (Table 2). All samples for Mo determination were powdered in an agate mortar and pestle, digested in a 2:2:1:1 acid solution of H_2O-HF-$HClO_4$-HNO_3, and subsequently analyzed using a PerkinElmer Elan 9000 mass spectrometer, with ± 2% analytical error.

3.4 Biomarkers for AOM

Four carbonate samples from a seep deposit at Hoodoo Dome (field station 10KRW001 in Table 1) were prepared and decalcified after methods described in Birgel et al. (2006a). After a saponification procedure with 6% KOH in methanol, the samples were extracted with a microwave extraction system (CEM Discovery) at 80 °C and 250 W with a dichloromethane-methanol (3:1) mixture. The total extracts were pre-cleaned by a separation into an *n*-hexane soluble and dichloromethane-soluble fraction. The *n*-hexane fraction was further separated by column chromatography into four fractions of increasing polarity (Birgel et al., 2008b). Only the hydrocarbon and the carboxylic acid fractions were found to contain compounds to constrain carbonate precipitation. Other than the hydrocarbon and carboxylic acid fractions, the alcohol fraction, was affected by thermal maturation and biodegradation and did not contain genuine

lipid biomarker signatures. This peculiarity is observed also in other ancient methane seep carbonates experiencing low to intermediate maturity (e.g. Birgel et al., 2006b; Birgel et al., 2006a; Little et al., 2015; Natalicchio et al., 2015). The carboxylic acids were found to contain two octadecenoic acids, which are rather unlikely to be preserved, especially when looking at the low to intermediate mature hydrocarbons (see also section 4.5). Both the hydrocarbons and the carboxylic acids were analyzed by coupled gas chromatography–mass spectrometry (GC-MS) with an Agilent 7890 A GC system, coupled to an Agilent 5975 C inert MSD mass spectrometer at the Department for Geodynamics and Sedimentology, University of Vienna. The carrier gas was helium. The GC temperature program used was as follows: 60 °C (1 min); from 60 °C to 150 °C at 10 °C/min; from 150 °C to 320 °C at 4 °C/min, 25 min isothermal. This temperature program was used for both fractions. Identification of individual compounds was based on retention times and published mass spectral data in comparison with other samples. Compound-specific carbon isotope analysis of molecular fossils was performed with a Thermo Fisher Trace GC Ultra connected via a Thermo Fisher GC Isolink interface to a Thermo Fisher Delta V Advantage spectrometer at the Department of Terrestrial Ecosystem Research, University of Vienna. Conditions chosen for the gas chromatograph were identical to those described above. Carbon isotopes are given as δ values in per mil relative to the Vienna Peedee belemnite (V-PDB) standard. Each measurement was calibrated using several pulses of CO_2 with known isotopic composition at the beginning and end of the run. Instrument precision was checked with a mixture of *n*-alkanes (C_{14} to C_{38}) of known isotopic composition. Analytical standard deviation was below 0.7‰.

4.0 RESULTS

4.1 Seep deposit distribution

One hundred and thirty-seven new seep deposits were discovered during the 2009 to 2011 field campaign. The addition of the two previously reported sites by Beauchamp et al. (1989) makes an extensive occurrence of 139 seep deposits on Ellef Ringnes Island. These occurred in 75 sites exclusively within the Christopher Formation. The majority were exposed along the periphery of the salt diapirs: around Hoodoo Dome (104), Dumbells Dome (17), Helicopter Dome (6), Haakon Dome (2), and Isachsen Dome (2). No seep deposits were found associated with Contour or Malloch domes (Fig. 2). Eight seep deposits were not spatially associated with an exposed salt diapir. Of these, seven were found near the centre of the island, 11.5 km or more away from the edge of the closest diapir (Fig. 2). These seep deposits were not associated with any mapped faults or fractures. One additional seep deposit was found 7.5 km southwest of Hakkon Dome, where the Christopher Formation is truncated by three large normal faults. All the seep deposits were located within the upper portion of the Invincible Point Member, below the regional volcanogenic marker unit dated at 105.4 +/-0.22 Ma (Evenchick et al., 2015) (Fig. 3). At any one site, seep deposits occur at the same defined stratigraphic level. Exposure made it difficult to trace this level between sites, but where it was possible seep mounds all appeared on the same stratigraphic horizon indicated by a siltstone bed. The stratigraphic position of the seep sites, measured relative to the regional marker unit, varied from a maximum of 42 m at Helicopter Dome to a minimum of 8 m at Dumbells Dome, a 34 m stratigraphic range. Geopetal fabrics, and bedding of a few larger deposits, have the same dip as the bedding of Christopher Formation strata that hosts them.

4.2 Seep deposit morphology

Based on their appearance in outcrop, we classified the seep deposits on Ellef Ringnes Island into three morphological categories: 1) carbonate mounds, 2) carbonate beds, and 3) carbonate crusts. These categories do not necessarily reflect the original morphology of the carbonates during formation. The carbonate mounds (Fig. 4b) were the most common observed morphology (125 individual mounds). However, this may have been partly a function of the ease of their observation from aerial helicopter survey. The carbonate mounds ranged in size from 0.2 to 6.7 m in width, 0.3 to 39.0 m in length, and 0.2 to 3.1 m in height (Table 1). The mounds were always greater in length than height, having an average length:height ratio of 4:1. They comprised large cohesive blocks of carbonate rocks up to ~1 m^3, to mounds completely composed of carbonate rubble.

The five carbonate beds were distinguished from the mounds based on their broad lateral extent and limited relief. They ranged in size from 0.1 to 4.2 m in width, 12.4 to 62.0 m in length, and 0.1 to 0.3 m in height (Table 1), with a length:height ratio greater than 15:1 (Fig. 4C). The carbonate beds were composed of carbonate rubble, rarely containing pieces of carbonate rocks larger than 10 cm^3. Three of the beds contained highly cemented calcareous siltstone, giving these beds slightly more relief than those without any siliciclastic content.

The three carbonate crusts had no relief, being composed of a thin layer (<0.2 m) of carbonate rubble. They ranged in size from 0.5 to 1.7 m in width, and 0.6 to 2.2 m in length (Table 1).

290 **4.3 Carbonate cements**

291 All the morphological variants of the seep deposits contained a number of distinct carbonate
292 cement phases (Fig. 4D), including 1) dark grey to grey-brown micrite, 2) banded calcite, 3)
293 botryoidal calcite, 4) yellow calcite, and 5) blocky calcite. The micrite formed circular to
294 elongated band-shaped areas that range from millimeters to 10s of cms in thickness. In thin
295 section, the micrite was beige to dark brown and contained grains including peloids, opaque
296 minerals, wood debris, and mollusc shells; it was also burrowed and fractured (Fig. 4E). Micrite
297 and associated peloids showed dissolution along corrosion surfaces.

298 The banded calcite was composed of beige (botryoidal calcite) and brown (yellow calcite)
299 layers that alternated in bands from 1 to 6 cm in thickness. The botryoidal calcite was composed
300 of well-developed, radiating, fibrous clusters of prismatic crystals separated into growth bands
301 indicated by opaque, submillimeter cement horizons (Fig. 4F,G). Petrographically, the botryoidal
302 calcite comprised clean and dirty phases. Clean botryoidal calcite was composed of clear
303 sweeping bands of prismatic calcite devoid of any inclusions or organic material. Individual
304 calcite crystals could only be distinguished in cross-polarised light and reached over 1mm in
305 length but were usually shorter than crystals of the other two botryoidal phases. Dirty botryoidal
306 calcite was similar to clean calcite except that it contained microcrystalline grey to brown
307 'debris' within the calcite. The dark material was dispersed throughout the whole botryoids, or
308 concentrated in horizons 70 to 200 μm apart. The submillimeter horizons were either very
309 smooth or had wavy 'V' shaped tips. Both the clean and dirty botryoidal calcite contained clear,
310 square-tipped needles that were over 2.3 mm in length and ~3.5 to 14 μm in width, observed
311 both petrographically and with SEM. Previous work has interpreted these as remnant acicular

312 aragonite needles (Savard et al., 1996). Corrosion surfaces within the botryoidal calcite layers
313 were typically associated with 10 to 50 μm dark bands of impurities.
314 The yellow calcite had a clotted, euhedral texture and did not display distinct crystal
315 boundaries. In thin section, yellow calcite was yellow to orange in colour and anhedral to clotted
316 in appearance. Individual layers of yellow calcite ranged in thickness from 150 μm to over 3 mm
317 and alternated with botryoidal calcite. The yellow calcite contained many opaque fragments,
318 between 10 to 500 μm in diameter. The amount of opaque debris within the yellow calcite was
319 variable among seep deposits, with some areas of yellow calcite containing no visible opaque
320 material. Corrosion surfaces were common at the basal contact between the yellow calcite and
321 the underlying cements it overgrew.
322 The blocky calcite was yellow to white and had a sparry texture with excellent crystal
323 development, and filled voids up to 8 cm (Fig. 4H). Petrographically, the blocky calcite crystals
324 ranged in size from < 30 μm to over 2 mm, with crystal boundaries that were typically sharp and
325 linear, but at times had a slight curve. The blocky calcite always nucleated from what was once a
326 free surface, with crystals that grew progressively larger as the void space filled in towards the
327 center. The blocky calcite did not contain organic matter or opaque grains, and was homogenous
328 in appearance. Microfractures as well as large void spaces, including centers of worm tubes,
329 were filled with blocky calcite that typically, but not always completely, filled the once-open
330 void.
331
332 **4.4 Carbon and oxygen isotopes**
333 Carbon and oxygen stable isotope data from the seep deposits fell into two distinct clusters (Fig.
334 5). Group 1 represented fossil fill, calcified wood, carbonate cement in calcareous sandstones,

335 and all of the cement phases (with the exception of the blocky calcite) and was characterised by
336 very negative $\delta^{13}C$ values ($\delta^{13}C$= -52.6‰ to -32.3‰) and $\delta^{18}O$ values around zero ($\delta^{18}O$= -7.1‰
337 to +1.3‰) (Fig. 5).
338 The blocky calcite cement formed a second sample cluster (Group 2) characterised by less
339 negative $\delta^{13}C$ values ($\delta^{13}C$ = -24.8‰ to -7.5‰) and more negative $\delta^{18}O$ values ($\delta^{18}O$= -21.3‰ to
340 -11.0‰) compared to Group 1. This included blocky cement that occurs as fill of a worm tube,
341 sample C-540385-1 in Table 2 and indicated as 'fossil fill' within the Group 2 cluster on Figure
342 5.
343 The $\delta^{13}C_{org}$ values of organic matter from the four carbonate deposit samples from site
344 10KRW001 had values of -47.9, -41.1, -38.1, and -31.7‰. In comparison, those of the
345 Christopher Formation mudstone samples ranged from -25.9 to -17.1‰ $\delta^{13}C_{org}$ (Table 3; Fig. 6).
346 There was no vertical trend in $\delta^{13}C_{org}$ values going up section towards the zone of seep deposits.
347
348 **4.5 Lipid biomarkers**
349 Biomarker analysis showed that the predominant compounds in the hydrocarbon fraction were
350 isoprenoidal hydrocarbons indicative of AOM, including head-to-tail linked phytane, tail-to-tail
351 linked crocetane and 2, 6, 10, 15, 19-pentamethylicosane (PMI), and three head-to-head linked
352 biphytanes comprising no, one, or two cyclopentane rings (Fig. 7A). Further, minor amounts of
353 pseudohomologue series of head-to-tail linked isoprenoids with 18 to 24 carbons were found.
354 Apart from the predominant isoprenoids, only minor *n*-alkanes with 16 to 23 carbon atoms and
355 terminally-branched alkanes with 16 to 19 carbon atoms were found in the hydrocarbons,
356 possibly derived from biodegraded organic matter or oil (e.g. Peters and Moldowan, 1993), and
357 entombed within the carbonate matrix. However, when comparing the high amounts of

358 isoprenoids with the only minor biodegraded lipid biomarkers, oil migration and the presence of
359 oil-derived lipid biomarkers was of minor importance for the studied samples.
360 All isoprenoids were significantly ^{13}C depleted, with the lowest $\delta^{13}C$ value of = -118 ‰
361 and -113 ‰ for PMI and phytane/crocetane, respectively. Somewhat less ^{13}C depleted values
362 were found for biphytanes (-109 ‰), and pristine (the least ^{13}C depleted compound). Most n-
363 alkane and branched alkane contents were too low to determine compound-specific isotopes, but
364 *n*-C_{23} was found, as well with a very low $\delta^{13}C$ value of -101 ‰, as reported in other methane
365 seep deposits (e.g. Chevalier et al., 2013; Peckmann et al., 2009; Thiel et al., 2001). The fatty
366 acids results (Fig. 7B) need to be used carefully, since pristine signatures were amalgamated with
367 secondary signatures, which most likely were introduced after the carbonate precipitation and the
368 entombment of the original signature of the microbial seep community. The most reliable
369 biomarkers of SRB, partners in AOM, are terminally-branched iso- and anteiso-C15 fatty acids,
370 which are prominent and also displayed the characteristic isotopic fingerprint of SRB of modern
371 sites, with $\delta^{13}C$ values of -92 ‰ and -87 ‰, respectively. Interestingly, the short-chain fatty
372 acids *n*-C_{14} and *n*-C_{16} still seem to represent original signals as well, since they were only
373 slightly less ^{13}C depleted than the branched fatty acids. This finding is rather unusual for
374 Mesozoic samples, where most often only the terminally-branched fatty acids carried the AOM
375 signature (e.g. Birgel et al., 2006b). In contrast, the two prominent octadecenoic acids probably
376 represent contaminants, since the pattern preserved in this sample is not like any patterns
377 recorded in modern seep sites (Elvert et al., 2003).
378

4.6 Paleontology

Of the 137 Ellef Ringnes Island seep deposits mapped in this study, 62 had macrofossils, 75 did not contain visible macrofossils, and two were not recorded for fossil content (Table 1). The carbonate mound morphologies typically had the highest density of macrofossils, but mounds with abundant calcareous siltstone contained few or no macrofossils. Four of the carbonate beds and two of the crusts contained macrofossils (Table 1). Fossil abundances were related to the amount of siliciclastics and size of each deposit, with high siliciclastic content and small deposits containing fewer fossils (Table 1).

The invertebrate macrofossil assemblage comprised molluscs, brachiopods and worm tubes belonging to at least 20 species (Table 4; Fig. 8). Only the nuculid bivalves and wood were previously identified from the two original Ellef Ringnes seep deposits (Beauchamp et al. 1989). The molluscs and brachiopods were preserved in the micrite cements, typically with their original shell material still present, such as nacre of nuculid bivalve and ammonite fossils. There was little evidence for transportation of shells on the seafloor, as few shells were fragmented and most of the bivalves and brachiopods were still articulated. The mollusc fauna was dominated by bivalves (n = 396; belonging to eight species; Fig. 8A-I), with smaller numbers of gastropods (n = 62; at least four species; Fig. 8K), ammonites (n = 32; at least four species; Fig. 8J) and scaphopods (n = 14). Brachiopods were rare, with only four specimens being found. The bivalve fauna included taxa with a variety of inferred paleoecologies (Table 4), the majority of which potentially had chemoautotrophic bacterial symbionts (n = 353; four species), together with small numbers of infaunal deposit feeders (n = 27; three species) and epifaunal filter feeders (n = 16; one species).

401 The ammonite fossils were frequently incomplete and were found only within large
402 mounds that contained very little siliciclastic material. Well preserved specimens were identified
403 as *Arcthoplites* (common), including *A.*(?) cf. *belli* (McLearn), as well as rarer *Cleoniceras* aff.
404 *canadense* Jeletzky, *Beudanticeras*(?), *Puzosia* aff. *sigmoidalis* Donovan (Fig. 8J) and
405 *Freboldiceras* aff. *irenense* (McLearn). Taken together, these indicate the *Beudanticeras affine*
406 regional ammonite zone (Jeletzky, 1964), and the *Douvilleiceras mammillatum* international
407 standard zone (Jeletzky, 1968), which is latest Early Albian in age, from 113 to 107 Ma (Ogg
408 and Hinnove, 2012).
409 Worm tube fossils were abundant (>500 specimens) in the Ellef Ringnes seep deposits
410 (Fig. 8L). Some of the tubes were enclosed by micrite, others by botryoidal calcite and yellow
411 calcite cements. The tubes had a variety of infillings, including micrite with peloids (sometimes
412 geopetal), botryoidal yellow calcite (Fig. 8M), and, less commonly, blocky calcite cement. The
413 worm tubes were preserved in various orientations and tended to cluster in groups that were
414 aligned parallel or sub-parallel to one another, oriented from sub-horizontal to vertical within a
415 given deposit (Fig. 8L). The tubes reached up to 12 cm in length and ~0.5 cm in diameter. The
416 tube walls were formed of concentrically laminated layers of calcite that ranged from 40 μm to
417 330 μm thick (Fig. 8M). These are the same fossils that previously had been described as large
418 serpulid tubes and smaller possible serpulid tubes in Beauchamp et al. (1989; Fig. 2) and
419 Beauchamp and Savard (1992; Fig. 5). However, they lack the chevron tube wall microstructure
420 typical of this group of polychaetes, and instead appear identical to tubular fossils from other
421 Cenozoic and Mesozoic seep deposits that have been identified as calcified probable
422 vestimentiferan worm tubes (e.g. Haas et al., 2009; Hilário et al., 2011).

Many of the Ellef Ringnes seep deposits contained fossilized wood, ranging from microscopic particles to pieces over 30 cm in length (Fig. 8N). The wood pores were cemented with calcite that preserved much of the original wood structure, including growth rings and knots. Some pieces of wood had fractures filled with late blocky calcite, but the majority of the wood was filled with early marine calcite with $\delta^{13}C$ values consistent with that of the seep carbonates (Fig. 5). The size and abundance of wood debris correlated with the size of the seep deposit, with small deposits containing microscopic or small fragments of wood and large deposits hosting large, isolated and very well preserved pieces of wood. Wood is also present in the enclosing Christopher Formation mudstone.

4.7 Paleo-redox geochemistry

The Christopher Formation mudstone contained between 0.22 to 2.29 ppm of Mo (Table 3), with the exception of one high value of 9.9 ppm immediately above the zone of seep deposits. Most samples had Mo concentrations (average of 0.98) that are below the PAAS concentration of 1.0 ppm (Turekian and Wedepohl, 1961) (Fig. 6). The Ellef Ringnes Island seep carbonates had values between 0.05 and 10.95 ppm (Table 2; Fig. 6).

5.0 INTERPRETATIONS AND DISCUSSION

5.1 Ellef Ringnes Island carbonates as seep deposits

The morphological variation, petrography, stable isotope values and biomarker contents of the carbonate deposits found on Ellef Ringnes Island are typical of both modern and other fossil seep carbonates, confirming that they were formed by the seepage of methane into the Sverdrup Basin in the Lower Cretaceous. We show that rather than being isolated occurrences, the original two

446 seep mounds described by Beauchamp et al. (1989) are part of a previously unrecognised
447 widespread occurrence of seep deposits on the island.
448 The very negative $\delta^{13}C_{VPDB}$ values of the Group 1 carbonates $\delta^{13}C$= -52.6‰ to -32.3‰) are
449 characteristic of carbonates that obtained carbon from AOM (Whiticar, 1999). The associated
450 $\delta^{18}O_{VPDB}$ values between -3‰ to 1‰, suggest carbonate precipitation near ambient seawater
451 temperature (O'Neil et al., 1969). As such the Group 1 carbonates are interpreted to represent
452 early diagenetic phases precipitated in the subsurface close to the seafloor via methane oxidation.
453 In contrast to Group 1, the Group 2 cements are exclusively blocky calcite and are the latest
454 phase of cements within the deposits. The range of $\delta^{13}C$ values are less negative than Group 1,
455 but still significantly more depleted in ^{13}C-depleted than normal marine carbonates (Fig. 5),
456 suggesting they formed from either an alternative carbon source, or different degree of mixing
457 between methane and seawater alkalinity. The $\delta^{18}O_{VPDB}$ values of group 2 cements are much
458 more ^{18}O-depleted than the early diagenetic material, suggesting precipitation under warmer
459 temperatures (O'Neil et al., 1969). Using the relationship that ~0.2‰ in $\delta^{18}O$ equates to a shift of
460 1 °C from the ambient seawater temperature (Faure, 1987), the late burial cements of the Ellef
461 Ringnes seep deposits would have precipitated in waters between 65 and 95 °C. This range is
462 higher than the maximum burial temperatures of the Christopher Formation, ~55 °C, as derived
463 from a burial history model for Hoodoo Dome (Dewing et al., 2016). This discrepancy may
464 relate to localized high geothermal gradients associated with the salt diapirs. High heat flow in
465 salt can create local thermal anomalies above and adjacent to salt structures. This was shown to
466 be the case in the Hazen F-54 well to the southwest of Ellef Ringnes Island, where geothermal
467 gradients affected by a salt diapir are significantly higher than background (Chen et al., 2010;

Lerche and O'Brien, 1987). There may have also been localized hydrothermal circulation near the diapirs (Grasby et al., 2012) that could have formed the Group 2 cements.

The newly discovered seep deposits are similar in scale to the two previously described seep mounds on Ellef Ringnes Island (Beauchamp et al., 1989; Beauchamp and Savard, 1992; Savard et al., 1996), but the identification of carbonate beds and crusts increases the range of morphologies of seep deposits in the Sverdrup Basin. Compared to other Phanerozoic seep deposits, the Ellef Ringes examples are fairly typical in size (e.g. Campbell, 2006; Campbell et al., 2002; Hammer et al., 2011; Kauffman et al., 1996; Kelly et al., 2000; Majima et al., 2005; Peckmann et al., 1999), although very large deposits of 100s metres diameter and 10s metres thickness, such as the Miocene Rocky Knob deposit from New Zealand (Campbell et al., 2008) or the Devonian Hollard Mound (Peckmann et al., 2005) are not present. In general, the greater the duration of seepage and/or increased methane flux will lead to larger deposits. Whether any genetic information can be gleaned from the morphologies of the Ellef Ringes deposits is equivocal, because of the eroded nature of the deposits at outcrop, particularly those with a mound shape. Nevertheless, the presence of beds and crusts is mirrored by the common occurrence of exhumed methane derived crusts and pavements at modern seeps (e.g. Campbell 2006; Himmler et al., 2015). The overall small sizes may also indicate a relatively short duration of growth.

The potential growth rate of individual seep deposits can be estimated by comparison with modern examples. A study using U-Th analyses found seep carbonate crusts, similar to our crust morphology, have growth rates that vary from 0.4 cm/kyr during early seep development, and 5.0 cm/kyr during late stages of seep development (Bayon, 2009), suggesting that carbonate crusts we observe that up to 50 cm thick could represent 10,000 to 125,000 years growth. Much

491 higher growth rates of 47 cm/kyr have been reported for columnar structures in the Arabian Sea
492 (Himmler et al., 2016). Using these growth rates the seep mound morphologies exposed on Ellef
493 Ringnes Island, from 20 to 310 cm high, would require methane seepage duration of only ~ 425
494 to 5,200 years to form – but upwards of 775,000 years at the lowest growth rate for modern
495 crusts. The shorter growth rates are more consistent with age dates of modern seep deposits,
496 however (Bayon, 2009; Berndt et al., 2014; Crémière et al., 2013). Given that there are 139 seep
497 deposits over an area of ~10,000 km^2, and that each deposit probably was active for thousands to
498 10s thousands of years, at least some may have been active at any time. Alternatively, the entire
499 system may have shut off and on during the period of formation, as suggested by the presence of
500 corrosion surfaces within individual seep deposits. Regardless, the density of seep deposits
501 exposed on Ellef Ringnes Island rivals that of other areas with abundant ancient seep deposits
502 (see section 1).
503 The areal extent of the seep deposits on Ellef Ringnes Island (10,000 km^2) is likely even
504 great than documented here. The methane seep mound reported on Prince Patrick Island, ~ 500
505 km distant, is of the same age (Beauchamp et al., 1989). Other nearby islands with Cretaceous
506 exposure (e.g. Amund Ringnes, Axel Heiberg, northern Ellesmere) have more steeply dipping
507 strata that would make carbonate mounds more difficult to observe without a focused search. In
508 general though the areal of extent of methane seepage is of the same order of magnitude as that
509 observed in regions of extensive modern methane seeps, such as the Gulf of Mexico (e.g.
510 Roberts and Aharon, 1994)
511

512 **5.2 Biomarkers of the anaerobic methane oxidation consortium**

513 The $\delta^{13}C_{org}$ values of organic matter within the Group 1 carbonates (-47.9 to -31.7 ‰) is
514 consistent with the cements, and distinct from $\delta^{13}C_{org}$ of organic matter in the surrounding
515 Christopher Formation mudstone (-25.9 to -17.1‰). This indicates a distinct habitat associated
516 with the seep mounds, with C metabolised from methane oxidation. This is seen further by the
517 significantly ^{13}C depleted isoprenoids ($\delta^{13}C$ of = -118 ‰ and -113 ‰ for PMI and
518 phytane/crocetane respectively), and *iso*- and *anteiso*-C_{15} fatty acids ($\delta^{13}C$ of = -92 ‰ and -87 ‰
519 respectively), that clearly indicate the incorporation of methane-derived carbon into the
520 preserved lipids by anaerobic methane oxidation and their syntrophic partners, sulphate-reducing
521 bacteria. The isoprenoids found in the hydrocarbon fraction indicate high thermal overprint of
522 the sample, as short-chain head-to-tail linked isoprenoids are present as pseudohomologues (c.f.
523 Birgel et al., 2008a; Heindel et al., 2015; Saito et al., 2015), consistent with the overall burial
524 history of the region (Dewing et al., 2016). Apart from the high temperature pseudohomolgue
525 signatures, the tail-to-tail linked isoprenoids crocetane and PMI are preserved unchanged in the
526 sample. Head-to-tail linked phytane is produced from the degradation of archaeol and/or
527 hydroxyarchaeol, former diether-bond membrane lipids of archaea, characteristic for Anaerobic
528 Methanotrophic archaea (ANME) (Niemann and Elvert, 2008). Further isoprenoids are
529 biphytanes, degradation products of former glycerol dialkyl glycerol tetraethers (GDGTs)(Liu et
530 al., 2016). Interestingly, biphytanes did not contain pseudohomologues, as found for the short
531 head-to-tail linked isoprenoids, and as described from other studies (e.g. Birgel et al., 2008a;
532 Heindel et al., 2015; Saito et al., 2015).

533 The substantial amounts of GDGTs with a predominance of the monocyclic homologue
534 accompanied by strong ^{13}C depletion, are indicative of ANME-1 (Niemann and Elvert, 2008). In

535 old samples, high biphytane contents still bearing rings with strong ^{13}C depletions were present
536 in Cretaceous and Eocene seep sites (Natalicchio et al., 2015; Peckmann et al., 2009), and may
537 also be used as ANME-1 indicators. However, ANME-2 also produces GDGTs, but with lower
538 contents (e.g. Blumenberg et al., 2004). The occurrence of abundant crocetane, which makes up
539 around 50% of the mixed phytane/crocetane peak in the sample, is a reliable indicator that
540 ANME-2 were present during carbonate formation. Crocetane and their unsaturated homologues
541 are found in most recent methane seep microbial communities dominated by ANME-2; in
542 contrast they are found only in minor concentrations in ANME-1 systems (Niemann and Elvert,
543 2008). Only a few Mesozoic seep deposits have so far been found to contain such high biphytane
544 contents (Peckmann et al., 2009; Sandy et al., 2012). The distribution found here suggests a
545 mixed community of both ANME-1 and ANME-2 consortia, although proving these were coeval
546 is problematic because biomarker data in ancient seep carbonates represents a time averaged
547 view of microbial activity during carbonate formation, which may have recorded seepage
548 activity and changes in the intensity over longer time periods (cf., Feng et al., 2014).
549
550 **5.3 Macrofauna**
551 Fossils collected from the new seep deposits indicate that the macrofossil diversity was
552 considerably greater than previously reported in Beauchamp et al. (1989) and Beauchamp and
553 Savard (1992) from the original two Ellef Ringnes sites. The macrofaunal assemblage is,
554 however, similar to that of other Mesozoic seeps, particularly those with a Boreal distribution,
555 with nektic and passively floating elements (ammonites and wood respectively), and benthic
556 elements (brachiopods, gastropods, scaphopods, bivalves and probable vestimentiferan tubes)
557 (e.g. Hammer et al., 2011; Hryniewicz et al., 2014; Hryniewicz et al., 2015). The abundance of

558 infaunal bivalves with probable chemosymbionts (e.g. lucinids, thyasirids, solemyids and
559 nucinellids) is also typical of Mesozoic seep communities, and these taxa are also common in
560 Cenozoic seeps, although not confined to this habitat (Hryniewicz et al., 2014). Their presence is
561 a good indication that there were high concentrations of sulfides within the sediment during the
562 early stage of formation of the seep deposits (micrite cement), and therefore a strong redox zone
563 at the time, linked to AOM reactions.

564 Probable vestimentiferan tubes are common in Mesozoic and Cenozoic methane seep
565 deposits and are typically associated with botyroidal cements (Haas et al., 2009 and references
566 therein), that represent slightly later stages of cement formation related to higher fluid flow (e.g.
567 Campbell et al., 2008) than micrites with peloids, infaunal molluscs, etc. This association can be
568 explained by comparison with the ecology of living vestimentiferans at methane seeps, where
569 firstly some hard substrate (i.e. exhumed seep carbonate and/or mollusc shells) is required for
570 initial settlement of juveniles, and secondly during subsequent growth the posterior part of the
571 tube projects into the substrate and is used to 'mine' sulphides (Dattagupta et al., 2008). Posterior
572 tube portions often co-occur to form bundles between nodules and crusts formed of early-stage
573 micrite cement (Haas et al., 2009; 2010). These ecological conditions are increasingly present
574 during more mature stages of seepage, when seep fluid flux is more concentrated into discrete
575 channels, often closer to the sediment-water interface. This is when slightly later stage cements,
576 such as fibrous aragonite crusts form (e.g. Haas et al., 2010). The geopetal sediment and various
577 cement phase infillings in the Ellef Ringnes seep worm tubes show that they often remained
578 open after the death of the animals and then acted as traps for subsequent sediment influx, and/or
579 channels for seep fluid flow, and/or late stage diagenetic cements (e.g. blocky calcite). This is

illustrated by tube worm fill with stable isotope values consistent with our Group 2 late-stage cements (Fig. 5).

5.4 Methane source

The seep deposits on Ellef Ringnes Island display very high isoprenoid hydrocarbon contents, released from intact ether-lipids, but no obvious overprint by oil, as indicated by high amounts of *n*-alkanes or a pronounced unresolved complex mixture (e.g. Feng et al., 2014; Naehr et al., 2009; Sandy et al., 2012), in the seep fluids. This implies a dominant methane source with little input of higher hydrocarbons, and as such an oil seep can be ruled out.

The $\delta^{13}C$ values of carbonate minerals depend on the contributions of carbon obtained from methane, non-methane higher hydrocarbons (oil), organic matter, biogenic methane, and normal marine inorganic carbon (DIC). Carbon derived from petroleum (oil) has $\delta^{13}C_{VPDB}$ values between -25‰ and -35‰, organic matter between -24‰ to -30‰, and DIC ~0‰ (Schoell, 1982). However, as lipid extracts from the Ellef Ringnes samples do not have an oil signature, this is an unlikely carbon source. The range of $\delta^{13}C$ values of the Group 1 cements ($\delta^{13}C$= -52.6‰ to -32.3‰; average -42.6‰) is similar to thermogenic gas recovered from petroleum wells in the Sverdrup Basin ($\delta^{13}C$ average of -42.0‰) (Grasby et al., 2012; Leythaeuser, 1986). However, seep carbonates generally have higher $\delta^{13}C$ values than the source methane due to contribution from DIC (Peckmann and Thiel, 2004). Using the average stable carbon isotope values of the thermogenic gas recovered from the Sverdrup Basin (-42.5‰) and the values from the seep carbonates (-42.6‰), the ratio of carbon in the seep deposits derived from a thermogenic methane source would have to have been nearly 100%, with no other carbon mixing, a process unknown to occur in modern seep environments. In the northern Gulf of

603 Mexico, seep fluids dominated by biogenic gas ($\delta^{13}C$= -88.9‰) produce seep carbonates with
604 $\delta^{13}C_{VPDB}$ values between -45.1‰ to -43.8‰ (Roberts et al., 2010). Similarly, seeps located
605 offshore southern Californian have biogenic methane ($\delta^{13}C$= -81.1 to -73.1 ‰) that forms
606 carbonates with $\delta^{13}C$values of -58 to -46 ‰ (Hein et al., 2006). These modern carbonate $\delta^{13}C$
607 values are similar to the seep carbonates preserved on Ellef Ringnes Island, and based on this we
608 argue that the stable isotope values of the Ellef Ringnes seeps reflect a shallow biogenic, rather
609 than a deeper thermogenic, methane source. This is supported by our biomarker analyses. The
610 carbon isotope fractionation between methane and the membrane lipids formed from that carbon
611 source is usually around -50 ‰. Given the values observed in seep mound biomarkers (PMI,
612 crocetane/phytane; biphytane), the $\delta^{13}C$ value of the methane gas would be estimated at -68 to -
613 59 ‰, again consistent with a biogenic methane source (Formolo et al., 2004).
614 The high density of seep deposits on the immediate flanks of the salt diapirs on Ellef
615 Ringnes Island may suggest there was some linkage between the development of seep deposits
616 and salt diapirism in the Sverdrup Basin. Geological maps of Ellef Ringnes Island (Evenchick
617 and Embry, 2012a; Evenchick and Embry, 2012b) show that both the Isachsen and Christopher
618 formations exposed on the flanks of salt diapirs are highly fractured and faulted, especially when
619 compared to the same formations away from the diapirs. However, while mobile, the salt diapirs
620 which are now exposed at the surface of the island had not penetrated the seafloor during the
621 Early Cretaceous (Dewing et al., 2016). The seep carbonate geopetal fabrics and seep deposit
622 bedding are also both parallel to the Christopher Formation strata, indicating that the seeps most
623 likely formed above the salt domes prior to tilting. We also note that seep deposits were found at
624 the same stratigraphic level away from any diapirs. Given this we argue that the apparent density
625 of methane seeps near diapirs is likely an exposure bias, where Christopher Formation strata are

626 preferentially exposed in uplifted beds surrounding the diapirs, making seep deposits more
627 readily observable in those locations.
628
629 **5.5 Environment of formation**
630 The profile of both $\delta^{13}C$ of organic carbon and Mo in the Christopher Formation shales does not
631 show any trend leading up to the zone of carbonate deposits, and only one data point shows high
632 Mo values immediately above the zone (Fig. 6). These low values suggest that methane seepage
633 occurred in normal, well-oxygenated bottom water conditions (Tribovillard et al., 2006) in the
634 Sverdrup Basin. This is consistent with the abundance and diversity of benthic fauna in the Ellef
635 Ringnes seep deposits. Modern methane seeps forming in anoxic basins and oxygen minimum
636 zones are devoid of large animals, and instead contain only microbial mats only (e.g. Himmler et
637 al., 2015).
638 The sequestration of Mo into pure carbonate phases involves the incorporation of low
639 reactive molybdate ions (MoO_4^{2-}) into the crystal lattice of carbonates in a similar fashion to
640 carbonate associated sulphate (CAS). The low chemical reactivity of molybdate ions in seawater
641 results in a long ocean residency time but also in the limited incorporation of molybdate into
642 carbonate minerals (Lyons et al., 2009). However, under anoxic conditions Mo can be reduced
643 and then sequestered by sulfides, such that Mo enrichments can be an indicator of locally anoxic
644 environments (Tribovillard et al., 2006). The Mo values of the Ellef Ringnes seep deposits vary
645 widely, from 0.05 to 10.95 ppm, in comparison with typical Mo concentration in shallow marine
646 carbonates of 0.4 ppm (Turekian and Wedepohl, 1961). In general, Mo values of the seep
647 carbonates are significantly higher than average shallow carbonate deposits, indicating local
648 anaerobic conditions associated with methane oxidation and carbonate precipitation. Consistent

649 with this, lipids found within the carbonate deposits are almost only associated with the
650 anaerobic oxidation of methane. No indications of prevailing oxic or microaerophilic conditions
651 were found, since no ^{13}C-depleted hopanoids or lanostanes have been identified (Birgel and
652 Peckmann, 2008; Natalicchio et al., 2015; Sandy et al., 2012). These results suggest the presence
653 of a strong redox zone within the sediment during carbonate precipitation, as also indicated by
654 the presence of infaunal bivalve species with potential chemosymbionts and the probable
655 vestimentiferan worm tubes (Table 4).
656
657 **5.6 Climate conditions during the methane seep event**
658 During Cretaceous time, the high paleolatitude setting of the Sverdrup Basin (72° N) was
659 characterised by numerous "cold-snaps" that punctuated otherwise warm Early Cretaceous
660 conditions (Grasby et al., in press; Herrle et al., 2015). Evidence for cold climate intervals during
661 the late Aptian-early Albian, including wood and pollen data, has been demonstrated for the
662 circum-Arctic region, as well as for lower latitudes (e.g. Galloway et al., 2015; Harland et al.,
663 2007; Kemper, 1987; Maurer et al., 2013; McAnena et al., 2013; Mutterlose et al., 2009).
664 Possible ice-rafted debris are also observed in Aptian strata of Spitsbergen (Dalland, 1977). A
665 cool climate in the Sverdrup Basin during Albian time is further supported by the widespread
666 occurrence of glendonites (an ikaite pseudomorph) in the Invincible Point Member (Grasby et
667 al., in press; Kemper, 1987; Kemper and Schmitz, 1975). Ikaite formation is considered to
668 require near freezing temperatures (~1.9 to 7.0 °C) coupled with a ten-fold increase in alkalinity
669 relative to normal seawater to form (Bischoff et al., 1993; Marland, 1975).
670 Previous studies have shown the co-occurrence of glendonites within active modern
671 methane seeps, related to organic matter decomposition driven by the initial stages of methane

672 seepage (Greinert and Derkachev, 2004; Teichert and Luppold, 2013). However, in our study
673 area, the glendonite bearing horizon in the Invincible Point Member occurs as discrete layers
674 over a 100 m stratigraphic range, ~200 m below the stratigraphic level of the seep carbonates.
675 This horizon is equivalent to the glendonite occurrence observed in the Invincible Point Member
676 of the Christopher Formation exposed at Glacier Fiord, Axel Heiberg Island (Herrle et al., 2015).
677 Based on their composite $\delta^{13}C$ stratigraphy they date the youngest glendonite bed at ca. 112.8
678 Ma (lowermost Albian), and they indicate that the loss of glendonites is coincident with a return
679 to warm temperatures at the end of a ~6 m.y. late Aptian to early Albian cooling event. A
680 transition from cold to warm climate at this time is also recorded in the subtropical Atlantic
681 Ocean, possibly related to volcanism (~114-110 Ma) of the Kerguelen LIP (McAnena et al.,
682 2013). Given these constraints, the possible age range for the methane release event that formed
683 the widespread carbonate seep deposits on Ellef Ringnes Island (between 113 to 107 Ma; section
684 4.6), corresponds closely to this climate warming event.
685
686 **5.7 Controls on methane release**
687 Where possible to observe, carbonate seep deposits occur at one single stratigraphic horizon,
688 below the regional marker unit dated at 105.4 Ma (section 4.1) and above the last occurrence of
689 glendonites dated at 112.8 Ma. The overall stratigraphic range of seep deposits across the island,
690 relative to the regional marker unit, is quite narrow (34 m) and could easily reflect variations in
691 seafloor topography and sedimentation rates. As such, the widespread occurrence of seep
692 deposits on Ellef Ringnes reflects a transient event. This raises the question as to what may have
693 influenced the short term, but aerially extensive, release of biogenic methane into the seafloor of
694 the Sverdrup Basin during Albian time. One possible mechanism is diaper mobilisation causing

695 gas migration along fracture systems. However, diapirs record a movement history for over 100
696 Ma (Dewing et al., 2016), suggesting that if this was the case than evidence of methane seepage
697 should be more common in the basin. The observation of carbonate mounds not associated with
698 diapirs further indicates that they do not directly control formation of methane seep deposits.
699 As described in section 5.6, the occurrence of widespread seep mounds in the Sverdrup
700 Basin is coincident with a period of Albian climate warming. This is similar to modern high
701 latitude methane seeps, where studies have shown that biogenic gas leakage occurred in response
702 to post ice-age climate warming, with continuous methane flux for 3 to 10 kyr in response to
703 post-glacial gas hydrate release (Berndt et al., 2014; Chand et al., 2016; Crémière et al., 2016).
704 We hypothesize that a similar event may have occurred in the Sverdrup Basin, whereby the early
705 Albian cold snap indicated by widespread glendonite occurrence would likely have been
706 associated with development of gas hydrates, trapping biogenic gas formed during
707 biodegradation of marine organics in the overall oxic depositional environment. Subsequent
708 climate warming may have caused hydrate dissociation and the regional, but transient,
709 occurrence of methane seep deposits. Modeled rates for hydrate melting shows gas release would
710 occur on ~100 ky time scale (Majorowicz et al., 2014), consistent with the potential age range of
711 the deposits.
712

713 # 6.0 CONCLUSIONS

714 The discovery of 137 new Early Cretaceous carbonate deposits on Ellef Ringnes Island,
715 Canadian Arctic Archipelago demonstrate the presence of an extensive field of methane seeps
716 that occur over an area greater than 10,000 km^2, throughout the central to southern part of the
717 island. This discovery greatly increases the overall area of methane seep sites recognised on the

718 island, the diversity of seep deposit types, and the macrofossil diversity present. Seep deposits
719 range in size from less than 1 m^3 to over 2.4 m high by 7.6 m long and are exposed within a
720 narrow stratigraphic range within the uppermost portion of the Invincible Point Member of the
721 Christopher Formation. While predominately exposed on the flanks of four exposed salt diapirs,
722 they are not restricted to this setting. The association of seep carbonates with diapirs most likely
723 reflects an exposure bias related to uplifted beds along the diapir flanks.
724 Each seep deposit is composed of various portions of six major carbonate phases as well
725 as moderately diverse fossil assemblages, including chemosymbiotic species. The carbonates
726 have very low $\delta^{13}C_{VPDB}$ values, as well as $\delta^{13}C$ values of biomarkers, indicative of carbonate
727 formed in the subsurface from the anaerobic oxidation of primarily biogenic methane.
728 The large geographical occurrence of seep deposits suggests widespread venting of
729 biogenic methane rich fluids. However, methane seepage was a transient event that occurred
730 during the latest Early Albian. We hypothesize that this may relate to possible gas hydrate
731 dissociation related to climate warming during that time.
732

733 ## ACKNOWLEDGEMENTS

734 Gennyne McCune contributed valuable information regarding the environments of the formation
735 and assistance in the field. Field work was supported by the Polar Continental Shelf Project.
736 Discussions with A. Embry on Cretaceous stratigraphy of the Sverdrup Basin are greatly
737 appreciated. ESS Contribution xxxx.
738

739 **REFERENCES**

740

741 Aloisi, G., Pierre, C., Rouchy, J.-M., Foucher, J.-P., and Woodside, J., 2000, Methane-related
742 authigenic carbonates of eastern Mediterranean Sea mud volcanoes and their possible
743 relation to gas hydrate destabilisation: Earth and Planetary Science Letters, v. 184, no. 1,
744 p. 321-338.
745 Balkwill, H. R., 1978, Evolution of the sverdrup basin: The american Association of Petroleum
746 Geologists Bulletin, v. 62, no. No.6, p. 1004-1028.
747 Bayon, G. H., G.M. and Bohn, M, 2009, U–Th stratigraphy of a cold seep carbonate crust:
748 Chemical Geology, v. 260, p. 47-56.
749 Beauchamp, B., Harrison, J. C., Nassichuk, W. W., Krouse, H. R., and Eliuk, L. S., 1989,
750 Cretaceous Cold-Seep Communities and Methane-Derived Carbonates in the Canadian
751 Arctic: Science, v. 244, no. 4900, p. 53-56.
752 Beauchamp, B., and Savard, M., 1992, Cretaceous chemosynthetic carbonate mounds in the
753 Canadian Arctic: PALAIOS, v. 7, no. 4, p. 434-450.
754 Berndt, C., Feseker, T., Treude, T., Krastel, S., Liebetrau, V., Niemann, H., Bertics, V. J.,
755 Dumke, I., Dünnbier, K., Ferré, B., Graves, C., Gross, F., Hissmann, K., Hühnerbach, V.,
756 Krause, S., Lieser, K., Schauer, J., and Steinle, L., 2014, Temporal Constraints on
757 Hydrate-Controlled Methane Seepage off Svalbard: Science, v. 343, no. 6168, p. 284-
758 287.
759 Birgel, D., Elvert, M., Han, X., and Peckmann, J., 2008b, 13C-depleted biphytanic diacids as
760 tracers of past anaerobic oxidation of methane: Organic Geochemistry, v. 39, no. 1, p.
761 152-156.
762 Birgel, D., Himmler, T., Freiwald, A., and Peckmann, J., 2008a, A new constraint on the
763 antiquity of anaerobic oxidation of methane: Late Pennsylvanian seep limestones from
764 southern Namibia: Geology, v. 36, no. 7, p. 543-546.
765 Birgel, D., and Peckmann, J., 2008, Aerobic methanotrophy at ancient marine methane seeps: A
766 synthesis: Organic Geochemistry, v. 39, no. 12, p. 1659-1667.
767 Birgel, D., Peckmann, J., Klautzsch, S., Thiel, V., and Reitner, J., 2006b, Anaerobic and Aerobic
768 Oxidation of Methane at Late Cretaceous Seeps in the Western Interior Seaway, USA:
769 Geomicrobiology Journal, v. 23, no. 7, p. 565-577.
770 Birgel, D., Thiel, V., Hinrichs, K.-U., Elvert, M., Campbell, K. A., Reitner, J., Farmer, J. D., and
771 Peckmann, J., 2006a, Lipid biomarker patterns of methane-seep microbialites from the
772 Mesozoic convergent margin of California: Organic Geochemistry, v. 37, no. 10, p. 1289-
773 1302.
774 Bischoff, J. L., Fitzpatrick, J. A., and Rosenbauer, R. J., 1993, The solubility and stabilization of
775 ikaite (CaCO3 · 6H20) from 0° to 25°C; environmental and paleoclimatic implications
776 for thinolite tufa: Journal of Geology, v. 101, p. 21-33.
777 Blumenberg, M., Seifert, R., Reitner, J., Pape, T., and Michaelis, W., 2004, Membrane lipid
778 patterns typify distinct anaerobic methanotrophic consortia: Proceedings of the National
779 Academy of Sciences of the United States of America, v. 101, no. 30, p. 11111-11116.
780 Boetius, A., Ravenschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R.,
781 Jorgensen, B. B., Witte, U., and Pfannkuche, O., 2000, A marine microbial consortium
782 apparently mediating anaerobic oxidation of methane: Nature, v. 407, no. 6804, p. 623-
783 626.

784 Boutelier, J., Cruden, A., Brent, T., and Stephenson, R., 2011, Timing and mechanisms
785 controlling evaporite diapirism on Ellef Ringnes Island, Canadian Arctic Archipelago:
786 Basin Research, v. 23, p. 478–498.
787 Campbell, K. A., 2006, Hydrocarbon seep and hydrothermal vent paleoenvironments and
788 paleontology: Past developments and future research directions: Palaeogeography,
789 Palaeoclimatology, Palaeoecology, v. 232, no. 2–4, p. 362-407.
790 Campbell, K. A., Farmer, J. D., and Des Marais, D., 2002, Ancient hydrocarbon seeps from the
791 Mesozoic convergent margin of California: carbonate geochemistry, fluids and
792 palaeoenvironments: Geofluids, v. 2, no. 2, p. 63-94.
793 Campbell, K. A., Francis, D. A., Collins, M., Gregory, M. R., Nelson, C. S., Greinert, J., and
794 Aharon, P., 2008, Hydrocarbon seep-carbonates of a Miocene forearc (East Coast Basin),
795 North Island, New Zealand: Sedimentary Geology, v. 204, no. 3–4, p. 83-105.
796 Chand, S., Crémière, A., Lepland, A., Thorsnes, T., Brunstad, H., and Stoddart, D., 2016, Long-
797 term fluid expulsion revealed by carbonate crusts and pockmarks connected to subsurface
798 gas anomalies and palaeo-channels in the central North Sea: Geo-Marine Letters, p. 1-13.
799 Chen, Z., Grasby, S. E., and Dewing, K., 2010, Temperature-depth plots for selected petroleum
800 exploration wells, Canadian Arctic Islands: Geological Survey of Canada, Open File, v.
801 6567, p. 1-16.
802 Chevalier, N., Bouloubassi, I., Birgel, D., Taphanel, M. H., and López-García, P., 2013,
803 Microbial methane turnover at Marmara Sea cold seeps: a combined 16S rRNA and lipid
804 biomarker investigation: Geobiology, v. 11, no. 1, p. 55-71.
805 Cochran, J. K., Landman, N. H., Larson, N. L., Meehan, K. C., Garb, M., and Brezina, J., 2015,
806 Geochemical evidence (C and Sr isotopes) for methane seeps as ammonite habitats in the
807 Late Cretaceous (Campanian) Western Interior Seaway: Swiss Journal of Palaeontology,
808 v. 134, no. 2, p. 153-165.
809 Crémière, A., Bayon, G., Ponzevera, E., and Pierre, C., 2013, Paleo-environmental controls on
810 cold seep carbonate authigenesis in the Sea of Marmara: Earth and Planetary Science
811 Letters, v. 376, p. 200-211.
812 Crémière, A., Lepland, A., Chand, S., Sahy, D., Condon, D. J., Noble, S. R., Martma, T.,
813 Thorsnes, T., Sauer, S., and Brunstad, H., 2016, Timescales of methane seepage on the
814 Norwegian margin following collapse of the Scandinavian Ice Sheet: Nature
815 Communications, v. 7, p. 11509.
816 Dalland, A., 1977, Erratic clasts in lower Tertiary deposits of Svalbard – Evidence of transport
817 by winter ice: Norsk Polarinstitutt Årbok, p. 151-166.
818 Dattagupta, S., Arthur, M. A., and Fisher, C. R., 2008, Modification of sediment geochemistry
819 by the hydrocarbon seep tubeworm Lamellibrachia luymesi: A combined empirical and
820 modeling approach: Geochimica et Cosmochimica Acta, v. 72, no. 9, p. 2298-2315.
821 Dewing, K., Springer, A., Guest, B., and Hadlari, T., 2016, Geological evolution and
822 hydrocarbon potential of the salt-cored Hoodoo Dome, Sverdrup Basin, Arctic Canada:
823 Marine and Petroleum Geology, v. 71, p. 134-148.
824 Dubilier, N., Bergin, C., and Lott, C., 2008, Symbiotic diversity in marine animals: the art of
825 harnessing chemosynthesis: Nat Rev Micro, v. 6, no. 10, p. 725-740.
826 Elvert, M., Boetius, A., Knittel, K., and Jørgensen, B. B., 2003, Characterization of Specific
827 Membrane Fatty Acids as Chemotaxonomic Markers for Sulfate-Reducing Bacteria

828 Involved in Anaerobic Oxidation of Methane: Geomicrobiology Journal, v. 20, no. 4, p.
829 403-419.
830 Embry, A. F., 1985, Stratigraphic subdivision of the Isachsen and Christopher formations (Lower
831 Cretaceous), Arctic Islands: Current Research, Part B, Geologic Survey of Canada, v.
832 Paper 85-1B, no. 239-246.
833 Embry, A. F., and Beauchamp, B., 2008, Sedimentary Basins of the World: The Sedimentary
834 Basins of the United States and Canada, Elsevier.
835 Emerson, S. R., and Huested, S. S., 1991, Ocean anoxia and the concentrations of molybdenum
836 and vanadium in seawater: Marine Chemistry, v. 34, no. 3–4, p. 177-196.
837 Evenchick, C. A., Davis, W. J., Bédard, J. H., Hayward, N., and Friedman, R. M., 2015,
838 Evidence for protracted High Arctic large igneous province magmatism in the central
839 Sverdrup Basin from stratigraphy, geochronology, and paleodepths of saucer-shaped
840 sills: Geological Society of America Bulletin.
841 Evenchick, C. A., and Embry, A. F., 2012a, Geology, Ellef Ringnes Island north, Nunavut:
842 Natural Resources Canada Canadian Geoscience Map 86 (Preliminary), scale 1: 125,000.
843 Evenchick, C. A., and Embry, A. F., 2012b, Geology, Ellef Ringnes Island south, Nunavut:
844 Natural Resources Canada Canadian Geoscience Map 87 (Preliminary), scale 1: 125,000.
845 Faure, G., 1987, New developments and applications in isotope geoscience: Chemical Geology;
846 Isotope Geoscience Section, v. 66, no. 1-2, p. 1-177.
847 Feng, D., Birgel, D., Peckmann, J., Roberts, H. H., Joye, S. B., Sassen, R., Liu, X.-L., Hinrichs,
848 K.-U., and Chen, D., 2014, Time integrated variation of sources of fluids and seepage
849 dynamics archived in authigenic carbonates from Gulf of Mexico Gas Hydrate Seafloor
850 Observatory: Chemical Geology, v. 385, p. 129-139.
851 Formolo, M. J., Lyons, T. W., Zhang, C., Kelley, C., Sassen, R., Horita, J., and Cole, D. R.,
852 2004, Quantifying carbon sources in the formation of authigenic carbonates at gas
853 hydrate sites in the Gulf of Mexico: Chemical Geology, v. 205, no. 3–4, p. 253-264.
854 Galloway, J. M., Tullius, D. N., Evenchick, C. A., Swindles, G. T., Hadlari, T., and Embry, A.,
855 2015, Early Cretaceous vegetation and climate change at high latitude: Palynological
856 evidence from Isachsen Formation, Arctic Canada: Cretaceous Research, v. 56, p. 399-
857 420.
858 Gould, D. B., and DeMille, G., 1964, Piercement Structures in Canadian Arctic Islands: Bulletin
859 of Canadian Petroleum Geology, v. 12, no. 3, p. 719.
860 Grasby, S. E., Chen, Z., and Dewing, K., 2012, Formation water geochemistry of the Sverdrup
861 Basin: Implications for hydrocarbon development in the High Arctic: Applied
862 Geochemistry, v. 27, p. 1623-1632.
863 Grasby, S. E., McCune, G. E., Beauchamp, B., and Galloway, J. M., in press, Lower Cretaceous
864 cold snaps led to widespread glendonite occurrences in the Sverdrup Basin, Canadian
865 High Arctic: Geological Society of America Bulletin.
866 Greinert, J., and Derkachev, A., 2004, Glendonites and methane-derived Mg-calcites in the Sea
867 of Okhotsk, Eastern Siberia: implications of a venting-related ikaite/glendonite
868 formation: Marine Geology, v. 204, no. 1–2, p. 129-144.
869 Haas, A., Little, C. T. S., Sahling, H., Bohrmann, G., Himmler, T., and Peckmann, J., 2009,
870 Mineralization of vestimentiferan tubes at methane seeps on the Congo deep-sea fan:
871 Deep Sea Research Part I: Oceanographic Research Papers, v. 56, no. 2, p. 283-293.

872 Haas, A., Peckmann, J., Elvert, M., Sahling, H., and Bohrmann, G., 2010, Patterns of carbonate
873 authigenesis at the Kouilou pockmarks on the Congo deep-sea fan: Marine Geology, v.
874 268, no. 1–4, p. 129-136.
875 Hammer, Ø., Nakrem, H. A., Little, C. T. S., Hryniewicz, K., Sandy, M. R., Hurum, J. H.,
876 Druckenmiller, P., Knutsen, E. M., and Høyberget, M., 2011, Hydrocarbon seeps from
877 close to the Jurassic–Cretaceous boundary, Svalbard: Palaeogeography,
878 Palaeoclimatology, Palaeoecology, v. 306, no. 1–2, p. 15-26.
879 Harland, M., Francis, J. E., Brentnall, S. J., and Beerling, D. J., 2007, Cretaceous (Albian–
880 Aptian) conifer wood from Northern Hemisphere high latitudes: Forest composition and
881 palaeoclimate: Review of Palaeobotany and Palynology, v. 143, no. 3–4, p. 167-196.
882 Hein, J. R., Normark, W. R., McIntyre, B. R., Lorenson, T. D., and Powell, C. L., 2006,
883 Methanogenic calcite, 13C-depleted bivalve shells, and gas hydrate from a mud volcano
884 offshore southern California: Geology, v. 34, no. 2, p. 109-112.
885 Heindel, K., Richoz, S., Birgel, D., Brandner, R., Klügel, A., Krystyn, L., Baud, A., Horacek,
886 M., Mohtat, T., and Peckmann, J., 2015, Biogeochemical formation of calyx-shaped
887 carbonate crystal fans in the subsurface of the Early Triassic seafloor: Gondwana
888 Research, v. 27, no. 2, p. 840-861.
889 Herrle, J. O., Schröder-Adams, C. J., Davis, W., Pugh, A. T., Galloway, J. M., and Fath, J., 2015,
890 Mid-Cretaceous High Arctic stratigraphy, climate, and Oceanic Anoxic Events: Geology.
891 Hilário, A., Capa, M., Dahlgren, T. G., Halanych, K. M., Little, C. T. S., Thornhill, D. J., Verna,
892 C., and Glover, A. G., 2011, New Perspectives on the Ecology and Evolution of
893 Siboglinid Tubeworms: PLoS ONE, v. 6, no. 2, p. e16309.
894 Himmler, T., Bayon, G., Wangner, D., Enzmann, F., Peckmann, J., and Bohrmann, G., 2016,
895 Seep-carbonate lamination controlled by cyclic particle flux: Scientific Reports, v. 6, p.
896 37439.
897 Himmler, T., Birgel, D., Bayon, G., Pape, T., Ge, L., Bohrmann, G., and Peckmann, J., 2015,
898 Formation of seep carbonates along the Makran convergent margin, northern Arabian Sea
899 and a molecular and isotopic approach to constrain the carbon isotopic composition of
900 parent methane: Chemical Geology, v. 415, p. 102-117.
901 Hinrichs, K.-U., Hayes, J. M., Sylva, S. P., Brewer, P. G., and DeLong, E. F., 1999, Methane-
902 consuming archaebacteria in marine sediments: Nature, v. 398, no. 6730, p. 802-805.
903 Hoehler, T. M., Alperin, M. J., Albert, D. B., and Martens, C. S., 1994, Field and laboratory
904 studies of methane oxidation in an anoxic marine sediment: Evidence for a methanogen-
905 sulfate reducer consortium: Global Biogeochemical Cycles, v. 8, no. 4, p. 451-463.
906 Hryniewicz, K., Little, C. T. S., and Nakrem, H. A., 2014, Bivalves from the latest Jurassic-
907 earliest Cretaceous hydrocarbon seep carbonates from central Spitsbergen, Svalbard:
908 Zootaxa, v. 3859, p. 1-66.
909 Hryniewicz, K., Nakrem, H. A., Hammer, Ø., Little, C. T. S., Kaim, A., Sandy, M. R., and
910 Hurum, J. H., 2015, The palaeoecology of the latest Jurassic–earliest Cretaceous
911 hydrocarbon seep carbonates from Spitsbergen, Svalbard: Lethaia, v. 48, no. 3, p. 353-
912 374.
913 Jeletzky, J. A., 1964, Lower Cretaceous marine index fossils of the sedimentary basins of
914 western and Arctic Canada. Illustrations of Canadian Fossils: Geological Survey of
915 Canada, Paper 64-11.

-, 1968, Macrofossil zones of the marine Cretaceous of the western Interior of Canada and their correlation with the zones and stages of Europe and the western Interior of the United States: Geological Survey of Canada, Paper 67-72.
Judd, A., and Hovland, M., 2007, Seabed Fluid Flow: The Impact on Geology, Biology and the Marine Environment, Cambridge University Press, 492 p.:
Kauffman, E. G., Arthur, M. A., Howe, B., and Scholle, P. A., 1996, Widespread venting of methane-rich fluids in Late Cretaceous (Campanian) submarine springs (Tepee Buttes), Western Interior seaway, U.S.A: Geology, v. 24, no. 9, p. 799-802.
Kelly, S. R. A., Blanc, E., Price, S. P., and Whitham, A. G., 2000, Early Cretaceous giant bivalves from seep-related limestone mounds, Wollaston Forland, Northeast Greenland: Geological Society, London, Special Publications, v. 177, no. 1, p. 227-246.
Kemper, E., 1987, Das Klima der Kreide-Zeit: Geol. Jb. Reiche. A., v. 96, p. 5-185.
Kemper, E., and Schmitz, H. H., 1975, Stellate nodules from the Upper Deer Bay Formation (Valanginian) of Arctic Canada: Geological Survey of Canada Paper 75-1C, p. 109-119.
Kulm, L. D., Suess, E., Moore, J. C., Carson, B., Lewis, B. T., Ritger, S. D., Kadko, D. C., Thornburg, T. M., Embley, R. W., Rugh, W. D., Massoth, G. J., Langseth, M. G., Cochrane, G. R., and Scamman, R. L., 1986, Oregon Subduction Zone: Venting, Fauna, and Carbonates: Science, v. 231, no. 4738, p. 561-566.
Lerche, I., and O'Brien, J. J., 1987, Modeling of Bouyant Salt Diapirism, in Dynamical Geology of salt and related Structures: Academic Press, Inc., p. 129-162.
Levin, L., A., 2005, Ecology of Cold Seep Sediments, Oceanography and Marine Biology, CRC Press, p. 1-46.
Leythaeuser, D., Stewart, K., 1986, Generation, migration and accumulation of hydrocarbons in the Sverdrup Basin, Canada. A geochemical basin study: Unpublished report to Panarctic Oils, GSC Catalogue No. QE196.9 L4 v. 2.
Little, C. T. S., Birgel, D., Boyce, A. J., Crame, J. A., Francis, J. E., Kiel, S., Peckmann, J., Pirrie, D., Rollinson, G. K., and Witts, J. D., 2015, Late Cretaceous (Maastrichtian) shallow water hydrocarbon seeps from Snow Hill and Seymour Islands, James Ross Basin, Antarctica: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 418, p. 213-228.
Liu, X.-L., Birgel, D., Elling, F. J., Sutton, P. A., Lipp, J. S., Zhu, R., Zhang, C., Könneke, M., Peckmann, J., Rowland, S. J., Summons, R. E., and Hinrichs, K.-U., 2016, From ether to acid: A plausible degradation pathway of glycerol dialkyl glycerol tetraethers: Geochimica et Cosmochimica Acta, v. 183, p. 138-152.
Lyons, T. W., Anbar, A. D., Severmann, S., Scott, C., and Gill, B. C., 2009, Tracking Euxinia in the Ancient Ocean: A Multiproxy Perspective and Proterozoic Case Study: The Annual Review of Earth and Planetary Science, v. 37, p. 507-534.
Macauley, J., Cruden, A., Brent, T., and Stephenson, R., 2000, Analogue modelling of evaporite diapirs on ellef ringnes island, canadian arctic archipelago: First results American Geophysical Union.
Majima, R., Nobuhara, T., and Kitazaki, T., 2005, Review of fossil chemosynthetic assemblages in Japan: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 227, no. 1–3, p. 86-123.

959 Majorowicz, J., Grasby, S. E., Safanda, J., and Beauchamp, B., 2014, Gas hydrate contribution to
960 Late Permian global warming: Earth and Planetary Science Letters, v. 393, no. 0, p. 243-
961 253.
962 Marland, G., 1975, The stability of CaCO3·6H2O (ikaite): Geochimica et Cosmochimica Acta,
963 v. 39, no. 1, p. 83-91.
964 Maurer, F., van Buchem, F. S. P., Eberli, G. P., Pierson, B. J., Raven, M. J., Larsen, P.-H., Al-
965 Husseini, M. I., and Vincent, B., 2013, Late Aptian long-lived glacio-eustatic lowstand
966 recorded on the Arabian Plate: Terra Nova, v. 25, no. 2, p. 87-94.
967 McAnena, A., Flogel, S., Hofmann, P., Herrle, J. O., Griesand, A., Pross, J., Talbot, H. M.,
968 Rethemeyer, J., Wallmann, K., and Wagner, T., 2013, Atlantic cooling associated with a
969 marine biotic crisis during the mid-Cretaceous period: Nature Geosci, v. 6, no. 7, p. 558-
970 561.
971 Milucka, J., Ferdelman, T. G., Polerecky, L., Franzke, D., Wegener, G., Schmid, M.,
972 Lieberwirth, I., Wagner, M., Widdel, F., and Kuypers, M. M. M., 2012, Zero-valent
973 sulphur is a key intermediate in marine methane oxidation: Nature, v. 491, no. 7425, p.
974 541-546.
975 Mutterlose, J., Bornemann, A., and Herrle, J., 2009, The Aptian-Albian cold snap: Evidence for
976 “mid” Cretaceous icehouse interludes: Neues Jahrbuch für Geologie und Paläontologie –
977 Abhandlungen, v. 252, p. 217-225.
978 Naehr, T. H., Birgel, D., Bohrmann, G., MacDonald, I. R., and Kasten, S., 2009, Biogeochemical
979 controls on authigenic carbonate formation at the Chapopote “asphalt volcano”, Bay of
980 Campeche: Chemical Geology, v. 266, no. 3–4, p. 390-402.
981 Naehr, T. H., Eichhubl, P., Orphan, V. J., Hovland, M., Paull, C. K., Ussler Iii, W., Lorenson, T.
982 D., and Greene, H. G., 2007, Authigenic carbonate formation at hydrocarbon seeps in
983 continental margin sediments: A comparative study: Deep Sea Research Part II: Topical
984 Studies in Oceanography, v. 54, no. 11–13, p. 1268-1291.
985 Nassichuk, and Roy, 1975, Mound-like carbonate rocks of early Cretaceous (Albian) age
986 adjacent to Hoodoo Dome, Ellef Ringnes Island, District of Franklin: Paper - Geological
987 Survey of Canada, v. 75-1, p. 565-569.
988 Nassichuk, W. W., and Davies, G. R., 1980, Stratigraphy and sedimentation of the Otto Fiord
989 Formation, Ottawa, Geological Survey of Canada, Bulletin 286, 87 p.:
990 Natalicchio, M., Peckmann, J., Birgel, D., and Kiel, S., 2015, Seep deposits from northern Istria,
991 Croatia: a first glimpse into the Eocene seep fauna of the Tethys region: Geological
992 Magazine, v. 152, no. 03, p. 444-459.
993 Niemann, H., and Elvert, M., 2008, Diagnostic lipid biomarker and stable carbon isotope
994 signatures of microbial communities mediating the anaerobic oxidation of methane with
995 sulphate: Organic Geochemistry, v. 39, p. 1668-1677.
996 O'Neil, J., Clayton, R., and TK, M., 1969, Oxygen isotope fractionartion in divalentmetal
997 carbonates: Journal of Chemical Physics, v. 51, p. 5547-5558.
998 Ogg, G. M., and Hinnove, L. A., 2012, Cretaceous, *in* Gradstein, F. M., Ogg, J. M., Schmitz, M.
999 D., and Ogg, G. M., eds., The Geologic Time Scale, Volume 2: Boston, Massachusetts,
1000 Elsevier, p. 1144.
1001 Orphan, V. J., House, C. H., Hinrichs, K.-U., McKeegan, K. D., and DeLong, E. F., 2001,
1002 Methane-Consuming Archaea Revealed by Directly Coupled Isotopic and Phylogenetic
1003 Analysis: Science, v. 293, no. 5529, p. 484-487.

1004 Paull, C. K., Hecker, B., Commeau, R., Freeman-Lynde, R. P., Neumann, C., Corso, W. P.,
1005 Golubic, S., Hook, J. E., Sikes, E., and Curray, J., 1984, Biological Communities at the
1006 Florida Escarpment Resemble Hydrothermal Vent Taxa: Science, v. 226, no. 4677, p.
1007 965-967.
1008 Peckmann, J., Birgel, D., and Kiel, S., 2009, Molecular fossils reveal fluid composition and flow
1009 intensity at a Cretaceous seep: Geology, v. 37, no. 9, p. 847-850.
1010 Peckmann, J., Little, C. T. S., Gill, F., and Reitner, J., 2005, Worm tube fossils from the Hollard
1011 Mound hydrocarbon-seep deposit, Middle Devonian, Morocco: Palaeozoic seep-related
1012 vestimentiferans?: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 227, no. 1–3,
1013 p. 242-257.
1014 Peckmann, J., and Thiel, V., 2004, Carbon cycling at ancient methane–seeps: Chemical Geology,
1015 v. 205, no. 3–4, p. 443-467.
1016 Peckmann, J., Thiel, V., Michaelis, W., Clari, P., Gaillard, C., Martire, L., and Reitner, J., 1999,
1017 Cold seep deposits of Beauvoisin (Oxfordian; southeastern France) and Marmorito
1018 (Miocene; northern Italy): microbially induced authigenic carbonates: International
1019 Journal of Earth Sciences, v. 88, no. 1, p. 60-75.
1020 Peters, K. E., and Moldowan, J. M., 1993, The biomarker guide: Interpreting molecular fossils in
1021 petroleum and ancient sediments, Englewood Cliffs, NJ (United States); Prentice Hall;
1022 None, Medium: X; Size: Pages: (363 p) p.:
1023 Ritger, S. D., Carson, B., and Suess, E., 1987, Methane-derived authigenic carbonates formed by
1024 subduction-induced pore-water expulsion along the Oregon/Washington margin:
1025 Geological Society of America Bulletin, v. 98, no. 2, p. 147-156.
1026 Roberts, H. H., and Aharon, P., 1994, Hydrocarbon-derived carbonate buildups of the northern
1027 Gulf of Mexico continental slope: A review of submersible investigations: Geo-Marine
1028 Letters, v. 14, no. 2, p. 135-148.
1029 Roberts, H. H., Feng, D., and Joye, S. B., 2010, Cold-seep carbonates of the middle and lower
1030 continental slope, northern Gulf of Mexico: Deep Sea Research Part II: Topical Studies in
1031 Oceanography, v. 57, no. 21–23, p. 2040-2054.
1032 Saito, R., Kaiho, K., Oba, M., Fujibayashi, M., Tong, J., and Tian, L., 2015, Predominance of
1033 archaea-derived hydrocarbons in an Early Triassic microbialite: Organic Geochemistry,
1034 v. 85, p. 66-75.
1035 Sandy, M. R., 1990, A New Early Cretaceous Articulate Brachiopod from the Northwest
1036 Territories, Canada, and Its Paleobiogeographic Significance: Journal of Paleontology, v.
1037 64, no. 3, p. 367-372.
1038 Sandy, M. R., Lazăr, I., Peckmann, J., Birgel, D., Stoica, M., and Roban, R. D., 2012, Methane-
1039 seep brachiopod fauna within turbidites of the Sinaia Formation, Eastern Carpathian
1040 Mountains, Romania: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 323–325,
1041 p. 42-59.
1042 Saumur, B. M., Dewing, K., and Williamson, M. C., 2016, Architecture of the Canadian portion
1043 of the High Arctic Large Igneous Province and implications for magmatic Ni–Cu
1044 potential: Canadian Journal of Earth Sciences, p. 1-15.
1045 Savard, M. M., Beauchamp, B., and Veizer, J., 1996, Significance of Aragonite Cements Around
1046 Cretaceous Marine Methane Seeps: Journal of Sedimentary Research, v. 66, no. 3, p.
1047 430-438.

Schoell, M., 1982, Applications of isotope analysis of petroleum and natural gas research: Spectra, v. 8, p. 32-41.
Schwartz, H., Sample, J., Weberling, K. D., Minisini, D., and Moore, J. C., 2003, An ancient linked fluid migration system: cold-seep deposits and sandstone intrusions in the Panoche Hills, California, USA: Geo-Marine Letters, v. 23, no. 3, p. 340-350.
Sibuet, M., and Olu, K., 1998, Biogeography, biodiversity and fluid dependence of deep-sea cold-seep communities at active and passive margins: Deep Sea Research Part II: Topical Studies in Oceanography, v. 45, no. 1, p. 517-567.
Stott, D. F., 1969, Ellef Ringnes Island, Canadian Arctic Archipelago: Geological Survey of Canada Paper 68-16.
Suess, E., 2014, Marine cold seeps and their manifestations: geological control, biogeochemical criteria and environmental conditions: International Journal of Earth Sciences, v. 103, no. 7, p. 1889-1916.
Taylor, S. R., and McLennan, S. M., 1985, The continental crust: Its composition and evolution, Blackwell Scientific Pub.,Palo Alto, CA; None, Medium: X; Size: Pages: 328 p.:
Teichert, B. M. A., and Luppold, F. W., 2013, Glendonites from an Early Jurassic methane seep — Climate or methane indicators?: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 390, p. 81-93.
Thiel, V., Peckmann, J., Schmale, O., Reitner, J., and Michaelis, W., 2001, A new straight-chain hydrocarbon biomarker associated with anaerobic methane cycling: Organic Geochemistry, v. 32, no. 8, p. 1019-1023.
Thorsteinsson, R., 1974, Carboniferous and Permian stratigraphy of Axel Heiberg Island and western Ellesmere Island, Canadian Arctic Archipelago: Canadian Geological Survey Bulletin, v. 224, p. 115.
Tribovillard, N., Algeo, T. J., Lyons, T., and Riboulleau, A., 2006, Trace metals as paleoredox and paleoproductivity proxies: An update: Chemical Geology, v. 232, no. 1–2, p. 12-32.
Turekian, K. K., and Wedepohl, K. H., 1961, Distribution of the elements in some major units of the Earth's crust: Geological Society of America Bulletin v. 72, no. 2, p. 175-191.
Voegelin, A. R., Nägler, T. F., Samankassou, E., and Villa, I. M., 2009, Molybdenum isotopic composition of modern and Carboniferous carbonates: Chemical Geology, v. 265, no. 3–4, p. 488-498.
Whiticar, M. J., 1999, Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane: Chemical Geology, v. 161, p. 291-314.
Wynne, P. J., Irving, E., and Osadetz, K. G., 1988, Paleomagnetism of cretaceous volcanic rocks of the sverdrup basin; magnetostratigraphy, paleolatitudes, and rotations: Canadian Journal of Earth Sciences, v. 25, no. 8, p. 1220-1239.

FIGURE CAPTIONS

Figure 1 Map showing location of the study area. The inset (Canadian Arctic Archipelago) shows the location of the Sverdrup Basin (outlined by dashed black line). Ellef Ringnes Island, the focus of this study, is located near the centre of the basin.

Figure 2 Map of Ellef Ringnes Island, composed of Lower Jurassic rocks in the northwest and upper most Cretaceous rocks in the southeast. The Island is pierced by seven salt diapirs, composed of the Pennsylvanian Otto Fiord Formation. Within the Christopher Formation, 139 early Cretaceous methane seep deposits have been discovered. The majority of the seep deposits are concentrated on the periphery of salt diapirs; with the exception of 8 seep deposits, not closely associated with diapirs. Geology is derived from (Evenchick and Embry, 2012a; Evenchick and Embry, 2012b).

Figure 3 Stratigraphic column of Cretaceous and Paleogene strata of the Sverdrup Basin. Stratigraphy is modified from (Embry and Beauchamp, 2008). Ancient methane seep deposits on Ellef Ringnes Island occur within the upper portion of the Invincible Point Member of the Christopher Formation, an Upper Aptian to Upper Albian marine shale. The horizon with occurrence of abundant glendonites is also shown.

Figure 4 Field photographs and thin sections of carbonate deposits. A) Distribution of carbonate deposits (marked by red arrows) as positive relief features on the arctic tundra, B) large carbonate deposit with relief up to 3 m, C) Carbonate bed with a

significantly greater length than height, D) Close up of outcrop showing characteristic banded cement texture, E) Photomicrograph of micritic cements within seep deposits, F) Photomicrograph under plain light showing banded cements, G) cross-polarised light image of banded yellow and botryoidal calcite, H) Close up of outcrop showing void filling blocky calcite. BC=blocky calcite, YC = yellow calcite, PL = peloid, OM = opaque mineral.

Figure 5 $\delta^{18}O_{VPDB}$ and $\delta^{13}C_{VPDB}$ plot of individual seep carbonate phases. All of the phases, excluding blocky calcite are very ^{13}C depleted, suggesting derivation from methane. The same phases have $\delta^{18}O_{VPDB}$ values close to zero, suggesting precipitation near ambient seawater temperature. Blocky calcite has much higher $\delta^{13}C_{VPDB}$ values and lower $\delta^{18}O_{VPDB}$ values, likely forming at increased temperature (depth) during burial diagenesis.

Figure 6 Geochemical plots of shale hosting carbonate seep deposits. Litholog of the Christopher formation shows section from the base of the formation to the resistant marker bed at the top of the Invincible Point Member. A) Plot of carbon isotopes of organic matter contained within the Christopher Formation as well as the seep carbonates (thought in part to be preserved biofilms). B) Plot of Mo concentration in both the Christopher Formation and seep carbonates. The Christopher shale contains less Mo than normal Post Archean Average Shale (Taylor and McLennan, 1985) suggesting precipitation under oxic conditions, whereas seep carbonates sequester much more Mo than normal carbonates, implying anoxic formation.

Figure 7 Chromatograms showing the lipid biomarkers found in the hydrocarbons and fatty acids recovered from seep carbonates. Stable carbon isotopes of the lipid biomarkers are also included.

Figure 8 Representative seep macrofossils. A) Solemyid bivalve; articulated specimen, internal mould lateral view. B) Lucinid bivalve; articulated specimen, internal mould lateral view; arrow points to anterior adductor muscle scar. C) Thyasirid bivalve: articulated specimen, internal mould lateral view. D) Same specimen as C), posterior view; arrow points to weak posterior sulcus. E) *Nucinella* sp.; articulated specimen, internal mould anterior view. F) Same specimen as E), lateral view; arrow points to lateral tooth. G) Malletid bivalve; partially articulated specimen, internal mould lateral view. H) Nuculid bivalve; articulated specimen, internal mould lateral view. I) Pectinid bivalve; articulated specimen, internal mould lateral view. J) Ammonite *Puzosia* aff. *sigmoidalis* Donovan; internal mould. K) Vetigastropod; internal mould lateral view. L) Cluster of probable vestimentiferan tubes. M) Photomicrograph under plain light of an oblique cross section through a probable vestimentiferan tube. Arrow points to 'delamination' structures. Note infilling of botryoidal banded calcite cement. N) Large pieces of wood preserved in seep carbonate at outcrop. Scale bars for A-K = 1 cm.