



UNIVERSITY OF LEEDS

This is a repository copy of *Development and reliability of a preliminary Foot Osteoarthritis Magnetic Resonance Imaging Score*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/113779/>

Version: Accepted Version

---

**Article:**

Halstead, J, Martín-Hervás, C, Hensor, EMA et al. (4 more authors) (2017) Development and reliability of a preliminary Foot Osteoarthritis Magnetic Resonance Imaging Score. *Journal of Rheumatology*, 44 (8). pp. 1257-1264. ISSN 0315-162X

<https://doi.org/10.3899/jrheum.160617>

---

© 2017. This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *The Journal of Rheumatology* following peer review. The definitive publisher-authenticated version Halstead, J, Martín-Hervás, C, Hensor, EMA et al. (4 more authors) (2017) Development and reliability of a preliminary Foot Osteoarthritis Magnetic Resonance Imaging Score. *Journal of Rheumatology*, 44 (8). pp. 1257-1264 is available online at: <http://dx.doi.org/10.3899/jrheum.160617>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Figure 2**



**Figure 2:** A and B show T1w and fat saturated T2w sagittal images. White arrow shows the talo-navicular joint was scored as a grade 0 (normal joint space and signal) joint space narrowing (JSN), grade 2 (moderate) osteophyte, the presence of effusion/synovitis, the presence of cysts, navicular bone were scored as grade 2 (34%-66%) bone marrow lesion (BML).

White arrow head shows the navicular-medial cuneiform was scored as grade 1 (increased signal in the joint space) JSN, grade 1 (mild) osteophytes, the presence of effusion/synovitis, the presence of cysts, and medial cuneiform bone were scored as grade 2 (34%-66%) bone marrow lesion (BML).