

Support and Assessment for Fall Emergency Referrals (SAFER) 2: a cluster randomised trial and systematic review of clinical effectiveness and cost-effectiveness of new protocols for emergency ambulance paramedics to assess older people following a fall with referral to community-based care when appropriate

Helen A Snooks, Rebecca Anthony, Robin Chatters, Jeremy Dale, Rachael Fothergill, Sarah Gaze, Mary Halter, Ioan Humphreys, Marina Koniotou, Phillipa Logan, Ronan Lyons, Suzanne Mason, Jon Nicholl, Julie Peconi, Ceri Phillips, Judith Phillips, Alison Porter, A Niroshan Siriwardena, Graham Smith, Alun Toghill, Mushtaq Wani, Alan Watkins, Richard Whitfield, Lynsey Wilson and Ian T Russell



**National Institute for
Health Research**

Support and Assessment for Fall Emergency Referrals (SAFER) 2: a cluster randomised trial and systematic review of clinical effectiveness and cost-effectiveness of new protocols for emergency ambulance paramedics to assess older people following a fall with referral to community-based care when appropriate

Helen A Snooks,^{1*} Rebecca Anthony,¹ Robin Chatters,²
Jeremy Dale,³ Rachael Fothergill,⁴ Sarah Gaze,¹
Mary Halter,⁵ Ioan Humphreys,⁶ Marina Koniotou,¹
Phillipa Logan,⁷ Ronan Lyons,¹ Suzanne Mason,²
Jon Nicholl,² Julie Peconi,¹ Ceri Phillips,⁶ Judith Phillips,⁸
Alison Porter,¹ A Niroshan Siriwardena,⁹
Graham Smith,¹⁰ Alun Toghill,¹⁰ Mushtaq Wani,¹¹
Alan Watkins,¹ Richard Whitfield,¹² Lynsey Wilson¹
and Ian T Russell¹

¹Patient and Population Health and Informatics, Swansea University Medical School, Swansea, UK

²School of Health and Related Research (SchARR), University of Sheffield, Sheffield, UK

³Warwick Medical School, University of Warwick, Coventry, UK

⁴Clinical Audit and Research Unit, London Ambulance Service NHS Trust, London, UK

⁵Faculty of Health and Social Care Sciences, St George's University Hospital, London, UK

⁶Swansea Centre for Health Economics, Swansea University, Swansea, UK

⁷Community Health Sciences, University of Nottingham, Nottingham, UK

⁸Centre for Innovative Ageing, Swansea University, Swansea, UK

⁹School of Health and Social Care, University of Lincoln, Lincoln, UK

¹⁰Service user for the SAFER 2 trial

¹¹Department of Geriatric and Stroke Medicine, Morriston Hospital, Swansea, UK

¹²Pre-hospital Emergency Research Unit (PERU), Welsh Ambulance Services NHS Trust, Cardiff, UK

*Corresponding author

Declared competing interests of authors: Helen Snooks is a member of the NIHR Journals Library Editorial Group.

Published March 2017

DOI: 10.3310/hta21130

This report should be referenced as follows:

Snooks HA, Anthony R, Chatters R, Dale J, Fothergill R, Gaze S, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER) 2: a cluster randomised trial and systematic review of clinical effectiveness and cost-effectiveness of new protocols for emergency ambulance paramedics to assess older people following a fall with referral to community-based care when appropriate. *Health Technol Assess* 2017;**21**(13).

Health Technology Assessment is indexed and abstracted in *Index Medicus/MEDLINE*, *Excerpta Medica/EMBASE*, *Science Citation Index Expanded (SciSearch®)* and *Current Contents®/Clinical Medicine*.

ISSN 1366-5278 (Print)

ISSN 2046-4924 (Online)

Impact factor: 4.058

Health Technology Assessment is indexed in MEDLINE, CINAHL, EMBASE, The Cochrane Library and the ISI Science Citation Index.

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics (COPE) (www.publicationethics.org/).

Editorial contact: journals.library@nhr.ac.uk

The full HTA archive is freely available to view online at www.journalslibrary.nhr.ac.uk/hta. Print-on-demand copies can be purchased from the report pages of the NIHR Journals Library website: www.journalslibrary.nhr.ac.uk

Criteria for inclusion in the *Health Technology Assessment* journal

Reports are published in *Health Technology Assessment* (HTA) if (1) they have resulted from work for the HTA programme, and (2) they are of a sufficiently high scientific quality as assessed by the reviewers and editors.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

HTA programme

The HTA programme, part of the National Institute for Health Research (NIHR), was set up in 1993. It produces high-quality research information on the effectiveness, costs and broader impact of health technologies for those who use, manage and provide care in the NHS. 'Health technologies' are broadly defined as all interventions used to promote health, prevent and treat disease, and improve rehabilitation and long-term care.

The journal is indexed in NHS Evidence via its abstracts included in MEDLINE and its Technology Assessment Reports inform National Institute for Health and Care Excellence (NICE) guidance. HTA research is also an important source of evidence for National Screening Committee (NSC) policy decisions.

For more information about the HTA programme please visit the website: <http://www.nets.nhr.ac.uk/programmes/hta>

This report

The research reported in this issue of the journal was funded by the HTA programme as project number 07/01/21. The contractual start date was in April 2009. The draft report began editorial review in May 2014 and was accepted for publication in September 2015. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The HTA editors and publisher have tried to ensure the accuracy of the authors' report and would like to thank the reviewers for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this report.

This report presents independent research funded by the National Institute for Health Research (NIHR). The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, NETSCC, the HTA programme or the Department of Health. If there are verbatim quotations included in this publication the views and opinions expressed by the interviewees are those of the interviewees and do not necessarily reflect those of the authors, those of the NHS, the NIHR, NETSCC, the HTA programme or the Department of Health.

© Queen's Printer and Controller of HMSO 2017. This work was produced by Snooks *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health. This issue may be freely reproduced for the purposes of private research and study and extracts (or indeed, the full report) may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Published by the NIHR Journals Library (www.journalslibrary.nhr.ac.uk), produced by Prepress Projects Ltd, Perth, Scotland (www.prepress-projects.co.uk).

Health Technology Assessment Editor-in-Chief

Professor Hywel Williams Director, HTA Programme, UK and Foundation Professor and Co-Director of the Centre of Evidence-Based Dermatology, University of Nottingham, UK

NIHR Journals Library Editor-in-Chief

Professor Tom Walley Director, NIHR Evaluation, Trials and Studies and Director of the EME Programme, UK

NIHR Journals Library Editors

Professor Ken Stein Chair of HTA Editorial Board and Professor of Public Health, University of Exeter Medical School, UK

Professor Andree Le May Chair of NIHR Journals Library Editorial Group (EME, HS&DR, PGfAR, PHR journals)

Dr Martin Ashton-Key Consultant in Public Health Medicine/Consultant Advisor, NETSCC, UK

Professor Matthias Beck Chair in Public Sector Management and Subject Leader (Management Group), Queen's University Management School, Queen's University Belfast, UK

Dr Tessa Crilly Director, Crystal Blue Consulting Ltd, UK

Dr Eugenia Cronin Senior Scientific Advisor, Wessex Institute, UK

Ms Tara Lamont Scientific Advisor, NETSCC, UK

Dr Catriona McDaid Senior Research Fellow, York Trials Unit, Department of Health Sciences, University of York, UK

Professor William McGuire Professor of Child Health, Hull York Medical School, University of York, UK

Professor Geoffrey Meads Professor of Health Sciences Research, Health and Wellbeing Research Group, University of Winchester, UK

Professor John Norrie Chair in Medical Statistics, University of Edinburgh, UK

Professor John Powell Consultant Clinical Adviser, National Institute for Health and Care Excellence (NICE), UK

Professor James Raftery Professor of Health Technology Assessment, Wessex Institute, Faculty of Medicine, University of Southampton, UK

Dr Rob Riemsma Reviews Manager, Kleijnen Systematic Reviews Ltd, UK

Professor Helen Roberts Professor of Child Health Research, UCL Institute of Child Health, UK

Professor Jonathan Ross Professor of Sexual Health and HIV, University Hospital Birmingham, UK

Professor Helen Snooks Professor of Health Services Research, Institute of Life Science, College of Medicine, Swansea University, UK

Professor Jim Thornton Professor of Obstetrics and Gynaecology, Faculty of Medicine and Health Sciences, University of Nottingham, UK

Professor Martin Underwood Director, Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, UK

Please visit the website for a list of members of the NIHR Journals Library Board:
www.journalslibrary.nihr.ac.uk/about/editors

Editorial contact: journals.library@nihr.ac.uk

Abstract

Support and Assessment for Fall Emergency Referrals (SAFER) 2: a cluster randomised trial and systematic review of clinical effectiveness and cost-effectiveness of new protocols for emergency ambulance paramedics to assess older people following a fall with referral to community-based care when appropriate

Helen A Snooks,^{1*} Rebecca Anthony,¹ Robin Chatters,² Jeremy Dale,³ Rachael Fothergill,⁴ Sarah Gaze,¹ Mary Halter,⁵ Ioan Humphreys,⁶ Marina Koniotou,¹ Phillipa Logan,⁷ Ronan Lyons,¹ Suzanne Mason,² Jon Nicholl,² Julie Peconi,¹ Ceri Phillips,⁶ Judith Phillips,⁸ Alison Porter,¹ A Niroshan Siriwardena,⁹ Graham Smith,¹⁰ Alun Toghill,¹⁰ Mushtaq Wani,¹¹ Alan Watkins,¹ Richard Whitfield,¹² Lynsey Wilson¹ and Ian T Russell¹

¹Patient and Population Health and Informatics, Swansea University Medical School, Swansea, UK

²School of Health and Related Research (SchARR), University of Sheffield, Sheffield, UK

³Warwick Medical School, University of Warwick, Coventry, UK

⁴Clinical Audit and Research Unit, London Ambulance Service NHS Trust, London, UK

⁵Faculty of Health and Social Care Sciences, St George's University Hospital, London, UK

⁶Swansea Centre for Health Economics, Swansea University, Swansea, UK

⁷Community Health Sciences, University of Nottingham, Nottingham, UK

⁸Centre for Innovative Ageing, Swansea University, Swansea, UK

⁹School of Health and Social Care, University of Lincoln, Lincoln, UK

¹⁰Service user for the SAFER 2 trial

¹¹Department of Geriatric and Stroke Medicine, Morriston Hospital, Swansea, UK

¹²Pre-hospital Emergency Research Unit (PERU), Welsh Ambulance Services NHS Trust, Cardiff, UK

*Corresponding author h.a.snooks@swansea.ac.uk

Background: Emergency calls are frequently made to ambulance services for older people who have fallen, but ambulance crews often leave patients at the scene without any ongoing care. We evaluated a new clinical protocol which allowed paramedics to assess older people who had fallen and, if appropriate, refer them to community-based falls services.

Objectives: To compare outcomes, processes and costs of care between intervention and control groups; and to understand factors which facilitate or hinder use.

Design: Cluster randomised controlled trial.

Participants: Participating paramedics at three ambulance services in England and Wales were based at stations randomised to intervention or control arms. Participants were aged 65 years and over, attended by a study paramedic for a fall-related emergency service call, and resident in the trial catchment areas.

Interventions: Intervention paramedics received a clinical protocol with referral pathway, training and support to change practice. Control paramedics continued practice as normal.

Outcomes: The primary outcome comprised subsequent emergency health-care contacts (emergency admissions, emergency department attendances, emergency service calls) or death at 1 month and 6 months. Secondary outcomes included pathway of care, ambulance service operational indicators, self-reported outcomes and costs of care. Those assessing outcomes remained blinded to group allocation.

Results: Across sites, 3073 eligible patients attended by 105 paramedics from 14 ambulance stations were randomly allocated to the intervention group, and 2841 eligible patients attended by 110 paramedics from 11 stations were randomly allocated to the control group. After excluding dissenting and unmatched patients, 2391 intervention group patients and 2264 control group patients were included in primary outcome analyses. We did not find an effect on our overall primary outcome at 1 month or 6 months. However, further emergency service calls were reduced at both 1 month and 6 months; a smaller proportion of patients had made further emergency service calls at 1 month (18.5% vs. 21.8%) and the rate per patient-day at risk at 6 months was lower in the intervention group (0.013 vs. 0.017). Rate of conveyance to emergency department at index incident was similar between groups. Eight per cent of trial eligible patients in the intervention arm were referred to falls services by attending paramedics, compared with 1% in the control arm. The proportion of patients left at scene without further care was lower in the intervention group than in the control group (22.6% vs. 30.3%). We found no differences in duration of episode of care or job cycle. No adverse events were reported. Mean cost of the intervention was £17.30 per patient. There were no significant differences in mean resource utilisation, utilities at 1 month or 6 months or quality-adjusted life-years. In total, 58 patients, 25 paramedics and 31 stakeholders participated in focus groups or interviews. Patients were very satisfied with assessments carried out by paramedics. Paramedics reported that the intervention had increased their confidence to leave patients at home, but barriers to referral included patients' social situations and autonomy.

Conclusions: Findings indicate that this new pathway may be introduced by ambulance services at modest cost, without risk of harm and with some reductions in further emergency calls. However, we did not find evidence of improved health outcomes or reductions in overall NHS emergency workload. Further research is necessary to understand issues in implementation, the costs and benefits of e-trials and the performance of the modified Falls Efficacy Scale.

Trial registration: Current Controlled Trials ISRCTN60481756 and PROSPERO CRD42013006418.

Funding: This project was funded by the National Institute for Health Research (NIHR) Health Technology Assessment programme and will be published in full in *Health Technology Assessment*; Vol. 21, No. 13. See the NIHR Journals Library website for further project information.

Contents

List of tables	xiii
List of figures	xvii
List of abbreviations	xix
Plain English summary	xxi
Scientific summary	xxiii
Chapter 1 Introduction	1
Structure of this report	1
Background	1
<i>Falls in older people</i>	1
<i>Emergency department overcrowding and potentially avoidable attendances</i>	2
<i>Conveyance rate of older adults who fall</i>	2
<i>Service models: community falls services</i>	3
Theoretical underpinning for intervention	4
Research aim and objectives	5
<i>Aim</i>	5
<i>Objectives</i>	5
Chapter 2 Systematic review of literature on effects of interventions by emergency medical staff for older people who fall	7
Abstract	7
<i>Background</i>	7
<i>Objective</i>	7
<i>Method</i>	7
<i>Results</i>	7
<i>Discussion</i>	7
<i>Study registration</i>	7
Introduction	7
<i>Background and rationale for the review</i>	8
Methods	8
<i>Protocol and registration</i>	8
<i>Eligibility criteria</i>	8
<i>Sources and search strategy</i>	9
<i>Study selection</i>	11
<i>Data extraction</i>	11
<i>Assessment of the risk of bias (quality)</i>	12
<i>Data synthesis</i>	12
Findings	12
<i>Inclusion</i>	12
<i>Study characteristics</i>	12
<i>Methodological quality</i>	13
<i>Study findings</i>	13

Discussion	22
<i>Strengths and limitations of this review</i>	23
<i>Review findings as context for Support and Assessment for Fall Emergency Referrals 2</i>	23
Chapter 3 Methods	25
Overview	25
<i>Aim</i>	25
<i>Objectives</i>	25
Cluster randomised controlled trial	25
<i>Trial design</i>	25
<i>Trial management</i>	25
<i>Setting</i>	26
<i>Participants</i>	26
<i>Randomisation</i>	26
<i>Recruitment</i>	26
<i>Interventions</i>	27
<i>Outcomes</i>	27
<i>Data collection</i>	29
<i>Data entry</i>	29
<i>Blinding</i>	29
<i>Sample size and power</i>	29
<i>Statistical methods</i>	30
<i>Missing data</i>	31
<i>Adverse events</i>	31
Economic evaluation	31
<i>Aim</i>	31
<i>Method</i>	31
<i>Health-related quality-of-life outcome</i>	32
<i>Lost to follow-up/missing data</i>	32
<i>Discounting</i>	32
<i>Uncertainty</i>	32
Qualitative methods	32
<i>Aim</i>	32
<i>Participants</i>	32
Data collection	33
<i>Patient interviews</i>	33
<i>Paramedic focus groups and interviews</i>	33
<i>Stakeholder focus groups and interviews</i>	34
<i>Qualitative data management</i>	34
<i>Qualitative data analysis</i>	34
<i>Involvement of service users</i>	35
Support and Assessment for Fall Emergency Referrals 2 trial progress	36
Summary of changes to the project protocol	38
Ethics and research governance	39
<i>Trial registration</i>	39
Chapter 4 Clinical effectiveness results	41
Participant flow	41
Recruitment	41
Baseline characteristics	41
<i>Individual patient level</i>	41
<i>Cluster level</i>	41
Numbers analysed	44

Outcomes and estimation	44
<i>Contamination</i>	55
<i>Patients ineligible for referral to the falls service</i>	55
<i>Patterns of referral to the falls service</i>	55
<i>Comparison of outcome measures</i>	58
Analysis of non-response	62
Harms	62
Chapter 5 Cost-effectiveness analysis	63
Results: summary of demographics	63
Costs of Support and Assessment for Fall Emergency Referrals 2 trial intervention	63
Resource use	63
Health-related quality of life	72
Cost-effectiveness analysis	72
Conclusion	72
Chapter 6 Qualitative results	75
Introduction	75
How the intervention was experienced by patients	76
<i>The fall in context</i>	76
<i>Patient experience of the ambulance service</i>	78
<i>Patient experience: those referred to a falls service</i>	81
<i>Patient experience: those who attended hospital</i>	82
<i>Patient health and quality of life since fall</i>	83
<i>The patient in context: other health and social care provision</i>	84
How the intervention was delivered in practice: paramedic and stakeholder views	85
<i>Paramedic and stakeholder expectations of the new intervention (pre trial)</i>	85
<i>Paramedics' and other stakeholders' experiences and views on context and existing practice</i>	86
<i>Experience of implementing the Support and Assessment for Fall Emergency Referrals 2 trial intervention</i>	88
<i>Factors which enabled/encouraged the use of the Support and Assessment for Fall Emergency Referrals 2 trial intervention</i>	88
<i>Factors which hindered the use of the Support and Assessment for Fall Emergency Referrals 2 trial intervention</i>	89
<i>Impact of the Support and Assessment for Fall Emergency Referrals 2 trial intervention on paramedic practice</i>	91
<i>Impact of the new pathway on falls services</i>	92
<i>Suggestions for modifications or developments of the Support and Assessment for Fall Emergency Referrals 2 trial intervention</i>	93
Summary of qualitative findings	94
<i>Patient experience</i>	94
<i>Paramedic and stakeholder experience</i>	95
Limitations of qualitative methods	96
<i>Patient experience</i>	96
<i>Paramedic and stakeholder experience</i>	96
Chapter 7 Discussion, conclusions and recommendations	97
Summary of key findings	97
Strengths and limitations	98
Implications (practice, policy and research)	99
Conclusions and recommendations for research	100

Acknowledgements	103
References	107
Appendix 1 Support and Assessment for Fall Emergency Referrals 2 clinical decision protocol	115
Appendix 2 Full data extraction and quality assessment tables	117
Appendix 3 Excluded studies	129
Appendix 4 Support and Assessment for Fall Emergency Referrals 2 organisational flow chart	137
Appendix 5 Station location maps	139
Appendix 6 Support and Assessment for Fall Emergency Referrals 2 postal consent pack	143
Appendix 7 Support and Assessment for Fall Emergency Referrals 2 1-month questionnaire	151
Appendix 8 Final definition of the intervention	159
Appendix 9 Support and Assessment for Fall Emergency Referrals 2 paramedic training session programme	165
Appendix 10 Support and Assessment for Fall Emergency Referrals 2 6-month questionnaire	167
Appendix 11 Support and Assessment for Fall Emergency Referrals 2 data analysis plan	173
Appendix 12 Patient semistructured interviews schedule	183
Appendix 13 Support and Assessment for Fall Emergency Referrals 2 paramedic pretrial focus group interview schedule	187
Appendix 14 Support and Assessment for Fall Emergency Referrals 2 paramedic trial-end interview schedule	189
Appendix 15 Stakeholder trial-end interview schedules	191
Appendix 16 Support and Assessment for Fall Emergency Referrals 2 framework for analysis of interviews and focus groups	195
Appendix 17 Support and Assessment for Fall Emergency Referrals 2 Involving People case study	199
Appendix 18 Site Consolidated Standards of Reporting Trials flow charts	201
Appendix 19 Further clinical effectiveness results	205

List of tables

TABLE 1 Inclusion and exclusion criteria	9
TABLE 2 Sources of literature used	10
TABLE 3 Search carried out in MEDLINE, the Allied and Complementary Medicine Database, and Cumulative Index to Nursing and Allied Health Literature Plus	11
TABLE 4 Summary of study characteristics	14
TABLE 5 Results from studies assessing outcomes at the time of the incident	15
TABLE 6 Results from studies assessing outcomes after the incident	16
TABLE 7 Summary of quality of included studies	18
TABLE 8 Studies reporting number of referrals made to a community service	20
TABLE 9 Studies reporting patient acceptance of referrals to a community service	20
TABLE 10 Studies reporting number of patients conveyed to the ED as an outcome	21
TABLE 11 Definition of the SAFER 2 trial intervention	28
TABLE 12 Periods of recruitment	43
TABLE 13 Baseline individual and cluster characteristics by treatment allocated	43
TABLE 14 Questionnaire response rates by treatment allocated	45
TABLE 15 Primary outcome and components at 1 month analysed by treatment allocated	46
TABLE 16 Primary outcome and components at 6 months analysed by treatment allocated	47
TABLE 17 Primary outcome and components at 1 month at each site analysed by treatment allocated	48
TABLE 18 Primary outcome and components at 6 months at each site analysed by treatment allocated	49
TABLE 19 Secondary outcomes at index incident analysed by treatment allocated	51
TABLE 20 Secondary outcomes at 1 month analysed by treatment allocated	53
TABLE 21 Secondary outcomes at 6 months analysed by treatment allocated	56
TABLE 22 Numbers of control group participants subsequently attended by an intervention group paramedic	57

TABLE 23 Falls referrals per intervention paramedic by site	58
TABLE 24 Falls referrals at index call for intervention patients by personal and incident characteristics	58
TABLE 25 Predictors of falls referral (full effects model)	59
TABLE 26 Reasons identified for non-referral of patients left at scene by intervention group paramedics	59
TABLE 27 Comparison of outcome measures at 1 and 6 months	59
TABLE 28 Comparison of characteristics of responders and non-responders to questionnaires at 1 month	62
TABLE 29 Serious adverse events within 2 days following index incident	62
TABLE 30 Intervention costs	64
TABLE 31 Published average unit costs	68
TABLE 32 Health-care resource use over 6 months by group	69
TABLE 33 Summary of health-care resource use over 6 months by group	71
TABLE 34 Mean SF-6D utilities over 6 months by group	72
TABLE 35 Sensitivity analysis	73
TABLE 36 Number of patients participating in the qualitative interviews	76
TABLE 37 Summary of paramedic and stakeholder interviews	77
TABLE 38 Data extraction table	118
TABLE 39 Quality assessment table (cohort studies)	125
TABLE 40 Quality assessment table (RCT)	126
TABLE 41 Quality assessment table (cost-effectiveness)	126
TABLE 42 Quality assessment table (qualitative)	127
TABLE 43 Final definition of the intervention	160
TABLE 44 Primary outcomes at 1 month, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral	206
TABLE 45 Components of the primary outcome at 1 month, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral	208

TABLE 46 Primary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral	210
TABLE 47 Components of the primary outcome at 6 months, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral	212
TABLE 48 Secondary outcomes at 1 month, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral	214
TABLE 49 Secondary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral	217

List of figures

FIGURE 1 The PRISMA flow chart	13
FIGURE 2 Flow of clusters and individuals through RCT for all participants	42
FIGURE 3 Short Form questionnaire-12 items PCS and mFES scores at 1 month: correlation = 0.546	60
FIGURE 4 Short Form questionnaire-12 items MCS and mFES scores at 1 month: correlation = 0.553	60
FIGURE 5 Short Form questionnaire-12 items PCS and mFES scores at 6 months: correlation = 0.627	61
FIGURE 6 Short Form questionnaire-12 items MCS and mFES scores at 6 months: correlation = 0.547	61
FIGURE 7 Cost-effectiveness plane for the SAFER 2 trial intervention	73

List of abbreviations

A&E	accident and emergency	PCS	physical component summary
AE	adverse event	PCT	primary care trust
AR	adverse reaction	PICOCS	population, intervention, comparator, outcomes, context and study design
CI	confidence interval	PRF	patient report form
CLRN	comprehensive local research network	PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
CONSORT	Consolidated Standards of Reporting Trials	ProFaNE	Prevention of Falls Network Europe
C-RCT	cluster randomised controlled trial	PSS	Personal Social Services
DMEC	Data Monitoring and Ethics Committee	PSSRU	Personal Social Services Research Unit
ECP	emergency care practitioner	QALY	quality-adjusted life-year
ED	emergency department	R&D	research and development
EMS	emergency medical service	RCT	randomised controlled trial
ePRF	electronic patient report form	SAE	serious adverse event
EQ-5D	European Quality of Health-5 Dimensions	SAFER	Support and Assessment for Fall Emergency Referrals study
GP	general practitioner	SAIL	Secure Anonymised Information Linkage database
HRA CAG	Health Research Authority Confidentiality Advisory Group	SD	standard deviation
HSCIC	Health and Social Care Information Centre	SF-12	Short Form questionnaire-12 items
HTA	Health Technology Assessment	SF-6D	Short Form questionnaire-6 Dimensions
ICC	intracluster correlation	SOP	standard operating procedure
LIT	local implementation team	TMG	Trial Management Group
MCS	mental component summary	TSC	Trial Steering Committee
MeSH	medical subject heading	WWORTH	West Wales Organisation for Rigorous Trials in Health and social care
mFES	modified Falls Efficacy Scale		
NICE	National Institute for Health and Care Excellence		
OR	odds ratio		

Plain English summary

The Support and Assessment for Fall Emergency Referrals (SAFER) 2 study aimed to assess the costs and benefits of new protocols for paramedics to assess older people following a fall, with an option to leave them at home with a referral to a community falls service. In three UK ambulance services, we compared what happened to patients attended by paramedics with the new protocols (intervention group) with what happened to patients attended by paramedics delivering usual care (control group). We interviewed a small sample of patients, paramedics and other staff about their experiences of the new model of care.

A total of 4655 patients were included in the trial. There were no differences in the number of further emergency health-care contacts or deaths between groups, but patients in the intervention group were less likely to make further emergency service calls. Although only 8% of patients were referred directly to falls services by paramedics, overall this meant that fewer patients were left at home without further care. The intervention was as safe as usual practice, and we did not find any differences in how long paramedics spent on each job or in patients' health and quality of life. Patients were generally happy with the care they received and paramedics found that the protocol increased their confidence.

The SAFER 2 study findings indicate that ambulance services may introduce this new pathway safely and at low cost, and expect reductions in further emergency service calls. However, we did not find any evidence of improved quality of life for patients or reductions in overall NHS emergency workload.

Scientific summary

Background

Emergency calls to ambulance services are frequently made for older people who have fallen, but ambulance crews often leave patients at the scene without any ongoing care. Subsequent falls and emergency episodes of care are common in this group. In the Support and Assessment for Fall Emergency Referrals (SAFER) 2 trial, we evaluated a new clinical protocol that allows paramedics to assess older people who had fallen and refer those who do not need to be taken to the emergency department (ED) to community-based falls services for continuing care.

Aim and objectives

Aim

To assess the benefits and costs of a complex intervention comprising training and clinical protocols enabling paramedics to assess older people who have fallen and refer them to falls services when appropriate.

Objectives

- To compare outcomes, processes and costs of care between intervention and control groups:
 - patient outcomes: rate and pattern of subsequent emergency health-care contacts or deaths; health-related quality of life; falls efficacy (fear of falling); and change in place of residence
 - processes of care: pathway from index incident; subsequent health-care contacts; ambulance service operational indicators and protocol compliance including clinical documentation
 - costs of care to the NHS.
- To understand how patients experience the new intervention.
- To identify factors which facilitate or hinder the use of the intervention.
- To inform the development of methods for falls research.

Methods

We undertook a systematic review of published evidence related to the effectiveness of emergency care interventions by ambulance crews for older people who had fallen. We searched the following electronic databases: The Cochrane Library, Allied and Complementary Medicine Database (AMED), Applied Social Sciences Index and Abstracts (ASSIA), British Nursing Index (BNI), Cumulative Index to Nursing and Allied Health Literature (CINAHL) Plus, Internurse, MEDLINE, PsycINFO, PubMed, Scopus, The British Library, UK Institutional Library Search, GreyNet, Conference Proceedings Citation Index (Web of Science), OpenGrey and The British Library's Electronic Table of Contents (Zetoc) between 1990 and July 2013.

The SAFER 2 study was a multicentre, cluster randomised controlled trial with economic evaluation and qualitative component. We undertook the trial in defined areas within three ambulance services in which falls services were set up but direct referral from the ambulance service did not take place. We allocated ambulance stations, and the paramedics who volunteered to take part, to intervention and control groups. We included patients in the trial if they were aged ≥ 65 years, resident in the catchment area of participating falls services, and attended by a study paramedic following an emergency call to the

ambulance service that was coded by a dispatcher as a fall without priority symptoms. Patients were recruited for their first incident within the trial period.

The complex intervention that we evaluated comprised assessment protocol, referral pathway, falls service, and training, clinical and operational support to paramedics. We asked paramedics at control stations to continue their usual practice, comprising routine assessment, initial care, assistance in moving and conveyance to ED unless the patient refused.

The primary outcome comprised subsequent emergency health-care contacts (death, emergency admissions, ED attendances and emergency service calls) at 1 month and 6 months.

Secondary outcomes from routine data:

- referral to falls service or other provider
- compliance with clinical documentation guidelines
- durations of ambulance service job cycle and episode of care
- duration of subsequent inpatient episodes
- subsequent fractures
- costs of care.

Secondary outcomes from patient questionnaires:

- health-related quality of life [Short Form questionnaire-12 items (SF-12)]
- patient satisfaction (Quality of Care Monitor, at 1 month only)
- 'fear of falling' [modified Falls Efficacy Scale (mFES)]
- self-reported falls.

Routine outcomes were retrieved from the ambulance services and matched to central NHS registers by the National Welsh Informatics Service and the Health and Social Care Information Centre (HSCIC) in to link them to NHS and Office for National Statistics data sets. We linked self-reported outcomes from postal questionnaires to the routine data by trial identifiers and analysed them in the Secure Anonymised Information Linkage (SAIL) databank without identifiers.

We collected qualitative data from patients attended by intervention paramedics concerning their experience and satisfaction, and from ambulance service paramedics and managers on their views of implementation of the intervention.

We analysed quantitative data by treatment allocated except for those describing uptake of the intervention. Qualitative data were analysed using a framework approach.

We obtained ethical approvals from the Research Ethics Committee for Wales, the National Information Governance Board and each participating health board, NHS trust and primary care trust. We followed all patients who did not actively decline consent (dissent) through the anonymised route.

Results

Systematic review

Of 5355 papers identified by searches, 138 appeared potentially eligible from title and abstract; we included 13 papers (12 studies) after detailed reading of full text. Studies from the USA, the UK, Australia and New Zealand comprised two randomised controlled trials, nine cohort studies and one qualitative study. These yielded limited and weak evidence of the most effective ways to deliver alternative treatments to older people who fall. A minority identified benefits – reductions in subsequent emergency calls and hospital admissions –

for patients referred to a falls prevention service or attended by a paramedic with additional skills. However, several studies did not explore such outcomes and the quality of most of the studies was low. We concluded that robust evidence from well-designed and rigorous research is needed to inform service delivery.

Support and Assessment for Fall Emergency Referrals 2 cluster randomised controlled trial

Recruitment

Between March 2011 and June 2012, across the three study sites, 3073 eligible patients were attended by 105 paramedics based at 14 ambulance stations randomly allocated to the intervention group, and 2841 eligible patients were attended by 110 paramedics based at 11 stations allocated to the control group. After excluding dissenting and unmatched patients, we included 2391 from the intervention group and 2264 from the control group in primary outcome analyses at 1 and 6 months.

Baseline data

Recruitment of patients was higher at site 1 than at the other sites. There was little difference in age, sex, time of call, distance to ED or time to recruitment between trial arms. There was one more intervention station at each site, but fewer paramedics per station.

Missing data

We retrieved a number of routine data for all 4704 eligible patients who did not dissent. However, neither the SAIL database nor HSCIC enabled us to match 49 of these, or to retrieve data on their ED attendances, hospital admissions and death. Questionnaire response rates at 1 month varied between sites from 30% to 51%, with 1307 questionnaires returned; and at 6 months from 53% to 74%, with 678 questionnaires returned.

Outcomes and estimation

Clinical effectiveness

One-third of patients had suffered a further emergency episode or death by 1 month [870 out of 2391 (36.4%) intervention group patients and 843 out of 2264 (37.2%) control group patients; odds ratio (OR) 0.96, 95% confidence interval (CI) 0.85 to 1.08]. By 6 months this had risen to two-thirds [1701 out of 2391 (71.1%) intervention group patients and 1592 out of 2264 (70.3%) control group patients; OR 1.02, 95% CI 0.89 to 1.16].

We found evidence of fewer emergency service calls by 1 month [442 out of 2391 (18.5%) intervention group patients called vs. 493 out of 2264 (21.8%) control group patients; OR 0.82, 95% CI 0.70 to 0.94]; thus, the mean number of further calls per patient per day at risk was 0.020 and 0.025, respectively (difference -0.004, 95% CI -0.008 to 0.000). By 6 months, 1046 out of 2391 (43.7%) intervention group patients had called the emergency services, compared with 1046 out of 2264 (46.2%) control group patients (OR 0.90, 95% CI 0.80 to 1.01); the mean number of further calls per patient per day at risk was 0.013 and 0.017, respectively (difference -0.005, 95% CI -0.007 to -0.002). These differences were largely consistent across sites, and there was some evidence of fewer ED attendances at 6 months (mean number of further attendances per patient: intervention group, 0.84; control group, 0.91; event ratio 0.81, 95% CI 0.72 to 0.91).

The proportion of patients transported to the ED at the time of index incident was similar in the two groups [1579 out of 2420 (65.2%) patients in the intervention group compared with 1431 out of 2284 (62.7%) patients in the control group; OR 1.08, 95% CI 0.96 to 1.22]. However, this proportion varied considerably by site, from < 60% at site 1 to nearly 80% at site 2. Intervention paramedics referred 8% of patients to falls services, ranging from 7.5% at site 1 to 9.7% at site 3. Control paramedics referred only 1.1% of patients to falls services, with some variation between sites. The number of patients left at the

scene without any referral for further care was lower in the intervention group (547 out of 2420; 22.6%) than in the control group (692 out of 2284; 30.3%) (OR 0.69, 95% CI 0.60 to 0.78).

Completion of clinical documentation was high (> 90% on all key physiological indicators recorded on-scene) across sites, with no clear difference between trial groups. We also found no differences in operational indicators [the mean duration of episodes of care, from emergency service call until patient's emergency episode was complete, was 196.8 minutes in the intervention group and 192.8 minutes in the control group (difference 2.05 minutes, 95% CI –6.68 to 10.77 minutes); and the mean duration of the job cycle, from emergency service call until the ambulance was free, was 99.9 minutes in the intervention group and 97.8 minutes in the control group (difference 1.69 minutes, 95% CI –0.75 to 4.12 minutes)].

We did not find differences in the following measures:

Duration (mean) of inpatient stay, which at 1 month was 2.25 days in the intervention group versus 2.10 days in the control group (difference 0.14 days, 95% CI –0.21 to 0.49 days) and at 6 months was 11.18 days in the intervention group versus 11.62 days in the control group (difference –0.56 days, 95% CI –1.88 to 0.76 days).

SF-12 mental health component (mean score), which at 1 month was 39.80 in the intervention group versus 38.89 in the control group (difference 0.90, 95% CI –0.74 to 2.55) and at 6 months was 43.21 in the intervention group versus 42.82 in the control group (difference 0.46, 95% CI –1.72 to 2.64).

SF-12 physical health component (mean score), which at at 1 month was 29.07 in the intervention group versus 29.40 in the control group (difference –0.50, 95% CI –1.85 to 0.86) and at 6 months was 30.44 in the intervention group versus 31.88 in the control group (difference –1.30, 95% CI –3.28 to 0.68).

Fall-specific mFES (mean score), which at 1 month was 3.71 in the intervention group versus 3.82 in the control group (adjusted difference –0.06, 95% CI –0.39 to 0.28) and at 6 months was 4.55 in the intervention group versus 4.79 in the control group (adjusted difference –0.23, 95% CI –0.73 to 0.27).

Satisfaction with care: technical mean scores at 1 month were 62.82 in the intervention group versus 63.21 in the control group (adjusted difference –0.32, 95% CI –1.27 to 0.63); in contrast, interpersonal mean scores were significantly higher, that is 68.92 in the intervention group versus 68.04 in the control group (adjusted difference 3.13, 95% CI 1.59 to 4.68).

Fewer intervention patients reported further falls by 1 month: 413 out of 621 (66.5%) in the intervention group versus 409 out of 589 (69.4%) in the control group (OR 0.72, 95% CI 0.54 to 0.96). However, there was no significant difference in subsequent fractures by 1 month, being reported by 98 out of 2391 patients in the intervention group (4.1%) versus 91 out of 2264 (4.0%) patients in the control group (OR 1.00, 95% CI 0.74 to 1.35). By 6 months, however, significant interactions between intervention and site masked any generic effect on further falls or subsequent fractures.

Sixty per cent of intervention group paramedics used their protocols to refer patients to falls services (i.e. between one and 11 times). Patients' age, sex and distance to ED did not influence these referrals. However, only 19 of 40 paramedics (48%) at site 3 referred patients to falls services, compared with 26 of 39 (67%) at site 1 and 19 of 26 (73%) at site 2.

Cost-effectiveness

We estimated the mean cost of the SAFER 2 study intervention across the three ambulance services as £17.30 per patient, including generic and local set-up costs such as training package development, training delivery and clinical support for implementation.

We estimated mean resource use by 1 month to be £3740.00 in the intervention group and £3514.00 in the control group (adjusted difference £190.24, 95% CI –£13.83 to £394.31), with the cost of initial hospital stays (£2523.27 and £2329.79, respectively) a major cost driver. Estimated mean resource use by 6 months was £8816.41 in the intervention group and £8661.77 in the control group (adjusted difference £24.20, 95% CI –£468.01 to £516.40), with the cost of subsequent hospital stays (£3982.21 and £4111.10, respectively) a major cost driver.

The imputed Short Form questionnaire-6 Dimensions (health and quality-of-life outcomes) scores showed that mean utilities over 6 months were slightly higher in the control group; however, the adjusted difference of –0.0026 was not statistically significant (95% CI 0.0066 to 0.0014).

Qualitative findings

Almost all of the 58 patients interviewed were very satisfied with the processes of assessment and examination carried out by paramedics. Most of those able to compare their SAFER 2 study encounter with the ambulance service with a previous contact after a fall could see little difference in the assessment and care delivered. Many patients were already accessing a complex network of community-based services, from a range of providers (NHS, social services, third sector).

Twenty-four paramedics participated in focus groups or interviews before implementation, and 25 paramedics and 31 other stakeholders, including ambulance service managers, trainers and falls services staff, participated in focus groups or interviews after the trial.

Before the trial, paramedics reported that they expected making significant use of the intervention. Post trial, paramedics in all sites reported that the intervention had increased their confidence in leaving patients at home. The structured training on the SAFER 2 study was felt to be more helpful than previous and more informal approaches to introducing innovations. They suggested that a 'simpler' intervention would have sufficed, with a less detailed flow chart. Reported barriers to referral included, in all sites, issues relating to patients' social situation and autonomy, and, in one site, a lack of knowledge of the role of the falls team. Some paramedics were concerned about increased times on-scene, but managers did not see this as an issue.

Methodological findings

We compared the performance of the generic SF-12 outcome measure with the falls-specific outcome measure, the mFES, in the study population and found the moderate correlation between them ($r = 0.55$ to 0.63), with significant floor and ceiling effects for the condition-specific measure.

Conclusions

The SAFER 2 study intervention to facilitate referral to community-based falls services was inexpensive and safe. The number of referrals made to falls services was lower than expected, and there was considerable variation between paramedics. The intervention reduced the number of patients left at the scene by attending ambulance crews without ongoing care, but had little effect on other processes of care. Although the numbers of further emergency health-care contacts and deaths were unchanged overall, subsequent emergency service calls were significantly reduced by 1 and 6 months. There was little evidence of impact on self-reported health outcomes or satisfaction.

Retrieval of anonymised linked data outcomes was highly successful in this 'e-trial', and ensured that findings were generalisable to the whole study population. We plan further analysis trial findings to describe fully the benefits and risks of this new approach.

The SAFER 2 trial has shown that ambulance services can introduce this new clinical pathway for patients without risk of harm, at modest cost and with some reductions in further emergency contacts. However, we did not find evidence of improved health outcomes for patients.

Future work recommendations

Further research is necessary to understand ambulance service implementation and paramedic uptake issues related to the falls intervention tested in the SAFER 2 study; costs and benefits of using anonymised linked data outcomes in trials; and performance of the mFES in this population.

Trial registration

This trial is registered as ISRCTN60481756 and as PROSPERO number CRD42013006418.

Funding

Funding for this study was provided by the Health Technology Assessment programme of the National Institute for Health Research.

Chapter 1 Introduction

Structure of this report

This study was a cluster randomised controlled trial (C-RCT) of a protocol for emergency ambulance paramedics to assess, and to refer to community-based care, older adults who had fallen and incorporated an economic evaluation and qualitative research.

The report begins with an overview of the literature and policy relating to older adults falling, avoidable conveyances to hospital, current paramedic practice, community falls services and the use of protocols for paramedics. We describe the underpinning theoretical framework for the intervention, clinical effectiveness and cost-effectiveness results and the qualitative findings.

We report a systematic review of the literature relating to interventions in the ambulance setting with older people who fall, and then describe the methods used to conduct the main C-RCT. We report the results of the trial, followed by results from the economic analysis and the qualitative components.

In the final chapter, we summarise and synthesise the findings from all three components of the study, the systematic review, C-RCT and qualitative study, providing interpretation in the light of other studies. We discuss the strengths and limitations of the research and generalisability to the NHS. The conclusions are followed by recommendations for future research.

Background

Falls in older people

Falls in older people are recognised as an important issue internationally,^{1,2} with high human costs, for the individual and their carers, and high organisational costs. It is estimated that around 30% of home-dwelling people aged ≥ 65 years fall every year,^{3–8} and the rate of falls is even higher among care home residents.⁹ Studies suggest that falls in older people are the cause of 20–30% of mild to severe injuries and 10–15% of emergency department (ED) attendances in this group.^{10,11} Falls in older people are often due to more than one underlying cause or risk factor; as the number of risk factors rises, so does the risk of falling.¹² Risk factors include muscle weakness, balance and gait, low blood pressure (or drop in blood pressure on assuming upright posture) and cognitive decline. The severity of fall-related complications increases with age.^{13,14} Falls are a cause of substantial rates of mortality and morbidity, as well as considerable contributors to immobility.⁸ Reduction in quality of life and physical activity leads to social isolation and functional deterioration with a high risk of resultant dependency and nursing home institutionalisation.^{15–18} Recovery from fall injury is often delayed in older people, which increases the risk of subsequent falls.⁸ Recurrent falls are associated with greater physician contact and functional decline.^{19–21} An additional complication is post-fall anxiety, leading to a fear of falling, which in turn can further contribute to deconditioning and weakness and in the long run may actually increase risk of falls.⁸

The *About the National Health Interview Survey*²² indicates that falls are the largest single cause of restricted activity days among older adults. Most people who fall do not seek any medical advice, and those who do are more likely to report to their general practitioner (GP).^{23,24} The combination of high falls with a high susceptibility to injury makes a relatively mild fall dangerous. This is because of a high prevalence of clinical diseases such as osteoporosis and age-related changes, that is slowed protective reflexes.⁸ In the UK, it has been estimated that falls account for 3% of total NHS service expenditure,¹⁰ with additional costs incurred by social care providers.²⁵ The prevention of falls in older people has been highlighted as a priority.^{11,26}

Although prevention appears effective,¹¹ reducing falls and associated morbidity depends on early identification of people at high risk and delivery of interventions across traditional service boundaries;²⁷ these priorities are now reflected in national and international guidelines.^{28–30}

Emergency department overcrowding and potentially avoidable attendances

Population growth, the increasing burden of chronic disease and population ageing, and the shortage of health-care workers are affecting health-care systems in many countries.³¹ EDs are under considerable pressure and overcrowding is a significant international problem, with a negative impact on both patient care and providers.³² Department of Health data³³ show that from 2003/4 to 2012/3 the number of attendances in English NHS ED units increased by nearly 32% (from 16.5 million to 21.7 million), although the majority of the increase was in minor injuries units and walk-in centres. Many of the older people presenting to EDs have had a fall.³⁴ One study showed that older people presenting to EDs after a fall represented 20% of all attendances and 14% of all hospitalisations in people aged > 65 years,³⁵ and the size of the problem is likely to be underestimated as falls are poorly defined, coded and recorded.³⁶

Older people can be kept for a longer time in the ED than younger adults because of difficulties in assessment and management, that is, they undergo more tests than younger adults, they are more likely to have cognitive limitations³⁷ and potential underlying diseases or, if they have common diseases, atypical symptoms,³⁸ and they are also more likely to be admitted than younger adults.³⁹ Pressures on acute services to discharge patients as soon as possible may mean that there is little focus on fall prevention interventions.³⁶ Older people taken to EDs following a fall are at high risk of falling again in the next year, with a 30% chance of sustaining fracture or dislocation.³⁵

It has been argued that many admissions of older people are avoidable, and that the demands placed on the hospital system may be relieved if alternatives can be found.⁴⁰ The 2006 White Paper *Our Health, Our Care, Our Say* set the direction on 'improving patient experience and significantly reducing unnecessary admissions to hospital',⁴¹ with paramedics and enhanced/senior paramedics to play a role in providing care at the scene. The National Medical Director of NHS England recently proposed key areas of change in the 2013 Keogh report *Transforming Urgent and Emergency Care Services in England 2013*.⁴² The reports look at developing the ambulance service into a mobile urgent treatment service capable of treating more patients at the scene so they do not need to be conveyed to hospital to initiate care. The report highlights opportunities to shift care closer to home, stating that 40% of ED patients are discharged requiring no treatment, up to 1 million emergency admissions were avoidable in 2012 and up to 50% of calls could be managed at the scene.

Conveyance rate of older adults who fall

Although demand on UK ambulance services continues to increase steadily, a recent study found that only 10% of calls are life-threatening, and an estimated half of all emergency calls relate to patients who could be treated at home.⁴³

People aged ≥ 65 years commonly call an emergency ambulance (through the emergency services) following a fall. In London (UK), this group accounts for about 60,000 attendances (8% of emergency ambulance attendances) each year.⁴⁴ This is very similar to the proportion reported in an urban emergency medical services (EMS) system in the USA.⁴³

Non-conveyance to EDs is high in this group, at close to 40% in London,⁴⁴ elsewhere in the UK^{45,46} and the USA.⁴³ In most cases (90%), patients who fall and are not conveyed to an ED have fallen in the home.⁴⁶ Non-conveyance of patients is recognised internationally as a safety and litigation risk.⁴⁷ In some UK ambulance services, performance targets are set for individual enhanced paramedics and paramedics regarding the percentage of patients they should leave at home, and the enhanced paramedic or paramedic decides who can be safely left at home. In 2011/12, the Department of Health, in conjunction with the College of Emergency Medicine, introduced eight accident and emergency (A&E) clinical quality indicators as part of the NHS Outcomes Framework. These became mandatory from 1 April 2011.

Indicator 1 (ambulatory care) measures the proportion of patients who are able to be treated at home by improved pathways of care for patients to avoid hospital admission.⁴⁸

Little is known about how, in the absence of specific guidelines or training to leave older people who fall at home, paramedics make these decisions. A study acknowledged the pragmatic nature of negotiation with patients whether or not to go to hospital.⁴⁹ A UK study identified that the non-clinical factors affecting these decisions include experience and confidence of ambulance staff, time into the shift, presence of carers, quality of the accommodation, waiting times at the local ED and prior knowledge of the patient.⁵⁰

Ambulance services, with key stakeholders, have been required to develop alternative care pathways, some of which involve the direct referral of older patients who have fallen to primary⁵¹ or community services.⁵² This is particularly relevant at a time when pressures on EDs give rise to safety concerns and challenges to demand management;⁵³ general practice and wider community services are being required to support the ED directly in order to avoid secondary care, in a time of major system reorganisation.⁵⁴ Alternative care pathways (avoiding use of the ED) for ambulance clinicians have been incentivised through local targets in the Commissioning for Quality and Innovation payment framework.⁵⁵ This emphasis is likely to continue given the pressure on EDs.

The National Service Framework for Older People²⁶ advocates that ambulance staff/clinicians refer to community-based care older people who have fallen, although this reflects consensus rather than research evidence. Previous studies in this setting have found that change in practice is difficult to achieve and new pathways of care are difficult to exploit.⁵⁶

Service models: community falls services

A range of different types of trial performed in community-dwelling individuals (someone who lives among the general population and is not living in an institution),⁵⁷ care home residents and hospital patients has established the evidence base for falls interventions.¹¹ Recent clinical guidelines from the American Geriatrics Society, British Geriatrics Society and American Academy of Orthopaedic Surgeons panel on falls prevention²⁹ have strongly advocated preventative approaches based on multidimensional risk factor assessment, exercise programmes (which include balance, strength and endurance training) and environmental assessment and modification. A 2004 meta-analysis demonstrated a 37% risk reduction in the monthly rate of falling for community-dwelling individuals when multifactorial intervention was implemented.⁵⁸ Falls clinics are one approach by which falls and injuries can be managed multifactorially.⁵⁹ A falls clinic is a space where older adults prone to falling can receive holistic support and treatment from nurses, physiotherapists and other health professionals. A recent randomised controlled trial (RCT) showed that the Chaos Falls Clinic in Finland is effective in preventing falls in home-dwelling people over 70 years of age at high risk of falling.⁶⁰ Patients in this trial were guided towards the falls clinic by regional health-care professionals (physicians, nurses or GPs), but relatives of patients could also contact the clinic for an assessment for eligibility.

A recent study found that referral to a community-based falls prevention of older people who had fallen and been left at home by their attending ambulance clinicians, service reduced further falls and improved clinical outcomes,⁶¹ and achieved cost-effectiveness.⁶² Recent policy changes in the UK have encouraged the development and implementation of alternative models of care for the delivery by the ambulance service. The literature points to the development of extended skills paramedics in line with government policies and supported by protocols. The use of enhanced paramedics has the potential to reduce running costs of the service considerably, while improving quality and focus, although significant 'one-off' costs associated with training are also necessary. However, Newman⁶³ noted, the 'emphasis on the national standards plus local flexibility appeared to reflect supposed shifts in the role of state' but more particularly presented a range of models which resulted in 'confusing messages about the relationship between government, the professionals and the public'.

Theoretical underpinning for intervention

Explicit definition of the theoretical underpinning for an intervention to be evaluated is a necessary step in the process for developing both the intervention to be tested as well as the methods for its evaluation. The theoretical basis for the Support and Assessment for Fall Emergency Referrals (SAFER) 2 study intervention – the protocol for emergency ambulance paramedics to assess older patients attended for a fall, and allowing referral for those not needing immediate care at ED to a falls service for community-based care (see *Appendix 1*) – was built on work carried out previously by the research team and other published research in the two areas of emergency prehospital care paramedic practice and service delivery, and care of older people who fall.

The intervention is hypothesised to work by improving the decision-making of paramedics in terms of safe non-conveyance and referral to appropriate community-based services of older people who have had a fall. Through the protocol, pathway, training, support and feedback, the intervention provided a formalised framework for decision-making and referral for this patient group. This is hypothesised to make a difference by one or more of the following four mechanisms:

- increasing paramedics' clinical knowledge of how to make appropriate non-conveyance decisions
- increasing paramedics' knowledge of falls services and pathways for referral
- increasing paramedics' confidence about making a non-conveyance decision and reducing anxiety about risk
- increasing awareness and likelihood that paramedics will consider non-conveyance and referral pathways as an option in appropriate cases.

For the intervention to make a difference to practice and to patient outcomes, a number of factors needed to be in place, including:

- effective referral pathways, and sufficient capacity in falls teams to respond to referrals
- training and support provided to paramedics that is appropriate and effective
- motivation on the part of paramedics to use the new protocol and referral pathway, including them not finding it too onerous or time-consuming.

If improved decision-making can be achieved, based on our best knowledge, we can expect the following beneficial effects:^{61,62,64–66}

- better outcomes (fewer subsequent falls and emergency episodes; higher satisfaction with care) for patients, in particular for those who would have been left at home with no further care but who are now referred to falls services, and also for those who may have been taken to the ED unnecessarily, with related risks of inpatient admission and hospital-related adverse events (AEs)
- reduced costs to the NHS and patients because of fewer initial ED episodes, inpatient admissions and subsequent falls and related care
- improved paramedic skill sets, level of professionalisation and practitioner morale.

Uncertainties lie at the certain points on the required causal pathway in terms of the feasibility of implementation and uptake by paramedics; acceptability to patients of non-conveyance/referral; improvements to processes of care for patients; of any wider impact on the operation of the ambulance service, such as increased job cycle times; and of patient health outcomes, including their perception of their own health.

We designed the SAFER 2 trial to gather data about each of these elements of the pathway in order to assess not only outcomes but also processes that may lead to improved outcomes. Study objectives and outcomes reflect the theoretical underpinning and hypothesised causal pathway.

Research aim and objectives

Aim

To assess the benefits and costs to patients and the NHS of a complex intervention comprising education, clinical protocols and pathways enabling paramedics to assess older people who have fallen and refer them to community-based falls services when appropriate.

Objectives

- To compare outcomes, processes and costs of care between intervention and control groups:
 - patient outcomes: rate and pattern of subsequent emergency health-care contacts or deaths, for any reason; health-related quality of life; psychological status, especially fear of falling; and change in place of residence
 - processes of care: pathway of care at index incident; subsequent health-care contacts; ambulance service operational indicators; and protocol compliance including clinical documentation
 - costs of care: provided by NHS and personal social services (PSS); incurred by patients or carers in seeking care.
- To estimate wider system effects of the introduction of the intervention on ambulance service performance and costs.
- To understand how patients experience the new health technology.
- To identify factors which facilitate or hinder the use of the intervention.
- To inform the development of methods for falls research especially outcome measures recommended for trials of interventions for older people who fall.⁵⁹
- To carry out a systematic review on the implementation and impact of interventions by emergency medical staff who care for older people who fall.

Chapter 2 Systematic review of literature on effects of interventions by emergency medical staff for older people who fall

Abstract

Background

Since the publication of the *National Service Framework for Older People 2001*,²⁶ the NHS has prioritised older people who fall. As the overall use of EDs increases, policy is shifting to encourage ambulances to convey fewer patients and refer them to other services, when appropriate.

Objective

To review evidence about effects of interventions within EMSs for older people who fall, which aim to reduce demand for EDs.

Method

We undertook a systematic search of 18 electronic databases. Studies were eligible for inclusion if they included empirical data on novel interventions by EMS in the community for older people who fall. We extracted outcomes, assessed studies for methodological quality and used narrative synthesis to analyse results.

Results

Of the 5355 articles identified, we selected 12 studies reported in 13 papers. Identified studies were excluded for the following main reasons: not English language, not EMS, not falls, no or entirely insufficient empirical data, no intervention or the outcomes of falls were not differentiated. The 13 papers reported data from 7411 participants: 7399 patients and 12 paramedics. Interventions fell into two groups: prospective screening on-scene or at the time of the emergency call and retrospective screening and referral to falls prevention.

Up to half of patients screened on-scene were conveyed to hospital, and referrals to falls services were low except when automatic (retrospectively). The majority of studies were judged to be of poor quality and data were heterogeneous, limiting opportunities for comparison and meta-analysis. Of the higher-quality studies, one trial reported a reduction in ED attendance and hospital admission and increased patient satisfaction; the other reported a reduction in ambulance calls over 12 months, with patients reporting fewer falls and increased confidence.

Discussion

Despite national policy to prevent falls and reduce use of emergency services, studies were few and evidence was mainly weak on effective ways to provide emergency care to older people who fall. When high-quality trial data were available, positive impact on patient and service outcomes was reported.

Study registration

This systematic review is registered on PROSPERO number CRD42013006418.

Introduction

This chapter presents a systematic review of the literature on the effects of novel interventions by EMSs for older people who fall. It describes the background and rationale for the review in the context of the

SAFER 2 study, defines the methods used and presents findings to identify existing evidence and topics for further research.

Background and rationale for the review

Since the publication of the *National Service Framework for Older People 2001*,²⁶ the NHS has prioritised older people who fall. These people have reduced quality of life and levels of physical activity, leading to social isolation and functional deterioration with high risk of increased dependency and institutionalisation.¹⁵⁻¹⁷ Treatment of falls is a major and rising cost for health systems internationally. In the UK, falls cost the NHS more than £2.3B per annum,⁶⁷ about 3% of total expenditure.¹⁰ In the USA, costs will increase by almost 75% between 2010 and 2020⁶⁸ and in Australia by more than double between 2004 and 2021.⁶⁹ Older people who fall and need emergency attention make up a substantial part of the EMS workload. In London, for example, they generate over 60,000 attendances, about 8% of the workload.^{44,51} International proportions are similar.⁷⁰ Standard paramedic practice is to assess injury and immediate care needs, move patients from where they have fallen and convey them to an ED unless they refuse.⁷¹ In the face of rising ED attendances, policy is shifting to encourage emergency ambulance services to convey fewer patients to EDs in the UK and internationally.^{40,72}

Research has highlighted opportunities for alternative treatment for older people who fall that reflect their health status and risk.^{43,46,73,74} Some older people who fall and call the emergency services can be identified in the ambulance call centre, allowing rapid and targeted alternative responses.⁷⁵ The cost of EMS attendance to older people who fall is high.⁷⁶ Innovative approaches are necessary and feasible,^{77,78} but evidence about cost-effectiveness of alternative treatment is needed.⁷⁶

A Cochrane review¹¹ of 159 randomised trials of interventions to reduce incidence of falls among older people living in the community reported that exercise interventions reduce risk and rate of falls. However, very few studies reported interventions by EMS staff.^{61,62,71,79} Gates *et al.*⁶⁵ found limited evidence supporting fall prevention programmes in EDs, primary care or the community but did not evaluate any initiatives delivered by EMS. More recently a systematic review by Mikolaizak *et al.*⁷⁴ reported that one study found that 49% of non-transported patients who had fallen had unplanned health contact within 14 days;⁴⁴ another study reported that 33% of such patients were admitted to hospital within 28 days;⁸⁰ and two studies that found that attendance by specially trained paramedics reduced subsequent falls, hospital attendance and other AEs.^{61,66,74} However, this review did not assess the quality of included studies or observe Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines,⁸¹ thus limiting interpretation of findings. None of these reviews considered issues affecting implementation of interventions for older people who fall, notably the views of EMS staff delivering care.

Methods

Protocol and registration

We registered this review with PROSPERO and the All Wales Systematic Review Register. We adhered to the guidance for reviews of health care by the Centre for Reviews and Dissemination⁸² and PRISMA.⁸¹

Eligibility criteria

We defined inclusion and exclusion criteria in terms of the population, intervention, comparator, outcomes, context and study design (PICOCs) of included studies, which is an adaption of population, intervention, comparator, outcomes (PICO). The additional 'C' in the acronym recognises the importance of the context within which the intervention is delivered.^{82,83}

We included studies that evaluated interventions by EMS staff in the community to older people who had fallen and generated an emergency service call. We considered any treatment above the care routinely provided by a paramedic or other EMS staff. *Table 1* shows our inclusion and exclusion criteria.

TABLE 1 Inclusion and exclusion criteria

PICOCS and limits	Inclusion	Exclusion
Population	<ul style="list-style-type: none"> Older people (aged ≥ 60 years) who fall at home or in the community and call for an emergency ambulance $\geq 80\%$ of participants recruited sustained a fall 	<ul style="list-style-type: none"> Patients who fall from higher than standing Patients who sustain falls in sports Patients whose fall is identified by home-based technology (e.g. motion sensors), which alerts EMSs $> 20\%$ of participants recruited sustained an injury other than a fall
Intervention	<ul style="list-style-type: none"> Enhanced practice delivered to older people attended by EMS following falls in the community – at the time of treatment or to follow up the fall 	<ul style="list-style-type: none"> Normal practice – patient is treated at home by an ambulance technician or paramedic with standard skills using standard procedures (viz. usual care within the ambulance service in which the study takes place) Referral to a falls service by ED or health-care professional Care delivered by helicopter EMSs
Comparator	No comparator required since any study design included	
Outcome	<ul style="list-style-type: none"> Referrals to other services ED conveyance Any outcome affecting patient care Subsequent falls Subsequent emergency calls for falls Costs Acceptability to patients Use of screening tool usage Views of staff on acceptability and implementation of intervention Strategies to increase referral rates 	<ul style="list-style-type: none"> Mere description of demography, geographical location of patient or presenting injuries Outcomes of decisions to transport to specialist treatment centres
Context	<ul style="list-style-type: none"> Prehospital emergency care 	<ul style="list-style-type: none"> ED Primary and community care Secondary care
Study design and reporting format	<ul style="list-style-type: none"> Full papers reporting empirical data from any study design 	<ul style="list-style-type: none"> Letters, comments, conference abstracts or opinion pieces Papers presenting no empirical data
Limits	<ul style="list-style-type: none"> English language Publications between 1990 and 2013 UK and international research 	<ul style="list-style-type: none"> Reports published before 1990

Sources and search strategy

We searched published and grey literature and reference lists of included studies. We searched the electronic databases listed in *Table 2* during the first week of July 2013 using a search strategy focusing mainly on the target population and the setting of the intervention. We used medical subject headings (MeSHs) and keywords when available. *Table 3* shows the search strategy for MEDLINE, the Allied and Complementary Medicine Database and Cumulative Index to Nursing and Allied Health Literature Plus.

Our search strategy focused on two elements of PICOCS:

1. population, using terms such as 'aged', 'health services for the aged', 'frail elderly', 'age', 'accidental falls', 'trip' and 'slip'
2. context, using terms such as 'emergency medical services', 'ambulatory care', 'ambulances', 'emergency medical technicians', 'urgent care', 'emergency care' and 'prehospital'.

TABLE 2 Sources of literature used

Type of literature	Data sources
Reviews	The Cochrane Library (Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, NHS Economic Evaluation Database, Cochrane Central Register of Controlled Trials and Cochrane Register of Methodological Reviews) Reference lists of included articles
General journal articles	AMED ASSIA BNI CINAHL Plus EMBASE Internurse MEDLINE PsycINFO PubMed Scopus Reference lists of included articles
Grey literature	The British Library UK Institutional Repository Search GreyNet Conference Proceedings Citation Index (Web of Science) OpenGrey Zetoc Reference lists of included articles
Journal citation reports	Web of Science

AMED, Allied and Complementary Medicine Database; ASSIA, Applied Social Sciences Index and Abstracts; BNI, British Nursing Index; CINAHL, Cumulative Index to Nursing and Allied Health Literature.

The terms for each database varied slightly according to MeSH or other indexed terms. The key terms were included, as well as keywords in title, abstract or text.

The search strategy used in three of the databases (MEDLINE, EMBASE and PubMed) can be summarised as follows:

- population, using the MeSH terms of 'Aged', 'Health Services for the Aged', 'Frail Elderly', 'Accidental Falls' and the keywords of 'age*', 'trip' and 'slip'
- context, using the MeSH terms of 'Emergency Medical Services', 'Ambulatory care', 'Ambulances', 'Emergency Medical Technicians' and the keywords of 'urgent care', 'emergency care', 'prehospital', 'pre-hospital' and 'emergency medical service*'
- combination of elements, combining each of the population terms and each of the context terms with 'OR'; and combining the population and the context terms with 'AND'.

TABLE 3 Search carried out in MEDLINE, the Allied and Complementary Medicine Database, and Cumulative Index to Nursing and Allied Health Literature Plus

Search identification	Search terms	Search options
		Limiters – English language
S22	S4 AND S11 AND S20	Search modes – Boolean/Phrase
S21	S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19	Search modes – Boolean/Phrase
S20	(MH 'Ambulatory Care')	Search modes – Boolean/Phrase
S19	(MH 'Ambulances')	Search modes – Boolean/Phrase
S18	(MH 'Emergency Medical Services') OR (MH 'Emergency Medical Technicians')	Search modes – Boolean/Phrase
S17	urgent care	Search modes – Boolean/Phrase
S16	emergency care	Search modes – Boolean/Phrase
S15	pre hospital	Search modes – Boolean/Phrase
S14	pre-hospital	Search modes – Boolean/Phrase
S13	ambulance	Search modes – Boolean/Phrase
S12	S6 OR S7 OR S8 OR S9 OR S10	Search modes – Boolean/Phrase
S11	age*	Search modes – Boolean/Phrase
S10	(MH 'Aged')	Search modes – Boolean/Phrase
S9	(MH 'Health Services for the Aged')	Search modes – Boolean/Phrase
S8	(MH 'Frail Elderly')	Search modes – Boolean/Phrase
S7	old*	Search modes – Boolean/Phrase
S6	elder*	Search modes – Boolean/Phrase
S5	S1 OR S2 OR S3	Search modes – Boolean/Phrase
S4	(MH 'Accidental Falls')	Search modes – Boolean/Phrase
S3	trip*	Search modes – Boolean/Phrase
S2	slip*	Search modes – Boolean/Phrase
S1	fall*	Search modes – Boolean/Phrase

Study selection

We downloaded electronic search results into the EndNote bibliographic software (Thomson Reuters, CA, USA). One researcher (RC, MH or MK) screened the title, abstract and keywords of each study using the inclusion and exclusion criteria in *Table 1*. A second researcher, blind to the first researcher's decision, screened 1 in 10 references, to ensure consistent decision-making. We resolved differences in discussion with the third researcher. We retrieved the full text of papers whose abstract met the inclusion criteria. We compared final recommendations and again achieved consensus on eligible studies in discussion with the third researcher.

Data extraction

We developed and piloted a data extraction framework reflecting the objectives of the review in line with guidance.⁸² Two researchers independently extracted the following items:

- year of publication, country of origin, setting, study design
- aim or research question, outcome measure(s), method(s), study population, sample size
- description of intervention
- key findings.

When possible, we used the taxonomy developed by the Prevention of Falls Network Europe (ProFaNE),⁸⁴ from the work of Hauer *et al.*,⁸⁵ to describe these data, thus increasing consistency of extraction across the included studies and opportunity for synthesis. The full data extraction table can be found in *Appendix 2*. We compared completed forms and agreed on content in discussion with the third researcher.

Assessment of the risk of bias (quality)

We assessed quality of included studies using checklists recommended by Centre for Reviews and Dissemination⁸² to inform comparisons and guide interpretation.⁸⁵ We identified these checklists as potentially suitable in the Centre for Reviews and Dissemination guidance. We used the Scottish Intercollegiate Guidelines Network checklist⁸⁶ for all study methods except qualitative reports, for which we used the summary criteria of Walsh and Downe.⁸⁷ We assessed general methodological quality as high, acceptable or low, defined by Scottish Intercollegiate Guidelines Network⁸⁶ thus:

- high: almost all criteria met; low risk of bias; conclusions unlikely to change after further research
- acceptable: most criteria met; some flaws with an associated risk of bias; conclusions may change after further research
- low: either most criteria not met or clear flaws in study design; conclusions likely to change after further research.

Data synthesis

We assessed quantitative results for their potential to contribute to a meta-analysis. We used narrative synthesis to synthesise heterogeneous data.⁸⁸ We explored relationships within and between studies, seeking to explain similarities and differences between study findings.

Findings

Inclusion

We identified a total of 5355 references. Our review of titles and abstracts suggested that 138 articles might be eligible. After studying full texts, we confirmed that 13 papers^{61,62,66,80,89-97} reporting 12 studies met our inclusion criteria, as detailed in *Figure 1*. *Table 4* summarises the 12 included studies; *Tables 5* and *6* describe them in more detail and *Table 7* assesses their quality. *Appendix 3* lists excluded studies (at full-text screening stage) with reasons for exclusion.

Study characteristics

Of the 12 studies, six^{61,62,66,80,89,94} came from the UK, four⁹⁰⁻⁹³ came from the USA, one⁹⁷ came from Australia and one⁹⁵ came from New Zealand. Study designs comprised two RCTs^{61,66} (one of which⁶¹ published a separate paper evaluating cost-effectiveness element of the study),⁶² nine cohort studies (of which six did not include a comparator arm^{89,91-93,95,97} and three included intervention and control arms^{80,90,96}) and one qualitative study.⁹⁴ Of the nine cohort studies, three were retrospective,^{92,95,96} five were prospective^{80,89-91,97} and one had both prospective (screening) and retrospective (referral) elements.⁹³ These study characteristics are summarised in *Table 4*.

The one-armed nature of six^{89,91-93,95,97} of the studies restricted them to reporting descriptive outcomes. All but Logan *et al.*⁶¹ reported outcomes at the time of the intervention or soon after. The most common outcome measures were referrals made to a receiving unit or service,^{66,89,91-93,97} acceptance of referral by the patient^{66,89,91,92,97} and patient conveyance following attendance by EMSs.^{66,80,95} Five studies reported follow-up outcomes for patients receiving a novel intervention (see *Table 6*).^{61,66,80,90,97} Three^{66,80,90} used short follow-up periods (14 and 28 days) with limited inclusion of outcome measures in two,^{80,90} whereas another collected physical health, quality-of-life and mortality outcomes.⁶⁶ One written as two research papers reported outcomes over 12 months;^{61,62} but one did not report the period of follow-up.⁹⁷




FIGURE 1 The PRISMA flow chart.

Ten^{61,66,80,89,90,92,93,95-97} of these studies included a total of 7399 patients and the qualitative study interviewed 12 paramedics.⁹⁴ However, one study did not specify its sample size.⁹¹

Methodological quality

The methodological quality of the studies was variable, but was generally weak (see *Table 7*). The controlled trials,^{61,66,90} economic evaluation⁶² and qualitative study⁹⁴ were considered acceptable or high quality and scored highly for internal validity. The cohort studies were mostly of weak quality and did not report sufficient information to determine if their methods minimised risk of bias or confounding. Seven out of eight of the cohort studies^{80,89,90,92,93,96,97} failed to identify potential confounders in study design and analysis. Shah *et al.*⁹³ and Metcalfe⁸⁹ did not provide a focused research question. Gray and Walker⁸⁰ scored fairly low using the quality assessment tool, but reviewers defined this study as of acceptable quality as, compared with the other studies included in this review, it included a control arm, a clearly focused aim and clearly written methods and results sections. Kue *et al.*⁹² did not define outcomes or reliably assess exposure. However, reviewers considered all studies that reported data that were directly relevant to this review, and all were included.⁹⁸ The full quality assessment for each included study is presented in *Appendix 2*, with a brief summary of the internal validity score, risk of bias and noted limitations given in *Table 7*.

Study findings

Interventions within ambulance services to treat older people who fall

The 12 evaluated interventions fell broadly into two main types; 10 encouraged EMS staff on-scene to adopt a new approach when making their initial assessment of the older person who had

TABLE 4 Summary of study characteristics

First author	Year of publication	Country	Study design	Study population	Sample size	Description of intervention
Metcalfe ⁸⁹	2006	UK	Cohort	Patients aged ≥ 65 years, but not transported	49	Provision of clinical assessment tool to paramedics with follow-up referral to rapid response team
Shah et al. ⁹⁰	2006	USA	Cohort (with control group)	Patients aged ≥ 65 years	258	Screening tool used by paramedics
Mason et al. ⁶⁶	2007	UK	C-RCT	Individuals aged > 60 years calling the emergency services and presenting with minor acute conditions in the scope of practice of the paramedic practitioner	3018	Assessment by a paramedic with an enhanced scope of practice in on-scene assessment, treatment and referral (paramedic practitioner)
Shandro et al. ⁹¹	2007	USA	Cohort	Patients aged ≥ 65 years, but not transported	Not reported	Paramedic referral to falls prevention
Gray and Walker ⁸⁰	2008	UK	Cohort (with control group)	Patients aged ≥ 65 years	233 attended by ECPs; 772 attended by ED	Provision of care by ECP (paramedic with further training)
Kue et al. ⁹²	2009	USA	Retrospective cohort	Patients aged ≥ 60 years, but not transported	721	Paramedic referral to social services
^a Logan et al. ⁶¹	2010	UK	RCT	Patients aged ≥ 60 years living in defined area, but not transported	204	Retrospective referral to falls prevention team following attendance by paramedic
^a Sach et al. ⁶²	2012	UK	Cost-effectiveness	Patients aged ≥ 60 years and not transported	157	Retrospective referral to falls prevention service following attendance by paramedic
Shah et al. ⁹³	2010	USA	Cohort	Patients aged ≥ 60 years	814	Provision of screening question to paramedics; follow-up referrals
Halter et al. ⁹⁴	2011	UK	Qualitative	12 ambulance staff who were trained in, and used the, clinical assessment tool	12 paramedics	Provision of clinical assessment tool to paramedics
Hoyle et al. ⁹⁵	2012	New Zealand	Retrospective cohort	Average age 62 years; but not all had suffered from falls	131	Provision of treatment protocols to ECPs
Studnek et al. ⁹⁶	2012	USA	Retrospective cohort (with control group)	911 callers to whom medical priority dispatch system sent ambulance	Phase 1, n = 160; phase 2, n = 101	Triage led by despatch nurse
Comans et al. ⁹⁷	2013	Australia	Cohort	Patients aged ≥ 65 years	21	Paramedic referral to falls prevention

ECP, emergency care practitioner.
 a These papers report the same study.

TABLE 5 Results from studies assessing outcomes at the time of the incident

Study	Number of patients screened using tool/total number of patients	Number of referrals/number of patients screened for referral (%)	Number of referrals accepted/number of patients referred (%)	Number conveyed to ED/number screened (%)	Percentage admitted to hospital	Paramedics views of the intervention	Other outcomes
Metcalfe, 2006 ⁸⁹	–	53/89 (60%)	51/53 (96.2%)	–	–	–	–
Shah <i>et al.</i> , 2006 ⁹⁰	210/258 (79%, 95% CI 74% to 84%)	–	–	–	–	–	–
Mason <i>et al.</i> , 2007 ⁸⁶	–	–	–	ED attendance 0 to 28 days: intervention group, 62.6%; control group, 87.5% ($p < 0.001$)	Admittance to hospital 0 to 28 days: intervention group, 40.4%; control group, 46.4% ($p < 0.001$)	n/a	Total episode time: intervention group, 235 minutes; control group, 278 minutes ($p < 0.001$). Investigations received: intervention group, 49.7%; control group, 67.9% ($p < 0.001$). Treatment received: intervention group, 81.3%; control, group 72.8% ($p < 0.001$)
Shandro <i>et al.</i> , 2007 ⁹¹	–	17 (number screened not reported)	11/17 (64.7%)	–	–	–	–
Gray and Walker, 2008 ⁸⁰	–	–	–	62/233 (27%)	25% reduction in intervention group	–	–
Kue <i>et al.</i> , 2009 ⁹²	–	7/721 (1%)	–	–	–	–	–
Shah <i>et al.</i> , 2010 ⁹³	814 (total number not reported)	124/814 (15%)	–	–	–	–	–
Halter <i>et al.</i> , 2011 ⁹⁴	–	–	–	–	–	Paramedics: reluctant to use novel intervention	–
Hoyle <i>et al.</i> , 2012 ⁹⁵	–	–	–	44/131 (34%)	–	–	–
Studnek <i>et al.</i> , 2012 ⁹⁶	–	–	–	–	9.3% reduction in intervention group	–	–
Comans <i>et al.</i> , 2013 ⁹⁷	–	21/638 (3%)	13/17 (76.5%)	–	–	–	–

CI, confidence interval; n/a, not applicable.

TABLE 6 Results from studies assessing outcomes after the incident

Study	Further falls	EMS calls	Fear of falling	Costs and QALYs	Admission to hospital (outcome measure), (p-value)	Other outcomes (outcome measure: results)
Shah <i>et al.</i> , 2006 ³⁰	–	–	–	–	–	Discussion with physician to discuss risk of falls: ^a intervention, n = 8/82 (9.8%); control, n = 1/37 (2.7%); p = 0.30 Receipt of changes to home environment: ^a intervention, n = 12/82 (15%); control, n = 4/37 (11%); p = 0.91 Recollection of educational materials: n = 17/82 (21%); control not applicable
Mason <i>et al.</i> , 2007 ⁶⁶	–	–	–	–	See Table 5	Subsequent unplanned contact with secondary care services (up to 28 days): intervention, 21.3%; control, 17.6% (p < 0.001) Worsening of physical health: intervention, 21.7%; control, 25.6%; p = 0.13 EQ-5D: no significant difference (p-value, NR) Mortality: no significant difference (p-value, NR) Patient satisfaction: very satisfied 85.5% intervention group vs. 73.8% control group; p < 0.001

Study	Further falls	EMS calls	Fear of falling	Costs and QALYs	Admission to hospital (outcome measure), (p-value)	Other outcomes (outcome measure: results)
Gray and Walker, 2008 ⁸⁰	–	–	–	–	Avoided admission rate: intervention group, 44%; control group, 52%; $n = 1005$; $p = 0.05$. 17% reduction at 28 days	–
Logan <i>et al.</i> , 2010; ⁶¹ Sach <i>et al.</i> , 2012 ⁶²	55% reduction in intervention group (95% CI 42% to 65%; $p < 0.001$)	40% reduction in intervention group, 95% CI 8% to 60%; $p = 0.018$	22% reduction in intervention group (95% CI 13% to 31%; $p < 0.001$)	Mean NHS and PSS costs: –£1551.00 less in intervention group (95% CI –£5932.00 less to £2829.00 more) Mean QALYs: 0.070 more in the intervention group (95% CI –0.010 to 0.150; $p = 0.086$)	Number of hospital days: intervention group, 1257; control group, 1141; $p = 0.70$	Nottingham extended activities of daily living scale medians: intervention group, $n = 8$; control group, $n = 6$; $p < 0.001$
Comans <i>et al.</i> , 2013 ⁹⁷	–	–	–	–	–	Patients completing referral programme: 5/8 (62.5%)

CI, confidence interval; EQ-5D, European Quality of Health-5 Dimensions; NR, not reported; QALY, quality-adjusted life-year.
a Intended outcomes of the intervention.

TABLE 7 Summary of quality of included studies

Study	Internal validity score/total possible score (n n/a)	How well was the study done to minimise bias?	Particular limitations noted during quality appraisal
Controlled trials			
Shah <i>et al.</i> , 2006 ⁹⁰	4/5 (5 n/a)	Moderate quality	Potential bias from choosing intervention cluster from two without randomisation; confounding not controlled for
Mason <i>et al.</i> , 2007 ⁶⁶	8/9 (1 n/a)	High quality	Participants could not be blinded; as some outcomes were self-reported, there was potential for reporting bias; high level of attrition (50%)
^a Logan <i>et al.</i> , 2010 ⁶¹	7/9 (1 n/a)	High quality	Participants could not be blinded
^a Sach <i>et al.</i> , 2012 ⁶²	7/8 (1 n/a)	High quality	As outcomes were self-reported, there was potential for reporting bias
Qualitative study			
Halter <i>et al.</i> , 2011 ⁹⁴	10/12 (0 n/a)	High quality	Lacks detail about ethics, researchers and analysis
Cohort studies			
Kue <i>et al.</i> , 2009 ⁹²	3/7 (7 n/a)	Low quality	Data not presented clearly; limited by small study numbers and being in one site
Comans <i>et al.</i> , 2013 ⁹⁷	5/11 (3 n/a)	Low quality	Very high attrition (75%); limited data used as the basis for evaluation
Hoyle <i>et al.</i> , 2012 ⁹⁵	3/6 (8 n/a)	Low quality	Subjective assessment and possibly bias in dispatch, disparity in transportation rates; uncontrolled evaluation with limited follow-up data
Gray and Walker, 2008 ⁸⁰	3/10 (4 n/a)	Low quality	Nominally controlled but collected different data from intervention and control groups, limiting comparison
Studnek <i>et al.</i> , 2012 ⁹⁶	4/8 (6 n/a)	Low quality	Nominally controlled but study sites changed between phases without relevant comparator/baseline data; very limited in terms of outcomes measured
Shandro <i>et al.</i> , 2007 ⁹¹	4/8 (6 n/a)	Low quality	Uncontrolled evaluation; very limited presentation of EMS-specific data and outcomes of care; number of exposed patients not stated
Shah <i>et al.</i> , 2010 ⁹³	2/9 (5 n/a)	Low quality	Feasibility study without controls or patient outcome measures
Metcalfe, 2006 ⁸⁹	2/9 (5 n/a)	Low quality	Uncontrolled evaluation with poorly reported methods; authors' conclusions do not match results
n/a, not applicable. a These papers report the same study.			

fallen;^{66,80,89-92,94,95,97} two did so retrospectively^{61,96} and one study⁹³ did both through intervention on-scene by EMS staff and retrospective action by other staff. Of the others in the first group, eight used paramedics/paramedic practitioners to intervene at the time of attendance^{66,80,90-92,94,95,97} whereas the other did so within the despatch centre, also known as ambulance control.⁹⁶ The two other studies used individuals not participating in the initial EMS care retrospectively to screen patients for referral.^{61,89}

Seven interventions screened patients on-scene with a predefined tool to decide whether or not to make a referral with the option of referring to a community-based service,^{89-94,97} one used the expertise of health professionals other than paramedics to decide upon conveyance,⁹⁶ three used paramedic practitioners trained to provide community-based clinical assessment for patients aged > 60 years,^{66,80,95} and one referred all eligible patients who were not conveyed following attendance by EMSs.⁶¹ Screening tools ranged from a single question about the patient's risk of falling,⁹⁰ through several questions about eligibility for referral^{90-93,97} to a flow chart to guide the EMS staff through decision-making.⁹⁴ One study screened the patients off-scene, with a nurse who retrospectively screened ambulance service records to decide whether or not to refer patients to a community service.⁸⁹ Once the patient was successfully screened and a decision was made to refer the patient, the studies referred them to a falls prevention service,^{61,91,97} social services,^{66,92} the patient's GP,^{66,90,94} district nurse,⁶⁶ a rapid response service⁸⁹ or a case management service.⁹³

Of the four^{66,80,95,96} studies not using predefined screening tools, but using the expertise of paramedic practitioners or health professionals other than paramedics, three assessed patients to decide whether to convey patients to hospital or to leave them at home,^{66,80,95} Mason *et al.*⁶⁶ used paramedic practitioners, and Gray and Walker⁸⁰ and Hoyle *et al.*⁹⁵ both used emergency care practitioners (ECPs) on-scene, paramedics with extended skills but without specific screening questions or falls service referral routes. Studnek *et al.*⁹⁶ transferred patients with medical problems deemed of 'low acuity' to a nurse-led advice line while an ambulance was travelling to their location. The nurse provided advice on alternative modes of care or transportation. The ambulance response was stopped only if the patient requested it.

Effect of interventions by emergency medical services for older people who fall

The outcomes measures assessed by the included studies were heterogeneous. Only a minority of the studies followed up patient outcomes: five^{89,91-93,95} collected outcomes only from the initial contact with the patient, relating to the emergency service call itself or the ambulance service's care of the patient on-scene; the remaining five reported outcomes after initial contact with the patient; and four^{66,80,90,97} studies reported both initial and subsequent outcomes. Among those reporting later outcomes, follow-up intervals were varied. Comans *et al.*⁹⁷ did not report this interval; two studies used periods of ≤ 1 month^{80,90} and the randomised trial assessed effectiveness and cost-effectiveness over 1 year.^{61,62} To try to synthesise the effect of the novel interventions on patients, we stratify the findings of the included studies by the time point at which the outcome was collected.

Outcomes pertaining to initial patient contact

Referrals

The most widely reported outcome measure was numbers of patient referrals to a community service following screening prompted by a fall, as reported by five studies^{89,91-93,97} (Table 8). Referral rates were generally low (between 1% and 15% of screened patients). However, Metcalfe⁸⁹ reported that 53 (60%) out of 89 patients screened by an experienced nurse within 24 hours of their fall were referred to a rapid response team or another appropriate service.

Three studies^{89,91,97} reported numbers of patients accepting referrals (Table 9). The acceptance rate was high, reported by Comans *et al.*⁹⁷ and Shandro *et al.*⁹¹ to be 76% and 65%, respectively. Metcalfe⁸⁹ reported that just two patients (3.8%) refused the referral generated by the rapid response team, giving a 96% acceptance rate. Although we have presented these data together, it should be noted that these studies measured referral acceptance at different time points in the referral process, with Comans *et al.*⁹⁷ reporting patients who consented to the study and underwent initial assessments, Shandro *et al.*⁹¹ assessing the number of patients enrolling on the programme, and Metcalfe⁸⁹ assessing the number of patients who accepted referrals.

Referral mechanisms that required on-scene referrals by EMS staff^{91,92,97} were undertaken less frequently and had lower patient acceptance rate than retrospective referrals, which had high patient acceptance.⁸⁹ As all these studies were of low quality, however, findings are not robust.

TABLE 8 Studies reporting number of referrals made to a community service

Referral component	Study				
	Kue <i>et al.</i> , 2009 ⁹²	Comans <i>et al.</i> , 2013 ⁹⁷	Shandro <i>et al.</i> , 2007 ⁹¹	Shah <i>et al.</i> , 2010 ⁹³	Metcalfe, 2006 ⁸⁹
Intervention	Referral pathway from EMS to community service				
Outcome	Number of referrals	Number attempted to refer	Number of referrals	Number of patients referred because of falls	Number of referrals made for patients experiencing a fall
Number of referrals	7	21	17	124	53
Sample size (patients screened for referral)	721	638	Unknown	814	89
Percentage referred	1	3	Unknown	15	60

TABLE 9 Studies reporting patient acceptance of referrals to a community service

Referral component	Study		
	Comans <i>et al.</i> , 2013 ⁹⁷	Shandro <i>et al.</i> , 2007 ⁹¹	Metcalfe, 2006 ⁸⁹
Intervention	Referral pathway from EMS to community service		Provision of screening tool with referral to community service
Outcome	Consented to study and underwent initial assessment	Number of enrolments on falls prevention programme from EMS referrals	Number patients accepting referrals
Data	13	11	51
Sample size (number of patients referred)	17	17	53
Percentage consented/accepted	76.5	64.7	96.2

Conveyance to emergency department

In three studies,^{66,80,95} ECPs or PPs (paramedics with enhanced training) took the decision to convey a patient to ED. ECPs or PPs did not use any additional decision-making tool above clinical judgement. Conveyance to the ED can only be reported here for two of the studies^{80,95} as the third study combined reporting of conveyance at the time (0 days) with conveyance at 28 days⁶⁶ and was varied (*Table 10*).

Admission to hospital

Two studies^{80,96} assessed whether or not patients who received a novel EMS intervention after making the emergency service call experienced reduced hospital admissions. Studnek *et al.*⁹⁶ found that hospital admissions fell from 35% ($n = 56$) to 25.7% ($n = 26$) when emergency service callers who had sustained a fall were screened to receive nurse-led telephone advice, in addition to a standard EMS attendance. Studnek *et al.*⁹⁶ did not calculate individual significant values for patients who had fallen, but found that, for all patients (with all possible medical complaints), there was a statistically significant difference in the proportion of patients discharged home following attendance at the ED (i.e. not admitted to hospital) compared with the control period ($p = 0.03$; CIs not reported). Gray and Walker⁸⁰ assessed hospital admittance at 72 hours post EMS attendance. Admission rates decreased from 51% ($n = 396$) during the control period to 26% ($n = 62$) during the intervention period (significance levels not reported).

TABLE 10 Studies reporting number of patients conveyed to the ED as an outcome

Referral component	Study	
	Hoyle <i>et al.</i> , 2012 ⁹⁵	Gray and Walker, 2008 ⁸⁰
Intervention	Paramedics with extended training (with no referral pathway)	
Outcome	Percentage of patients transported to ED	Patients attending ED
Data	44	62
Sample size (number of patients screened/attended)	131	233
Percentage of patients transported to ED	34	27

Outcomes collected following initial contact with patients

Five studies, written as six research papers,^{61,62,66,80,90,97} reported follow-up outcomes. Five reported the impact on subsequent falls and health service utilisation^{61,66,80,90,97} and one reported the impact on costs to the NHS.⁶²

Subsequent falls and use of health care

Logan *et al.*⁶¹ reported that patients referred to a falls prevention team experienced 55% fewer falls over 12 months than the control group [95% confidence interval (CI) 0.35 to 0.58; $p < 0.001$]. Patients also had reduced fear of falling (assessed by the falls efficacy scale, mean difference -16.5 , 95% CI -23.2 to -9.8 ; $p < 0.001$), were more active [assessed by the Barthel Activities of Daily Living Index, odds ratio (OR) 2.91, 95% CI 1.18 to 7.20]. There was also a significant reduction in emergency ambulance calls for falls over 12 months (effect size 0.60, 95% CI 0.40 to 0.92; $p = 0.018$).

Mason *et al.*⁶⁶ reported several patient outcomes with a significant difference for the intervention group (those attended by a paramedic practitioner): a 25% decrease in attendance at an emergency department at 0 and 28 days following the incident (relative risk 0.72, 95% CI 0.68 to 0.75; $p < 0.001$), a 6.1% decrease in hospital admission during the same period (relative risk 0.87, 95% CI 0.81 to 0.94; $p < 0.001$), an increase in subsequent unplanned contact with secondary care services (relative risk 0.121, 95% CI 1.06 to 1.38; $p < 0.001$), and an 11.7% increase in those reporting being 'very satisfied with their care' (relative risk 1.16, 95% CI 1.09 to 1.23; $p < 0.001$).

The other three studies^{80,90,97} reporting follow-up outcomes were cohort studies collecting outcome measures associated with specific aspects of the novel intervention. Shah *et al.*⁹⁰ assessed whether or not patients had discussed their risk of falls with their doctor and if they had received changes to their home environment. Although increased numbers of patients had these attributes in the intervention arm, the difference was not significant.

Gray and Walker⁸⁰ reported a 17% reduction in admissions at 28 days following EMS attendance during the intervention period than in the control period when patients were attended by paramedics with usual training (reduction from 52% to 44%, $n = 1005$, $df = 1$; $p = 0.05$).

Comans *et al.*⁹⁷ reported the number of patients who completed the referral programme, with five patients who were referred to a falls prevention service completing the programme, out of the eight patients who initially enrolled on the programme.

Effect on cost

One paper⁶² found that the referral of patients who had been attended and left at home by EMS to a falls prevention service, carried out within a randomised trial,⁶⁰ was cost-effective. The mean difference in NHS and PSS costs between the intervention and control groups was $-\pounds 1551.00$ per patient over 1 year (95% CI $-\pounds 5932.00$ to $\pounds 2829.00$). The mean difference in quality-adjusted life-years (QALYs) was 0.070 (95% CI -0.010 to 0.150) in favour of the intervention group.

What is the impact on and views of emergency medical services staff of interventions designed to improve the care of and outcomes for older people who fall?

Staff appeared reluctant to use new interventions to treat older people who fall, according to two studies.^{94,97} Halter *et al.*⁹⁴ reported variable use of a clinical assessment tool to review whether to safely leave a patient at home or convey him or her to the ED. This qualitative study reported that paramedics were reluctant to use a systematic decision-making tool but instead relied on informal processes, showing the need for a consistent message and support for ambulance staff. Comans *et al.*⁹⁷ identified that repeated education sessions, good relations and regular contact between falls services and paramedics raised referral rates, but these were not sustained when awareness-raising ended.

Discussion

This systematic review identified limited and mainly weak evidence regarding the most effective ways to deliver alternative treatments to older people who fall. The 12 studies that were included in this review implemented variable novel interventions, limiting comparisons that can be made between studies. We classified the interventions within two groups: (1) on-scene screening during patient assessment or triage during the emergency service call to refer them to a community health service or hospital conveyance; or (2) retrospective screening and referral to a community health service.

The studies included in this review are heterogeneous in terms of interventions and outcomes, and, while this is a finding in itself, we have considered the studies to be similar enough, in a small research field, to analyse as a group because of the similarity of the context and patient group included in all the studies. We have not attempted meta-analysis for this reason, and it is the heterogeneity of outcome measures, and a lack of follow-up outcomes to assess the long-term effect of the novel interventions on patients, that means it is difficult to assess the impact of these interventions. The majority of outcome measures were descriptive, detailing the number of referrals that were made and the acceptance of these referrals to patients. Studies utilising these descriptive outcome measures identified that a low percentage of the overall number of screened patients were referred, except where the referral was made retrospectively externally to the ambulance service. Similarly, acceptance of the referral by patients increased when the referral was undertaken externally to the ambulance service. Although these outcome measures are useful for understanding the paramedics' and patients' adoption of the intervention, they do not allow understanding of the effect of the novel intervention on the patient.

Five studies, written as six research papers, undertook follow-up of patients beyond the initial contact with EMS,^{61,66,80,90,96} five of which undertook between-group analyses. They identified that, compared with the control groups, patients who were treated for a fall by EMS had fewer future falls, called the EMS less often and received cost-effective care when referred to a fall prevention service;⁶¹ were less likely to be admitted to hospital when being treated by a paramedic practitioner⁶⁶ or ECP;⁸⁰ and were more likely to discuss fall risk with their doctor when a referral was made.⁹⁰ The study not undertaking between-group analyses found that five out of eight patients who enrolled in a fall prevention programme completed it.⁹⁷

Emergency medical services staff's views of the novel interventions were seldom reported; the one qualitative study included in this review concluded that EMS staff were reluctant to use formalised assessment techniques, instead relying on their usual decision-making process.⁹⁴ One study did not directly report paramedics' views, but did suggest that use of the intervention increased with regular contact and good relations.⁹⁷ Although other included studies did not directly assess EMS staffs views of the novel interventions, the studies that reported referral rates reported a low percentage of referrals taking place. These findings highlight the difficulty of changing practice with EMS.

Assessment of study quality, using recommended tools, identified that the included papers were mostly poorly reported and overall quality of the evidence was not high. Most of the studies were single-arm cohort studies and many had methodological drawbacks leaving them open to bias. The majority of the

included studies were single-group cohorts and only one RCT was identified. Seven of the included papers were deemed to be of low quality, two of acceptable quality and three of high quality.

Previous systematic reviews undertaken to assess the effect of referral mechanisms on patients who fall have found varying level of benefits for patients. A review looking to assess the effect of multidisciplinary falls prevention service treatment on patients referred from community settings found limited evidence to support a reduction in falls or fall-related injuries for referred patients.⁶⁵ Another review, undertaken to identify evidence regarding patients who are not conveyed following a fall, found that appropriate interventions can benefit patients who have fallen, including reducing AE and EMS calls.⁷⁴

The current review identified benefits within a minority of the included studies – one study reported significant decreases to future EMS attendances and hospital admissions when patients are referred retrospectively to a falls prevention service, while two studies described a reduction in admission to hospitals when patients are attended by an ECP. These positive findings on patient and service outcomes are, however, set in the context of the number of studies for which such outcomes are not explored and the low quality of the majority of the included studies.

High-quality evidence from well-designed and rigorously conducted research is needed to understand how effective services can be adopted and delivered which will provide appropriate and safe care for patients. Future research should focus on the ability of paramedics to make decisions regarding a patients' conveyance, paramedics' views of the intervention and the long-term health consequences for patients of such interventions.

Strengths and limitations of this review

We used a systematic process in line with good practice^{82,99} to identify international research and review reported data. In undertaking this review, we were unable either to consider papers in languages other than English or to request original data from study authors because of time constraints.

Heterogeneity of interventions and outcomes and low study quality mean data should be interpreted with caution. It also limits ability to generalise findings.

The quality assessment tools that were utilised were not tailored to single-group cohort studies. Therefore, many of the questions were not relevant to the majority of studies that were included in this review. Nevertheless, the inability to answer a question was regarded as a sign of low quality.

Review findings as context for Support and Assessment for Fall Emergency Referrals 2

The SAFER 2 trial selected the age of 65 years and older as its cut-off point for inclusion, as both the *National Service Framework for Older People 2001*²⁶ and the National Institute for Health and Care Excellence (NICE) guidelines for falls⁶⁷ use this cut-off point, although they recognise this as an arbitrary point that varies in relevance to individual health and disability.

In the systematic review, however, we chose to also include studies that stated their lower age for inclusion to be 60 years, as we were aware from our reading on the topic that two^{61,66} UK trials had used this slightly lower age breakpoint.

Although we recognise that this might suggest that the cut-off point of ≥ 60 years of age could or should have been used for the SAFER 2 trial, we are reassured that our results are comparable when we look at the mean age of participants in the 10 of the 12 studies that provide us with this demographic information about participants.^{61,66,89-93,95-97} 7 of these 10 studies report mean ages of participants in the 'frail older people' group, ranging from 77 to 83 years^{61,66,90-93,97} and one of the most common age group of participants as 80-89 years.⁸⁹ The two studies of exception were those where fall was one condition only amongst a range studied, and these reported average ages of 47⁹⁶ and 62 years.⁹⁵ We therefore feel

confident that the different lower age parameter for the systematic review and the trial did not impact on relevance of one to the other.

We suggest that the fact that this review found limited evidence on the topic of support and assessment for older people who fall and are attended by the emergency ambulance service, and even more limited evidence of high quality, is the context and justification for the SAFER 2 trial, the findings of which are now reported.

Chapter 3 Methods

Overview

The SAFER 2 study design comprised a C-RCT, with economic evaluation and qualitative component.

Aim

To assess the benefits and costs to patients and the NHS of a complex intervention comprising education, clinical protocols and pathways enabling paramedics to assess older people who have fallen and refer them to community-based falls services when appropriate.

Objectives

- To compare outcomes, processes and costs of care between intervention and control groups:
 - patient outcomes: rate and pattern of subsequent emergency health-care contacts or deaths; health-related quality of life; falls efficacy (fear of falling); and change in place of residence
 - processes of care: pathway of care at index incident; subsequent health-care contacts; ambulance service operational indicators and protocol compliance including clinical documentation
 - costs of care.
- To understand how patients experience the new health technology.
- To identify factors which facilitate or hinder the use of the intervention.
- To inform the development of methods for falls research.

Cluster randomised controlled trial

Trial design

The trial was conducted in geographically defined sites within three UK ambulance services. The unit of clustering at each site was an ambulance station. The clusters comprised the paramedics based at eligible stations who volunteered to participate in the trial. Once stations had been randomly allocated to intervention or control groups, participating paramedics based at those stations were asked to deliver care according to their group allocation, irrespective of their location on any particular shift.

We chose a cluster design, as opposed to a patient-level RCT, as the intervention included training and new assessment skills, and it would be impossible to switch on and off as would be required in a patient-level RCT. Other units of randomisation have been used in trials based in prehospital emergency care, with limited success.¹⁰⁰ We chose to randomly allocate stations in order to allow the hosting ambulance services to support change in practice for paramedics based at intervention stations while minimising contamination to practice by paramedics based at control stations.

Trial management

Following the Medical Research Council guidelines for good practice in clinical trials,¹⁰¹ the management structure for the trial (see *Appendix 4*) included an independent Trial Steering Committee (TSC) and Data Monitoring and Ethics Committee (DMEC), with an internal Trial Management Group (TMG), local implementation team (LIT) in each area, core research team, and task and finish groups for specific aspects of the trial such as data management. The TSC provided external oversight and advice to the chief investigator, the Health Technology Assessment (HTA) programme and the sponsor on all aspects. The DMEC had access to unblinded comparative data to monitor the data and make recommendations to

the TSC if there were ethical or safety reasons why the trial should not continue. The TMG managed the trial at an overarching level. The LITs dealt with operational issues at each site and provided liaison opportunities between participating services. The core team that managed the trial at a day-to-day operational level was smaller and included the chief investigator, site principal investigators and researchers.

Setting

We undertook the trial in prehospital emergency care, with paramedics delivering the intervention in partnership with community-based falls services (known as falls prevention services). We selected sites within three ambulance services in England and Wales in which a falls service was available, but no process was in place for paramedics to make direct referrals from the scene of emergency service call attendances. *Appendix 5* maps stations and hospitals with ED departments at each site.

Participants

Ambulance stations were eligible for inclusion in the study as a cluster if they were situated within a participating ambulance service and were a base for paramedics who regularly attended patients within a participating falls prevention service's catchment area.

We invited paramedics based at selected ambulance stations to participate in the trial before allocating those stations randomly between intervention and control groups. Paramedics were approached via e-mail, letter and station posters and were asked to return a reply slip if they were interested in participating. Paramedics who volunteered to participate were given a £50.00 voucher.

Patients received either intervention or control group care, depending on the group allocation of the attending paramedic. During training, intervention paramedics were instructed to lead on care if there was also a control group paramedic on-scene, and these patients were included in the intervention group for analysis.

Inclusion criteria:

- aged ≥ 65 years
- resident in the catchment area of participating falls services
- attended by a study paramedic following an emergency call to the ambulance service which was coded by a dispatcher as a fall without priority symptoms (Advanced Medical Priority Dispatch System Code 17)
- first eligible call during the trial period.

Randomisation

Randomisation of stations to groups was carried out in accordance with principles related to cluster randomisation and blinding of analysis team, as outlined in the West Wales Organisation for Rigorous Trials in Health and social care (WWORTH) standard operating procedure (SOP) on randomisation.¹⁰² The trial statistician (WYC) and manager (JP) undertook randomisation after recruitment of paramedics had closed, to avoid selection bias, using trial site, volume of calls, falls service and the number of participating paramedics as stratification variables.

Recruitment

Patients were identified as being potentially eligible for study inclusion from routine emergency service dispatch records through standardised queries written for each site. Site researchers confirmed eligibility of individual patients by retrieving corresponding patient report forms (PRFs), which are routinely completed by paramedics when they attend a patient. PRFs include patient identifiers and demographics; and operational, clinical assessment and treatment information.

In this cluster trial we did not ask paramedics to approach patients for consent to participate in the trial during the emergency episode. Instead, following identification of eligible patients as detailed in *Participants*, we sought consent from patients for follow-up through routine medical records and by postal questionnaire. Following considerable discussion with the Research Ethics Committee, our consent process included contacting patients by post and then, if necessary, by telephone or a home visit. Patients then had the opportunity to actively 'consent' or 'decline consent' (dissent). However, we were unable to contact some patients at all (no response by post or telephone). We had Health Research Authority Confidentiality Advisory Group (HRA CAG) permission to follow up this group anonymously (but they neither 'consented' nor 'declined consent'). The group we analysed consisted of those who had consented and those who we could not contact (follow-up was through an anonymised route). Only those who had actively declined consent (dissented) were excluded from this group. The postal consent pack (see *Appendix 6*) included:

- covering letter
- consent form
- patient information sheet
- 1-month questionnaire (see *Appendix 7*)
- Freepost return envelope
- £5.00 voucher, to thank participants for their time.

Interventions

Experimental: intervention group care

The core of the health technology we evaluated was a clinical protocol for the care of older people who have fallen, enabling emergency ambulance paramedics to assess and refer them to community-based falls services (see *Appendix 1*). During a workshop held to define the intervention, participants from across specialties and sites agreed common minimum standards for each component of the intervention, with flexibility allowed for local differences in processes such as referral and documentation. The complex intervention comprised the assessment protocol (the health technology being trialled) and six other components, as shown in *Table 11*. Local variations are shown in detail in *Appendix 8*.

Control group care

Control group paramedics did not receive training in the intervention. We asked paramedics based at control stations to continue their usual practice. Although we know that conveyance rates vary considerably among services, stations and paramedics, we did not seek to standardise practice, as we were unable to identify best practice. Current practice in the control group was therefore care as usual comprising assessment of injury or other condition requiring immediate care, assistance in moving and conveyance to ED unless the patient refused.

Outcomes

Outcome measures at 1 month and 6 months after the patient's index incident were consistent with recommendations of ProFaNE.¹⁰³

Primary outcome

A composite outcome of subsequent emergency health-care contacts (in order of severity, i.e. death, emergency admissions, ED attendances or emergency service calls), summarised by:

- proportion of patients who suffer these events
- interval to first event
- event rate.

TABLE 11 Definition of the SAFER 2 trial intervention

Component	Core minimum standard	Local variation permitted
Assessment protocol	Clinical protocol to guide paramedic through the decision-making process and assess the patient's risk of further falls and safety for non-conveyance	Consent process for patient agreement to be referred to falls service Protocol may be used as aide memoire or as a specific additional form to the PRF
Training	One full day, with training package including programme, written materials and DVD (see <i>Appendix 9</i>)	Background and role of trainers, including members of falls teams when possible
Referral pathway	A pathway to allow intervention paramedics to refer patients to a falls prevention service, including a system to allow auditing of receipt of referral forms	Method and process of referral, to minimise any delays on-scene Participation of GP practices across falls service ^a catchment area
Referral tool	A form recording information to be passed to the falls prevention service, including basic demographic, clinical and contact details	Other information included in referral form
Falls service response to referral	Telephone contact with patient, by professional with ability to recognise cases where urgent input is needed Initial face-to-face multifactorial assessment; multidisciplinary assessment if appropriate	Time frame for initial patient contact and assessment
Falls service provision	NICE guidance-adherent service: multidisciplinary treatment delivered by team of appropriately qualified professionals (may include, e.g. equipment, balance training, medical review)	Details of service and patient intervention
Clinical and operational support	Clinical support debrief with ED every 2–4 weeks plus feedback on outcomes for individual patients referred to falls service	Exact method and means of clinical support and feedback

DVD, digital versatile disc.

^a At site 1 not all GP practices within the trial catchment area were linked to the participating falls services. Intervention paramedics needed to consult a list of participating GP practices before offering referrals to patients. These patients were trial eligible, but were not eligible for the full intervention. Numbers recruited are noted in *Chapter 4*.

Secondary outcomes

At index incident:

- onward pathway of care (conveyed to the ED; referred to falls service; referred to other provider)
- compliance with guidelines for ambulance service clinical documentation
- durations of: ambulance service job cycle (from receipt of emergency service call to time that ambulance reports being free for next call); episode of care (from receipt of emergency service call to end of emergency episode, i.e. the patient is left at home, discharged from ED or admitted to hospital); or time to falls service response.

At 1 month and 6 months after patient's index incident:

- duration of subsequent inpatient episodes
- subsequent reported fractures
- health-related quality of life, as measured by the Short Form questionnaire-12 items (SF-12)¹⁰⁴
- patient satisfaction as measured by the Quality of Care Monitor¹⁰⁵ and 'fear of falling' as measured by the modified Falls Efficacy Scale (mFES)¹⁰⁶
- self-reported further falls

- costs of care to NHS and PSS
- self-reported costs incurred by patients and carers.

Qualitative:

- views of ambulance service paramedics, managers and partners on implementation of the intervention
- experience and satisfaction of patients receiving the intervention.

Data collection

- Ambulance service: routine clinical and operational management information about index and subsequent contacts from despatch records (completed by emergency service call takers) and PRFs completed by paramedics during or immediately after patient attendance.
- Falls services: information about referrals and falls service response.
- Participants (or their carers): questionnaire data, sent by post to eligible patients at 1 month (see *Appendix 7*) and those who consented to a further questionnaire at 6 months (see *Appendix 10*).
- Health and Social Care Information Centre (HSCIC) (England) and National Welsh Informatics Service (Wales): routine data about ED attendances, hospital admissions and deaths for all matched study patients over the study period.

At all three sites, patients who dissented by any means were excluded from all data collection. For all other patients, we collected anonymised data on index and subsequent health-care contacts. Identifying information collected at ambulance and falls services was split from clinical information by our trial partners within the three ambulance services. In order to improve the matching rate achieved by National Welsh Informatics Service and HSCIC, our ambulance service partners used imperfect emergency service call information such as name, address and date of birth to look up NHS numbers. Once patients were matched, clinical outcome data were retrieved and transferred for analysis into the Secure Anonymised Information Linkage (SAIL) gateway when it was joined up with questionnaire data.

Data entry

The method of data entry varied by source and site. Ambulance service despatch data were downloaded electronically to study databases. There were two types of PRF: paper and electronic. Sites 2 and 3 used only paper forms and site 1 used a mix of paper and electronic forms. Paper forms were input into study databases manually by the study team, whereas electronic forms were input automatically after they were downloaded from a central server. Questionnaires were input into study databases using Teleform software version 10 (Cardiff Software, Sunnyvale, CA, USA). A quality assurance check was carried out on all manually entered data.

Blinding

Blinding was carried out when possible. Paramedics, trial managers and site researchers were not blinded, as they needed to know the allocation of participating ambulance stations for operational reasons. The trial statistician and all other TMG members were kept blind to allocations in order to eliminate reporting bias. This was undertaken by coding the allocated groups in order to prevent group identification. Blinding statisticians is a technique recognised, both by statisticians in general and by the Wworth SOP on statistics,¹⁰⁷ as best practice in statistical analysis. Even though the Wworth SOP also requires that trials agree statistical analysis plans before starting analysis, these cannot cover all eventualities, with the result that unblinded statisticians could unconsciously take analytical decisions that favour intervention over control or vice versa. Unblinding occurred after completion of primary analyses.

Sample size and power

We estimated our trial sample size from our principal outcome: the proportion of participants who, within 6 months, die or contact emergency services. From previous trials of interventions for older people who have sustained a fall and presented for emergency treatment, summarised in a recent systematic review,¹⁰⁸

we made the conservative estimate that trial patients had about a 50% chance of making another emergency contact or dying within 6 months. We judged that a change of 5% in this proportion could be clinically and economically important. In the absence of clustering, a sample size of 4190 evaluable participants would have yielded 90% power to detect a change of at least 5% (from 50% to $\leq 45\%$ or $\geq 55\%$) when using a two-sided 5% significance level. As participants came from 25 clusters, we needed to adjust this sample size to allow for intraclass correlation (ICC). We estimated this ICC from the findings of the SAFER 1 trial,⁷⁹ which evaluated the clinical effectiveness and cost-effectiveness of computerised clinical decision support software for use by paramedics when attending older adults who had a fall. The SAFER 1 trial estimated the ICC for the same outcome, but over 1 month rather than 6 months, as 0 when clustering participants by station (as in the SAFER 2 trial), but 0.005 when clustering participants by paramedic (as in the SAFER 1 trial).⁷⁹ To be conservative, we allowed for an ICC of 0.002 by increasing the target evaluable sample to 6290, namely 25 clusters \times 251.6 participants per cluster. This sample would have more than 90% power to detect a change of 0.18 in the number of emergency contacts of 1.8 over 6 months, given a standard deviation (SD) of 1.5. Hence, the SAFER 2 trial was able to detect a difference of 1 emergency contact in 10 avoided (or induced) by the intervention.

We had originally postulated that patients recruited to the study would have a 40% chance of making an emergency contact or dying within 6 months and that the ICC could be as high as 0.03. Under those assumptions our target sample of 6290 would have yielded 80% power to detect a change of at least 10% (i.e. from 40% to $\leq 30\%$ or $\geq 50\%$) when using a two-sided 5% significance level. When the SAFER 1 trial showed that the assumed ICC was unduly pessimistic, recruitment was progressing well. Therefore, rather than finish the trial early, we decided, with the approval of both TSC and DMEC, to be less conservative in assuming a worst ICC of 0.02, thus yielding enough power to detect a change of only 5% in the emergency contact rate, still a clinically important difference in the view of our advisers.

Statistical methods

The full analysis plan can be found in *Appendix 11*. Primary analysis was by 'treatment allocated'. Analyses included logistic regression for binary outcomes; cross-tabulations and risk ratios for categorical outcomes; and survival analysis including Cox's proportional hazards models for times to events. We used multilevel modelling to estimate (random) station effects and (fixed) group effects and analysed repeated observations as such.

Our principal outcome is made up of a hierarchy of events. We therefore undertook analysis incrementally: first, deaths; second, emergency admissions plus deaths; third, ED attendances plus admissions and deaths; and, finally, emergency service calls plus attendances, admissions and deaths. Although we originally planned to undertake analysis for all events and for those coded as a fall, in practice data quality across the levels precluded analysis of falls only. We have presented results as follows: the proportion of patients who called the emergency services, attended ED, were admitted or died; survival analysis of the time to the first subsequent emergency contact; the mean number of further emergency contacts adjusted for time at risk, excluding days in hospital or after death; and recurrent event analysis when feasible. We also examined the effect of the intervention on patient satisfaction, health-related quality of life and costs (as described under *Economic evaluation*).

Potential predictors of triage decisions include the distance between the site of the index event and the ED; patients' age, sex and history of previous falls; type of presentation [specifically, a seasonality term capturing when, during the calendar year, the call was made, and an indicator of whether or not it was made out of (GP) hours]; and time since recruitment started (recruitment point). The list of adjusting factors (covariates) includes those suggested by the TSC and may be divided into three subsets: (1) group and site interactions, reflecting study design; (2) participant characteristics (age and gender); and (3) characteristics related to the timing and location of the index incident. We used the covariates in the analysis. It is possible that patients in the catchment area of one station may receive care for a subsequent event from a paramedic based at another station participating in the study but allocated to a different group. Nevertheless, analysis was still by treatment allocated. We have identified referrals made

to falls services at subsequent emergency service call attendances and are able to use this for censored analysis in order to minimise contamination, but this is not included in the current report.

We proposed to identify any wider system effects, by comparing response times during the trial period across the study catchment area and surrounding areas with pretrial response times and response times elsewhere. This analysis would only be necessary if effects were found on ambulance operational performance indicators – job cycle time within the trial population.

To inform the development of outcome measures for falls research as recommended by ProFaNE,¹⁰³ we compared SF-12 and derived Short Form questionnaire-6 Dimensions (SF-6D) scores with mFES scores to establish their construct validity. We also assessed their predictive validity by comparing scores with the number of further events and the time to the first subsequent event.

Missing data

We adopt a consistent approach to missing data relating to both clinical effectiveness and cost-effectiveness, except when individual outcome measures require some variation in that approach. For each variable, we consider the frequency of missing data; if there is no reason to suspect that data are not missing completely at random, we use appropriate imputation methods to mitigate the problem of missing data. Specifically, in addition to taking the SF-6D score as 0 when a participant is dead at the data collection point, we imputed missing values by regression (thus adjusting for other covariates) from all available values of that score at other data points.

Adverse events

As the study population had high mortality and morbidity, we did not routinely record or report AEs that were neither serious nor adverse reactions (ARs) in the sense of possibly being caused by the new clinical protocol for referring to falls services. The main potential AR is misdiagnosis, which could lead to an inappropriate pathway of care. As misdiagnosis is reliably identifiable only through patient complaints or coroner's inquest, we focus on these, and treat them as serious ARs. Any patient complaint or coroner's inquest at which the ambulance service was asked to supply information related to non-conveyance of a trial participant from the index incident triggered was investigated by the local principal investigator and chief investigator. We also investigated suspected ARs brought to our attention in any other way.

Death or emergency hospital admissions are serious adverse events (SAEs). As these form the primary outcome of this trial, and are not unusual or unexpected in the study population, we report them at the end of the trial. In particular, the imbalance between intervention and control groups in the occurrence of SAEs or serious ARs was the subject of statistical analysis at the end of the trial.

Economic evaluation

Aim

We undertook an economic evaluation alongside the RCT from the perspective of the UK NHS and PSS, in line with the approach recommended by NICE. Economic analysis estimated the costs of providing the intervention and the consequences of the scheme for the NHS and PSS in terms of inpatient admissions, ED attendances, GP consultations, out-of-hours GP contacts, NHS Direct contacts and use of social services.

Method

We estimated the costs of providing the new intervention by utilising data collected from financial reports and documents, relevant information logged as a part of routine practice and from resource utilisation recording sheets, together with reference to patient records and discussions with relevant finance staff. We collected data on participants' use of health service and social services resources from paramedic records, routine hospital records and patient-completed questionnaires. We estimated NHS resource use from routine data including duration of ambulance job cycles and episodes of care; records of resource

use; and patient records. We also estimated social services resource costs from discussion with relevant social services departments. Costs were then calculated using published unit costs such as the Personal Social Services Research Unit (PSSRU) Unit Costs of Health and Social Care 2013¹²³ and *NHS Reference Costs 2011–12*.¹⁰⁹

Health-related quality-of-life outcome

The SF-6D scores (derived from SF-12 scores) were used to estimate the QALYs gained from the intervention and an incremental cost per QALY, applying appropriate threshold incremental cost-effectiveness ratios, such as the £20,000.00 and £30,000.00 per QALY thresholds used by NICE. These ratios were presented along with their associated cost-effectiveness acceptability curve.

Lost to follow-up/missing data

If a participant was dead at the data collection point, the SF-6D score was taken as 0. To avoid outliers, SF-6D and mFES scores were taken to be the minimum value observed for that measure in the relevant treatment group. The problem of missing data was addressed using an appropriate regression-based imputation approach, as reported earlier.

Discounting

As the outcomes were assessed at 1 month and 6 months, no discounting was employed.

Uncertainty

We addressed uncertainty by applying bootstrapping for cost-effectiveness acceptability curves and CIs. Sensitivity analyses were undertaken to assess the robustness of results to changes in the configuration of the intervention and other health service costs.

Qualitative methods

Aim

The qualitative component of the trial was designed to address the following two of the study objectives:

1. Gain an in-depth understanding of how the intervention is experienced by patients.
2. Understand how the intervention is delivered in practice, identifying factors which enable or hinder its use.

Participants

Patients

We undertook qualitative interviews with a sample of trial patients who had experienced a fall, were seen by an intervention paramedic and agreed to the interview. Patients were selected from all three study sites, with a target number of 20 from each site. A mix of patients was identified for interview: those who had been taken to ED; those who had been referred to a falls service; and those who were neither taken to the ED nor referred to a falls service.

All eligible patients who had been referred to a falls service were selected for interview. The other two patient groups (taken to ED and left at home without referral) were much larger; sampling was therefore carried out so that only every 10th sequential patient was selected for interview. The selected patients were then telephoned to arrange an interview. Whenever possible, this occurred 6–8 weeks after the index incident, although in practice some interviews took place up to 4 months after the index incident. A log of the interviews undertaken and the characteristics of the patient's index incident (i.e. location, disposal, patient sex and patient age) was kept in order to ensure that all patient groups were interviewed.

Paramedics

We invited all intervention paramedics who volunteered to take part in the study to discuss their expectations and experiences of the intervention. Whenever possible, paramedics were invited to take part in a focus group,¹¹⁰ bringing together between four and eight of their peer group in a local venue within their ambulance service area. When paramedics were not available to take part in a focus group, we invited them to take part in a face-to-face interview.

We carried out interviews with paramedics before they started to carry out the intervention (pre trial but after having received the training) and after the patient recruitment phase (post trial).

Other stakeholders

In each study site, data were gathered after the end of the trial period from other relevant staff involved in delivering the intervention: training staff and management staff from the ambulance service, and those involved in managing and delivering the falls service. Participants were selected purposively and were invited to take part in focus groups; those who were not able to attend a focus group were invited to take part in an interview instead.

In each of site 2 and 3, one focus group was held of staff involved in delivering the falls service, and one of staff in the ambulance service with a training or supervisory role. In addition to this, in site 3 two one-to-one interviews were conducted with stakeholders who were not able to attend a focus group. In site 1, stakeholders took part in four one-to-one interviews, but no focus group.

Data collection

We interviewed patients and paramedics only in the intervention arm (not in the control group) because we were interested in how the intervention was used and experienced in practice, and the mechanisms by which it may or may not have worked.

Patient interviews

A schedule (see *Appendix 12*) for face-to-face, semistructured patient interviews was developed by the research team, drawing on the aims and objectives of the study, in order to explore:

- patients' experience of the ambulance service
- patients' experience of those seen by a falls service
- patients' health since fall
- patients' satisfaction with treatment.

Face-to-face semistructured interviews were conducted by experienced researchers from the study team (RA, RC, MK, LW). In some cases, a carer or other person who had been present at time of fall also took part in the interview or was present to support the participant. Interviews took place in patients' own homes, and were recorded and later transcribed in full.

Paramedic focus groups and interviews

Data collection with paramedics took place at two points in the study:

1. at baseline, after paramedics had been trained in the SAFER 2 trial intervention but before they started using it in practice (see *Appendix 13*)
2. after the end of the patient recruitment period (see *Appendix 14*).

Separate topic guides were developed by the research team for each phase of data collection, drawing on the aims and objectives of the study. The topic guides were reviewed by members of the LITs, which

included ambulance service personnel at different clinical, operational and management levels. The focus group topic guides were adapted for use in one-to-one interviews. The topic guide covered:

- paramedics' views and attitudes towards the new intervention
- any preconceptions about the new way of working
- factors which enabled the use of the new referral pathway
- factors which hindered the use of the new referral pathway.

Each focus group was led by one researcher, with a second member of the team acting as support and taking notes to enable the linkage of texts to speakers, and noting other details, such as points of consensus or disagreement, issues that drew strong emotional responses such as anger, fear or anxiety. All focus groups and interviews, with the permission of participants, were recorded and transcribed in full.

Stakeholder focus groups and interviews

Topic guides (see *Appendix 15*) were developed by the research team for use with ambulance service personnel and wider stakeholders, and covered:

- perception of how well the intervention worked
- impact on the stakeholder's organisation
- cross-organisational working
- views on the process of implementing the intervention.

Stakeholder focus groups took place before the main results of the trial were known. Each was led by one researcher, with a second member of the team acting as support and taking notes to enable the linkage of texts to speakers, and noting other details, such as points of consensus or disagreement, issues that drew strong emotional responses such as anger, fear or anxiety. Each interview was carried out face to face, and was conducted by one member of the research team. All focus groups and interviews, with the permission of participants, were recorded and transcribed in full.

Qualitative data management

As this was a team-based project involving collection of data in multiple sites, maintaining consistency in data collection and transcription was crucial. The co-ordinator set up a data management system consisting of instructions on converting raw data to computer files (in the form of a transcription protocol), on organising data storage and on data archiving steps and a data management checklist. The transcription protocol ensured that standard conventions were adopted throughout the transcription process, and that a standard presentation format was used. All transcripts were anonymised and stored securely. Transcripts of interviews with stakeholders were sent back to the participant for checking, to ensure that they were accurate.

Qualitative data analysis

The data were analysed thematically using a modified version of the framework analysis approach for applied policy research.¹¹¹ This is a systematic, dynamic and transparent method of analysis, which generates themes from the original accounts of participants.¹¹²

Two separate analysis tasks were carried out: data from patient interviews were analysed to address objective 3; and data from focus groups and interviews with paramedics and stakeholders were analysed to address objective 4. Each task consisted of the following stages:

- The transcripts were distributed for reading among members of the research team (RA, RC, SG, MK, AP, LW and AT, the patient representative) so that each had at least five to read, with each transcript being read by two people.
- Each member of the research team made notes on themes and ideas, guided by the relevant study objective.

- The research team met to discuss and agree a coding frame.
- Two researchers independently coded up two transcripts each using the coding frame, discussed how well it worked, and then revised the coding frame. The final coding frames are shown in *Appendix 16*.
- RA, RC, MK and AP used the revised coding frame on one transcript, then compared coding to check consistency.
- Transcripts were then shared among RA, RC, MK and AP for coding. RA then entered all codes into an NVivo database (QSR International, Warrington, UK) (one database relating to objective 3 and one relating to objective 4).
- RC, MK and AP met to discuss overall themes and findings relating to each objective, drawing on the NVivo database and the coding frame.

The qualitative findings in *Chapter 6* present anonymised quotations from participants. In line with the guidance from Corden and Sainsbury,¹¹³ quotations have been used to present evidence, to support explanation of complex ideas, to provide illustration, and to deepen understanding, particularly when respondents had an emotional response. Although many quotations were selected to illustrate the majority view, some show exceptional viewpoints, as noted. In some cases, quotations were selected to illustrate complexity or ambiguity of perspective, as noted.

Involvement of service users

Involving service users in research is encouraged to improve relevance, quality and accountability of research.¹¹⁴⁻¹¹⁷ The SAFER 2 trial followed the principles and procedures outlined in the WWORTH SOP for user inclusion¹¹⁸ to ensure that members of the public were actively involved throughout the research process. This SOP provided a starting point for developing a model of user involvement in the SAFER 2 trial. Moreover, the SAFER 2 trial aimed to address the challenges of involving older people, who are often in poor health because of their falls history or falls risk. The model took into account the multiple layers of a multisite trial and proposed three tiers of involvement in different forums at the strategic, site and local levels where overall or specific input could be made. We followed best practice guidance by involving two service users in each of the strategic trial committees.^{114,119,120} We also involved service users at site ($n = 3$) and local levels ($n = 18$) of the trial to provide a range of forums and opportunities for involvement by older people with risk of falling. Our involvement model received the Involving People Award in 2014 as an example of best practice.

Strategic level

Two service user members were sought for each of the TMG, TSC and DMEC. These strategic level meetings involved trial co-applicants and independent members. They were responsible for trial oversight and took strategic level decisions.

Site level

Local implementation team meetings took place at each site to oversee the delivery of the trial, and service users were sought to participate in these. In addition, in each of the three sites a Service User Reference Group was formed to provide forums for service users to discuss and contribute to the study. They were co-ordinated by the site researchers and held meetings at universities and community venues. Membership, meeting arrangements and frequency varied, reflecting local interest and individual needs.

Local level

Service user representatives took part in task-and-finish groups to develop the patient questionnaire and associated paperwork and to find ways to improve questionnaire response rates. Finally, towards the end of the study period, service users took part in qualitative analysis meetings.

Service users were recruited by various routes:

- previous experience of participating in research studies or membership of a standing user involvement group connected to a university

- membership of a third-sector organisation bringing together members of the target population (e.g. the Princess Royal Trust for Carers)
- Involving People Network providing links to service users
- personal contact from the research team.

All service users contributing to the trial were provided with training and support (including expenses payments) in line with guidelines set by the public involvement organisation INVOLVE.¹¹⁴

Through the methods adopted by the trial, service users were involved in a range of trial processes, including developing and planning research; overseeing the management of the trial; refining data collection methods; and contributing to analysis, particularly of the qualitative data. *Appendix 17* is a case study from one of our service users about his involvement in the SAFER 2 trial.

Support and Assessment for Fall Emergency Referrals 2 trial progress

This section details the ‘story’ of the study, charting progress from the awarding of funding to the completion of the project, highlighting the main events leading to the successful completion of the study and events that led to unforeseen setbacks to the project timetable.

The SAFER 2 study received confirmation of funding from the HTA programme in January 2009. The trial ethics application was submitted to the Multicentre Research Ethics Committee for Wales in April 2009. This highlighted the proposed consent process (the same used in the SAFER 1 trial), in which patients were invited to ‘opt out’ if they did not wish to take part in the study. The committee approved this process subject to permission from the then National Information Governance Board Ethics and Confidentiality Committee (now the HRA CAG), as it would involve accessing patient information without their explicit written consent. The Ethics and Confidentiality Committee did not give approval on the basis an active opt-in consent process should be used. This decision had implications for the trial as it required potential participants to return signed consent forms, which would be likely to lead to a low inclusion rate with the more elderly, frail and vulnerable patients under-represented. After consultation with our research team members, we proposed an alternative process for recruitment, combining both informed consent (gained either postally or via telephone) and anonymous follow-up, initially via the SAIL system in Wales. Ethical approval was gained in February 2010, 11 months behind schedule. In 2012, a further application to the HRA CAG for anonymised follow-up of patients from England who had not declined consent (dissented) through the HSCIC was approved.

Alongside this, the study team needed to confirm the participation of ambulance and falls services; negotiation was required with service providers to agree arrangements for participation.

- In site 1, two of the six falls prevention services declined to participate in the trial because of concerns with capacity.
- In site 2, one of the two health boards implemented a ‘Frailty Programme’ during the same time period the trial was scheduled to run. Under the Frailty Programme, all older people who fell and called the emergency services were referred on to be seen at home immediately. This compromised the trial methods, as there would be no control group. Negotiations were ongoing for several months as to the best way to run the two programmes alongside each other, but the health board eventually decided to not participate in the study because of this conflict with local service developments.
- Owing to delays setting up the study, the commencement of the SAFER 2 trial coincided with the ambulance service that contained site 3 introducing a financial reimbursement for referrals to GPs of people who had fallen. This was incompatible with the RCT design because of the elimination of the control group, which would lead to significant issues at this site. Following the chief investigator’s attendance at a senior-level strategic meeting, site 3 reaffirmed its commitment to full participation in the

trial, with the GP referral programme introduced across the service except in the trial catchment area. Research and development (R&D) permissions were signed off, with a collaborator's agreement in place.

Owing to risks around the participation of site 3 and falls services in the site 1 and site 2, in August 2010 we held discussions with a fourth ambulance service that expressed an interest in joining the trial. After a period of negotiation and consideration across local partner services, it decided that it could not participate in the trial at this time.

By early 2010, participating services were confirmed. In total, the study involved three ambulance services, with 26 NHS partners based within five comprehensive local research networks (CLRNs) and NHS trusts. Research governance processes commenced across the three trial sites, later than scheduled, in March 2010. Although English ambulance services, which each span several CLRNs, had a lead CLRN identified, at the time it was not clear how this process (i.e. research processes, cost attribution and support to carry out studies) worked for the participating acute trusts and primary care trusts (PCTs) in the trial areas. On advice from these organisations, principal investigators were identified and approached for sign-up at the 26 NHS organisations. Site-specific information forms were prepared, but these could neither be completed nor a formal sign-up of NHS organisations be achieved without agreement of the attribution of costs across the trial. Although NHS costs had been estimated at the outset of the planning process, these were affected by changes to the study design (in particular the recruitment and data retrieval processes) and lack of clarity about cost attribution. Discrepancies between CLRN and NHS research site views on cost attribution led to the set-up of a meeting with the Department of Health in May 2010, in which the attribution of costs was finally agreed.

Alongside the processing of research governance approvals, the study team were developing the SAFER 2 trial intervention. Undertaken using the Medical Research Council guidance,⁶⁴ the intervention was established through the use of previous literature, a stakeholder workshop, modelling and patient involvement (see *Appendix 8*).

It was very difficult to predict how long it would take to complete R&D processes at global and local levels across the research sites. R&D approvals for all three sites (with the exception of some acute trusts in site 3, where full R&D approvals were not gained until March 2012) were finally gained in October 2010, allowing the trial to commence.

The delays outlined above had a knock-on effect on the overall timescale of the project. In mid-2010, we revised the project timetable, proposing that the full pilot with recruitment of patients for the study started in January 2011 followed by the main trial in February. Owing to the study team receiving additional data regarding the number of older fallers seen by paramedics, we reassessed the expected recruitment rate in order to meet recruitment targets, resulting in a proposed extension to the recruitment phase of 2 months (until the end of October 2011 instead of the end of August 2011).

As the trial gained momentum, we looked to recruit two new research support officers, funded through service support costs (CLRN/National Institute for Social Care and Health Research) to undertake administrative tasks associated with patient recruitment. These ambulance service-based roles (which were undertaken by paramedics in site 2 and site 3 and an administrative assistant within site 1) were key to paramedic recruitment, as our researchers had restricted access to ambulance stations and identifiable data. Full funding agreements had to be in place before the ambulance services would proceed with recruitment to these posts, which led to delays in advertising and filling them. Further delays occurred in site 3 because of issues releasing paramedic staff to undertake these roles. Inevitably, this had a knock-on effect on the paramedic recruitment schedule.

Paramedic recruitment to the trial commenced in November 2010 at the three trial sites. Recruitment of paramedics was initially slow; some showed unwillingness to be involved in the trial when there was a 50% chance of being randomly allocated to the control group. In December 2010, we received approval

from the ethics committee to provide a £50.00 voucher to paramedics for taking part (in London the money was pooled into a station training budget); this allowed paramedic recruitment to increase.

Paramedic training was undertaken once stations (clusters) were randomly allocated to either control or intervention arms of the trial. Paramedics based at intervention stations underwent training delivered by ambulance service trainers, falls service representatives and the study team, using a cascade method originating with the training team who developed the clinical protocol. In early 2011, following the successful completion of paramedic training, patient recruitment commenced. Site 1 was first to start in March 2011, with sites 2 and 3 following in April and July, respectively.

Identification of eligible patients required a high level of resources. In two of the trial sites (1 and 2), PRFs containing patient details needed to confirm eligibility into the trial were particularly difficult to locate. These forms were completed by paramedics on-scene and were stored at ambulance stations before being centrally collated. Considerable effort was put into locating the forms, which were often difficult to locate because of the issues with data reliability. Ambulance service senior managers were aware of the issue and were helpful in terms of supporting the location and retrieval of the outstanding forms.

The study team wrote to patients to gain informed consent once study eligibility had been confirmed. Informed consent is particularly challenging in prehospital care research and in the SAFER 2 trial this was even more so as the participants were frail and elderly. Initially, patient consent rates were low, at around 10%. In August 2011, following a review of the consent procedures, the study team proposed an alteration to the consent process, including simplification of the consent form, sending of the questionnaire with the initial patient letter and inclusion of a £5.00 voucher to thank patients for their time spent considering the study. A substantial amendment was submitted to and approved by the Multicentre Research Ethics Committee, allowing the process to be altered. The consent rate to identifiable data follow-up increased to 30%. In early 2012, a front cover was added to the existing questionnaire to improve response rates.

In October 2011, the trial was temporarily suspended at site 1, because of internal governance concerns regarding the processes of verbal consent of patients over the telephone. As a consequence, we revised the way in which verbal consent was recorded across the trial. From the start of the patient recruitment phase, it was clear that fewer eligible patients were being attended by study paramedics within sites 2 and 3 than estimated from the pilot data received at the start of the study: this was mainly because of paramedics working outside the study catchment area. This, coupled with the previously mentioned issues around ethical approvals, information governance and R&D permissions, meant the study fell behind its recruitment schedule. Originally, patient recruitment was scheduled to finish in December 2011. In early 2012, the HTA funded an extension to the study, allowing patient recruitment to carry on until the end of June 2012.

Summary of changes to the project protocol

The following changes were made to the original protocol after it was funded:

- The process for recruiting patients to the trial was amended following ethical approval from an opt-out process to an active opt-in consent process including follow-up telephone call.
- Participating ambulance stations at the London site were changed from South to North Central London stations, following set-up of a falls service in the original proposed study area.
- The paramedic recruitment process was amended to include a £50.00 voucher for those who volunteered.
- The number of paramedics recruited at each site was updated to reflect actual numbers recruited, which were greater than originally stated.
- The European Quality of Health-5 Dimensions (EQ-5D) was not used. In order to be consistent with outcome measures used on the SAFER 1 trial, we instead used the derived SF-6D.

- Patient recruitment was further amended after the trial commenced to include a simpler process and the addition of a £5.00 voucher with the invitation pack. This was in order to improve response rates.
- An anonymised data follow-up process for the two English sites was added.
- We originally planned to compare test–retest reliability and responsiveness of the instruments, by sending a retest questionnaire to 100 randomly selected patients from each site. We did not proceed with this because of the evident difficulties our frail and elderly patients were experiencing completing essential study documentation.
- Administration of the Patient Generated Index was also originally included as part of the qualitative patient interviews. Owing to time and resource pressures this was not carried out.
- The study received two extensions. The first extended the study by 15 months, from 30 June 2012 to 30 September 2013. Owing to difficulties in gaining ethical permission at the commencement of the study, there had been a delay to the start, which had a knock-on effect to the rest of the trial. Recruitment was also slower than expected, and the extension allowed for us to reach our projected recruitment target. The second (unfunded) extension was for a further 4 months, to 31 January 2014. This was to allow time for full analysis of the data, following difficulties in retrieving anonymised outcome data.

Ethics and research governance

Ethical approval was obtained from the Research Ethics Committee for Wales, Information Governance approval from the National Information Governance Board, and NHS R&D approval from each participating health board, NHS trust and PCT.

Trial registration

The trial is registered as Current Controlled Trials ISRCTN60481756 and UKCRN 6801. The trial protocol has been published.⁷¹

Chapter 4 Clinical effectiveness results

Participant flow

Between March 2011 and June 2012 a total of 5914 eligible patients were attended by 215 paramedics based at 25 ambulance stations across the three study sites [Consolidated Standards of Reporting Trials (CONSORT) flow chart in *Figure 2*; site CONSORT flow charts in *Appendix 18*]. Six of 31 eligible stations withdrew after randomisation but before the start of patient recruitment because of a conflicting 'Frailty Programme' intervention being introduced throughout one participating health board area at site 2. Participating paramedics based at 14 stations were randomly allocated to the intervention group and those based at the remaining 11 stations were randomly allocated to the control group. After 1210 (20%) dissenting patients were excluded, 4704 (80%) were available for follow-up: 2420 in the intervention group and 2284 in the control group. As all but 49 patients were subsequently matched at SAIL or HSCIC, we included 4655 patients in analysis of anonymised linked data primary outcome.

Recruitment

Patients were recruited to the trial between 14 March 2011 and 30 June 2012. Start dates varied between March and July, as sites were ready to begin recruiting patients. All sites finished recruitment at the same time point (*Table 12*).

We followed up patients for 6 months after the index incident.

Baseline characteristics

Individual patient level

Although we expected recruitment of patients to be similar across ambulance service sites, and similar numbers of ambulance stations participated in the trial at each site, recruitment of patients was much higher at site 1 as a result of several factors including:

- longer recruitment period at site 1
- operational practice which took paramedics out of the trial catchment area, particularly in site 3
- loss of six stations and related paramedics at site 2.

There was also variation in recruitment between trial arms across sites, with a higher proportion of control group patients recruited at site 1 than at either site 2 or 3. There was very little difference between groups in age, overall or at any site. There were more women than men recruited to the trial, although the proportion of men was slightly higher in the intervention group than the control group, with about 3% more at each site and overall. There was little difference between trial arms in the proportion of calls made outside usual GP operating hours was similar between trial arms, distance to ED or time to recruitment (*Table 13*). There were some differences between sites, for example site 3 had a lower proportion of female patients, a lower proportion of calls made 'out of hours' and shorter distances to ED. This was our most urban site, which may have accounted for these differences in study population.

Cluster level

We randomly allocated one more station to the intervention group than the control group at each site. However, we recruited fewer paramedics recruited per station in the intervention arm than in the control arm at each site.




FIGURE 2 Flow of clusters and individuals through RCT for all participants. Adapted from Snooks et al.¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 12 Periods of recruitment

Site	Recruitment period		
	Start	End	Length in days
Site 1	14 March 2011	30 June 2012	475
Site 2	4 April 2011	30 June 2012	454
Site 3	6 July 2011	30 June 2012	361

TABLE 13 Baseline individual and cluster characteristics by treatment allocated

Variables	Group	
	Intervention	Control
Individual variables		
<i>Number of participants</i>		
All sites	2420	2284
Site 1	1329	1352
Site 2	544	436
Site 3	547	496
<i>Age (years), mean (SD) [n]</i>		
All sites	82.54 (7.97) [2414]	82.14 (8.11) [2275]
Site 1	82.99 (7.81) [1329]	82.57 (7.91) [1352]
Site 2	82.79 (7.97) [538]	81.79 (8.60) [427]
Site 3	81.20 (8.21) [547]	81.28 (8.16) [496]
<i>Female, proportion (%)</i>		
All sites	1480/2419 (61.2)	1477/2284 (64.7)
Site 1	816/1329 (61.4)	882/1352 (65.2)
Site 2	351/543 (64.6)	292/436 (67.0)
Site 3	313/547 (57.2)	303/496 (61.1)
<i>Emergency service index call out of hours, proportion (%)</i>		
All sites	1012/2419 (41.8)	954/2282 (41.8)
Site 1	624/1328 (47.0)	626/1352 (46.3)
Site 2	221/544 (40.5)	174/436 (39.9)
Site 3	167/547 (30.5)	154/494 (31.2)
<i>Distance to ED (miles), mean (SD) [n]</i>		
All sites	4.77 (3.43) [2406]	4.65 (3.08) [2270]
Site 1	5.03 (3.09) [1329]	5.04 (2.61) [1352]
Site 2	7.33 (3.38) [542]	6.75 (3.65) [434]
Site 3	1.55 (0.74) [535]	1.71 (0.72) [484]

continued

TABLE 13 Baseline individual and cluster characteristics by treatment allocated (*continued*)

Variables	Group	
	Intervention	Control
Recruitment point (days), mean (SD) [n]		
All sites	221.8 (129.0) [2420]	220.8 (129.2) [2284]
Site 1	239.4 (136.4) [1329]	238.1 (132.9) [1352]
Site 2	217.4 (128.8) [544]	227.0 (128.7) [436]
Site 3	183.4 (98.6) [547]	168.3 (102.9) [496]
Cluster variables		
<i>Number of stations (number of paramedics)</i>		
All sites	14 (105)	11 (110)
Site 1	5 (39)	4 (38)
Site 2	5 (26)	4 (26)
Site 3	4 (40)	3 (46)
<i>Recruited paramedics per station, mean (SD)</i>		
All sites	7.5 (5.185)	10 (6.708)
Site 1	7.8 (4.97)	9.5 (4.203)
Site 2	5.2 (3.033)	6.5 (2.082)
Site 3	10 (7.348)	15.33 (11.15)

Reprinted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Numbers analysed

Table 14 shows questionnaire response rates for all 4704 eligible patients who did not dissent and for whom we sought to retrieve linked data outcomes. However, SAIL and HSCIC could not match the demographic data for 29 intervention patients (18 in site 1, 4 in site 2 and 7 in site 3) and 20 control patients (15 in site 1, 3 in site 2 and 2 in site 3). Therefore, we could not retrieve data on ED attendances, hospital admissions or death for these 49 patients.

Accounting for people who did not consent to questionnaires and those who died within 1 month, the overall response rate was 36.5%. Although there was little difference between intervention and control groups, response rates ranged from 30.0% at site 3 to 49.5% at site 2.

At 6 months the overall response rate was 58.7% (60.1% in the intervention group and 57.2% in the control group). Again there was variation in response rate between sites from 54.1% at site 1 to 67.4% at site 3. Furthermore, in site 3, 73.8% of the intervention group returned a 6-month questionnaire compared with 60.3% of the control group.

Outcomes and estimation

We did not find any significant differences between groups in our composite primary outcome at 1 month or 6 months. One-third of patients had suffered a further emergency episode or death by 1 month, rising to over two-thirds by 6 months. When primary outcome components were analysed separately we found

TABLE 14 Questionnaire response rates by treatment allocated

Variable	Group								
	Intervention				Control				Total
	Site 1	Site 2	Site 3	Total	Site 1	Site 2	Site 3	Total	
Total number of eligible participants	1329	544	547	2420	1352	436	496	2284	4704
No consent to 1-month questionnaire (before new consent process), <i>n</i>	330	117	n/a	447	301	93	n/a	394	841
Died within 1 month, <i>n</i>	86	31	30	147	79	26	32	137	284
Participants sent 1-month questionnaire, <i>n</i>	913	396	517	1826	972	317	464	1753	3579
Valid 1-month questionnaires returned, <i>n</i>	322	191	155	668	337	162	140	639	1307
1-month questionnaire response rate, %	35.3	48.2	30.0	36.6	34.7	51.1	30.2	36.5	36.5
No consent to 6-month questionnaire, <i>n</i>	759	299	313	1371	743	261	295	1299	2670
Died within 6 months, <i>n</i>	249	116	93	458	270	75	75	420	878
Participants sent 6-month questionnaire, <i>n</i>	321	129	141	591	339	100	126	565	1156
Valid 6-month questionnaires returned, <i>n</i>	172	79	104	355	185	62	76	323	678
6-month questionnaire response rate, %	53.6	61.2	73.8	60.1	54.6	62.0	60.3	57.2	58.7
n/a, not applicable.									

evidence of significantly lower demand in the intervention group. *Tables 15 and 16* show fewer further emergency service calls at both 1 and 6 months, with differences largely consistent across sites, and some indication of fewer ED attendances at 6 months, albeit with significant variations across the three sites (*Tables 17 and 18*).

Table 19 shows that rate of conveyance to ED at the index incident was similar between groups overall and at each site, although the proportion conveyed varied considerably by site, from < 60% at site 1 to nearly 80% at site 2. Eight per cent of trial eligible patients were referred to falls services by their attending paramedic in the intervention arm, varying from 7.5% at site 1 to 9.7% at site 3. Very few patients were referred to falls services in the control group; the rate was 1% overall, but higher at site 3, where there was a service-wide initiative to increase referrals of this patient group. More patients were left at scene without referral for further care in the control group than in the intervention group, 30% versus 23% overall; a proportion which varied widely by site and was highest in the control group at site 1 (37%). Completion of clinical documentation was high across groups and sites with no clear difference between trial arms. We also found no differences in the operational indicators of duration of episode of care (time from emergency service call until patient's emergency episode was complete) or job cycle (time from emergency service call until ambulance free).

No differences were found in the secondary outcomes of duration of inpatient stay, or self-reported quality of life or fall-related self-efficacy (fear of falling) at 1 month or 6 months (*Tables 20 and 21*). There was evidence of differences between groups in the proportion reporting further falls at 1 and 6 months: lower in the intervention group at 1 month, but higher at 6 months amid significant variations across sites. There were also significant differences between groups in the interpersonal aspect of patient satisfaction at 1 month

TABLE 15 Primary outcome and components at 1 month analysed by treatment allocated

Primary outcome	Raw data		Adjusted comparison ^{a,b}		ICC	
	Intervention	Control	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Overall composite outcome						
Proportion of patients with further emergency service call, ED attendance, emergency admission or death, ^c <i>n/N</i> (%)	870/2391 (36.4)	843/2264 (37.2)	OR = 0.956; <i>p</i> = 0.461	0.848 to 1.077	0.0009	0 to 0.0048
Primary outcome components						
Proportion of patients dying (any cause), ^d <i>n/N</i> (%)	147/2391 (6.1)	136/2264 (6.0)	OR = 0.994; <i>p</i> = 0.960	0.780 to 1.266	0	n/a
Proportion with further emergency admission, ^e <i>n/N</i> (%)	517/2391 (21.6)	475/2264 (21.0)	OR = 1.039; <i>p</i> = 0.595	0.903 to 1.196	0.0019	0.0002 to 0.0060
Proportion of patients with further ED attendance, ^f <i>n/N</i> (%)	463/2391 (19.4)	418/2264 (18.5)	OR = 1.067; <i>p</i> = 0.392	0.920 to 1.237	0.0070	0.0021 to 0.0151
Further ED attendances per patient, ^g mean (SD) [<i>n</i>]	0.2631 (0.6162) [2197]	0.2609 (0.7954) [2093]	Λ = 1.104 (<i>p</i> = 0.219)	0.943 to 1.293	0.0058	0.0017 to 0.0128
Further ED attendances/patient/day at risk, ^h mean (SD) [<i>n</i>]	0.0236 (0.1018) [2197]	0.0223 (0.0833) [2093]	Δ = 0.0011; <i>p</i> = 0.710 Δ_L = 0.0436; <i>p</i> = 0.413	-0.0045 to 0.0066 -0.0609 to 0.1481	0.0068 0.0100	0.0020 to 0.0150 0.0035 to 0.0204
Proportion of patients with further emergency service call, ⁱ <i>n/N</i> (%)	442/2391 (18.5)	493/2264 (21.8)	OR = 0.815; <i>p</i> = 0.006	0.705 to 0.943	0.0056	0.0018 to 0.0119
Further emergency service calls per patient, ^j mean (SD) [<i>n</i>]	0.2981 (0.7758) [2197]	0.3378 (0.7823) [2093]	Λ = 0.883; <i>p</i> = 0.049	0.780 to 1.000	0.0038	0.0011 to 0.0086
Further emergency service calls/patient/day at risk, ^k mean (SD) [<i>n</i>]	0.0204 (0.0641) [2197]	0.0245 (0.0814) [2093]	Δ = -0.0040; <i>p</i> = 0.071 Δ_L = -0.1354; <i>p</i> = 0.013	-0.0083 to 0.0003 -0.2418 to -0.0290	0.0043 0.0046	0.0011 to 0.0101 0.0013 to 0.0103

n/a, not applicable.

- a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender; and whether or not the index call was made out of (GP) hours.
 - b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a multiplicative event ratio (Λ) from negative binomial regression models for count data; and an additive group effect [Δ , in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(y + 0.001)$ in place of y] from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.
 - c Gender (*p* < 0.001); seasonality (*p* = 0.002); out of hours (*p* = 0.008).
 - d Gender (*p* < 0.001); age (*p* < 0.001); seasonality (*p* = 0.038).
 - e Site 3 (*p* = 0.001); out of hours (*p* = 0.010).
 - f Site 1 (*p* = 0.003); site 3 (*p* < 0.001); age (*p* = 0.036); age² (*p* = 0.020); seasonality (*p* = 0.002).
 - g Site 1 (*p* = 0.006); site 3 (*p* < 0.001); interaction between site 3 and group (*p* = 0.031); age (*p* < 0.001); age² (*p* = 0.001); seasonality (*p* = 0.004); days at risk (*p* < 0.001).
 - h For Δ : site 3 (*p* < 0.001); for Δ_L : site 1 (*p* = 0.013), site 3 (*p* < 0.001); seasonality (*p* = 0.003).
 - i Site 1 (*p* < 0.001); site 3 (*p* = 0.002); age (*p* < 0.001); seasonality (*p* = 0.004); out of hours (*p* < 0.001); recruitment point (*p* < 0.001).
 - j Site 1 (*p* < 0.001); age (*p* = 0.029); distance to ED (*p* = 0.001); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); days at risk (*p* < 0.001).
 - k For Δ : age (*p* < 0.001); distance to ED (*p* = 0.004); recruitment point (*p* = 0.012); out of hours (*p* = 0.025); for Δ_L : site 1 (*p* = 0.001); site 3 (*p* = 0.009); age (*p* < 0.001); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); seasonality (*p* = 0.008).
- Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 16 Primary outcome and components at 6 months analysed by treatment allocated

Primary outcome	Raw data		Adjusted comparison ^{a,b}		ICC	
	Intervention	Control	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Overall composite outcome						
Proportion of patients with further emergency service call, ED attendance, emergency admission or death, ^c <i>n/N</i> (%)	1701/2391 (71.1)	1592/2264 (70.3)	OR = 1.018; <i>p</i> = 0.789	0.895 to 1.157	0	<i>n/a</i>
Primary outcome components						
Proportion of patients dying (any cause), ^d <i>n/N</i> (%)	458/2391 (19.2)	419/2264 (18.5)	OR = 1.187; <i>p</i> = 0.094	0.971 to 1.451	0	<i>n/a</i>
Proportion with further emergency admission, ^e <i>n/N</i> (%)	1153/2391 (48.2)	1084/2264 (47.9)	OR = 1.001; <i>p</i> = 0.984	0.891 to 1.125	0.0052	0.0016 to 0.0113
Proportion of patients with further ED attendance, ^f <i>n/N</i> (%)	1079/2391 (45.1)	1021/2264 (45.1)	OR = 0.999; <i>p</i> = 0.986	0.888 to 1.123	0.0191	0.0085 to 0.0342
Further ED attendances per patient, ^g mean (SD) [<i>n</i>]	0.844 (1.392) [2380]	0.913 (2.738) [2257]	Λ = 0.810; <i>p</i> < 0.001	0.722 to 0.909	0.0144	0.0062 to 0.0261
Further ED attendances/patient/day at risk, ^h mean (SD) [<i>n</i>]	0.0169 (0.0907) [2380]	0.0144 (0.0686) [2257]	Δ = 0.0025; <i>p</i> = 0.292	-0.0021 to 0.0071	0.0041	0.0011 to 0.0098
			Δ_L = -0.0163; <i>p</i> = 0.711	-0.1024 to 0.0699	0.0264	0.0127 to 0.0451
Proportion of patients with further emergency service call, ⁱ <i>n/N</i> (%)	1046/2391 (43.7)	1046/2264 (46.2)	OR = 0.899; <i>p</i> = 0.076	0.799 to 1.011	0.0030	0.0006 to 0.0079
Further emergency service calls per patient, ^j mean (SD) [<i>n</i>]	1.136 (2.506) [2380]	1.251 (2.672) [2257]	Λ = 0.931; <i>p</i> = 0.076	0.860 to 1.007	0.0040	0.0012 to 0.0090
Further emergency service calls/patient/day at risk, ^k mean (SD) [<i>n</i>]	0.0125 (0.0363) [2380]	0.0172 (0.0599) [2257]	Δ = -0.0045; <i>p</i> = 0.002	-0.0073 to -0.0017	0.0029	0.0004 to 0.0069
			Δ_L = -0.1183; <i>p</i> = 0.010	-0.2079 to -0.0286	0.0036	0.0009 to 0.0086

n/a, not applicable.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (in days since start of study); Seasonality; indicators of gender; and whether or not the index call was made during out of (GP) hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a multiplicative event ratio (Λ) from negative binomial regression models for count data; and an additive group effect [Δ , in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(y + 0.001)$ in place of y] from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* < 0.001); recruitment point (*p* = 0.001); seasonality (*p* = 0.022).

d Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* = 0.001); site 3 (*p* = 0.003); interaction between site 1 and group (*p* = 0.023); distance to ED (*p* = 0.037).

e Site 1 (*p* = 0.019); gender (*p* = 0.003); age (*p* = 0.015); recruitment point (*p* = 0.031); distance to ED (*p* < 0.001).

f Site 1 (*p* = 0.009); site 3 (*p* < 0.001); out of hours (*p* = 0.020); distance to ED (*p* = 0.003); age² (*p* < 0.001); seasonality (*p* = 0.019).

g Site 3 (*p* < 0.001); interaction between site 1 and group (*p* < 0.001); gender (*p* < 0.001); seasonality (*p* < 0.001); recruitment point (*p* = 0.033); distance to ED (*p* < 0.001); days at risk (*p* < 0.001).

h For Δ : seasonality (*p* = 0.012); recruitment point (*p* = 0.018); site 3 (*p* = 0.037). For Δ_L : site 1 (*p* = 0.036); site 3 (*p* < 0.001); age (*p* < 0.001); distance to ED (*p* = 0.001); out of hours (*p* = 0.023); seasonality (*p* = 0.003).

i Site 1 (*p* = 0.016); site 3 (*p* = 0.035); age (*p* < 0.001); recruitment point (*p* < 0.001); out of hours (*p* < 0.001).

j Site 1 (*p* < 0.001); site 3 (*p* = 0.031); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); distance to ED (*p* = 0.001).

k For Δ : out of hours (*p* < 0.001); seasonality (*p* = 0.003); age (*p* = 0.006); recruitment point (*p* = 0.020). For Δ_L : out of hours (*p* < 0.001); distance to ED (*p* = 0.002); age (*p* < 0.001); recruitment point (*p* < 0.001).

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 17 Primary outcome and components at 1 month at each site analysed by treatment allocated

Primary outcomes	Site	Raw data		Site effects ^{a,b}	
		Intervention	Control	Main	Interaction
Proportion of patients with further emergency service call, ED attendance, emergency admission or death, <i>n/N</i> (%)	Site 1	481/1311 (36.7)	509/1337 (38.1)	OR = 1.136; <i>p</i> = 0.268	OR = 1.091; <i>p</i> = 0.583
	Site 2	172/540 (31.9)	152/433 (35.1)	<i>n/a</i>	OR = 0.864; <i>p</i> = 0.285
	Site 3	217/540 (40.2)	182/494 (36.8)	OR = 1.078; <i>p</i> = 0.582	OR = 1.333; <i>p</i> = 0.125
Proportion of patients dying (any cause), <i>n/N</i> (%)	Site 1	86/1311 (6.6)	79/1337 (5.9)	OR = 0.983; <i>p</i> = 0.941	OR = 1.173; <i>p</i> = 0.616
	Site 2	31/540 (5.7)	26/433 (6.0)	<i>n/a</i>	OR = 0.953; <i>p</i> = 0.862
	Site 3	30/540 (5.6)	31/494 (6.3)	OR = 1.048; <i>p</i> = 0.864	OR = 0.922; <i>p</i> = 0.830
Proportion with further emergency admission, <i>n/N</i> (%)	Site 1	271/1311 (20.7)	272/1337 (20.3)	OR = 1.001; <i>p</i> = 0.993	OR = 1.091; <i>p</i> = 0.644
	Site 2	104/540 (19.3)	88/433 (20.3)	<i>n/a</i>	OR = 0.935; <i>p</i> = 0.679
	Site 3	142/540 (26.3)	115/494 (23.3)	OR = 1.190; <i>p</i> = 0.278	OR = 1.257; <i>p</i> = 0.291
Proportion of patients with further ED attendance, <i>n/N</i> (%)	Site 1	256/1311 (19.5)	244/1337 (18.2)	OR = 1.311; <i>p</i> = 0.078	OR = 1.080; <i>p</i> = 0.711
	Site 2	79/540 (14.5)	63/433 (14.5)	<i>n/a</i>	OR = 1.006; <i>p</i> = 0.972
	Site 3	128/540 (23.7)	111/494 (22.5)	OR = 1.702; <i>p</i> = 0.002	OR = 1.065; <i>p</i> = 0.788
Further ED attendances per patient, mean (SD) [<i>n</i>]	Site 1	0.2642 (0.6574) [1230]	0.2337 (0.5449) [1258]	$\Lambda = 1.224$; <i>p</i> = 0.161	$\Lambda = 1.103$; <i>p</i> = 0.613
	Site 2	0.2017 (0.5017) [466]	0.1901 (0.4824) [384]	<i>n/a</i>	$\Lambda = 1.014$; <i>p</i> = 0.938
	Site 3	0.3174 (0.6042) [501]	0.3969 (1.374) [451]	$\Lambda = 2.050$; <i>p</i> < 0.001	$\Lambda = 0.777$; <i>p</i> = 0.240
Further ED attendances/patient/day at risk, mean (SD) [<i>n</i>]	Site 1	0.0217 (0.0775) [1230]	0.0201 (0.0753) [1258]	$\Delta = 0.0063$; <i>p</i> = 0.245	$\Delta = -0.0018$; <i>p</i> = 0.811
				$\Delta_L = 0.1933$; <i>p</i> = 0.058	$\Delta_L = -0.2631$; <i>p</i> = 0.851
	Site 2	0.0172 (0.0611) [466]	0.0138 (0.0661) [384]	<i>n/a</i>	$\Delta = 0.0034$; <i>p</i> = 0.600
				<i>n/a</i>	$\Delta_L = 0.0803$; <i>p</i> = 0.504
	Site 3	0.0341 (0.1648) [501]	0.0357 (0.1115) [451]	$\Delta = 0.0219$; <i>p</i> = 0.001	$\Delta = -0.0050$; <i>p</i> = 0.572
				$\Delta_L = 0.4744$; <i>p</i> < 0.001	$\Delta_L = -0.0955$; <i>p</i> = 0.563
Proportion of patients with further emergency service call, <i>n/N</i> (%)	Site 1	257/1311 (19.6)	320/1337 (23.9)	OR = 1.689; <i>p</i> < 0.001	OR = 0.855; <i>p</i> = 0.442
	Site 2	78/540 (14.4)	68/433 (15.7)	<i>n/a</i>	OR = 0.906; <i>p</i> = 0.585
	Site 3	107/540 (19.8)	105/494 (21.3)	OR = 1.449; <i>p</i> = 0.031	OR = 1.010; <i>p</i> = 0.966

TABLE 17 Primary outcome and components at 1 month at each site analysed by treatment allocated (*continued*)

Primary outcomes	Site	Raw data		Site effects ^{a,b}	
		Intervention	Control	Main	Interaction
Further emergency service calls per patient, mean (SD) [n]	Site 1	0.3130 (0.8111) [1230]	0.3831 (0.8763) [1258]	$\Lambda = 1.682$; $p < 0.001$	$\Lambda = 0.824$; $p = 0.278$
	Site 2	0.2318 (0.6508) [466]	0.2292 (0.5686) [384]	n/a	$\Lambda = 0.987$; $p = 0.933$
	Site 3	0.3234 (0.7920) [501]	0.3034 (0.6384) [451]	$\Lambda = 1.311$; $p = 0.078$	$\Lambda = 1.077$; $p = 0.721$
Further emergency service calls/ patient/day at risk, mean (SD) [n]	Site 1	0.0196 (0.0595) [1230]	0.0247 (0.0728) [1258]	$\Delta = 0.0073$; $p = 0.088$ $\Delta_L = 0.3538$; $p = 0.001$	$\Delta = -0.0068$; $p = 0.241$ $\Delta_L = -0.2094$; $p = 0.142$
	Site 2	0.0191 (0.0642) [466]	0.0174 (0.0733) [384]	n/a n/a	$\Delta = 0.0017$; $p = 0.736$ $\Delta_L = 0.0196$; $p = 0.873$
	Site 3	0.0238 (0.0743) [501]	0.0301 (0.1067) [450]	$\Delta = 0.0127$; $p = 0.012$ $\Delta_L = 0.2877$; $p = 0.020$	$\Delta = -0.0080$; $p = 0.248$ $\Delta_L = -0.1077$; $p = 0.525$

n/a, not applicable.

a Site effects are calculated using a base model which includes only indicators for group, site and interactions from the list of 'core' covariates, and are not adjusted for other covariates in that list. Main effects use site 2 as the reference category.

b The main and interaction effects reflect the variable under consideration; specifically, we report OR from logistic regression models for binary variables; a multiplicative event ratio (Λ) from negative binomial regression models for count data; and additive effects [Δ , in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(y + 0.001)$ in place of y] from linear models for measurement variables.

TABLE 18 Primary outcome and components at 6 months at each site analysed by treatment allocated

Primary outcomes	Site	Raw data		Site effects ^{a,b}	
		Intervention	Control	Main	Interaction
Proportion of patients with further emergency service call, ED attendance, emergency admission or death, n/N (%)	Site 1	933/1310 (71.2)	948/1337 (70.9)	OR = 1.214; $p = 0.101$	OR = 0.918; $p = 0.598$
	Site 2	372/540 (68.9)	289/433 (66.7)	n/a	OR = 1.103; $p = 0.476$
	Site 3	396/540 (73.3)	355/494 (71.9)	OR = 1.273; $p = 0.092$	OR = 0.976; $p = 0.901$
Proportion of patients dying (any cause), n/N (%)	Site 1	249/1311 (19.0)	270/1337 (20.2)	OR = 1.208; $p = 0.190$	OR = 0.710; $p = 0.073$
	Site 2	116/540 (21.5)	75/433 (17.3)	n/a	OR = 1.306; $p = 0.105$
	Site 3	93/540 (17.2)	74/494 (15.0)	OR = 0.841; $p = 0.333$	OR = 0.904; $p = 0.671$

continued

TABLE 18 Primary outcome and components at 6 months at each site analysed by treatment allocated (*continued*)

Primary outcomes	Site	Raw data		Site effects ^{a,b}	
		Intervention	Control	Main	Interaction
Proportion with further emergency admission, <i>n/N</i> (%)	Site 1	597/1311 (45.5)	620/1337 (46.4)	OR = 0.953; <i>p</i> = 0.663	OR = 0.959; <i>p</i> = 0.782
	Site 2	258/540 (47.8)	206/433 (47.6)	<i>n/a</i>	OR = 1.008; <i>p</i> = 0.950
	Site 3	298/540 (55.2)	258/494 (52.2)	OR = 1.205; <i>p</i> = 0.158	OR = 1.117; <i>p</i> = 0.537
Proportion of patients with further ED attendance, <i>n/N</i> (%)	Site 1	592/1311 (45.2)	584/1337 (43.7)	OR = 1.235; <i>p</i> = 0.062	OR = 1.170; <i>p</i> = 0.310
	Site 2	196/540 (36.3)	167/433 (38.6)	<i>n/a</i>	OR = 0.908; <i>p</i> = 0.467
	Site 3	291/540 (53.9)	270/494 (54.7)	OR = 1.920; <i>p</i> < 0.001	OR = 1.068; <i>p</i> = 0.717
Further ED attendances per patient, mean (SD) [<i>n</i>]	Site 1	0.813 (1.362) [1310]	0.790 (1.355) [1336]	Λ = 1.172; <i>p</i> = 0.067	Λ = 1.259; <i>p</i> = 0.055
	Site 2	0.561 (0.912) [535]	0.678 (1.083) [429]	<i>n/a</i>	Λ = 0.820; <i>p</i> = 0.058
	Site 3	1.204 (1.743) [535]	1.453 (5.296) [492]	Λ = 2.163; <i>p</i> < 0.001	Λ = 1.009; <i>p</i> = 0.945
Further ED attendances/patient/day at risk, mean (SD) [<i>n</i>]	Site 1	0.0155 (0.0859) [1310]	0.0117 (0.0461) [1336]	Δ = -0.0036; <i>p</i> = 0.426	Δ = 0.0046; <i>p</i> = 0.455
				Δ_i = 0.1424; <i>p</i> = 0.086	Δ_i = 0.0942; <i>p</i> = 0.405
	Site 2	0.0145 (0.1033) [535]	0.0153 (0.1117) [429]	<i>n/a</i>	Δ = -0.0008; <i>p</i> = 0.874
				<i>n/a</i>	Δ_i = -0.0690; <i>p</i> = 0.477
	Site 3	0.0231 (0.0886) [535]	0.0209 (0.0701) [492]	Δ = 0.0056; <i>p</i> = 0.297	Δ = 0.0030; <i>p</i> = 0.677
				Δ_i = 0.5620; <i>p</i> < 0.001	Δ_i = 0.0353; <i>p</i> = 0.793
Proportion of patients with further emergency service call, <i>n/N</i> (%)	Site 1	590/1311 (45.0)	635/1337 (47.5)	OR = 1.413; <i>p</i> = 0.002	OR = 0.763; <i>p</i> = 0.077
	Site 2	233/540 (43.1)	169/433 (39.0)	<i>n/a</i>	OR = 1.186; <i>p</i> = 0.195
	Site 3	223/540 (41.3)	242/494 (49.0)	OR = 1.500; <i>p</i> = 0.002	OR = 0.618; <i>p</i> = 0.008
Further emergency service calls per patient, mean (SD) [<i>n</i>]	Site 1	1.260 (2.917) [1310]	1.412 (3.163) [1336]	Λ = 1.684; <i>p</i> < 0.001	Λ = 0.883; <i>p</i> = 0.252
	Site 2	0.849 (1.497) [535]	0.839 (1.456) [429]	<i>n/a</i>	Λ = 1.011; <i>p</i> = 0.910
	Site 3	1.120 (2.183) [535]	1.173 (1.882) [492]	Λ = 1.398; <i>p</i> < 0.001	Λ = 0.944; <i>p</i> = 0.656

TABLE 18 Primary outcome and components at 6 months at each site analysed by treatment allocated (*continued*)

Primary outcomes	Site	Raw data		Site effects ^{a,b}	
		Intervention	Control	Main	Interaction
Further emergency service calls/patient/day at risk, mean (SD) [n]	Site 1	0.0118 (0.0308) [1310]	0.0177 (0.0604) [1336]	$\Delta = 0.0038$; $p = 0.165$	$\Delta = -0.0045$; $\rho = 0.228$
				$\Delta_L = 0.2946$; $p = 0.001$	$\Delta_L = -0.2524$; $\rho = 0.034$
	Site 2	0.0125 (0.0376) [535]	0.0139 (0.0620) [429]	n/a	$\Delta = -0.0015$; $\rho = 0.646$
				n/a	$\Delta_L = 0.1088$; $\rho = 0.286$
	Site 3	0.0144 (0.0460) [535]	0.0184 (0.0565) [492]	$\Delta = 0.0045$; $p = 0.170$	$\Delta = -0.0025$; $\rho = 0.573$
				$\Delta_L = 0.3329$; $p = 0.001$	$\Delta_L = -0.3508$; $\rho = 0.013$

n/a, not applicable.

a Site effects are calculated using a base model which includes only indicators for group, site and interactions from the list of 'core' covariates, and are not adjusted for other covariates in that list. Main effects use site 2 as the reference category.

b The main and interaction effects reflect the variable under consideration; specifically, we report OR from logistic regression models for binary variables; a multiplicative event ratio (Δ) from negative binomial regression models for count data; and additive effects [Δ , in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(y + 0.001)$ in place of y] from linear models for measurement variables.

TABLE 19 Secondary outcomes at index incident analysed by treatment allocated

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, p-value	95% CI	Estimate	95% CI
Conveyed to ED, ^{c,d} n/N (%)	All	1579/2420 (65.2)	1431/2284 (62.7)	OR = 1.082; $p = 0.205$	0.958 to 1.223	0.0512	0.0266 to 0.0832
	Site 1	779/1329 (58.6)	755/1352 (55.8)				
	Site 2	426/544 (78.3)	344/436 (78.9)				
	Site 3	374/547 (68.4)	332/496 (66.9)				
Referred to falls service by emergency service crew, ^{c,e} n/N (%)	All	204/2420 (8.4)	26/2284 (1.1)	OR = 51.730; $p < 0.001$	16.46 to 162.54	0.0400	0.0204 to 0.0652
	Site 1	100/1329 (7.5)	3/1352 (0.2)				
	Site 2	51/544 (9.4)	0/436 (0)				
	Site 3	53/547 (9.7)	23/496 (4.6)				

continued

TABLE 19 Secondary outcomes at index incident analysed by treatment allocated (*continued*)

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Left at scene without referral, ^f <i>n/N</i> (%)	All	547/2420 (22.6)	692/2284 (30.3)	OR = 0.686; <i>p</i> < 0.001	0.600 to 0.784	0.0375	0.0195 to 0.0612
	Site 1	379/1329 (28.5)	496/1352 (36.7)				
	Site 2	72/544 (13.2)	84/436 (19.3)				
	Site 3	96/547 (17.6)	112/496 (22.6)				
Key physiological indicators recorded at scene							
Respiratory rate, ^g <i>n/N</i> (%)	All	2318/2420 (95.8)	2165/2284 (94.8)	OR = 1.278; <i>p</i> = 0.090	0.963 to 1.695	0.0449	0.0215 to 0.0764
	Site 1	1281/1329 (96.4)	1288/1352 (95.3)				
	Site 2	495/544 (91.0)	386/436 (88.5)				
	Site 3	542/547 (99.1)	491/496 (99.0)				
Pulse rate, ^h <i>n/N</i> (%)	All	2319/2420 (95.8)	2173/2284 (95.1)	OR = 1.216; <i>p</i> = 0.186	0.910 to 1.624	0.0511	0.0254 to 0.0851
	Site 1	1296/1329 (97.5)	1309/1352 (96.8)				
	Site 2	489/544 (89.9)	378/436 (86.5)				
	Site 3	534/547 (97.6)	486/496 (98.0)				
Level of consciousness, ⁱ <i>n/N</i> (%)	All	2327/2420 (96.2)	2189/2284 (95.8)	OR = 1.058; <i>p</i> = 0.704	0.790 to 1.418	0.0016	0.0001 to 0.0055
	Site 1	1262/1329 (95.0)	1285/1352 (95.0)				
	Site 2	533/544 (98.0)	427/436 (97.9)				
	Site 3	532/547 (97.3)	477/496 (96.2)				
Length of episode of care (minutes), ^j mean (SD) [<i>n</i>]	All	196.8 (153.9) [2410]	192.8 (152.8) [2273]	Δ = 2.048; <i>p</i> = 0.645	-6.68 to 10.77	0.0180	0.0074 to 0.0336
	Site 1	185.4 (130.9) [1328]	181.8 (135.8) [1352]				
	Site 2	205.4 (211.0) [540]	209.1 (217.9) [431]				
	Site 3	216.2 (135.5) [542]	208.7 (122.4) [490]				

TABLE 19 Secondary outcomes at index incident analysed by treatment allocated (*continued*)

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Length of job-cycle time ^k (minutes), mean (SD) [<i>n</i>]	All	99.9 (41.9) [2416]	97.8 (43.9) [2277]	$\Delta = 1.685$; <i>p</i> = 0.174	-0.746 to 4.117	0.0167	0.0070 to 0.0308
	Site 1	97.5 (41.4) [1328]	97.2 (44.9) [1352]				
	Site 2	108.8 (47.7) [543]	102.2 (48.2) [435]				
	Site 3	96.9 (35.0) [545]	95.8 (36.3) [490]				

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender; and whether or not the index call was made during out of (GP) hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables and an additive group effect (Δ , in the same units as the dependent variable from linear models for measurement variables).

c Some patients were conveyed and referred (intervention group *n* = 34, control group *n* = 18). Statistically significant covariates are listed in further footnotes.

d Out of hours (*p* < 0.001); site 1 (*p* < 0.001); site 3 (*p* < 0.001); recruitment point (*p* = 0.001).

e Out of hours (*p* = 0.001); site 3 (*p* < 0.001); interaction between group and site 3 (*p* < 0.001); seasonality (*p* = 0.028); age (*p* = 0.032).

f Out of hours (*p* < 0.001); site 1 (*p* < 0.001); recruitment point (*p* = 0.001); site 3 (*p* = 0.028).

g Site 1 (*p* < 0.001); site 3 (*p* < 0.001); recruitment point (*p* = 0.009); age (*p* = 0.039).

h Site 1 (*p* < 0.001); site 3 (*p* < 0.001); age (*p* = 0.008); age² (*p* = 0.047).

i Site 1 (*p* < 0.001).

j Out of hours (*p* < 0.001); recruitment point (*p* < 0.001); site 1 (*p* < 0.001); seasonality (*p* = 0.002); age (*p* = 0.018).

k Distance to ED (*p* < 0.001); out of hours (*p* < 0.001); recruitment point (*p* < 0.001); seasonality (*p* < 0.001); site 1 (*p* < 0.001).

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 20 Secondary outcomes at 1 month analysed by treatment allocated

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Duration of subsequent inpatient episodes (nights in hospital, truncated) at 30 days, ^c mean (SD) [<i>n</i>]	All	2.25 (6.14) [2391]	2.10 (6.05) [2264]	$\Delta = 0.141$; <i>p</i> = 0.426	-0.207 to 0.490	0.0012	0.0001 to 0.0045
	Site 1	1.98 (5.64) [1311]	1.97 (5.73) [1337]				
	Site 2	2.51 (6.72) [540]	2.42 (6.77) [433]				
	Site 3	2.64 (6.67) [540]	2.17 (6.20) [494]				
Proportion with further reported fractures, ^d <i>n/N</i> (%)	All	98/2391 (4.1)	91/2264 (4.0)	OR = 1.002; <i>p</i> = 0.987	0.744 to 1.351	0.0049	0.0013 to 0.0114
	Site 1	38/1311 (2.9)	42/1337 (3.1)				
	Site 2	41/540 (7.6)	27/433 (6.2)				
	Site 3	19/540 (3.5)	22/494 (4.5)				

continued

TABLE 20 Secondary outcomes at 1 month analysed by treatment allocated (*continued*)

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Self-reported outcomes							
Quality of life SF-12 MCS, ^e mean (SD) [<i>n</i>]	All	39.80 (12.47) [447]	38.89 (12.16) [410]	$\Delta = 0.902$; $p = 0.282$	-0.744 to 2.547	0.0080	0.0004 to 0.0306
	Site 1	40.68 (12.85) [214]	40.30 (13.04) [217]				
	Site 2	37.76 (11.59) [126]	35.63 (10.69) [104]				
	Site 3	40.44 (12.53) [107]	39.26 (10.88) [89]				
Quality of life SF-12 PCS, ^f mean (SD) [<i>n</i>]	All	29.07 (9.97) [447]	29.40 (10.28) [410]	$\Delta = -0.495$; $p = 0.472$	-1.847 to 0.856	0	n/a
	Site 1	28.92 (9.81) [214]	29.62 (11.47) [217]				
	Site 2	28.13 (9.66) [126]	28.39 (7.99) [104]				
	Site 3	30.47 (10.55) [107]	30.03 (9.58) [89]				
Patient satisfaction QCM Technical, ^g mean (SD) [<i>n</i>]	All	62.82 (7.98) [563]	63.21 (8.16) [551]	$\Delta = -0.320$; $p = 0.506$	-1.265 to 0.625	0.0102	0.0018 to 0.0273
	Site 1	63.13 (7.45) [274]	63.79 (7.97) [295]				
	Site 2	61.37 (9.78) [163]	61.64 (8.63) [137]				
	Site 3	64.01 (6.07) [126]	63.58 (7.91) [119]				
Patient satisfaction QCM Interpersonal, ^h mean (SD) [<i>n</i>]	All	68.92 (8.66) [563]	68.04 (9.12) [551]	$\Delta = 3.132$; $p < 0.001$	1.587 to 4.678	0.0123	0.0017 to 0.0357
	Site 1	67.59 (9.63) [274]	67.94 (9.93) [295]				
	Site 2	71.18 (7.42) [163]	68.25 (8.51) [137]				
	Site 3	68.88 (7.28) [126]	68.06 (7.68) [119]				
Fall-related self-efficacy (fear of falling), ⁱ mean (SD) [<i>n</i>]	All	3.714 (3.040) [634]	3.815 (3.117) [600]	$\Delta = -0.055$; $p = 0.743$	-0.385 to 0.275	0.0062	0.0004 to 0.0224
	Site 1	3.836 (3.103) [302]	4.052 (3.273) [321]				
	Site 2	3.148 (2.866) [182]	3.194 (2.666) [151]				
	Site 3	4.156 (3.035) [150]	3.955 (3.133) [128]				

TABLE 20 Secondary outcomes at 1 month analysed by treatment allocated (*continued*)

Secondary outcome	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, p-value	95% CI	Estimate	95% CI
Proportion of patients who reported ≥ 1 further fall, ^j n/N (%)	All	413/621 (66.5)	409/589 (69.4)	OR = 0.723; $p = 0.025$	0.544 to 0.961	0.0102	0.0016 to 0.0285
	Site 1	211/296 (71.3)	215/314 (68.5)				
	Site 2	112/178 (62.9)	105/149 (70.5)				
	Site 3	90/147 (61.2)	89/126 (70.6)				

MCS, mental component summary; n/a, not applicable; PCS, physical component summary; QCM, Quality of Care Monitor.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; an additive group effect (Δ , in the same units as the dependent variable) from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Age ($p = 0.004$); site 1 ($p = 0.014$).

d Site 1 ($p < 0.001$); distance to ED ($p = 0.001$); gender ($p = 0.009$); age ($p = 0.020$).

e Site 1 ($p < 0.001$); site 3 ($p = 0.019$); age ($p = 0.037$); out of hours ($p = 0.038$).

f Age² ($p < 0.001$); out of hours ($p = 0.014$).

g Site 1 ($p < 0.001$); site 3 ($p < 0.001$).

h Interaction between site 1 and group ($p < 0.001$); interaction between site 3 and group ($p = 0.029$).

i Age ($p < 0.001$); out of hours ($p < 0.001$); site 1 ($p < 0.001$); site 3 ($p = 0.004$).

j Interaction between site 1 and group ($p = 0.016$).

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

arising from higher scores from intervention patients in site 2, but not in the technical aspect of patient satisfaction. There was evidence of a difference in the proportion of patients with further fractures at 6 months (but not at 1 month), with pronounced differences between groups across the three sites (approximately equal at site 1, higher in intervention patients in site 2 and lower in intervention patients at site 3).

Contamination

Table 22 shows the percentage of control patients subsequently attended within the trial period by a paramedic from the intervention group, < 5% at 1 month rising to about 15% at 6 months, with broadly similar rates at each site.

Patients ineligible for referral to the falls service

A subgroup of participants in site 1 were registered at GP practices without agreement for referral of patients to falls services and were less able to benefit from the full SAFER 2 intervention. Appendix 19 therefore presents the main results after excluding these patients. Results in these tables are consistent with those presented in the preceding sections.

Patterns of referral to the falls service

Sixty per cent of designated intervention group paramedics referred trial eligible patients to a falls service (Table 23). Paramedics referred up to 11 patients, although most made referrals once or twice. Patient age, sex, distance to ED did not appear to influence referral (Table 24); however, patients were more likely to be referred to falls services out of hours and less likely to be referred at site 1 than at the other sites (Table 25).

For most patients left at the scene by intervention group paramedics without referral to a falls service, no reason was identified. At site 1, 231 patients were not eligible for referral because, although they lived in the study catchment area, they were not registered with participating GPs. Other reasons for non-referral included refusal, not appropriate for referral or information that the patient was already under the care of a falls team or other service (Table 26).

TABLE 21 Secondary outcomes at 6 months analysed by treatment allocated

Outcome	Site	Raw data		Adjusted comparison ^{a,b}			
		Intervention (A)	Control (B)	Estimate, p-value	95% CI significance level	Estimate	95% CI
Routine data outcomes							
Duration of subsequent inpatient episodes (nights in hospital, truncated at 180 days), ^c mean (SD) [n]	All	11.18 (22.80) [2391]	11.62 (23.52) [2264]	$\Delta = -0.563$; $p = 0.403$	-1.884 to 0.757	0.0065	0.0021 to 0.0138
	Site 1	9.48 (19.29) [1311]	9.95 (19.95) [1337]				
	Site 2	14.48 (29.56) [540]	14.75 (30.01) [433]				
	Site 3	12.03 (22.52) [540]	13.42 (25.53) [494]				
Proportion with further reported subsequent fractures, ^d n/N (%)	All	228/2391 (9.5)	222/2264 (9.8)	OR = 1.449; $p = 0.015$	1.076 to 1.952	0.0172	0.0073 to 0.0318
	Site 1	93/1311 (7.1)	107/1337 (8.0)				
	Site 2	92/540 (17.0)	59/433 (13.6)				
	Site 3	43/540 (8.0)	56/494 (11.3)				
Self-reported outcomes							
Quality of life SF-12 MCS, ^e mean (SD) [n]	All	43.21 (12.57) [258]	42.82 (12.28) [241]	$\Delta = 0.463$; $p = 0.677$	-1.717 to 2.643	0	n/a
	Site 1	44.00 (12.50) [127]	43.45 (12.55) [144]				
	Site 2	41.84 (12.42) [53]	38.95 (12.77) [43]				
	Site 3	42.84 (12.84) [78]	44.24 (10.67) [54]				
Quality of life SF-12 PCS, ^f mean (SD) [n]	All	30.44 (11.33) [258]	31.88 (11.67) [241]	$\Delta = -1.300$; $p = 0.198$	-3.282 to 0.682	0	n/a
	Site 1	29.83 (11.29) [127]	32.36 (11.72) [144]				
	Site 2	31.61 (12.89) [53]	30.77 (12.79) [43]				
	Site 3	30.64 (10.31) [78]	31.49 (10.70) [54]				
Fall-related self-efficacy (fear of falling), ^g mean score (SD) [n]	All	4.547 (3.328) [341]	4.792 (3.393) [310]	$\Delta = -0.230$; $p = 0.368$	-0.729 to 0.270	0.0018	0 to 0.0219
	Site 1	4.643 (3.402) [167]	5.091 (3.477) [176]				
	Site 2	4.217 (3.254) [72]	4.010 (3.423) [60]				
	Site 3	4.624 (3.273) [102]	4.716 (3.090) [74]				

TABLE 21 Secondary outcomes at 6 months analysed by treatment allocated (*continued*)

Outcome	Site	Raw data		Adjusted comparison ^{a,b}			
		Intervention (A)	Control (B)	Estimate, p-value	95% CI significance level	Estimate	95% CI
Proportion with change in place of residence, ^h n/N (%)	All	32/319 (10.0)	19/280 (6.8)	OR = 1.497; p = 0.183	0.826 to 2.713	0.0032	0 to 0.0235
	Site 1	15/156 (9.6)	9/163 (5.5)				
	Site 2	5/73 (6.8)	4/50 (8.0)				
	Site 3	12/90 (13.3)	6/67 (9.0)				
Proportion of patients who reported ≥ 1 further fall, ⁱ n/N (%)	Site 1	110/159 (69.2)	105/172 (61.0)	OR = 1.423; p = 0.134	0.897 to 2.258	–	–
	Site 2	44/74 (59.5)	43/56 (76.8)	OR = 0.365; p = 0.019	0.157 to 0.848	–	–
	Site 3	74/96 (77.1)	44/68 (64.7)	OR = 1.835; p = 0.084	0.922 to 3.652	–	–

MCS, mental component summary; n/a, not applicable; PCS, physical component summary; QCM, Quality of Care Monitor.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a multiplicative event rate ratio (Λ , scalar) from negative binomial regression models for count data; an additive group effect (Δ in the same units as the dependent variable), from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Site 1 ($p < 0.001$); age ($p = 0.001$); age² ($p = 0.016$).

d Gender ($p < 0.001$); site 1 ($p < 0.001$); interaction between site 1 and group ($p = 0.022$); interaction between site 3 and group ($p < 0.001$).

e Age ($p = 0.024$).

f Age ($p < 0.001$).

g Age ($p < 0.001$); out of hours ($p = 0.043$).

h Age² ($p = 0.030$).

i Analysis of this variable reveals not only five significant covariates [age ($p = 0.004$); site 1 ($p = 0.010$); recruitment point ($p = 0.015$); distance to ED ($p = 0.014$); and gender ($p = 0.036$)] but also two major interactions [between site 3 and group ($p = 0.001$); and between site 1 and group ($p = 0.003$)]. In these unusual circumstances, we have omitted the misleading sub-row covering all participants and extended the three heterogeneous site-specific rows to summarise each site separately. Significant covariates are site 1, age ($p = 0.003$); site 2, distance to ED ($p = 0.001$); and seasonality ($p = 0.021$); and site 3, none.

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 22 Numbers of control group participants subsequently attended by an intervention group paramedic

Site	Count, n/N (%)	
	Within 1 month	Within 6 months
All	98/2264 (4.3)	356/2264 (15.7)
Site 1	59/1337 (4.4)	224/1337 (16.8)
Site 2	15/433 (3.5)	60/433 (13.9)
Site 3	24/494 (4.9)	72/494 (14.6)

TABLE 23 Falls referrals per intervention paramedic by site

Number of referrals to falls service ^a	Site 1 paramedics, n (%)	Site 2 paramedics, n (%)	Site 3 paramedics, n (%)	Total paramedics, n (%)
0	13 (33.3)	7 (26.9)	21 (52.5)	41 (39.0)
1	8 (20.5)	9 (34.6)	8 (20.0)	25 (23.8)
2	6 (15.4)	2 (7.7)	6 (15.0)	14 (13.3)
3	1 (2.6)	3 (11.5)	0 (0.0)	4 (3.8)
4	0 (0.0)	0 (0.0)	4 (10.0)	4 (3.8)
5	2 (5.1)	3 (11.5)	1 (2.5)	6 (5.7)
6	5 (12.8)	0 (0.0)	0 (0.0)	5 (4.8)
7	2 (5.1)	1 (3.8)	0 (0.0)	3 (2.9)
8	1 (2.6)	0 (0.0)	0 (0.0)	1 (0.9)
9	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
10	0 (0.0)	1 (3.8)	0 (0.0)	1 (0.9)
11	1 (2.6)	0 (0.0)	0 (0.0)	1 (0.9)
Total paramedics recruited to study	39	26	40	105

a When there was more than one intervention paramedic attended to a patient, we classified all such paramedics as referring the patient.

TABLE 24 Falls referrals at index call for intervention patients by personal and incident characteristics

Characteristics	Referred patients ^a		Not referred patients	
	n (%)	SD	n (%)	SD
Mean age (years)	83.88	7.15	82.14	8.11
Men	61 (35.1)	–	807 (35.3)	–
Women	113 (64.9)	–	1477 (64.7)	–
Mean distance to nearest ED (miles)	4.57	3.27	4.65	3.08
Out of hours	91 (52.3)	–	954 (41.8)	–
Site 1	81(46.6)	–	1352 (59.2)	–
Site 2	49 (28.2)	–	436 (19.1)	–
Site 3	44 (25.3)	–	496 (21.7)	–
Total	174	–	2284	–

a Excluding two control participants.

Comparison of outcome measures

To meet our methodological objective we compared the performance of the generic SF-12 with the mFES ('Fear of Falling'), a falls-specific outcome measure, in this population (Table 27 and Figures 3–6). The SF-12 mental component summary (MCS) and physical component summary (PCS) scores at 1 month both showed mean values at the lower end of the scale (lower scores indicate worse health), indicating mental and physical scores below the population norms. Similarly, the mFES at 1 month showed a clustering of scores at the lower end of the scale, indicating worse health. At 6 months, both scales again showed mean score values at the lower end of the scale, indicating worse health. There was, however, a small increase in MCS, PCS and mFES scores, indicating a slight improvement in patient-reported outcomes since the 1-month follow-up.

TABLE 25 Predictors of falls referral (full effects model)

Predictor	B	p-value	Exp(B)	95% CI for Exp (B)	
				Lower	Upper
Site 1	-0.448	0.015	0.639	0.446	0.917
Site 3	0.204	0.389	1.226	0.771	1.951
Gender	0.101	0.487	1.107	0.832	1.472
Age	0.079	0.025	1.082	1.010	1.160
Age ²	-1.702	0.085	0.182	0.026	1.264
Distance to ED	-0.041	0.137	0.960	0.910	1.013
Out of hours	0.456	0.001	1.578	1.202	2.072
Recruitment point	-0.217	0.426	0.805	0.472	1.373
Seasonality	0.187	0.067	1.206	0.987	1.474
Constant	-3.586	-	-	-	-

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 26 Reasons identified for non-referral of patients left at scene by intervention group paramedics

Reason	All, n (%)	Site 1, n (%)	Site 2, n (%)	Site 3, n (%)
No reason identified	377 (68.9)	241 (63.6)	60 (83.3)	76 (79.2)
Not registered with an eligible GP	68 (12.4)	68 (17.9)	0 (0)	0 (0)
Recorded by paramedic as				
Not appropriate for referral	38 (6.9)	25 (6.6)	4 (5.6)	9 (9.4)
Refused referral	31 (5.7)	20 (5.3)	6 (8.3)	5 (5.2)
Care plan in place	13 (2.4)	6 (1.6)	2 (2.8)	5 (5.2)
Already under a falls team	20 (3.7)	19 (5.0)	0 (0)	1 (1.0)
Total	547	379	72	96

TABLE 27 Comparison of outcome measures at 1 and 6 months

Outcome measure	n ^a	Mean score	SD
1 month			
SF-12 MCS	1160	43.84	9.36
SF-12 PCS	1160	35.03	8.01
mFES	1252	3.47	3.07
6 months			
SF-12 MCS	595	44.95	9.04
SF-12 PCS	595	38.23	7.72
mFES	663	4.55	3.45

PROM, patient-reported outcome measure.

a This number represents those for whom a PROM score could be generated at each time point.




FIGURE 3 Short Form questionnaire-12 items PCS and mFES scores at 1 month: correlation = 0.546.




FIGURE 4 Short Form questionnaire-12 items MCS and mFES scores at 1 month: correlation = 0.553.




FIGURE 5 Short Form questionnaire-12 items PCS and mFES scores at 6 months: correlation = 0.627.




FIGURE 6 Short Form questionnaire-12 items MCS and mFES scores at 6 months: correlation = 0.547.

There was moderate correlation between the SF-12 MCS and PCS scores and the mFES scores at both 1 and 6 months (range 0.55–0.63). Unusually, there were significant floor and ceiling effects for the condition-specific measure, mFES, indicating less sensitivity to detect small changes in patient outcomes than the SF-12 disease-specific measure.

Analysis of non-response

We have not analysed dissenters, as the National Information Governance Board withdrew consent to use their data.

Questionnaire response rates for those who did not dissent did not vary between trial arms (Table 28). Responders were slightly younger and the significant difference in recruitment point shows that the response rate improved over the course of the trial.

Harms

We defined a SAE as an emergency service call, ED attendance, emergency hospital admission or death identified as occurring within 2 days of the index incident. These are reported in Table 29. There is little difference in the rate of occurrence of these events between the intervention group and the control group, or in the total number of events between groups ($n = 331$ in intervention group vs. $n = 334$ in control group). During the trial there was no serious ARs in which a SAE was reported as possibly related to use of the intervention, including referral to a falls service.

TABLE 28 Comparison of characteristics of responders and non-responders to questionnaires at 1 month

Variables	Responders to 1-month questionnaire	Non-responders to 1-month questionnaire	Adjusted comparison	
			Estimate, p -value	95% CI
Proportion in intervention group, n/N (%)	668/1307 (51.1)	1752/3397 (51.6)	OR = 0.982; $p = 0.775$	0.864 to 1.115
Age (years), mean (SD) [n]	81.33 (8.06) [1303]	82.74 (8.00) [3384]	$\Delta = -1.408$; $p < 0.001$	-1.920 to -0.896
Proportion of females, n/N (%)	828/1307 (63.4)	2129/3396 (62.7)	OR = 1.029; $p = 0.675$	0.901 to 1.174
Proportion with index call out of hours, n/N (%)	503/1305 (38.5)	1462/3395 (43.1)	OR = 0.829; $p = 0.005$	0.728 to 0.945
Distance to ED (miles), mean (SD) [n]	4.87 (3.39) [1296]	4.66 (3.22) [3380]	$\Delta = 0.209$; $p = 0.050$	0.000 to 0.419
Recruitment point (days), mean (SD) [n]	246.1 (125.1) [1306]	211.8 (129.4) [3396]	$\Delta = 34.30$; $p < 0.001$	26.11 to 42.48

TABLE 29 Serious adverse events within 2 days following index incident

SAE	Intervention ($N = 2420$), n (%)	Control ($N = 2284$), n (%)	Total ($N = 4704$), n (%)
Emergency service call	101 (4.2)	117 (5.1)	218 (4.6)
ED attendance	78 (3.2)	92 (4.0)	170 (3.6)
Emergency admission	133 (5.5)	109 (4.8)	242 (5.1)
Death	19 (0.8)	16 (0.7)	35 (0.7)
Total	331 (13.7)	334 (14.6)	665 (14.1)

Chapter 5 Cost-effectiveness analysis

Results: summary of demographics

We included 4704 eligible participants in the analysis: 2681 from site 1, 980 from site 2 and 1043 from site 3. Of these, 2420 patients were randomly allocated to the intervention group and 2284 to the control group. Males accounted for 38.8% of the intervention group and 35.3% of the control group. The mean age was 82.5 years in the intervention group and 82.1 years in the control group.

Costs of Support and Assessment for Fall Emergency Referrals 2 trial intervention

The costs of the SAFER 2 trial intervention amounted to £41,854.00, which translates to a mean cost of £17.30 per eligible patient, as shown in *Table 30*. They include costs associated with training and implementation of the scheme, with a distinction made between costs associated with establishing the scheme (its 'roll-out' to other sites), which relate to development of the scheme and training in the respective sites, and the costs of enabling the delivery of the scheme in each area.

Resource use

The costs of health care used by trial participants, in the care of trial paramedics, were derived by multiplying items of resource use by the published unit costs in *Table 31*.

All but 49 patients were subsequently matched at SAIL or the HSCIC and, therefore, 4655 patients (2391 intervention, 2264 control) were included in the analysis of the anonymised linked data primary outcome. We combined two types of resource use. We calculated costs relating to emergency service calls, ED attendances and hospital stays for all matched participants, using routinely collected data for 1 month and 6 months following the index call, and then combined self-reported costs, obtained via questionnaires, over 1 and 6 months. *Table 32* shows that the main differences at 6 months are in the mean costs of initial hospital stays (£2523.27 in the intervention group vs. £2329.79 for control), and of subsequent hospital stays (£3982.21 for intervention vs. £4111.10 for control). At 6 months, mean total costs were £7386.39 in the intervention group and £7301.31 in the control group. Consistent with the clinical effectiveness analysis in *Chapter 4*, there are statistically significant differences in the mean costs of subsequent emergency service calls over 1 month, and of ED attendances over 6 months. These may warrant further investigation, although such differences are generally obscured when aggregated with larger (but statistically non-significant) differences in other elements of resource use – notably, hospital stays (see *Table 32* and its footnotes).

As questionnaire response rates were low at both 1 month and 6 months, self-reported resource use costs were used to impute unbiased estimates of such costs for all matched participants. *Table 33* summarises all resource costs. Again using the linear modelling method outlined in *Chapter 4*, the estimated intervention effect (adjusted for statistically significant covariates) at 1 month is £190.24 (95% CI –£13.83 to £394.31; $p = 0.068$), and at 6 months is £24.20 (95% CI –£468.01 to £516.40; $p = 0.923$); see *Table 33* and its footnotes.

TABLE 30 Intervention costs

Task	Site 1		Site 2		Site 3		Data sources	Total
	Basis of estimate	Actual cost	Basis of estimate	Actual cost	Basis of estimate	Actual cost		
Central set-up costs								
Intervention meeting in Derby to define common core and local optional components of intervention	One meeting (5 hours) x five medical consultants at £50.00 per hour; three falls team leaders at £40.00 per hour; one nurse consultant at £40.00 per hour; one research champion (paramedic) at £26.00 per hour; one ECP at £32.25 per hour; two clinical trainers at £32.25 per hour; one clinical effectiveness manager at £32.25 per hour	£2825.00	-	-	-	-	Ambulance service	£2825.00
Intervention training subgroup to design DVD	Four meetings (1 hour each) + 1 day development work (7 hours) x one ECP at £32.25 per hour; one nurse consultant at £40.00 per hour	£795.00	-	-	-	-	Ambulance service	£795.00
Production of DVD for paramedic training	Falls specialists involvement 10 hours at £40.00 per hour	£400.00	-	-	-	-	Ambulance service	£400.00

Task	Site 1		Site 2		Site 3		Data sources	Total
	Basis of estimate	Actual cost	Basis of estimate	Actual cost	Basis of estimate	Actual cost		
Local set-up costs								
Training of clinical educators	One session (4 hours) x two trainers at O/T rates (£32.25 per hour)	£333.00	Two sessions (including practice sessions) x one trainer at £187.50 daily rate	£375.00	Two sessions x one trainer at £187.24 daily rate	£374.00	Ambulance service	£1082.00
Clinical educators provision of training to paramedics	Four sessions (4 hours) x two clinical educators at O/T rates (£32.25 per hour)	£1032.00	Three sessions (7.5 hours each) x one trainer at £187.50 daily rate	£562.50	Seven sessions (7.5 hours each) x two trainers at £187.24 daily rate	£2621.00	Ambulance service	£4215.50
Others provision of training to paramedics	Two sessions (4 hours) x one ECP at £32.25 per hour; one nurse consultant at £40.00 per hour	£578.00	n/a	–	n/a	–	Ambulance service	£578.00
Training of paramedics	40 paramedics x 7.5 hours at £26.00 per hour (O/T included in the rate) Travel: 10 miles at £0.45 per mile	£8175.00	26 paramedics x 1 day training (12-hour day) at daily rate £262.00 (O/T included in the rate) (1.5 x) £21.83 Travel costs: 80 miles at £0.45 per mile	£7748.00	40 paramedics x 8 hours session + extra O/T payment of 4 hours = 12 hours at £21.00 per hour Travel: 40 x two zone 1 + two tube ticket travel cards at £8.00	£10,400.00	Ambulance service	£26,323.00
Cost of catering for paramedic training	Four sessions x £25.00 per group per session	£100.00	26 paramedics at £5.00 meal allowance per head	£130.00	40 paramedics at £8.34 per head	£334.00	Ambulance service	£564.00
Falls service leads delivering training to paramedics	Four sessions (2 hours) x one falls service lead at £40.00 per hour Travel: £4.00 per session	£336.00	n/a	–	Four sessions (1 hour each) x one falls service lead at £15.60 hourly rate Travel: four x two zone 1 + two tube ticket travel cards at £8.00	£94.00	Ambulance service	£430.00

continued

TABLE 30 Intervention costs (continued)

Task	Site 1		Site 2		Site 3		Total	
	Basis of estimate	Actual cost	Basis of estimate	Actual cost	Basis of estimate	Actual cost		
Partners and falls service set-up and implementation management costs	Six ambulance service-specific meetings (15 minutes each) x one operational support manager (£32.25 per hour); one head of audit and research (£49.90 per hour); one clinical tutor (£32.25 per hour); one lead nurse (£32.25 per hour). Two falls service meetings (60 minutes each) x three falls service team leader (£40.00 per hour). Two LIT meetings (15 minutes each) x AE consultant £50.00 per hour	£438.00	Six LIT and other meetings (15 minutes to discuss referral pathway) x two clinical trainers at £21.00 per hour; four falls leads at £15.60; one emergency bed service manager at £21.90 per hour; one medical director at £48.90 per hour; one area operations manager at £30.00 per hour; one team leader at £18.30 per hour	£335.00	Six LIT and other meetings (15 minutes to discuss referral pathway) x two clinical trainers at £21.00 per hour; four falls leads at £15.60; one emergency bed service manager at £21.90 per hour; one medical director at £48.90 per hour; one area operations manager at £30.00 per hour; one team leader at £18.30 per hour	£335.00	Ambulance service	£1108.00
	Two operational support managers (£32.25 per hour); one head of audit and research; one clinical tutor (£32.25 per hour); one lead nurse (£32.25 per hour); three falls service team leader (£40.00 per hour)							

Task	Site 1		Site 2		Site 3		Data sources	Total
	Basis of estimate	Actual cost	Basis of estimate	Actual cost	Basis of estimate	Actual cost		
Delivery costs								
RSO/paramedic time providing feedback and clinical support to paramedics	2 days per week at £14.70 per hour	£206.00	2 days per week x RSO day rate £187.50	£375.00	2 days per week at RSO day rate £252.00	£504.00	Ambulance service	£1085.00
Travel for ongoing clinical support	20 miles per week in lease car	£303.00	n/a	£0.00	n/a	–	Ambulance service	£303.00
Feedback forms from falls service to paramedics	91 referrals x 30 minutes per referral undertaken by one falls therapist at £30.00 per hour	£1365.00	n/a	–	100 referrals (not all were eligible) x 30 minutes by falls therapist at £15.60 per hour	£780.00	Ambulance service	£2145.00
Total intervention costs								
Intervention costs total	£16,886.00	£16,886.00	Intervention costs total	£9525.50	Intervention costs total	£15,442.00	–	£41,854.00
Number of eligible patients	1329	1329	Number of eligible patients	544	Number of eligible patients	547	–	2420
Cost per eligible patient	£12.71	£12.71	Cost per eligible patient	£17.51	Cost per eligible patient	£28.23	–	£17.30
DVD, digital versatile disk; n/a, not applicable; O/T, overtime; RSO, research support officer.								

TABLE 31 Published average unit costs

Health service resource	Average unit cost	Range of unit costs	Range of annualised equipment costs (3.5% discount)	Average annualised equipment cost (3.5% discount)	Source
Ambulance attendance (patient conveyed)	£230.00	–	–	–	NHS Reference Costs 2012–13 ¹²²
Ambulance attendance (patient not conveyed)	£174.00	–	–	–	NHS Reference Costs 2012–13 ¹²²
ED attendance cost	£115.00	–	–	–	NHS Reference Costs 2012–13 ¹²²
Non-elective inpatient long stay (4.0 days)	£2752.00	£531.00–9657.00	–	–	NHS Reference Costs 2012–13 ¹²²
Non-elective inpatient long stay (excess days)	£230.00	£67.00–556.00	–	–	NHS Reference Costs 2012–13 ¹²²
Non-elective inpatient short stay	£521.00	£133.00–1376.00	–	–	NHS Reference Costs 2012–13 ¹²²
Falls service referrals	£119.00	–	–	–	NHS Reference Costs 2012–13 (from 'Hospital at Home/Early Discharge Schemes') ¹²²
GP surgery visit (average 11.7 minutes)	£43.00	–	–	–	PSSRU (2013) ¹²³
GP telephone consultation (average 7.1 minutes)	£26.00	–	–	–	PSSRU (2013) ¹²³
GP home visit (average 23.4 minutes)	£110.00	–	–	–	PSSRU (2013) ¹²³
NHS Direct	£25.00	–	–	–	BMA (2011) ¹²⁴
Community nurse home visit (district nursing sister, district nurse)	£70.00	–	–	–	PSSRU (2013) ¹²³
Outpatient attendance (same as non-elective inpatient stay average short stay cost)	£467.00	–	–	–	NHS Reference Costs 2012–13 ¹²²
Nursing/residential care weekly (private) (average weekly cost)	£758.00	–	–	–	PSSRU (2013) ¹²³
Nursing/residential care weekly (local authority) (average weekly cost)	£796.00	£415.00–1331.00	–	–	PSSRU (2013) ¹²³
Local authority day care (average cost)	£40.00	–	–	–	PSSRU (2013) ¹²³
Extra care package for older people (average weekly cost)	£428.00	–	–	–	PSSRU (2013) ¹²³
Adjustable shower stools and chairs	£81.00	£14.00–148.00	£1.80–18.00	£9.90	PSSRU (2013) ¹²³
Perching stool with arms or back	£23.00	–	£2.70	£2.70	PSSRU (2013) ¹²³
Toilet frame and seat	£30.00	–	£3.60	£3.60	PSSRU (2013) ¹²³
Mobile shower chair	£55.00	–	£6.60	£6.60	PSSRU (2013) ¹²³

TABLE 31 Published average unit costs (continued)

Health service resource	Average unit cost	Range of unit costs	Range of annualised equipment costs (3.5% discount)	Average annualised equipment cost (3.5% discount)	Source
Bath step	£20.00	–	£2.50	£2.50	PSSRU (2013) ¹²³
Standard bath lift, two types	£317.00	£303.00–330.00	£36.00–40.00	£38.00	PSSRU (2013) ¹²³
Linked bed raisers, pair	£32.00	–	£3.80	£3.80	PSSRU (2013) ¹²³
Adjustable trolley	£34.00	–	£4.10	£4.10	PSSRU (2013) ¹²³
High-back chair	£121.00	–	£14.50	£14.50	PSSRU (2013) ¹²³
Variety of indoor and outdoor grab rails	£47.00	£3.40–91.00	£0.40–11.00	£5.70	PSSRU (2013) ¹²³
Walking sticks	£38.00	£22.00–54.00	£2.60–6.40	£4.50	PSSRU (2013) ¹²³
Commodes	£57.00	£29.00–85.00	£3.40–10.30	£6.85	PSSRU (2013) ¹²³
Wheelchair (per year)	£89.00	–	–	–	PSSRU (2013) ¹²³
Home care	£42.00	–	–	–	PSSRU (2013) ¹²³

BMA, British Medical Association.

TABLE 32 Health-care resource use over 6 months by group

Index	Intervention (n = 2391)			Control (n = 2264)		
	Observed	Total cost	Mean cost (SD)	Observed	Total cost	Mean cost (SD)
Routine data						
Proportion conveyed at index emergency services call n/N (%)	1562/2391 (65.3%)	£503,506.00	£210.58 (£26.66)	1424/2264 (62.9%)	£473,680.00	£209.22 (£27.06)
Proportion attending ED following index call n/N (%)	1419/2391 (59.3%)	£163,185.00	£68.25 (£56.50)	1311/2264 (57.9%)	£150,765.00	£66.59 (£56.79)
Proportion admitted to hospital following index call n/N (%)	906/2391 (37.9%)	–	–	843/2264 (37.2%)	–	–
Index hospital stays (nights in hospital, truncated at 1 month)	12735	£4,628,754.00	£1935.91 (£3096.33)	10857	£4,057,019.00	£1791.97 (£2954.50)
Index hospital stays (nights in hospital, truncated at 6 months)	18068	£6,033,134.00	£2523.27 (£5221.25)	15703	£5,274,639.00	£2329.79 (£4981.03)
Index falls service referrals	204	£24,276.00	£10.15 (£33.25)	26	£3,094.00	£1.37 (£12.68)
1-month follow-up routine data						
Subsequent emergency service calls ^a	662	£133,724.00	£55.93 (£153.64)	710	£143,420.00	£63.35 (£153.06)
Subsequent ED attendances ^b	591	£67,965.00	£28.43 (£68.94)	552	£63,480.00	£28.04 (£88.41)

continued

TABLE 32 Health-care resource use over 6 months by group (continued)

Index	Intervention (n = 2391)			Control (n = 2264)		
	Observed	Total cost	Mean cost (SD)	Observed	Total cost	Mean cost (SD)
Subsequent hospital stays (nights in hospital, truncated at 1 month) ^c	5375	£2,287,737.00	£956.81 (£2208.39)	4758	£2,042,487.00	£902.16 (£2168.19)
1-month follow-up self-reported questionnaire data						
GP surgery visit	331	£14,233.00	£21.50 (£53.51)	359	£15,437.00	£24.20 (£58.99)
GP telephone consultation	404	£10,504.00	£15.87 (£34.67)	408	£10,608.00	£16.63 (£37.50)
GP home visit	325	£35,750.00	£54.00 (£119.76)	355	£39,050.00	£61.21 (£139.02)
NHS Direct	169	£4,225.00	£6.38 (£26.20)	158	£3,950.00	£6.19 (£26.45)
Community nurse (district nursing sister, district nurse) visit	588	£41,160.00	£62.18 (£145.55)	554	£38,780.00	£60.78 (£150.56)
Outpatients	420	£196,140.00	£296.28 (£692.51)	382	£178,394.00	£279.61 (£697.57)
Social service equipment (adjustable shower stools, walking frames, etc.)	–	£1,686.00	£2.55 (£10.48)	–	£1,432.00	£2.25 (£9.29)
Further home care/carer	–	£588.00	£0.89 (£6.05)	–	£840.00	£1.32 (£7.32)
Local authority day care/ nursing/residential care weekly/extra care package for older people	–	£0.00	£0.00 (£0.00)	–	£0.00	£0.00 (£0.00)
1-month self-reported sub-total	662	£304,286.00	£459.65 (£809.09)	638	£288,491.00	£452.18 (£823.77)
6-month follow-up routine data						
Subsequent emergency services calls ^d	2712	£547,824.00	£229.12 (£505.58)	2825	£570,650.00	£252.05 (£539.13)
Subsequent ED attendances ^e	2014	£231,610.00	£96.87 (£159.99)	2063	£237,245.00	£104.79 (£314.41)
Subsequent hospital stays (nights in hospital, truncated at 6 months) ^f	26739	£9,521,458.00	£3982.21 (£6840.75)	26311	£9,307,534.00	£4111.10 (£6963.00)
6-month follow-up self-reported questionnaire data						
GP surgery visit	682	£29,326.00	£83.55 (£175.15)	602	£25,886.00	£80.14 (£161.92)
GP telephone consultation	700	£18,200.00	£51.85 (£106.10)	528	£13,728.00	£42.50 (£88.15)
GP home visit	493	£54,230.00	£154.50 (£372.27)	410	£45,100.00	£139.63 (£321.09)
NHS Direct	393	£9,825.00	£27.99 (£93.32)	270	£6,750.00	£20.90 (£73.78)
Community nurse (district nursing sister, district nurse) visit	1100	£77,000.00	£219.37 (£418.91)	1078	£75,460.00	£233.62 (£447.88)
Outpatients	864	£403,488.00	£1149.54 (£2204.78)	736	£343,712.00	£1064.12 (£2053.32)

TABLE 32 Health-care resource use over 6 months by group (*continued*)

Index	Intervention (<i>n</i> = 2391)			Control (<i>n</i> = 2264)		
	Observed	Total cost	Mean cost (SD)	Observed	Total cost	Mean cost (SD)
Social service equipment (adjustable shower stools, walking frames, etc.)	–	£1,985.00	£5.63 (£18.80)	–	£1,288.00	£3.99 (£12.85)
Further home care/carer	–	£462.00	£1.32 (£7.33)	–	£630.00	£1.95 (£8.85)
6-month self-reported subtotal	351	£594,506.20	£1693.75 (£2759.19)	323	£512,553.95	£1586.85 (£2527.89)
Intervention	2391	£41,352.00	£17.30 (£0.00)	2264	£0.00	£0.00 (£0.00)
Totals for 1 month	–	£8,154,797.55	£3410.62 (£3597.28)	–	£7,222,436.70	£3190.12 (£3526.34)
Totals for 6 months	–	£17,660,863.50	£7386.39 (£8619.93)	–	£16,530,160.95	£7301.31 (£8594.00)

a Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = -10.42$ (95% CI -20.04 to -0.79 ; $p = 0.034$).

b Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = -4.85$ (95% CI -10.81 to 1.11 ; $p = 0.110$).

c Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = 58.46$ (95% CI -66.86 to 183.78 ; $p = 0.361$).

d Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = -17.02$ (95% CI -47.01 to 12.97 ; $p = 0.266$).

e Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = -21.97$ (95% CI -40.89 to -3.06 ; $p = 0.023$).

f Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = -183.06$ (95% CI -577.72 to 211.61 ; $p = 0.363$).

Adapted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TABLE 33 Summary of health-care resource use over 6 months by group

Index	Intervention (<i>n</i> = 2391)		Control (<i>n</i> = 2264)	
	Total cost	Mean cost	Total cost	Mean cost
1-month follow-up				
Cost of index call up to 1 month	£5,361,085	£2,242.19	£4,684,558	£2,069.15
Routine costs by 1 month	£2,489,426	£1,041.17	£2,249,387	£993.55
Self-reported costs at 1 month (imputed)	£1,091,880	£456.66	£1,022,581	£451.67
Total costs by 1 month ^a	£8,942,391	£3,740.02	£7,956,526	£3,514.37
6-month follow-up				
Cost of index call up to 6 months	£6,765,465	£2,829.55	£5,902,178	£2,606.97
Routine costs by 6 months	£10,300,892	£4,308.19	£10,115,429	£4,467.95
Self-reported costs at 6 months (imputed)	£4,013,668	£1,678.66	£3,592,630	£1,586.85
Total costs by 6 months ^b	£21,080,025	£8,816.41	£19,610,237	£8,661.77

a Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = 190.24$ (95% CI -13.83 to 394.31 ; $p = 0.068$).

b Estimate of the intervention effect (in £, from fitted linear model, adjusting for statistically significant covariates) is $\Delta = 24.20$ (95% CI -468.01 to 516.40 ; $p = 0.923$).

Health-related quality of life

We estimated SF-6D utilities from questionnaires received at 1 and 6 months. However, the average proportion of missing data items was 5.5% at 1 month and 10.1% at 6 months; therefore, we imputed these missing items from the predicted SF-6D utilities adjusted for the number of reported deaths. The resulting imputed SF-6D utilities were higher in the control group at 6 months, although this did not achieve statistical significance. As the utilities were collected at 6 months, the QALYs gained by the intervention group was -0.0026 (95% CI -0.0066 to 0.0014), that is more QALYs were gained by the control group (Table 34).

Cost-effectiveness analysis

In summary, as there is no difference between groups in relation to resource use, the net cost of the intervention is £17.30. There is also no statistically significant difference in the number of QALYs generated between the two groups.

To test the extent to which this conclusion depends on the various assumptions we made, we conducted sensitivity analyses based on the upper and lower bounds of the 95% CI for both net cost (using the adjusted comparisons in Table 33) and net QALY (Table 35).

The sensitivity analysis highlights that the extent to which the SAFER 2 trial intervention can be regarded viewed as being cost-effective is inconclusive. This is further evidenced by the cost-effectiveness plane in Figure 7, generated from bootstrapping (5000 resamples). There is a net mean cost of £17.30, some clinically important statistically significant differences in the number of subsequent emergency service calls (at 1 month) and ED attendances (at 6 months), but no statistically significant difference in QALYs gained.

Conclusion

The relative cost-effectiveness of the SAFER 2 trial intervention is inconclusive, and further investigation is warranted to establish whether or not it represents value for money. It has resulted in important difference in the number of subsequent emergency service calls at 1 month and ED attendances at 6 months, but no difference between intervention and usual care in relation to QALYs gained.

TABLE 34 Mean SF-6D utilities over 6 months by group

Index	Group		Adjusted comparison	95% CI
	Intervention (A)	Control (B)		
1 month				
Derived SF-6D, mean (SD) [n]	0.5565 (0.1264) [480]	0.5547 (0.1278) [452]	$\Delta = -0.0011$; $p = 0.893$	-0.0173 to 0.0151
6 months				
Derived SF-6D, mean (SD) [n]	0.5885 (0.1377) [280]	0.5970 (0.1391) [257]	$\Delta = -0.0078$; $p = 0.509$	-0.0311 to 0.0155
SF-6D (imputed), mean (SD) [n]	0.4695 (0.2351) [2376]	0.4838 (0.2376) [2249]	$\Delta = -0.0107$; $p = 0.114$	-0.0240 to 0.0026
QALYs mean (SD) [n]	0.2093 (0.0709) [2375]	0.2133 (0.0713) [2245]	$\Delta = -0.0026$; $p = 0.202$	-0.0066 to 0.0014

TABLE 35 Sensitivity analysis

Parameter	Incremental cost of intervention	Incremental QALY	Cost per QALY gained
Baseline (95% CI)	£17.30 (-£475.01 to £509.40)	-0.0026 (-0.0066 to 0.0014)	Usual care is dominant. North-west quadrant of cost-effectiveness plane
Upper 95% bound of net cost	£509	-	£363,857. North-east quadrant of cost-effectiveness plane
Upper 95% bound of net QALY	-	0.0014	
Upper 95% bound of net cost	£509	-	Usual care is dominant. North-west quadrant of cost-effectiveness
Lower 5% bound of net QALY	-	-0.0066	
Lower 5% bound of net cost	-£475	-	£71,971. South-west quadrant of cost-effectiveness plane
Lower 5% bound of net QALY	-	-0.0066	
Lower 5% bound of net cost	-£475	-	Intervention is dominant. South-east quadrant of cost-effectiveness plane
Upper 5% bound of net QALY	-	0.0014	




FIGURE 7 Cost-effectiveness plane for the SAFER 2 trial intervention.

Chapter 6 Qualitative results

Introduction

The qualitative component of the study was designed to find out how the intervention was experienced both by patients and by paramedics and other stakeholders involved in delivering it. In particular, it aimed to identify factors which facilitate or hinder use of the protocol and referral pathway. The SAFER 2 trial intervention was hypothesised to work by improving the decision-making of paramedics in terms of safe non-conveyance and referral of older people who have had a fall to appropriate community-based services, leading to better outcomes for patients, reduced costs and improved paramedic skill sets. Through the protocol, pathway, training, support and feedback, the intervention provided a formalised framework for decision-making and referral for this patient group, in comparison with the informal approaches that characterise existing practice. This is hypothesised to make a difference by one or more of the following four mechanisms:

1. increasing paramedics' clinical knowledge of how to make appropriate non-conveyance decisions
2. increasing paramedics' knowledge of falls services and pathways for referral
3. increasing paramedics' confidence about making a non-conveyance decision and reducing anxiety about risk
4. increasing awareness of and likelihood that paramedics will consider non-conveyance and referral pathways as an option in appropriate cases.

In order for the intervention to make a difference to practice and to patient outcomes, a number of factors needed to be in place, including:

- effective referral pathways, and sufficient capacity in falls teams to respond to referrals
- training and support provided to paramedics that is appropriate and effective
- motivation on the part of paramedics to use the new protocol and referral pathway.

The qualitative component of the study aimed to examine the perspective of paramedics and other clinical stakeholders on these hypothesised mechanisms and necessary factors. It also aimed to gather data from patients to assess whether or not they were satisfied with the new intervention and whether or not, if relevant, the referral pathways were effective.

To address how the intervention was experienced by patients, we interviewed 58 patients attended by paramedics trained in the use of the new protocol from across the three sites, of whom 36 had been transferred to hospital, 12 were referred to a falls service and 10 remained at home without a referral (*Table 36*). Full details of sampling and data collection processes are given in *Chapter 4*. In this chapter, respondents are identified by identification numbers: numbers beginning with 1 are from site 1, those beginning with 2 are from site 2, and those beginning with 3 are from site 3, then participants in each site are numbered sequentially in the order in which they were interviewed. An 'H' after the number indicates that the patient was transferred to hospital, an 'F' shows that the patient was referred to a falls service and an 'N' indicates that the patient had neither outcome and was simply left at home. Findings from all three sites are written up together, structured by theme, with variations between sites noted when relevant. In the majority of cases, the respondent is the patient. In 18 cases, a family member with caring responsibilities also took part in the interview; where relevant in the text, quotations from carers are indicated by a 'c' appended to the end of the patient identification number.

TABLE 36 Number of patients participating in the qualitative interviews

Site	Transported to hospital	Referred to falls service	Neither transported nor referred	Total
Site 1	9	4	4	17
Site 2	15	2	2	19
Site 3	12	6	4	22
Total	36	12	10	58

As discussed in the qualitative data analysis section of *Chapter 3*, evidence is presented as anonymised quotations to support explanation of complex ideas, provide illustration and deepen understanding. Most quotations were selected to illustrate the majority view; however, some do convey exceptional viewpoints, as noted. In some cases, quotations were selected to illustrate complexity or ambiguity of perspective, as noted.

To address the experience of those delivering the intervention, we carried out focus groups and interviews with paramedics who were randomly allocated to the intervention group across the three sites at two time points: before the implementation of the intervention (24 paramedics) and after the completion of the live phase of the trial during which patients were recruited (25 paramedics). After the completion of the patient recruitment phase, we carried out focus groups and interviews with 31 stakeholders who had a role in delivering the intervention (e.g. ambulance service managers, training officers and staff of falls services). This chapter has been structured around themes related to the research questions and has been informed by the theoretical underpinning for the study. It brings together findings from the different staff groups and different sites, noting differences between sites when relevant. *Table 37* summarises the paramedic and stakeholder focus groups and interviews that took place.

How the intervention was experienced by patients

The fall in context

Every patient we interviewed described having a fall. Fifty-one respondents talked about the circumstances of the index incident and what they felt caused it. The majority (28) of the falls were described as taking place within the home, many of them trips and slips when moving around or using steps, and eight were cases of respondents sliding off chairs or the bed. A further 11 of the respondents described falls in the garden or outside the house, while hedge-trimming or watering plants, shovelling coal, putting the bins out, and so on. The remainder of the falls happened while the respondent was out and about. A minority of respondents described injuries resulting from the fall, ranging from fractures and being knocked unconscious to cuts and bruises.

Some falls were described as having a clear mechanical cause, such as 2–6H missing his footing on a stepladder while cutting the hedge. However, for many more, both inside and outside the home, the cause seemed unclear, as in the case of one respondent who fell in her kitchen:

All of a sudden, I never felt it, it just . . . just . . . like it just . . . my mind went blank. I just felt myself falling and when I looked around I was on the floor. So I thought to myself ‘silly bugger’.

3–21N

Many of the respondents explicitly stated that they did not know why they fell, while for a striking number the fall was something which ‘just’ happened: ‘my leg just gave way’ (1–1H); ‘I just dropped’ (1–7F); ‘I just slid down’ (1–2N), and so on.

TABLE 37 Summary of paramedic and stakeholder interviews

Site	Data collection	Participants
Site 1	Pretrial paramedic focus group	Four paramedics
	Pretrial paramedic interviews	Two paramedics
	Post-trial paramedic focus group	Four paramedics
	Post-trial stakeholder focus group	Four stakeholders working for ambulance service: head of research and audit; research project manager; operational support manager; and clinical quality manager
	Post-trial stakeholder interviews	Four stakeholders: ambulance service clinical team leader; and three falls team leaders
Site 2	Pretrial paramedic focus group	Four paramedics
	Post-trial paramedic focus group	Seven paramedics
	Post-trial stakeholder focus group 1	Three stakeholders working for ambulance service: R&D manager; clinical effectiveness manager; and a research assistant
	Post-trial stakeholder focus group 2	13 stakeholders: community staff nurse; acute trust head of strategy; team leader; community deputy team leader; team leader; district nurse team leader; district nurse; team leader district nurse; district nurse; district nurse team leader; district nurse team leader; district nurse team leader; and community sister
Site 3	Pretrial paramedic focus group 1	Six paramedics, one clinical tutor
	Pretrial paramedic focus group 2	Eight paramedics
	Post-trial paramedic focus group 1	Seven paramedics
	Post-trial paramedic focus group 2	Seven paramedics
	Post-trial stakeholder focus group 1	Three stakeholders working for ambulance service: research manager; deputy information manager; and clinical tutor
	Post-trial stakeholder focus group 2	Three stakeholders: senior ED clinical research nurse; falls co-ordinator and interim head of falls team; and falls lead physiotherapist
	Post-trial stakeholder interviews	Two stakeholders: ambulance service medical director and library housebound service manager

Others seemed to be seeking an explanation for why they fell:

But you know being in the middle of the night, waking up out of a sleep, you're not very steady, are you? I mean in the best of times.

3-9F

The falls happen like somebody's switching off the light switch according to the doctor. It's not a stumble or a trip, it-, this is just a fall which happens [clicks fingers], which is related to probably blood pressure or whatever. They are looking at the fact an epileptic fit may be involved here.

2-7H

In some cases this was expressed in terms of wanting to find an external cause for the fall:

But as I say the fact that I fell was simply something mechanical. I took a step back and I'd got a saucepan in one hand and a spoon in the other so I wasn't holding onto things as I usually do. It's the sort of thing that can happen to anybody, I think.

3-20N

Forty-three of the respondents described having had one or more previous falls, before the index incident, although not all of these incidents would have been attended by an ambulance. Some seemed habituated to falling:

Cause I've fell loads of times before.

1-11N

I have falls . . . I'm a faller.

3-18F

I can fall over at a drop of a hat, can't I?

1-5H

However, while some of those who had previously fallen had said that they had managed to do so without damage, and without professional intervention to get back on their feet, for others there was past history of significant damage sustained in falling:

I said, 'Oh my goodness, not again [wife's name], 'cause she'd broke the other hip 12 months ago.

1-4Hc

Thirteen of the respondents described having fallen again between the index incident and the interview. For the majority of them, this was a single fall, or a series of minor falls dismissed as 'little trips' (3-5H), but three described having had two or even three more major falls for which an ambulance had been called.

Patient experience of the ambulance service

Care

Respondents were asked to describe what happened when the paramedics arrived following the fall. In terms of immediate care or intervention, much of the largest group of responses ($n = 29$) concerned the patient being physically lifted, simply by hand, with the help of inflating equipment ($n = 7$) or in one case with the help of a towel. A small number of respondents compared the experience of being lifted by the paramedics with previous occasions when family, friends or neighbours had lifted them, sometimes resulting in hurt to themselves.

A minority of respondents described other forms of treatment administered by the paramedic, such as the dressing of wounds and the administration of pain relief, including morphine. In some cases, paramedics advised the patient to contact the GP for further care needs.

More generally, respondents talked about the way in which paramedics delivered care and built rapport with their patients. These comments were entirely positive across all three sites, and across all patient groups. The very process of examination and asking questions was described by some patients as having a positive effect:

I was very impressed, they gave me confidence, I felt safe with them, I felt they'd looked at what they needed to look at . . . yes they said to take painkillers and yes it was perfectly all right to have a drink because I might be a bit dehydrated by now and they said-, and I thought, yes now you've seen me I can do what I need to do.

1-6N

I thought they were very good. I mean they dealt with it efficiently, the questions gave me confidence . . . It was the confidence, you know you're going to get a bit of pain but it's the confidence they give you that you can work at it.

3–11H

Paramedics were described as treating respondents with respect, making them feel safe, and generally removing anxiety:

They put me at my ease lovely. I relaxed, I thought they were wonderful.

2–18H

Others used the words 'fantastic', 'marvellous', 'efficient', 'caring', 'kindly', 'respectful', 'kind', 'nice', 'brilliant and a good laugh' and 'the nicest of all the public services'. This respondent reflected the overall tone of the interviews:

I can't praise them enough.

2–16H

Assessment and examination

Respondents were asked to describe the processes of assessment and physical examination which they underwent after their fall. Recall of the details was frequently imprecise, not surprisingly, as the interviews were done some months after the incident, and stress or memory problems may have clouded their recall. Almost all showed patience with the assessment process and were positive in their reporting of it.

Respondents described in detail the various checks which paramedics went through:

They took me downstairs onto the ambulance and I was there for about half an hour. They were checking me over . . . and they were very very thorough and they were very good as well . . . Yes they checked my heart, blood pressure, they looked me over where I hurt myself and they were writing it all down which I suppose goes to the hospital. They written down all the drugs and they knew what they were.

3–16H

Thorough 100% medical . . . They phoned up the GP and checked my medication and took my blood pressure. They gave me a walking test and helped me . . . I was surprised at the thoroughness of the test. I just took it as the normal procedure.

3–7F

From what I can remember, she was as thorough as any doctor, with the things that she did, didn't she?

1–14F

One respondent was pleased with the manner and confidence with which the paramedics went through the assessment, without referring to a 'crib sheet':

Oh, they were natural with it. They weren't reading it up as we went, no . . . That would've affected my confidence if they had done.

1–10H

The only respondent who expressed reservations about the assessment and examination process was 1–6Hc, whose wife was attended after a fall:

Don't get out to a patient's house when everybody's frustrated and uptight and everything about somebody being–, being very, very poorly and then start playing about with computers and, you know, this sort of this nonsense . . . Initially I get a little bit worked up about the fact that they ask so

many what-, seems to me to be extraneous questions . . . But they seem to ask so many questions and you seem to be there forever and you think well, is all this relevant to taking somebody into hospital that's sick? You ring an ambulance in the first place because you think it's urgent that they need hospital attention, how can they burn all this time up?

1-6Hc

Those with previous experience of being attended by paramedics after a fall were asked to compare this experience (index incident) with any other falls. Three said they felt that the assessment as part of the SAFER 2 trial was more thorough, but the majority ($n = 11$) of respondents who made a comparison did not think there was a difference.

It was pretty much the same . . . Pretty consistent and they always use the same sort of procedures.

2-17H

Well I didn't know they were following a new process, it seemed exactly the same as the last time they came.

3-19N

Decision-making

A minority of respondents described the process of decision-making which was informed by the assessment and examination. These descriptions were almost exclusively in terms of the decision about whether or not to attend hospital, rather than about referral to the falls team. One respondent, describing how a decision was reached by the lead paramedic, seemed very positive about the use of a decision-making tool:

And then eventually the young lady said, 'I think you should go into hospital', and . . . they took her by ambulance to the hospital. It was a very straight cut simple sort of uncomplicated affair. Much different to a lot of previous times we've called the ambulance.

1-4Hc

Not all the respondents were so clear about the process of reasoning which took place:

I didn't know, they knew I had to get into hospital but I don't think they knew what I had done, I'm not sure about this, not that they missed anything, but I'm not sure, but they certainly were very sure that I had to go to hospital.

3-9H

A few respondents described how the paramedics drew them into the decision-making process, offering them the choice of whether or not to go to hospital: some decided to stay home, while others chose to go in for additional examinations:

They gave me an option to go for a check-up in the hospital and I didn't think it was necessary because the hospital would have just done the same treatment I received from the paramedics.

3-7F

I wanted to go, I wanted to make sure if there was anything broken they would fix it.

3-16H

A small number, although advised by paramedics to attend hospital, exercised their right to refuse, one of them in fairly strong terms:

They said it's not just that we want you to go to hospital; we don't know if a thorough enough examination has been done. We want to feel sure that everything has been done and we haven't

done it. PLEASE go. And I said 'On my head be it, I won't go and if anyone wants to know – you begged me to go . . . I don't want to go to that bloody awful hospital.'

3–18F

But more common was a sense that the respondent put their confidence in the clinical skills and decision-making of paramedics:

I mean, you rely on their judgement, don't you?

2–11H

Patient experience: those referred to a falls service

Twelve of the respondents had been identified as having been referred to a falls service by the attending ambulance crew. Respondents described being contacted within 2 weeks of referral at most, some within a week, and receiving a range of interventions, such as 6-week courses in falls avoidance at a day centre, home visits with exercise training, the installation of grab rails in the home, and equipment such as bed levers and perching stools.

Among the respondents who had been referred to the falls service and had received input, satisfaction was generally high:

Oh it's the best thing that's ever happened in my life.

1–8F

They're tops. They really are. Not one of them you could fault. Very, very helpful . . . They're marvellous people. And so patient with you.

1–12F

I had wonderful care from [local authority]. Absolutely remarkable.

3–4F

Being identified as suitable for a falls service did not necessarily equate to receiving the intervention. One respondent felt that the falls service offered nothing helpful, for another the intervention was stopped for medical reasons and a third was identified as suitable for referral to a falls service, but refused it altogether because:

I thought it was a waste of time really.

3–21N

Three respondents recorded as having been referred could not recall any contact, which may reveal a fault with the recording system or with the referral system.

Although respondents were forthcoming about their views on the falls services themselves, it was hard to disentangle any opinion on the *process* of being referred to the falls service as part of the SAFER 2 trial intervention. One respondent was unusual in commenting on this, indicating that she felt some confusion about the referral process:

Well, they just said to me when we–, 'You know we have to refer you now to the falls team' and so I said, 'Right'. But I wondered what's the falls team? What it was all about.

1–12F

More generally, it was not always clear whether the intervention the respondent was talking about was one which was delivered by a falls service associated with the SAFER 2 trial, or was actually a similar service

delivered by another team or accessed through a different route. Respondents themselves were in some cases unclear about this:

Well, I feel as though, yes, I mean I got this appointment at the falls, although the doctor, obviously, did it, but I think it was through the SAFER 2 that they did it.

1-7F

Oh, sorry . . . I don't know who arranged that then.

1-14F

Patient experience: those who attended hospital

Thirty-six of the respondents had been taken to hospital as part of the SAFER 2 trial. The care they received in hospital varied greatly, from being 'sent home with some paracetamol' (1-1H), to being admitted for surgery. Others were radiographed, stitched or were put in a cast, and a small number ended up with a lengthy stay in hospital.

The majority of respondents had a positive evaluation of their visit to hospital. Generally, this was expressed in terms of praise for the care provided by individual clinicians:

[The nurse] did fill me with confidence, you know, he was very, very good.

1-10H

The hospital staff, they were all absolutely wonderful, can't rate them enough.

1-15H

They were very comforting, you know, they gave me confidence in them and they were thoughtful and kind and helpful. I can't praise them enough, nor the nurse that did the stitching.

2-16H

But, yeah, the doctor was marvellous, the people who done the X-ray, great. No problems.

2-6H

Two respondents stated explicitly that the decision to go in to hospital had been the right one, the wife of 3-6Hc suggesting that it 'saved your life'.

However, for some the experience of attending hospital proved to be very frustrating. 2-6H was radiographed for suspected broken ribs, although even after the radiograph, 'they couldn't say for sure if there was a break or not'. Another respondent was reported by a carer as receiving even less in the way of an intervention:

We went into [ED] in the [hospital], we were there till quarter to three when a doctor eventually came to see her. She had a quick look at her, asked me to get her off the bed and walk her, and because she walked, to take her home . . . not even an X-ray to see whether she had broken her leg.

2-3Hc

Other negative aspects of attending hospital reported by respondents included delays in handover and admission:

We were all lined up, it was like a production line, waiting to be handed over to the doctor.

2-10H

Another respondent mentioned the difficulty of getting only a one-way journey in the ambulance:

But, the only thing I have against all that, they'll get you to the hospital but they don't see about you getting home again.

1-1H

Patient health and quality of life since fall

Respondents were asked how their health had been since the fall, leading to extensive discussions not just of health, but also of mobility and quality of life. Some patients made a comparison between their current state and their prefall status, while others compared how they were at the time of the interview with their immediate post-fall status; a few talked through the whole sequence of fall and recovery. A great range of experience was described, across all three outcome groups, and all three study sites. It was not possible to discern a clear pattern of difference between the three outcome groups, except that the group of patients who had been taken to hospital, as might be expected, included some people with more significant injuries.

A minority of respondents had very positive stories of recovery after the fall. These were most striking in the case of people who had sustained fractures or other significant injury in the fall:

Getting better all the time.

2–10H

Mum is amazingly better. She walks around the whole house on her own. When she came back from hospital she was actually moved with a hoist, she had no movement whatsoever and that has changed dramatically.

2–17Hc

Among those who had been referred to a falls service, a minority described very clearly how the intervention they were receiving was supporting an improvement in their well-being:

And now with the exercises . . . they show you to do, which I keep doing. I mean I couldn't have walked from here to the table at one time, now I can walk to the table . . . Walk round the house with confidence, you know . . . not frightened of falling.

1–12F

Well I'm walking better and I'm feeling better in myself because I've gotten over the fall. And I try to do little things. I haven't quite got enough balance at the moment . . . and that's what they're trying to work on.

3–9F

Although there were positive stories of recovery, a larger number of respondents talked about the increased limitations they experienced since the fall. This was a case of adapting what they did, taking more care and keeping clear of risky places such as the garden:

I've been very careful . . . No I haven't fallen since, I haven't done any gardening though you see and I've only done very little housework so I've tended not to do anything where I might fall.

1–6N

For some, having to accept these limitations was distressing:

I haven't been up that road on my own since that happened . . . I used to go out once a week. Yeah, I loved it. One leg wasn't like 100% but I could do it. I can't do it now, it breaks my heart . . . I definitely can't walk like I used to.

2–4H

For many of the respondents, it was a loss of confidence and a fear of falling that prevented them from going out and about:

So I'll freeze for fear of falling . . . Hoping to tell me legs where I want to go.

2–13H

No but I won't go out in the street on my own. I can't, I just can't, I just freeze. I go to that door, I can't even go out to the street door . . . Oh yeah. I used to go shopping and everything. I am frightened I am gonna fall.

3-7H

Some expanded on this theme, talking about the fall as something which had pushed them a little further along the inevitable journey into old age:

I have a loss of confidence, I am old but suddenly I feel much older. I feel a great sense of loss of freedom, of a loss of independence . . . Somebody said to me what do I most miss and I said 'being out in the air' I do stand at the back door, but I'm frightened of the stairs.

3-9H

I don't get up around quite so much as I did. And it's hard 'cause I now, which I now, which I never felt before, feel old. I am a bit nervous about doing physical things and where I'm going and should I take my walking stick and things like that.

3-15H

I'm walking now and I start crying. Then they say why and I say all this things I don't feel it before and it's my age.

3-1N

A large proportion of the respondents described their health, following the fall, as much the same as before. However, many of these then went on to report pain, reduced mobility or lack of confidence since the fall, presumably because they were thinking of these as being distinct from health. For example, 2-15H reported his health as being 'just the same' since the fall but went on to describe having to take tablets for pain relief and now finding it hard to get up after sitting on the settee.

The patient in context: other health and social care provision

Nearly half of the respondents described the networks of health and social care provision which supported their lives, much of which had direct relevance to preventing falls or managing their impact.

The most commonly reported form of support was an emergency alarm (also known as a Lifeline or Linkline), mentioned by 12 respondents across all three study sites. These alarms feature a pendant carried around the neck, which the user presses to call for help after a fall. For some respondents, they featured as part of the story of the index incident; for others, they were mentioned more as a source of reassurance.

In all three areas, many respondents reported having previously been provided, through different routes, with the type of support offered by the falls services associated with the SAFER 2 trial intervention. This provision included the fitting of grab rails and other adaptations to the home, equipment such as commodes and walking frames, and physiotherapy. Routes to accessing services varied; for one respondent, the GP functioned as an effective gateway:

Well apparently if you fall more than a certain number of times in a year, it has to be reported which my GP did, so I had a occupational therapist came out to see me and a physiotherapist. And they sorted out all the things, the aids that they thought I would need, namely a trolley and the second hand rail on the stairs, and grip rails everywhere, which I'm thankful for.

1-5H

One respondent described how previous falls had acted as a trigger to push her up the waiting list:

Then when I had that fall, she got a bit worried so it moved up a bit to the next one, I moved up, and that's how it's been going on, moving up and moving up to the next one 'til they found time to come down and assess me.

3–9F

An alternative route to provision was on hospital discharge, which was cited as a way of accessing both practical home care for a limited period, rehabilitation (described by one as 'walking stick instructions' 3–13H) and a range of useful equipment such as trolleys which contributes to falls prevention. These routes to access were not always straightforward, with two respondents describing having gone through the assessment process but then being found not eligible for support.

They did send an [occupational therapist] out to me, but she wasn't very good, I didn't end up getting anything . . . But my husband has–, well, we've been trying to get a rail for the bath for years, each time he's had an assessment, but we never get it.

1–8F

Other respondents described other forms of health or social care that contributed to their well-being at home, including specialist falls clinic, foot clinic, pain clinic and visits from the district nurse. A small number of respondents described support which they paid for themselves, outside the range of public sector or third-sector provision. In two cases this was people employed to help with cleaning, laundry and gardening; in another three cases it was privately purchased equipment, such as a specialist bed, while two reported paying privately for physiotherapy.

How the intervention was delivered in practice: paramedic and stakeholder views

Paramedic and stakeholder expectations of the new intervention (pre trial)

Both paramedics and stakeholders expected significant usage of the new referral pathway. Paramedics thought that the new referral protocol would be utilised widely if it was simple to complete.

I do think it can't be too difficult to do that referral, that's going to be the biggy. It's whether it takes an hour to sort something out. It needs to be a 5/10-minute phone call and 'yeah, no problem'.

Site 1 paramedic

If you make it a 20-page document people are not going to go with it.

Site 3 paramedic

Paramedics believed that patient care would be improved through the intervention, through avoidance of ED and through referral to a falls prevention service.

Once they can determine why the patient's having the falls, it could be just poor housing, rugs, things like that, slippers, could be medical, could be down to their medicines what they're taking, . . . it's got to build their confidence, it's got to do.

Site 1 paramedic

Paramedics from all three study sites expected personal gains, including an increase in job satisfaction resulting from offering a more complete care intervention:

You feel like you're finishing a job. You're not leaving it half done. You know, it's 'Doris has fallen again, we'll go back to her tomorrow' [but] 'Oh Doris is fallen, she's being referred, someone is looking after her'.

Site 1 paramedic

You'll feel like you've actually achieved something for the patient because you're not actually just moving them along on a conveyer belt to hospital. You'll feel that you're actually benefiting them.

Site 3 paramedic

With the exception of one, paramedics believed that the intervention would increase time on-scene. Paramedics from sites 2 and 3 expressed anxiety on how the control room would respond to an increase in the time on-scene.

If you're leaving people at home you need to do the checks to make sure they're safe to stay, that's the thing isn't it, you can't just do a quick blood pressure, a quick pulse . . . then go. You know, in the real world of course it isn't like that and I think a lot of people up in like control, they think it is like that and it's not . . .

Site 3 paramedic

The referral bit will take time. I don't know how control will feel about that.

Site 2 paramedic

Despite paramedics' worries, ambulance service managers in sites 1 and 3 did not feel that increases in time on-scene would be an issue.

I have no problem with paramedics spending an extra half an hour, three-quarters of an hour on-scene doing a thorough assessment that's appropriate for the patient. And at the end of the day, the call cycle time will still be quicker than going into ED.

Site 1 stakeholder

I don't necessarily think that time constraints on-scene would be a problem, I mean most paramedics are quite adamant that time on-scene is my time, I'm not going to get any pressure from, you know, control, if I'm with a patient then that's the time that I need so I don't think that that would have any impact on them referring or not referring.

Site 3 stakeholder

Concern was raised by falls team leaders at the number of referrals that would be received the trial, and the capacity of the services to deal with this, although one spoke of plans to bring in locum staff in order to cover increases in demand for their service.

Paramedics' and other stakeholders' experiences and views on context and existing practice

The majority of paramedics felt that current practice does not provide appropriate care to elderly patients who fall. Paramedics from all three of the ambulance services stated that current practice involved assessing patients for injuries, and depending on the outcome of the assessment and patient preference, conveying the patient to an ED, leaving the patient at home with no onward referral, or leaving the patient at home and referring to their GP. In two of the three sites (sites 1 and 3), there had been previous attempts to set up alternative pathways for elderly patients who had fallen, referring them to, for example, intermediate care teams, rapid response teams, district nurses and physiotherapy services, although these pathways had fallen out of use.

Overall, paramedics felt that they are able to make decisions regarding leaving patients at home and onward referrals, but they lack the referral pathways to properly undertake this. This was stated explicitly by a paramedic within site 1, with paramedics from sites 2 and 3 stating that they routinely make decisions regarding patient conveyance.

But this has been happening for years hasn't it. We've never had anything put in place. I'd say we've got more access to more places now than we've ever had, and it's still not enough . . . because nine times out of 10 we know exactly what that patient needs, and we just can't access it.

Site 1 paramedic

The process of deciding whether a patient stays at home or is referred to hospital is not always a straightforward one, and paramedics gave examples from existing practice where patients and paramedics disagreed. These were most striking in cases in which a patient wished to stay at home, but the paramedic felt, based on clinical assessment, that they should go to hospital; in these reported cases the paramedic's view prevailed, in part because of anxieties about risk:

I had a lady the other day, she clearly needed to go in and she was arguing with us for half an hour. I think we just wore her down in the end, but she had to go.

Site 2 paramedic

You bully them, you do. You bully them to some extent 'cause you think, I'm not risking it.

Site 2 paramedic

A number of paramedics discussed their anxieties about the risks of leaving a patient at home following a fall, worrying both about the patient, and about any comeback on the paramedic if it were to turn out to be the wrong decision:

I always think back at the end of the shift, oh God, I hope they're all right.

Site 2 paramedic

At the back of our minds, it's always gonna be . . . 'am I gonna be wrong, am I gonna get in trouble', you know, not that anybody wants to do anything wrong, not that we'd purposely do it. We're always trying to do the right thing, but if you miss something for whatever reason, and we're only human, . . . what are the ramifications gonna be?

Site 1 paramedic

Paramedics within sites 1 and 3 discussed the concept of 'covering their backs' when attending an elderly patient at home following a fall, in order to mitigate the risk of action against them if something were to go wrong. In some cases, this resulted in the patient being conveyed to hospital even if the paramedic could not see a clear clinical need:

Sometimes we convey the patient when quite clearly they didn't need to go in. In essence you feel like you need to cover your own backside because nobody else will.

Site 3 paramedic

Referrals to GPs were seen as problematic in all three sites: paramedics were not confident that the GP would respond to a referral (owing to not realising the urgency or simply not receiving the referral) or that they would make an onward referral if appropriate:

They [GPs] just don't seem to understand they have options as well, that they can refer people on. I just don't trust the GP to make that choice, to know that there are more choices. It's alright for us, we know we can refer onto social services . . .

Site 3 paramedic

Paramedics' professional status was seen as an important factor in the ability of paramedics to consider alternatives to conveying patients to the ED. One paramedic appeared to express frustration that the potential of paramedics to act as the gateway to a range of care services was untapped:

I went to a fall, and there was a district nurse who turned up just after I got there, . . . and the questions she got asked, and the ease of [the district nurse] getting that person respite – I was just gobsmacked at how easy it was. All those questions I could have just easily done . . . I think that paramedics have generally been seen as just take them to hospital, they haven't been seen as having any clue.

Site 1 paramedic

Pay, and the responsibilities attached to it, were mentioned by paramedics (although not managers) as an issue in sites 1 and 2:

Yeah, people are like, 'Well I don't get paid to [undertake additional decision-making to leave patients at home] so why should I do it?' So then everybody goes to hospital, so then you've got an A&E then that is absolutely chocca full of people who could possibly have been treated elsewhere by a different pathway.

Site 2 paramedic

Experience of implementing the Support and Assessment for Fall Emergency Referrals 2 trial intervention

Post trial, paramedics and stakeholders reflected on the process of implementing SAFER 2 trial as part of the research study. Paramedics stated three main reasons for volunteering to take part in the study: persuasion by a colleague; opportunity for personal development and learning new skills; and dissatisfaction with existing practice, in terms of both the lack of referral options available and the frustration of multiple attendances to patients for falls.

And you're taking somebody in and you know you're going to be straight back out because, you know, so if there's something that you can stop from going in so it's not a . . . waste of resources.

Site 2 paramedic

In sites 2 and 3, parallel developments of other pathways, while the SAFER 2 trial was under development, were felt to have led to confusing messages for paramedics:

It made it quite complicated from a local point of view, so I had 30 paramedics who were trained as intervention paramedics and 60 who weren't and trying to tell them, 'No, you can't choose the [SAFER 2] one yet,' 'Oh but we have seen adverts everywhere and we're getting letters from operations directors saying we must refer patients,' 'Well no you can't.' So it was quite complicated getting the right message out to the right staff, I think we got there but it was quite challenging to do that.

Site 3 ambulance manager

In all three ambulance services, the majority of managers felt that implementation of the SAFER 2 trial intervention had an impact on their role and workload during the initial phases of the study, when paramedic training was being carried out, but less so in the latter parts of the study:

[My involvement] all seemed to be very front loaded and then I think I lost my way, if I'm perfectly honest with it, during the latter-, I'd still come to the meetings and stuff, but once the training was there and everything was set up, it was almost like, right, sort of sit back and watch what happens.

Site 1 manager

Factors which enabled/encouraged the use of the Support and Assessment for Fall Emergency Referrals 2 trial intervention

According to the paramedics, usage of the new referral pathway was encouraged by its simplicity.

Overall, paramedics were positive regarding the suitability of the training they received. One paramedic compared the benefits of the SAFER 2 trial training to previous training for novel ambulance service interventions, which (as at the other two sites as well) seemed to be limited to e-mail or notice board updates.

Well, there hasn't been any training apart from this [the SAFER 2 trial training], . . . I mean this has been superb really, you know. You get to sit and relax and get trained on it but the new pathways are reading a piece of paper and acting on it. So, we're not trained on it [previous pathways].

Site 1 paramedic

Only one stakeholder commented on the value of the SAFER 2 trial training, and in positive terms:

The [paramedics] have now had some training that's been approved by some specialists so they feel very confident now at making those decisions with the backing of this triage tool.

Site 3 stakeholder

Paramedics in all three trial sites stated that they were aware of the clinical support available to them throughout the trial, although the majority did not actively seek clinical support at any point during the trial. Paramedics in all sites thought newsletters and e-mail updates were important to provide feedback and so maintain the motivation to refer:

[The newsletters were] a kind of confirmation that you are doing the right thing. I can't remember any specific points but I remember one and I don't know what it was, I can remember reading it and thinking, oh yeah, OK, I'm doing that so I'm doing the right thing.

Site 3 paramedic

Factors which hindered the use of the Support and Assessment for Fall Emergency Referrals 2 trial intervention

Both stakeholders and paramedics reported aspects of paramedic practice and the organisation of the ambulance service which hindered use of the SAFER 2 trial intervention. Paramedics from all three sites stated that taking a patient to the ED was often the easier option, especially when nearing the end of a shift.

You've got a lot of paramedics who, anything for an easy life – 'I'll go back take them to [ED] . . . Some of them want to do the best so they get off on time, or don't have a late finish, or least path of resistance.

Site 1 paramedic

Time of day, coupled with the patient's social situation, was mentioned by paramedics as affecting their ability to make a referral. Paramedics from all sites stated that they would feel uncomfortable leaving an elderly patient at home late at night without support, even if they were not injured. A small number of paramedics stated that a more immediate intervention than SAFER 2 trial was required by paramedics in order to assure confidence when leaving a patient at home.

I think if we had, as you said earlier, a direct phone number to a person that we knew was going to come out here in 2 or 3 hours, we'd maybe feel a little bit more confident to leave her there.

Site 2 paramedic

Paramedics said they often forgot to refer when on-scene, this being influenced by the time of day and the amount of time they had already been on the road.

At different times of the day . . . you're more tired, so you do occasionally leave the scene and then think, ah, I haven't done that [the SAFER 2 trial decision tool].

Site 2 paramedic

Ambulance service managers were aware of this, one suggesting that the use of the SAFER 2 trial declined over time:

I just feel that–, possibly 'cause it's over 12 months they did it– and . . . there wasn't any other trigger mechanism to trigger them to participate actively, participate. And bit by bit, they started just to forget.

Site 1 clinical tutor

In contrast to examples described in *Paramedics' and other stakeholders' experiences and views on context and existing practice* of paramedics persuading patients that they needed to go to hospital, they also described examples of patients, or those who cared for them, putting pressure on paramedics for a hospital transfer, even where paramedics might propose that a patient be left at home with a referral to a falls team. Even though the SAFER 2 trial decision tool had been used, paramedics were frustrated in taking the recommended step to refer to a falls service:

You feel pressure. When you attend elderly fallers, you face pressure from the families, their next door neighbours, . . . generally speaking that patient will take the advice of their next door neighbour and they'll go to the hospital. So we face a lot of pressures from around us to take them in. The family will say, 'Oh, I'd prefer her to go in and see a doctor. I'd rather she went in and had an X-ray,' even if you explain that they don't X-ray people routinely. So there's a lot of pressure around taking them in.

Site 2 paramedic

You're dealing with a frail, elderly group and often with little support in the community . . . added to which I think there is still the 'you call, we haul' mentality . . . call for an ambulance, you expect to go to hospital, that's what we'll do then.

Site 3 stakeholder

Obtaining consent from patients to refer to the falls team was seen as an issue in all sites:

Patients didn't want to be referred, we meet a lot of obstinate people who [laughs]–, 'Get me off the floor, get me bed, I don't want anything else from you and I'll be calling you back tomorrow night'.

Site 3 paramedic

Paramedics and managers also reported that it could be difficult to ascertain from patients if they already have involvement from such a service, or from an equivalent service:

If you went to a couple of falls to the same person and in the meanwhile they'd had an intervention of some description from a falls team, they might not immediately recognise that as being the intervention . . . It's really quite hard, I think.

Site 1 ambulance service manager

And we just feel, are you going on overkill almost, you know, if you start referring to another little team, you know, and they've already got this, that and the other coming in? You feel you're just repeating things.

Site 1 paramedic

Paramedics reported particular issues to do with referrals of patients living in nursing homes, where the referral to the falls service could be seen by nursing home staff as threatening:

If you referred somebody in their own private address, it was welcomed because you explained the package . . . But in the nursing home environment, you felt as though you were actually putting them on the spot . . . they were being scrutinised because they weren't providing the care or the facilities as what was available.

Site 2 paramedic

Paramedics in sites 1 and 2 saw lack of information regarding the role of the falls teams as an issue, specifically the time frames for initial contact and the interventions provided. Paramedics suggested that increased knowledge of the intervention would have positively influenced the number of referrals.

Evident across all interviews with ambulance service managers were some gaps in communication between them and the study paramedics. Although paramedics spoke of the importance of clinical support in the initial stages of a new pathway, most managers stated that they had not spoken directly to paramedics about the study and had not provided clinical support to study paramedics, with one manager in site 1 describing the difficulty of 'catching' a paramedic in order to discuss progress. Paramedics in all sites told a similar story.

[Asked if management were supportive of paramedics being involved in the trial] No one ever said anything.

Site 1 paramedic

I don't think they [ambulance service managers] knew who was taking part and who wasn't.

Site 2 paramedic

Impact of the Support and Assessment for Fall Emergency Referrals 2 trial intervention on paramedic practice

The Clinical Decision Flow Chart, used by paramedics on-scene, was felt to be a useful aide-memoire, but its impact on practice was variable. Paramedics from all three sites discussed a process of 'settling in' to the new assessment and referral process after their training. Many paramedics stated that it took time to get used to using the Clinical Decision Flow chart, but once they had, it was only used to remind them of specific details on-scene.

The more you used it, the less you needed to refer to it, or only certain points or anything. And if you weren't sure about something then you'd—, oh, just let me go and check on that. So once you got used to it, as I say, it was very straightforward to use.

Site 2 paramedic

I found it useful as a memory jogger.

Site 3 paramedic

For those based in sites 1 and 3, the flow chart did not contain any new procedures that were not already undertaken on-scene, apart from the referral to the falls team:

I think it [the flow chart] just reinforced what we already know. I guess . . . we're all used to dealing with falls.

Site 1 paramedic

In site 2, by contrast, the flow chart was felt to bring new assessment techniques into practice, to which paramedics responded positively:

I never did two blood pressures on people, when they were on the floor and when they got up. That didn't cross my mind. So you know that was definitely a learning curve, so that's given us extra skills.

Site 2 paramedic

Paramedics from all three sites felt that the SAFER 2 trial did not bring about significant changes in decision-making on-scene; paramedics felt that the decision to convey or leave a patient at home was a decision that they often made with some confidence but the new referral protocol 'reinforced what they already knew'.

Paramedics in sites 1 and 2 stated that the SAFER 2 trial intervention increased their confidence to leave a patient at home through two mechanisms: the reassurance that the falls team will contact the patient and the confidence in the assessment of a patient who has fallen. Paramedics within site 3 only stated the

former. Opinion of ambulance service managers was split between those who thought the confidence had increased, and those that believed that confidence was still an issue.

People are very confident, they are happy making those decisions [not to convey a patient to A&E] I'd say in general.

Site 3 manager

One of the things that we've highlighted as a possible reason [for not referring patients] is the lack of confidence, which is meaning that people aren't leaving patients at home that could be possibly left at home.

Site 1 operational support manager

Paramedics from all three sites thought that the new pathway increased time on-scene, mainly through three mechanisms: increased time undertaking patient assessment, increased time recording patients' assessment on PRF, and time spent on-scene making the referral to the falls team. Paramedics within site 3 discussed 'welfare checks' (a call from the control room to ensure the welfare of the paramedic after a certain period of non-communication), which the paramedics perceived as 'hurry-up checks'. One paramedic realised a benefit to spending more time on-scene with the patient:

I think for the length of time that we would have been there before SAFER 2-, 'cause you know, obviously when you actually do the referral it takes up to an hour, so then you get to see the patient sort of in their own environment and possibly doing things that you wouldn't normally because you'd probably just take them down the hospital.

Site 3 paramedic

Impact of the new pathway on falls services

Falls service representatives from all three sites commented on the appropriateness of the referrals made from paramedics; the majority of patients were thought to have been referred appropriately, although one falls service representative suggested some were not:

... there were some who maybe should have gone into hospital and actually not being kept at home.

Site 3 stakeholder

Representatives stated that the referrals they received through the SAFER 2 trial were a mix of patients already known to the falls services, and patients of whom the falls services had not previously been aware. Falls service team leaders from sites 1 and 3 reported that the newly identified patients referred through the SAFER 2 trial were often more frail than those normally receiving treatment from falls teams; a number of patients referred had subsequent A&E attendances and emergency service calls between the referral being made and contact with the falls teams.

The impact of paramedic referrals on the falls service differed between trial sites. In site 2, falls service representatives reported the impact as substantial in terms of paperwork, time and difficulties with onward referral, although the assessment and onward referral process was not thought to be very different from normal practice.

It was a bit of a put off really, and I have to be honest, when the referral come through ... you were sort of like dreading it a little bit.

Site 2 falls service representative

Representatives from sites 1 and 3, however, felt that the new referral pathway did not impact significantly upon the service, with representatives in site 3 commenting on the positive impact of the new pathway on their falls service.

I think the response time was quite interesting 'cause ours was a community team, we had a response time and within 1 to 10 days or quicker . . . What SAFER 2 did was . . . it made us firm up a few systems in our single point of access to flag up these fallers, patients, with the SAFER 2 you know, they go on the top, they need to be prioritised so we had to do a bit of work around that which was I think good for the service to kick in a sort of more urgent response, which we hadn't had before, we weren't an urgent sort of rapid response team. But after SAFER 2 we had that ability.

Site 3 falls service representative

A number of representatives discussed receiving repeat referrals for the same patient; this was generally felt to be useful, helping to identify gaps in current patient care.

At a personal level, falls service representatives from sites 1 and 3 were positive about their involvement in the study, stating that it was 'worthwhile' and 'enjoyable'. Despite the difficulties, representatives within site 2 were also positive regarding their involvement.

Keeping people well and healthy in their own homes for longer, so I'm glad we did take part. And I think it is something that, you know, the health board will look to develop further in future, but you know, we do need to make sure we get it right, and address some of the issues that you've all raised.

Site 2 falls service representative

Suggestions for modifications or developments of the Support and Assessment for Fall Emergency Referrals 2 trial intervention

Participants suggested modifications in four categories: the project set-up, training, patient assessment and referral processes/post referral. Suggested modifications differed by study site, reflecting the variety of issues experienced across the sites.

In sites 1 and 3, ambulance service managers suggested that key ambulance service staff should be involved as soon as possible in the set-up of a study, to make important decisions on behalf of the trust and to have input into the study design and data collection methods.

Falls service representatives from site 1 identified training as an issue, although their equivalents in the other two sites did not, perhaps reflecting the more limited involvement falls service representatives had in training in site 1. A falls service representative from site 1 suggested that training in the SAFER 2 trial should be carried out at regular intervals, in order to include and update more paramedics:

Going back to the training really and maybe just doing some supplementary training and . . . some scenarios with them . . . a quick half hour would have been enough just to say, 'Thank you for your referrals, thank you for your interest', . . . they could think of patients that they thought of, 'oh yeah well I maybe could have referred that lady but she seemed quite fit . . . she seemed quite able but she'd had a couple of falls'.

Site 1 falls service representative

Paramedics in sites 2 and 3 felt that the clinical decision flow chart needed to be simplified, to include less routine information, and more about the causes of falls:

I think on the flow chart if you're looking at possibly improving it maybe something a bit more . . . shorter version, maybe tick box type of thing.

Site 3 paramedic

On the flow chart, the current one, sometimes there was a bit too much information.

Site 2 paramedic

One site 3 paramedic felt that a smaller A5-sized flow chart would be better as it would fit into paramedics' pockets, to allow improved access on-scene. By contrast, other stakeholders across all sites did not comment on the flow chart being an issue.

Paramedics within both sites 1 and 3 felt that the decision to convey a patient would be easier to make if they had access to patients' records on-scene.

A copy of what their blood pressure is normally like, their normal observations so then . . . we can see it's . . . abnormal and then it's going to help our decision-making whether we're going to leave them or not.

Site 3 paramedic

In site 1, electronic patient report forms (ePRFs) were implemented during the study period, and ambulance service managers suggested that patient assessment could be improved if the ePRF included prompts to remind paramedics to consider a referral to the falls service. An assessment checklist on the ePRF was also considered beneficial by stakeholders.

. . . putting something on to the electronic Toughbook[® Panasonic, Newark, NJ, USA], create some sort of mandatory check. You know, . . . do not pass go unless you have notified the falls team, etc., whatever it might be, so there is some sort of check there built into the system and-, so it's programmed and you've identified there's a fall . . . for me, that way there, you will ensure that you've actually triggered the response that you want, to trigger it, it's going to be referred.

Site 1 clinical trainer

Other suggested improvements included increasing the number of people on telephone lines receiving referrals, increasing the capacity to refer (suggested by a site 2 paramedic), and providing an information sheet to referred patients to help them understand 'what will happen next' (site 3 paramedic). Improvements to the feedback process were suggested by paramedics from all three study sites, and by stakeholders in site 1. Paramedics from sites 1 and 3 argued for prompt feedback following a falls service referral, while a paramedic from site 2 explained that confirmation that the patient received the intervention would increase their confidence in the referral process:

We weren't sure what happened after we left. There was a presumption that they were going to be seen. It would have been nice to have some feedback to say that Doris has been seen yesterday and this has been done, so that you know it's working rather than presuming it's working.

Site 2 paramedic

Summary of qualitative findings

Patient experience

We used the qualitative phase of research to assess whether or not patients were satisfied with the new intervention and whether or not, if relevant, the referral pathways by which we had hypothesised that the SAFER 2 trial intervention would make a difference were effective.

Almost all the respondents were very satisfied with the processes of assessment and examination carried out by paramedics and tended to see them as part of a relationship based on empathy and care. The majority of those who were in a position to make a comparison between the SAFER 2 trial encounter with the ambulance service and a previous contact after a fall could not see much difference in the assessment and care delivered. When patients talked about having a say in decision-making, it was generally in terms of whether or not they would travel to hospital, not in terms of whether or not they wished to have contact with the falls service.

Even when patients were referred to the falls service, this did not necessarily lead to them receiving an intervention. Patients referred to falls services may in practice have come by many and various routes. Many of the patients in the SAFER 2 study were already tapping into a complex, and potentially confusing, network of community-based services, delivered by a range of providers (NHS, social services or third sector).

Although some patients reported good recovery after their fall, many had lasting ill-effects, with loss of confidence a major issue. Many worked out their own ways of reducing risk of a future fall, but at some cost to their quality of life.

For the vast majority of patients, the index incident was not the first fall they had experienced, although previous falls may not have resulted in contact with the ambulance service, and many went on to have a subsequent fall. Some seemed quite habituated to falling. Many falls were reported by patients as having an unspecified or unclear cause.

Paramedic and stakeholder experience

Through the qualitative component of the study, we aimed to examine the perspective of paramedics and other clinical stakeholders on the hypothesised mechanisms by which the SAFER 2 trial intervention was meant to work, that is, improving the decision-making of paramedics in terms of safe non-conveyance and referral of older people who have had a fall to appropriate community-based services, and necessary factors to allow this to happen.

The intervention was hypothesised to increase paramedics' clinical knowledge of how to make appropriate non-conveyance decisions and their knowledge of falls services and pathways for referral. In sites 1 and 2 (but not site 3), paramedics reported that a lack of knowledge around the role of the falls team and the structure of their intervention limited their ability to refer.

It was also hypothesised that it would increase paramedics' confidence about making a non-conveyance decision and reducing anxiety about risk and would increase the likelihood that paramedics will consider non-conveyance and referral pathways as an option in appropriate cases. There were mixed messages about whether or not confidence had been increased from both paramedics and managers. Paramedics emphasised that the decision-making process remained complex, with patients and carers having a voice which might contradict the paramedic and the protocol.

We also examined the three factors that we felt needed to be in place: effective referral pathways and sufficient capacity in falls teams; effective training and support for paramedics; and motivation on the part of paramedics to use the intervention. Before introduction of the intervention, falls service stakeholders also felt that the new pathway would impact upon their role greatly, although after the intervention stakeholders from two of the three sites reported no significant change to workload. Falls team managers in sites 1 and 2 stated that many referred patients declined the falls service intervention, although this was not mentioned as an issue at site 3. Paramedics in all three sites described reasons for not referring to falls services relating to patients' social situation and patient autonomy.

After implementation, paramedics reported that support for implementing the new intervention (i.e. paramedic training, flow chart and clinical support) had a positive effect on their ability to refer and make decisions about patient conveyance. The structured training on the SAFER 2 trial was felt to be more helpful than previous, more informal, approaches to introducing innovations, especially in sites 1 and 2. In sites 1 and 2, the SAFER trial processes such as the get up and go test and taking sitting and standing blood pressures were new to paramedics, while paramedics in site 3 already undertook these.

In terms of motivation, before introduction of the intervention, paramedics generally saw it as an effective way to deal with elderly patients who fall, and potentially an improvement on current practice. Current practice was felt to be geared towards conveyance to hospital even when not appropriate for the patient, although paramedics from all sites stated that they routinely make the decision to leave a patient at home following a fall. Paramedics expected to make significant usage of the new referral pathway, resulting in increased time on-scene, but also personal gains in terms of their skills and job satisfaction.

Throughout the transcripts, there was a lack of evidence of paramedics and managers having 'useful' contact with each other. Paramedics from all three study sites stated that they had limited contact with their managers during the study period. This is reinforced by the fact that paramedics were 'worried' about increased times on-scene, but managers did not see this as an issue.

Suggested modifications differed by study site, reflecting the variety of issues experienced across the sites. Paramedics from all three sites agreed that a 'simpler' intervention would have sufficed, with a less detailed flow chart. This was not commented on by ambulance service managers.

Limitations of qualitative methods

Patient experience

The sampling of patients for the interviews was heavily skewed (36 out of 58) towards those who had been transported to hospital. This may have meant that the patients interviewed were more likely, on average, to have had a major fall, and/or were more likely to have wanted to be transported to hospital.

Some patients found it difficult to recall details of the index incident many months after the event. When patients had had a number of falls, it was not always entirely clear which fall they were recalling and describing in response to the interviewer's questions.

It was hard to disentangle patients' views on the SAFER 2 trial assessment protocol specifically from their views on the more general process of examination and assessment.

Paramedic and stakeholder experience

It proved difficult to interview an equivalent cross-section of ambulance service managers within each trust, which led to a small variation in the set of questions asked in each site.

Focus groups and interviews were undertaken at different times between May 2011 (pre-trial focus groups) and March 2013 (post-trial focus groups). The time of year may have affected the responses provided by both stakeholders and paramedics, as service delivery pressures and priorities might have differed; therefore, it could possibly have affected the issues at the forefront of the interviewees' minds.

Each site varied in the basic training provided to paramedics, the set-up of the service, service delivery pressures and priorities at the time of the study.

Chapter 7 Discussion, conclusions and recommendations

Summary of key findings

There is a limited evidence base on prehospital alternatives to conveyance to hospital for older people who fall. The systematic review included 12 papers, only two of which were RCTs. Studies were of variable quality, with the majority being rated as low quality. The review identified limited and weak evidence regarding the most effective ways to deliver alternative treatments to older people who fall. A minority of the included studies reported benefits: reductions in subsequent emergency calls and hospital admissions when patients who had been left at home by their attending ambulance crew were referred retrospectively to a falls prevention service or when patients were attended by a paramedic with additional skills (ECP). However, in the majority of studies such outcomes were not explored. The review concluded that high-quality evidence from well-designed and rigorously conducted research is needed to inform service delivery.

In our RCT, we did not find any differences between trial arms in the composite primary outcome. However, we found consistent evidence of a small reduction in subsequent emergency service calls, from 21.8% of participants in the control arm to 18.5% in the intervention arm at 1 month, and from 46.2% to 43.7% at 6 months. Six per cent of patients had died in each group by 1 month and 18% by 6 months. Nearly one-third of patients had suffered a further emergency episode or death by 1 month, rising to over two-thirds by 6 months. Referrals to falls services were low: 8% in the intervention group. Overall, and at each site, fewer patients were left at the scene without any referral for further care in the intervention group than in the control group. We did not find clear evidence of any differences in other secondary outcomes related to processes of care, further injuries or self-reported quality of life, satisfaction or fear of falling.

Uptake of the intervention was variable. Sixty per cent of intervention group paramedics made at least one referral, but most of these made only one or two in total.

The intervention cost £17.30 per patient and generated some clinically important, statistically significant differences in the number of subsequent emergency service calls (at 1 month) and ED attendances (at 6 months), but no statistically significant difference in QALYs gained.

Although the SAFER 2 trial intervention is relatively low cost, this was not offset over 6 months by reductions in health-care resource use, neither did it generate improved health-related quality of life among patients allocated at random to the intervention relative to those controls allocated to usual care.

Retrieval of anonymised linked data outcomes was highly successful in this 'e-trial', with analysable outcomes available for 80% of those eligible. Such a high rate of inclusion produces findings that are generalisable to the whole-study population. Further investigation and analysis of these trial findings is warranted to fully describe the benefits and risks of this new approach.

Through the qualitative component of the study, we examined the perspective of patients, paramedics and other clinical stakeholders to help us better understand how the SAFER 2 trial intervention might work. In terms of increasing paramedics' clinical knowledge of how to make appropriate non-conveyance decisions, and their knowledge of falls services and pathways for referral, there were mixed messages, with paramedics in two sites reporting that a lack of knowledge around the role of the falls team and the structure of their intervention limited their ability to refer. Similarly, there were mixed messages in relation

to the intervention's role in increasing confidence to make non-conveyance decision and reducing in anxiety about risk. Paramedics emphasised that the decision-making process remained complex, with patients and carers having a voice which might contradict the paramedic and the protocol. Paramedics in all three sites described reasons for not referring to falls services relating to patients' social situation and patient autonomy.

Patients and carers in the intervention arm were highly satisfied with the care received, although generally they did not notice differences in practice compared with previous contacts. Interviews with patients and carers revealed that the scope for the intervention to make a measurable impact is limited by a number of factors. For various reasons, the pathway may not have led to a falls service intervention. Many of the patients in the SAFER 2 study were already tapping into a complex, and potentially confusing, network of community-based services, delivered by a range of providers (NHS, social services or third sector). The index fall is not necessarily an isolated incident, but more often part of a bigger and more complex pattern of falling, so it may not be straightforward to restore confidence and mobility after a fall, even with an intervention.

Strengths and limitations

The SAFER 2 study was a large-scale, multicentre randomised trial in the challenging setting of prehospital emergency care. We met our recruitment target for this frail group and achieved, through retrieval of anonymised linked routine data outcomes, a very high rate of inclusion: we reported our primary outcome for 80% of eligible patients, a much higher proportion than usually achieved.¹²⁵ Completion of 'designed data sets' concerning compliance with treatment and trial protocols, including reasons for non-referral, was low. This may reflect the context; we carried out this trial in the emergency care environment in which crews are under pressure to not only meet the care needs of each patient they attend but also to meet operational requirements which largely consist of time-based performance standards. As a general rule in trials in this setting, including the SAFER 2 trial, we try to minimise the collection of any new data from practitioners and rely instead on routinely collected information. In this way we maximise the inclusion of patients and the completeness of key data items and minimise intrusion into the clinical and operational context of the trial. In the SAFER 2 trial our primary and many secondary outcomes were based on routinely collected clinical and operational information and the completeness of these data sets is very high. Data that relied on additional paperwork by crews or patients had low completion rates and a higher rate of missing data items.

With low ICC, there is little dilution from observed to effective sample sizes, so we can be confident in findings reported. Statistical analyses have attempted to assess the effect of the intervention while allowing for variations across the three sites; analyses with statistically significant interactions therefore require careful interpretation.

The SAFER 2 trial tested a complex intervention. We were able to clearly define the components of that intervention with our research and clinical partners across sites. We delineated each component, common minimum standards, and aspects that could vary locally. This allows our trial results to be transparent and repeatable, in a field in which interventions often lack clarity.¹²⁶

Some aspects of the trial limit the usefulness of the results, in particular the fact that the trial took place in a context of changing practice and competing service innovations to care for this patient group. At each trial site we lost geographical areas, GP practices, stations, paramedics and patients because of the changing environment in which the trial took place. In site 3, an alternative referral innovation was introduced service-wide, incentivised by local commissioners. In site 1, only some GP practices were able to access the falls service which was taking part in our study, and patients not registered to those practices were ineligible for the emergency service call referral; in addition, many patients who had fallen were diverted to a clinical desk in the ambulance control room for an alternative (non-paramedic) response.

In site 2, a Frailty Programme was introduced in one of the initially participating health board areas, and, as a result, stations in this area were withdrawn from the trial. Six of 31 stations and 49 of 264 paramedics withdrew following randomisation, as presented in the CONSORT flow chart in *Chapter 4*. The withdrawal of stations was similar across trial arms, but withdrawal of paramedics was higher in the intervention arm. We do not know whether or not these withdrawals affected implementation of the intervention. Overall, we overcame these challenges which are perhaps inevitable in a pragmatic trial in a dynamic service environment. We completed the trial with balanced groups and achieved a sample size sufficient to detect important effects related to the intervention.

Uptake of the intervention was low, at 8% of patients in the intervention group. We do not know what the appropriate rate should be, as the protocol was intended to be applied across the spectrum of older people who had fallen, and many may have had a clinical need to be taken to the ED. Approximately one-quarter of patients were left at the scene of their call without a referral. Although this varied between groups and across sites, it is probably the best indication of the 'pool' of patients who might benefit from referral to a community-based service. The referral rate was variable between individual paramedics, with over one-third never making a referral at all during the study period. Referral rates did not vary by age, sex or distance to ED. Patients were less likely to be referred at site 1, although some patients were ineligible for the referral pathway at that site; and patients were more likely to be referred out of hours, but we do not have any further insight into why this was the case. In addition, we found no difference between trial arms on conveyance rate, suggesting that practice did not change significantly during the trial.

We recruited volunteer paramedics to participate in the trial, with the agreement of the three collaborating ambulance services. With slow recruitment we agreed to the introduction of £50.00 vouchers for each participating ambulance service, although, in practice, services used the payments in different ways. This reflects the practical considerations of carrying out multicentre experimental research in a live health service delivery environment. It does mean that the trial paramedics may not be representative of all paramedics and, therefore, we need to be cautious about interpretation of our findings.

It was not possible to undertake planned analyses of fall-related subsequent events, as data quality was found to be poor, with different codes and levels of completeness between hospitals and sites. It is well known that falls are poorly categorised, as, for example, slips, trips, stumbles or collapses, which is why we made the decision to analyse all events. Any effects on falls-related events would have also been evident in the all cause event proportions and rates, so this further exploration became less important once we had completed analysis of the primary outcome.

Implications (practice, policy and research)

With three participating ambulance services, 25 stations, > 200 paramedics and > 10 community-based falls services, and with outcomes available for 80% of eligible patients, we are confident that findings are generalisable to the UK setting and other similar health systems.

The SAFER 2 trial intervention was associated with a small reduction in the proportion of patients making further emergency service calls, and in the number of further calls made. However, this effect was not carried through to other parts of the emergency care system. There were no clear effects in secondary care, on ED attendances or emergency admissions. This finding is difficult to interpret, as we would have expected any difference in one part of the system to be reflected in other parts. It may be that patients in the intervention arm are more confident to self-manage if they fall again; however, the lack of impact at ED suggests that the effect may be restricted to those who were not conveyed to ED. Even a modest reduction in emergency service calls in this population is a success for the intervention in terms of prehospital care in which operational pressures are so high. Ambulance services may wish to implement this intervention in order to tackle rising demand but it will not make a huge difference to emergency service workload.

Uptake was lower than expected, with < 10% of trial eligible patients attended by intervention group paramedics referred to falls services. There may be several reasons for this, including a reluctance to change practice, a risk-averse climate within emergency prehospital care, patients who were unwilling to be referred or a lower than expected number of eligible older people who were not already under the care of a falls or similar frailty team. Some of these reasons reflect the population and context of the trial. Within the trial, all reasonable efforts were made to support a change in clinical practice; however, variability in referral rates suggests that uptake of the new referral pathway could have been higher. We do not know if higher use of the falls referral pathway would further reduce emergency service calls.

The smaller than expected effect may be because the pool of people who might benefit is smaller than previously expected, because preventative measures are more likely to be effective than an emergency intervention once a fall has occurred, or it may be because with many initiatives in place to target this group, there simply is little remaining scope for impact by emergency service paramedics. These results contrast with previous findings,⁶¹ perhaps because of the larger sample size and increased number of centres in the SAFER 2 trial.

Direct referral by ambulance crews to falls services is being implemented across the UK,⁷⁸ alongside many other new clinical pathways. In terms of tackling rising demand and managing workload, particularly of patients who present repeatedly to the emergency service, we can be confident that direct referral is not going to revolutionise care. We have not found evidence of harm and have found evidence of limited benefit to patients and the NHS.

Conclusions and recommendations for research

The complex SAFER 2 trial intervention, with a protocol for paramedics to assess older people who had fallen and refer those without need for immediate clinical care to community-based falls services, was inexpensive and safe. We did not find any effect on our primary outcome, although, when broken down into its components, there was a small reduction in the occurrence and rate of further emergency service calls. We did not find any evidence of improved quality of life, although some aspects of satisfaction were higher in the intervention group. Referral to falls services was lower than expected and variable between paramedics, although fairly consistent between sites. Fewer patients were left at scene in the intervention group by their attending ambulance crews without ongoing care than in the control group; however, other processes of care were unaltered.

Retrieval of anonymised linked data outcomes was highly successful in this 'e-trial', with analysable outcomes available for 80% of those eligible. Such a high rate of inclusion produces findings that are generalisable to the whole study population.

In terms of research priorities:

- Understanding implementation issues further is a research priority in this area for older people who fall and other groups for whom the default emergency service response is conveyance to hospital.
- Further investigation and analysis of outcomes retrieved through anonymised data linkage is warranted to fully describe the benefits and risks of this new approach.
- Further research regarding the performance of the SF-12 and the mFES for older people who fall is warranted, including:
 - exploring whether or not any differences exist in the either of the two scales between baseline and 6 months
 - comparing differences in outcomes between the intervention and the control sites at each time point
 - comparing the completion rates of the scales in the intervention and control sites

- examining differences between scale scores in those patients who went on to have a further fall (using routine ED data)
- examining whether or not either scale had any predictive ability in terms of predicting the likelihood of future falls.

In conclusion, the SAFER 2 trial findings indicate that ambulance services may introduce this new clinical pathway for patients without risk of harm, and with some limited impact on emergency service call workload. We did not find evidence of improved health outcomes for patients or reductions in emergency episodes across the NHS.

Acknowledgements

We would like to acknowledge and thank the following people who have contributed to the SAFER 2 study and the preparation of this report.

Bridie Evans (research officer), who contributed to the systematic review chapter and provided expertise on service user involvement throughout the trial.

Wai-Yee Cheung (senior lecturer in statistics and co-applicant), who contributed to the protocol development and provided statistical input to the early part of the trial, including randomisation.

Chris Garnsworthy (service user), who contributed to writing the Plain English summary and provided feedback about the results from a service user perspective.

Other acknowledgements

We would also like to thank the following people for their enthusiasm, support and hard work throughout the duration of the study.

To all the patients gave their time to complete questionnaires and interviews.

To all members of the falls services teams: Marie Ward; Kate Roberston; Jenny Fantom; Lyndsey Richardson; Lyn Messer; Paul Crank; Sonia Jones; Anna McCartney; and Rachel King.

To all the paramedics who participated, and to all the other ambulance service staff who supported the trial: Glyn Thomas; David Thomas; Caroline Hogan; Matthew Turner; Anne Spaight; Helen Stook; Sue Poucher; Joy Graves; Liz Team; Nadya Essam; Stacey Knowles; Michael Fuller; Darren Coxon; Martin Sheehan; Duncan Booth; Terry Doherty; Wendy Hazard; Melanie Edwards; and Jane Turner.

To National Institute for Social Care and Health Research, which provided invaluable support to the trial in Wales.

We thank the Trial Steering Committee members: Professor Chris Salisbury (chairperson), Chris Hartley-Sharpe, Greg Morgan, Dr Suzanne Wyatt; Professor Jonathan Benger and the Data Monitoring and Ethics committee members: Dr Kerenza Hood (chairperson), Dr Antony Bayer, Dr John Belcher and Colin Thomson.

To our colleagues at Swansea University, Moira Morgan, Anne Surman, Sherry Jenkins and Claire Evans, who have provided unfailing administrative support to the study.

To Rohan D'Silva and the Health Informatics Research Unit team.

Contributions of authors

Helen A Snooks (Professor of Health Services Research), chief investigator, led the development of the research question, study design, study implementation and report submission.

Rebecca Anthony (trial manager) managed all aspects of the conduct of the trial and supervised research staff, wrote the first draft of *Chapter 1* and contributed to all other report chapters.

Robin Chatters (research associate), site researcher; managed day-to-day conduct of the trial at Nottingham site, wrote the first draft of *Chapter 2* and contributed to all other report chapters.

Jeremy Dale (Professor of Primary Care), co-applicant, provided expertise in primary and emergency care research.

Rachael Fothergill (Head of Clinical Audit and Research), principal investigator, provided prehospital care research expertise.

Sarah Gaze (trial manager) managed all aspects of the conduct of the trial, supervised research staff and contributed to all report chapters.

Mary Halter (senior research fellow), co-applicant, acted as site academic lead, provided expertise in emergency prehospital care, care of older people who have fallen and local research management, and conducted the systematic review.

Ioan Humphreys (research officer) contributed to the economic analysis and wrote the first draft of *Chapter 5*.

Marina Koniotou (research officer) managed the day-to-day conduct of the trial at the London site and contributed to all report chapters.

Philippa Logan (Professor of Rehabilitation Research), co-applicant, provided expertise in falls prevention and research.

Ronan Lyons (Professor of Public Health), co-applicant, provided expertise in public health, emergency care, injury prevention and service evaluation (SAIL-linked data).

Suzanne Mason (Professor of Emergency Medicine), co-applicant, provided specialist advice related to emergency care research and supervision of site researcher.

Jon Nicholl (Professor of Health Services Research), co-applicant, provided expertise in health service research, HTA and emergency care.

Julie Peconi (trial manager), co-applicant, managed all aspects of the conduct of the trial and supervised research staff at the early stage of the trial.

Ceri Phillips (Professor of Health Economics), co-applicant, provided health economics expertise and analysis and led the economic analysis.

Judith Phillips (Professor of Gerontology), co-applicant, provided expertise in the social care and needs of older people and the design of the qualitative components of the trial.

Alison Porter (senior research officer) led the qualitative analysis, wrote *Chapter 6* and contributed to all other report chapters.

A Niroshan Siriwardena (Professor of Primary and Pre-Hospital Health Care), principal investigator, gave advice related to development of protocol and training.

Graham Smith (service user), TMG member, gave advice throughout trial.

Alun Toghill (service user), TMG member, gave advice throughout trial.

Mushtaq Wani (consultant geriatrician), co-applicant, provided clinical expertise in the care of the elderly and falls.

Alan Watkins (senior statistician), provided analysis of trial data and wrote *Chapter 4*.

Richard Whitfield (development manager), principal investigator; provided prehospital care research expertise.

Lynsey Wilson (research assistant) managed day-to-day conduct of the trial at the Wales site.

Ian T Russell (Professor of Clinical Trials), co-applicant, gave methodological advice, gave mentorship of trials management, provided statistical analysis and contributed to all report chapters.

All authors contributed to the preparation of the final report and approved the final version.

Publications

Darnell G, Mason SM, Snooks H. Elderly falls: a national survey of UK ambulance services. *Emerg Med J* 2012;**29**:1009–10.

Snooks H, Anthony R, Chatters R, Cheung WY, Dale J, Donohoe R, *et al*. Support and assessment for fall emergency referrals (SAFER2) research protocol: Cluster randomised trial of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community-based care. *BMJ Open* 2012;**2**:e002169.

Snooks HA, Anthony R, Chatters R, Dale J, Fothergill RT, Gaze S, *et al*. Paramedic assessment of older adults after falls, including community care referral pathway: cluster randomized trial [Published online ahead of print March 13 2017]. *Ann Emerg Med* 2017.

Data sharing statement

Data are stored within the Secure Anonymised Information Linkage (SAIL) databank at the Health Information Research Unit (HIRU) at Swansea University. All proposals to use SAIL data sets must comply with HIRU's information governance policy.

References

1. Bergeron E, Clement J, Lavoie A, Ratte S, Bamvita JM, Aumont F, *et al.* A simple fall in the elderly: not so simple. *J Trauma* 2006;**60**:268–73. <http://dx.doi.org/10.1097/01.ta.0000197651.00482.c5>
2. Sikron F, Giveon A, Aharonson-Daniel L, Peleg K. My home is my castle! Or is it? Hospitalizations following home injury in Israel, 1997–2001. *Isr Med Assoc J* 2004;**6**:332–5.
3. Tinetti ME, Baker DI, McAvay G, Claus EB, Garrett P, Gottschalk M, *et al.* A multifactorial intervention to reduce the risk of falling among elderly people living in the community. *N Engl J Med* 1994;**331**:821–7. <http://dx.doi.org/10.1056/NEJM199409293311301>
4. Tinetti ME, Speechley M. Prevention of falls among the elderly. *N Engl J Med* 1989;**320**:1055–9. <http://dx.doi.org/10.1056/NEJM198904203201606>
5. Tinetti ME. Preventing falls in elderly persons. *N Engl J Med* 2003;**348**:42–9. <http://dx.doi.org/10.1056/NEJMc020719>
6. Bergland A, Wyller TB. Risk factors for serious fall related injury in elderly women living at home. *Inj Prev* 2004;**10**:308–13. <http://dx.doi.org/10.1136/ip.2003.004721>
7. Gillespie L. Preventing falls in elderly people. *BMJ* 2004;**328**:653–4. <http://dx.doi.org/10.1136/bmj.328.7441.653>
8. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* 2006;**35**(Suppl. 2):ii37–ii41. <http://dx.doi.org/10.1093/ageing/afl084>
9. McKay C, Anderson KE. How to manage falls in community dwelling older adults: a review of the evidence. *Postgrad Med J* 2010;**86**:299–306. <http://dx.doi.org/10.1136/pgmj.2009.093468>
10. Scuffham P, Chaplin S, Legood R. Incidence and costs of unintentional falls in older people in the United Kingdom. *J Epidemiol Community Health* 2003;**57**:740–4. <http://dx.doi.org/10.1136/jech.57.9.740>
11. Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, *et al.* Interventions for preventing falls in older people living in the community 2012. *Cochrane Database System Rev* 2012;**9**:CD007146. <http://dx.doi.org/10.1002/14651858.CD007146.pub3>
12. NIH Senior Health. *Falls and Older Adults. Causes and Risk Factors*. 2013. URL: www.nihseniorhealth.gov/falls/causesandriskfactors/01.html (accessed 30 January 2013).
13. O’Loughlin JL, Robitaille Y, Boivin J-F, Suissa S. Incidence of and risk factors for falls and injurious falls among the community-dwelling elderly. *Am J Epidemiol* 1993;**137**:342–54.
14. Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988;**319**:1701–7. <http://dx.doi.org/10.1056/NEJM198812293192604>
15. Keene GS, Parker MJ, Pryor GA. Mortality and morbidity after hip fractures. *BMJ* 1993;**307**:1248–50. <http://dx.doi.org/10.1136/bmj.307.6914.1248>
16. Gibson MJ, Andres RO, Kennedy TE, Coppard LC, Kellogg International Work Group on the Prevention of Falls by the Elderly. The prevention of falls in later life. A report of the Kellogg International Work Group on the Prevention of Falls by the Elderly. *Dan Med Bull* 1987;**34**:1–24.
17. Tinetti ME. Factors associated with serious injury during falls by ambulatory nursing home residents. *J Am Geriatr Soc* 1987;**35**:644–8. <http://dx.doi.org/10.1111/j.1532-5415.1987.tb04341.x>

18. Keil DP, O'Sullivan P, Teno JM, Mor V. Health care utilization and functional status in the aged following a fall. *Med Care* 1991;**29**:221–8. <http://dx.doi.org/10.1097/00005650-199103000-00004>
19. Wolinsky FD, Johnson RJ, Fitzgerald JF. Falling, health status, and the use of health services by older adults. A prospective study. *Med Care* 1992;**30**:587–97. <http://dx.doi.org/10.1097/00005650-199207000-00002>
20. Tinetti ME, Williams CS. The effect of falls and fall injuries on functioning in community-dwelling older persons. *J Gerontol A Biol Sci Med Sci* 1998;**53**:M112–19. <http://dx.doi.org/10.1093/gerona/53A.2.M112>
21. Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA* 1989;**261**:2663–8. <http://dx.doi.org/10.1001/jama.1989.03420180087036>
22. Centers for Disease Control and Prevention. *About the National Health Interview Survey*. 2010. URL: www.cdc.gov/nchs/nhis/about_nhis.htm (accessed 31 January 2014).
23. Carr AJ, Higginson IJ. Are quality of life measures patient centred? *BMJ* 2001;**322**:1357–60. <http://dx.doi.org/10.1136/bmj.322.7298.1357>
24. Graham HJ, Firth J. Home accidents in older people: role of primary health care team. *BMJ* 1992;**305**:30–2. <http://dx.doi.org/10.1136/bmj.305.6844.30>
25. Tian Y, Thompson J, Buck D, Sonola L. *Exploring the System-wide Costs of Falls in Older People in Torbay*. London: The King's Fund; 2013.
26. Department of Health. *National Service Framework for Older People 2001*. URL: www.gov.uk/government/uploads/system/uploads/attachment_data/file/198033/National_Service_Framework_for_Older_People.pdf (accessed 10 April 2014).
27. Close JC. Interdisciplinary practice in the prevention of falls—a review of working models of care. *Age Ageing* 2001;**30**:8–12. http://dx.doi.org/10.1093/ageing/30.suppl_4.8
28. Aoyagi K, Ross PD, Davis JW, Wasnich RD, Hayashi T, Takemoto T. Falls among community-dwelling elderly in Japan. *J Bone Miner Res* 1998;**13**:1468–74. <http://dx.doi.org/10.1359/jbmr.1998.13.9.1468>
29. American Geriatric Society, American Academy of Geriatrics Society, Orthopaedic Surgeons Panel On Falls Prevention. Guideline for the prevention of falls in older persons. *J Am Geriatr Soc* 2001;**49**:664–72. <http://dx.doi.org/10.1046/j.1532-5415.2001.49115.x>
30. National Institute for Health and Care Excellence. *Clinical Practice Guidelines for the Assessment and Prevention of Falls in Older People 2004*. URL: www.nice.org.uk/nicemedia/pdf/CG021fullguideline.pdf (accessed 17 April 2014).
31. NHS Confederation. *The Ambulance Service Network. A Vision for Emergency and Urgent Care: the Role of the Ambulance Services 2008*. URL: www.nhsconfed.org/Publications/Pages/Avisionfor.aspx (accessed 15 April 2014).
32. Winkin V, Pinckaers V, D'Orio V, Ghuysen A. Overcrowding estimation in the emergency department: is the simplest score the best? *Crit Care* 2013;**17**:258. <http://dx.doi.org/10.1186/cc12196>
33. The King's Fund. *Are Accident and Emergency Attendances Increasing 2013*. URL: www.kingsfund.org.uk/blog/2013/04/are-accident-and-emergency-attendances-increasing (accessed 15 April 2014).
34. Close JC, Lord SR, Antonova EJ, Martin M, Lensberg B, Taylor M, et al. Older people presenting to the emergency department after a fall: a population with substantial recurrent healthcare use. *Emerg Med J* 2012;**29**:742–7. <http://dx.doi.org/10.1136/emered-2011-200380>

35. Close J, Ellis M, Hooper R, Glucksman E, Jackson S, Swift C. Prevention of falls in the elderly trial (PROFET): a randomised controlled trial. *Lancet* 1999;**353**:93–7. [http://dx.doi.org/10.1016/S0140-6736\(98\)06119-4](http://dx.doi.org/10.1016/S0140-6736(98)06119-4)
36. Kawano T, Nishiyama K, Anan H, Tujimura Y. Direct relationship between aging and overcrowding in the ED, and a calculation formula for demand projection: a cross-sectional study. *Emerg Med J* 2014;**31**:19–23. <http://dx.doi.org/10.1136/emered-2012-202050>
37. The Fire Defense Agency. *Result of Research of the Acceptance of the Medical Institution in Emergency Conveyance in 2009*. In MolAa, Editor Communications Tokyo: The Fire Defense Agency; 2010.
38. Bridges J. *Listening Makes Sense: Understanding the Experiences of Older People and Relatives Using Urgent Care Services in England*. London: City University; 2008. URL: www.staff.city.ac.uk/~rbbc318/lms/resources/final_report.pdf (accessed 15 April 2014).
39. Miller S. *Unnecessary and Avoidable Hospital Admissions for Older People*. Canberra, ACT: Department of Health and Ageing; 2003.
40. Department of Health (DH). *Taking Healthcare to the Patient: Transforming NHS Ambulance Services*. London: DH; 2005. URL: <http://rcp.info/Portals/11/Future/NHS%20EMS%20Policy%20Recommendation2005.pdf> (accessed 7 May 2015).
41. Department of Health. *Our Health, Our Care, Our Say*. London: HMSO; 2006.
42. NHS England. *Transforming Urgent and Emergency Care Services in England*. Leeds: NHS England; 2013. URL: www.nhs.uk/NHSEngland/keogh-review/Documents/UECR.Ph1Report.FV.pdf (accessed 8 April 2014).
43. Weiss SJ, Chong R, Ong M, Ernst AA, Balash M. Emergency medical services screening of elderly falls in the home. *Prehosp Emerg Care* 2003;**7**:79–84. <http://dx.doi.org/10.1080/10903120390937148>
44. Snooks HA, Halter M, Close JC, Cheung WY, Moore F, Roberts SE. Emergency care of older people who fall: a missed opportunity. *Qual Saf Health Care* 2006;**15**:390–2. <http://dx.doi.org/10.1136/qshc.2006.018697>
45. Marks PJ, Daniel TD, Afolabi O, Spiers G, Nguyen-Van-Tam JS. Emergency (999) calls to the ambulance service that do not result in the patient being transported to hospital: an epidemiological study. *Emerg Med J* 2002;**19**:449–52. <http://dx.doi.org/10.1136/emj.19.5.449>
46. Close JC, Halter M, Elrick A, Brain G, Swift CG. Falls in the older population: a pilot study to assess those attended by London ambulance service but not taken to A&E. *Age Ageing* 2002;**31**:488–9. <http://dx.doi.org/10.1093/ageing/31.6.488>
47. Goldberg RJ, Zautcke JL, Koenigsberg MD, Lee RW, Nagorka FW, Kling M, *et al*. A review of prehospital care litigation in a large metropolitan EMS system. *Ann Emerg Med* 1990;**19**:557–61. [http://dx.doi.org/10.1016/S0196-0644\(05\)82189-4](http://dx.doi.org/10.1016/S0196-0644(05)82189-4)
48. Department of Health, College of Emergency Medicine. *A&E Clinical Quality Indicators 2011*. URL: www.imperial.nhs.uk/patients/ourstandards/emergency-care-quality-indicators/index.htm (accessed 26 March 2014).
49. Porter A, Snooks H, Youren A, Gaze S, Whitfield R, Rapport F, *et al*. 'Should I stay or should I go?' Deciding whether to go to hospital after a 999 call. *J Health Serv Res Policy* 2007;**12**:32–8. <http://dx.doi.org/10.1258/135581907780318392>
50. Snooks HA, Kearsley N, Dale J, Halter M, Redhead J, Foster J. Gaps between policy, protocols and practice: a qualitative study of the views and practice of emergency ambulance staff concerning the care of patients with non-urgent needs. *Qual Saf Health Care* 2005;**14**:251–7. <http://dx.doi.org/10.1136/qshc.2004.012195>

51. London Ambulance Service NHS Trust. *GP Newsletter February 2013 Edition 10 NHS 111 – Final Stages*. London: London Ambulance Service NHS Trust; 2013. URL: www.londonambulance.nhs.uk/health_professionals/gp_information/gp_news.aspx (accessed 10 April 2014).
52. East Midlands Ambulance Service NHS Trust. *Falls Team Shortlisted for National Award*. Nottingham: East Midlands Ambulance Service NHS Trust; 2013. URL: www.emas.nhs.uk/about-us/emas-aspect/emas-aspect-2013/emas-aspect-22/ (accessed 16 April 2014).
53. Hassan T, McMillan P, Walsh C, Higginson I. *The Drive for Quality. How to Achieve Safe, Sustainable Care in our Emergency Departments: System Benchmarks and Recommendations*. London: The College of Emergency Medicine; 2013. URL: <http://secure.collemergencymed.ac.uk/Shop-Floor/Professional%20Standards/Quality%20in%20the%20Emergency%20Department/> (accessed 16 April 2014).
54. Great Britain. *Health and Social Care Act 2012*. London: The Stationery Office; 2012.
55. NHS Institute for Innovation and Improvement. *Commissioning for Quality and Innovation (CQUIN) Payment Framework 2006–2013*. URL: www.institute.nhs.uk/commissioning/pct_portal/cquin.html (accessed 16 April 2014).
56. Cooper S, Barrett B, Black S, Evans C, Real C, Williams S, et al. The emerging role of the emergency care practitioner. *Emerg Med J* 2004;**21**:614–18. <http://dx.doi.org/10.1136/emj.2003.011247>
57. Anderson KN, Anderson LE, Glanze WD. *Mosby's Medical, Nursing, & Allied Health Dictionary*. St. Louis, MO: Mosby-Year Book. Inc; 1994.
58. Chang JT, Morton SC, Rubenstein LZ, Mojica WA, Maglione M, Suttrop MJ, et al. Interventions for the prevention of falls in older adults: systematic review and meta-analysis of randomised clinical trials. *BMJ* 2004;**328**:680. <http://dx.doi.org/10.1136/bmj.328.7441.680>
59. Hill KD, Moore KJ, Dorevitch MI, Day LM. Effectiveness of falls clinics: an evaluation of outcomes and client adherence to recommended interventions. *J Am Geriatr Soc* 2008;**56**:600–8. <http://dx.doi.org/10.1111/j.1532-5415.2007.01626.x>
60. Palvanen M, Kannus P, Piirtola M, Niemi S, Parkkari J, Jarvinen M. Effectiveness of the Chaos Falls Clinic in preventing falls and injuries of home-dwelling older adults: a randomised controlled trial. *Injury* 2014;**45**:265–71. <http://dx.doi.org/10.1016/j.injury.2013.03.010>
61. Logan PA, Coupland CA, Gladman JR, Sahota O, Stoner-Hobbs V, Robertson K, et al. Community falls prevention for people who call an emergency ambulance after a fall: randomised controlled trial. *BMJ* 2010;**340**:c2102. <http://dx.doi.org/10.1136/bmj.c2102>
62. Sach TH, Logan PA, Coupland CA, Gladman JR, Sahota O, Stoner-Hobbs V, et al. Community falls prevention for people who call an emergency ambulance after a fall: an economic evaluation alongside a randomised controlled trial. *Age Ageing* 2012;**41**:635–41. <http://dx.doi.org/10.1093/ageing/afs071>
63. Newman J. *Modernising Governance: New Labour, Policy and Society*. London: Sage; 2001.
64. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;**337**:a1655.
65. Gates S, Fisher JD, Cooke M, Carter Y, Lamb SE. Multifactorial assessment and targeted intervention for preventing falls and injuries among older people in community and emergency care settings: systematic review and meta-analysis. *BMJ* 2008;**336**:130–3. <http://dx.doi.org/10.1136/bmj.39412.525243.BE>

66. Mason S, Knowles E, Colwell B, Wardrope J, Dixon S, Gorringer R. Paramedic Practitioner Older People's Support Trial (PPOPS): a cluster randomised controlled trial. *BMJ* 2007;**335**:919–25. <http://dx.doi.org/10.1136/bmj.39343.649097.55>
67. National Institute for Health and Care Excellence (NICE). *Falls: Assessment and Prevention of Falls in Older People*. NICE guidelines CG161. London: NICE; 2013. URL: <http://publications.nice.org.uk/falls-assessment-and-prevention-of-falls-in-older-people-cg161> (accessed 8 November 2013).
68. Centers for Disease Control and Prevention. *Costs of Falls Among Older Adults 2013*. URL: www.cdc.gov/homeandrecreationsafety/falls/fallcost.html (accessed 23 April 2014).
69. Hendrie D, Hall SE, Arena G, Legge M. Health system costs of falls of older adults in Western Australia. *Aust Health Rev* 2004;**28**:363–73. <http://dx.doi.org/10.1071/AH040363>
70. Cone DC, Ahern J, Lee CH, Baker D, Murphy T, Bogucki S. A descriptive study of the 'lift-assist' call. *Prehosp Emerg Care* 2013;**17**:51–6. <http://dx.doi.org/10.3109/10903127.2012.717168>
71. Snooks H, Anthony R, Chatters R, Cheung W-Y, Dale J, Donohoe R, *et al*. Support and assessment for fall emergency referrals (SAFER 2) research protocol: cluster randomised trial of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community-based care. *BMJ Open* 2012;**2**:e002169. <http://dx.doi.org/10.1136/bmjopen-2012-002169>
72. Darzi A. *High Quality Care for All: NHS Next Stage Review Final Report*. London: Department of Health; 2009.
73. Tiedemann A, Mikolaizak AS, Sherrington C, Segin K, Lord SR, Close JC. Older fallers attended to by an ambulance but not transported to hospital: a vulnerable population at high risk of future falls. *Aust N Z J Public Health* 2013;**37**:179–85. <http://dx.doi.org/10.1111/1753-6405.12037>
74. Mikolaizak AS, Simpson PM, Tiedemann A, Lord SR, Close JC. Systematic review of non-transportation rates and outcomes for older people who have fallen after ambulance service call-out. *Australas J Ageing* 2013;**32**:147–57. <http://dx.doi.org/10.1111/ajag.12023>
75. Snooks H, Cheung WY, Gwini S, Humphreys I, Sánchez A, Siriwardena N. Can older people who fall be identified in the ambulance call centre to enable alternative responses or care pathways? *Emerg Med J* 2011;**28**:e1. <http://dx.doi.org/10.1136/emj.2010.108605.15>
76. Newton JL, Kyle P, Liversidge P, Robinson G, Wilton K, Reeve P. The costs of falls in the community to the North East Ambulance Service. *Emerg Med J* 2006;**23**:479–81. <http://dx.doi.org/10.1136/emj.2005.028803>
77. Mason S, Wardrope J, Perrin J. Developing a community paramedic practitioner intermediate care support scheme for older people with minor conditions. *Emerg Med J* 2003;**20**:196–8. <http://dx.doi.org/10.1136/emj.20.2.196>
78. Darnell G, Mason SM, Snooks H. Elderly falls: a national survey of UK ambulance services. *Emerg Med J* 2012;**29**:1009–10. <http://dx.doi.org/10.1136/emmermed-2011-200419>
79. Snooks H, Cheung WY, Close J, Dale J, Gaze S, Humphreys I, *et al*. Support and Assessment for Fall Emergency Referrals (SAFER 1) trial protocol. Computerised on-scene decision support for emergency ambulance staff to assess and plan care for older people who have fallen: evaluation of costs and benefits using a pragmatic cluster randomised trial. *BMC Emerg Med* 2010;**10**:2. <http://dx.doi.org/10.1186/1471-227X-10-2>
80. Gray JT, Walker A. Avoiding admissions from the ambulance service: a review of elderly patients with falls and patients with breathing difficulties seen by emergency care practitioners in South Yorkshire. *Emerg Med J* 2008;**25**:168–71. <http://dx.doi.org/10.1136/emj.2007.050732>



81. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLOS Med* 2009;**6**:e1000097. <http://dx.doi.org/10.1371/journal.pmed.1000097>
82. University of York. *Systematic Reviews CRD's Guidance for Undertaking Reviews in Healthcare CRD*. 2009. URL: www.york.ac.uk/inst/crd/pdf/Systematic_Reviews.pdf (accessed 23 April 2014).
83. Petticrew S, Roberts H. *Systematic Reviews in the Social Sciences, A Practical Guide*. Oxford: Blackwell Publishing; 2006. <http://dx.doi.org/10.1002/9780470754887>
84. Prevention of Falls Network Europe. *Manual for the Fall Prevention Classification System*. 2007. URL: www.profane.eu.org/documents/Falls_Taxonomy.pdf (accessed 23 April 2014).
85. Hauer K, Lamb SE, Jorstad EC, Todd C, Becker C. Systematic review of definitions and methods of measuring falls in randomised controlled fall prevention trials. *Age Ageing* 2006;**35**:5–10. <http://dx.doi.org/10.1093/ageing/afi218>
86. Scottish Intercollegiate Guidelines Network. *Sign 50 A Guideline Developer's Handbook 2011*. URL: www.sign.ac.uk/pdf/sign50.pdf (accessed 23 April 2014).
87. Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery* 2006;**22**:108–19. <http://dx.doi.org/10.1016/j.midw.2005.05.004>
88. Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M. *Guidance on the Conduct of Synthesis In Systematic Reviews*. 2006. URL: www.lancaster.ac.uk/shm/research/nssr/research/dissemination/publications/NS_Synthesis_Guidance_v1.pdf (accessed 23 April 2014).
89. Metcalfe J. Enhancing the care of older people in the community. *Nurs Stand* 2006;**21**:35–9. <http://dx.doi.org/10.7748/ns2006.10.21.7.35.c6360>
90. Shah MN, Clarkson L, Lerner EB, Fairbanks RJ, McCann R, Schneider SM. An emergency medical services program to promote the health of older adults. *J Am Geriatr Soc* 2006;**54**:956–62. <http://dx.doi.org/10.1111/j.1532-5415.2006.00736.x>
91. Shandro JR, Spain DA, Dicker RA. Recruitment strategies for a fall prevention program: if we build it, will they really come? *J Trauma* 2007;**63**:142–6. <http://dx.doi.org/10.1097/TA.0b013e318068428a>
92. Kue R, Ramstrom E, Weisberg S, Restuccia M. Evaluation of an emergency medical services-based social services referral program for elderly patients. *Prehosp Emerg Care* 2009;**13**:273–9. <http://dx.doi.org/10.1080/10903120802706179>
93. Shah MN, Caprio TV, Swanson P, Rajasekaran K, Ellison JH, Smith K, *et al*. A novel emergency medical services-based program to identify and assist older adults in a rural community. *J Am Geriatr Soc* 2010;**58**:2205–11. <http://dx.doi.org/10.1111/j.1532-5415.2010.03137.x>
94. Halter M, Vernon S, Snooks H, Porter A, Close J, Moore F, *et al*. Complexity of the decision-making process of ambulance staff for assessment and referral of older people who have fallen: a qualitative study. *Emerg Med J* 2011;**28**:44–50. <http://dx.doi.org/10.1136/emj.2009.079566>
95. Hoyle S, Swain AH, Fake P, Larsen PD. Introduction of an extended care paramedic model in New Zealand. *Emerg Med Australas* 2012;**24**:652–6. <http://dx.doi.org/10.1111/j.1742-6723.2012.01608.x>
96. Studnek JR, Thestrup L, Blackwell T, Bagwell B. Utilization of prehospital dispatch protocols to identify low-acuity patients. *Prehosp Emerg Care* 2012;**16**:204–9. <http://dx.doi.org/10.3109/10903127.2011.640415>
97. Comans TA, Currin ML, Quinn J, Tippet V, Rogers A, Haines TP. Problems with a great idea: referral by prehospital emergency services to a community-based falls-prevention service. *Inj Prev* 2013;**19**:134–8. <http://dx.doi.org/10.1136/injuryprev-2011-040076>

98. Fitzpatrick-Lewis D, Ciliska D, Thomas H. *The Methods for the Synthesis of Studies without Control Groups*. Hamilton, ON: National Collaborating Centre for Methods and Tools; 2009. URL: www.nccmt.ca/pubs/non-RCT2_EN.pdf (accessed 23 April 2014).
99. Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration; 2011. URL: www.cochrane-handbook.org (accessed 23 April 2014).
100. Snooks H, Foster T, Nicholl J. Results of an evaluation of the effectiveness of triage and direct transportation to minor injuries units by ambulance crews. *Emerg Med J* 2004;**21**:105–11. <http://dx.doi.org/10.1136/emj.2003.009050>
101. Medical Research Council (MRC). *Good Clinical Practice In Clinical Trials*. London: MRC; 1998. URL: www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC002416 (accessed 14 May 2014).
102. WWORTH. *SOP24 Standard Operating Procedure for Randomisation*. Swansea: Swansea University Medical School; 2012.
103. Lamb SE, Jorstad-Stein EC, Hauer K, Becker C. Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 2005;**53**:1618–22. <http://dx.doi.org/10.1111/j.1532-5415.2005.53455.x>
104. Ware J Jr, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;**34**:220–33. <http://dx.doi.org/10.1097/00005650-199603000-00003>
105. Carey RG, Seibert JH. A patient survey system to measure quality improvement: questionnaire reliability and validity. *Med Care* 1993;**31**:834–45. <http://dx.doi.org/10.1097/00005650-199309000-00008>
106. Hill KD, Schwarz JA, Kalogeropoulos AJ, Gibson SJ. Fear of falling revisited. *Arch Phys Med Rehabil* 1996;**77**:1025–9. [http://dx.doi.org/10.1016/S0003-9993\(96\)90063-5](http://dx.doi.org/10.1016/S0003-9993(96)90063-5)
107. WWORTH. *SOP28 Standard Operating Procedure for Statistics*. Swansea: Swansea University Medical School; 2012.
108. Aminzadeh F, Dalziel WB. Older adults in the emergency department: a systematic review of patterns of use, adverse outcomes, and effectiveness of interventions. *Ann Emerg Med* 2002;**39**:238–47. <http://dx.doi.org/10.1067/mem.2002.121523>
109. Department of Health (DH). *NHS Reference Costs 2011–12*. London: DH; 2012.
110. Barbour RS. *Are Focus Groups an Appropriate Tool for Studying Organisational Change. Developing Focus Group Research*. London: Sage; 1999. pp. 113–26.
111. Ritchie J, Spencer L. Qualitative Data Analysis for Applied Policy Research. In Huberman M, Miles M, editors. *The Qualitative Researcher's Companion*. London: Sage; 2002.
112. Srivastava A, Thomson S. Framework analysis: a qualitative methodology for applied policy research. *JOAAG* 2009;**4**.
113. Corden A, Sainsbury R. *Using Verbatim Quotations in Reporting Qualitative Social Research: Researchers' Views*. York: Social Policy Research Unit, University of York; 2006.
114. INVOLVE. *Briefing Notes for Researchers: Involving the Public in NHS, Public Health and Social Care Research*. Eastleigh: INVOLVE; 2012. URL: www.invo.org.uk/posttypepublication/involve-briefing-notes-for-researchers/ (accessed 14 May 2014).
115. Boote J, Telford R, Cooper C. Consumer involvement in health research: a review and research agenda. *Health Policy* 2002;**61**:213–36. [http://dx.doi.org/10.1016/S0168-8510\(01\)00214-7](http://dx.doi.org/10.1016/S0168-8510(01)00214-7)

116. Entwistle VA, Renfrew MJ, Yearley S, Forrester J, Lamont T. Lay perspectives: advantages for health research. *BMJ* 1998;**316**:463–6. <http://dx.doi.org/10.1136/bmj.316.7129.463>
117. Chalmers I. A guide to patient-led good controlled trials. *Lancet* 2000;**356**:774. [http://dx.doi.org/10.1016/S0140-6736\(05\)73681-3](http://dx.doi.org/10.1016/S0140-6736(05)73681-3)
118. Evans BA, Bedson E, Bell P, Hutchings H, Lowes L, Rea D, *et al.* Involving service users in trials: developing a standard operating procedure. *Trials* 2013;**14**:219. <http://dx.doi.org/10.1186/1745-6215-14-219>
119. Boote J, Baird W, Sutton A. Public involvement in the design and conduct of clinical trials: a review. *Int J Interdiscip Soc Sci* 2011;**5**:91–111.
120. INVOLVE. *Public Involvement in Clinical Trials: Supplement to the Briefing Notes for Researchers*. 2012. URL: www.nihr.ac.uk/get-involved/INVOLVEpublicinvolvementinclinicaltrialsBriefingnotes2012.pdf (accessed 23 February 2014).
121. Snooks HA, Anthony R, Chatters R, Dale J, Fothergill RT, Gaze S, *et al.* Paramedic assessment of older adults after falls, including community care referral pathway: cluster randomized trial [Published online ahead of print March 13 2017]. *Ann Emerg Med* 2017. <https://doi.org/10.1016/j.annemergmed.2017.01.006>
122. Department of Health (DH). *NHS Reference Costs 2012–13*. London: DH; 2013.
123. Curtis L. *Unit Costs of Health and Social Care 2013*. Canterbury: PSSRU, University of Kent; 2013.
124. Turner J, Snooks H, Youren A, Dixon S, Fall D, Gaze S, *et al.* *The Costs and Benefits of Managing some Low Priority 999 Ambulance Calls by NHS Direct Nurse Advisers. Final Report to the Service Delivery and Organisation R&D Programme*. Sheffield: SCHARR (School of Health and Related Research), Sheffield University; 2006.
125. Fletcher B, Gheorghe A, Moore D, Wilson S, Damery S. Improving the recruitment activity of clinicians in randomised controlled trials: a systematic review. *BMJ Open* 2012;**2**:e000496. <http://dx.doi.org/10.1136/bmjopen-2011-000496>
126. Hawe P, Shiell A, Riley T, Gold L. Methods for exploring implementation variation and local context within a cluster randomised community intervention trial. *J Epidemiol Community Health* 2004;**58**:788–93. <http://dx.doi.org/10.1136/jech.2003.014415>

Appendix 1 Support and Assessment for Fall Emergency Referrals 2 clinical decision protocol

Reprinted from Snooks *et al.*¹²¹ with permission from the American College of Emergency Physicians. Copyright © 2017 American College of Emergency Physicians. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Appendix 2 Full data extraction and quality assessment tables

TABLE 38 Data extraction table

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Metcalfe, 2006 (UK) ⁸⁸	Cohort study to examine the benefit to patients of a single point of access to other services	Patients aged ≥ 65 years being attended by an emergency ambulance within the PCT area and being left at home	Clinical assessment tool and training to support ambulance staff. External agency undertook clinical triage for referral to a rapid response team and further onward referrals	$n = 49$	Number of referrals made for patients experiencing a fall Number of onward referrals and destination of referrals	$n = 53$ $n = 51$: specialist nurse, $n = 1$: rapid response team, $n = 17$: community hospital, $n = 1$: geriatrician, $n = 2$: community physiotherapist, $n = 4$: GP, $n = 9$: district nurse, $n = 8$: community psychiatric nurse, $n = 3$: and social care, $n = 6$
Shah <i>et al.</i> , 2006 (USA) ⁹⁰	Cluster controlled trial to evaluate the effect of an EMS programme by screening at least 80% of community-dwelling older adults to identify those at risk for falls	Intervention group: residents aged ≥ 65 years ($n = 1011$ in 2000) Usual-care control group: residents aged ≥ 65 years ($n = 720$) Patients were excluded if they could not speak English, if they had been previously included in the study or if they were institutionalised (group home, nursing home or jail)	EMTs provided with training, participating in a 90-minute case-based discussion on caring for older adults. Training on the screening programme was provided. EMTs instructed to screen community-dwelling patients during emergency responses to evaluate risk of falling. Instructional materials were given if the patient was considered to be at risk	$n = 258$ (total patients attended by EMTs for all conditions). $n = 143$ control	Number patients refusing referrals Patients' risk status (falls) Recollection of educational materials Recollection of discussions with physicians regarding risks Receipt of changes to the home environment	$n = 2$ Successful for screening for falls history $n = 210/258$ (79%, 95% CI 74% to 84%) Poor recollection of falls brochure for intervention group, only $n = 17$ (21%) Infrequent discussions with PCPs about risks: intervention $n = 8$ (10%) and control $n = 1$ (3%), with no PCP visit in 35% intervention and 46% control cases No difference in changes to prevent falls: intervention: $n = 12/82$ (15%), control: $n = 4/37$ (11%); $p = 0.91$

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Mason <i>et al.</i> , 2007 (UK) ⁸⁶	C-RCT	Individuals aged 60 years and above calling the emergency services for a presenting complaint in the scope of practice of the paramedic practitioner	Paramedics with extended skills (paramedic practitioners)	<i>n</i> = 3018	ED attendance 0 to 28 days Admittance to hospital 0 to 28 days Total episode time	Intervention = 62.6%; control = 87.5% (<i>p</i> < 0.001) Intervention = 40.4%; control = 46.4% (<i>p</i> < 0.001) Intervention = 235 minutes; control = 278 minutes (<i>p</i> < 0.001)
Shandro <i>et al.</i> , 2007 (USA) ⁹¹	Cohort study to evaluate recruitment strategies and to outline challenges and solutions of the implementation of a fall prevention programme	Patients, who are living independently, were over 65 years of age and had an emergency service dispatch for a fall but were not transported to the ED	Provided with educated on the goals and design of a fall prevention programme. Voluntary referral mechanism for self-report or report of a medical provider for participants that had a recent fall and were able to participate in a home safety evaluation and strengthening programme	Eligible patient numbers not specified	Number of referrals by EMS staff Number of enrolments on falls prevention programme by EMS referrals	Intervention = 49.7%; control = 67.9% (<i>p</i> < 0.001) Intervention = 81.3%; control = 72.8% (<i>p</i> < 0.001) <i>n</i> = 17 referred to the prevention programme <i>n</i> = 11 (65%) enrolled onto the prevention programme

continued

TABLE 38 Data extraction table (continued)

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Gray and Walker, 2008 (UK) ⁸⁰	Time-series analysis cohort study with control arm to establish the impact of ECPs in terms of avoided admissions for falls	All patients aged ≥ 65 years who had called the emergency services after a fall and were seen by the ECP service; or who had attended ED following attendance by non-ECP ambulance personnel	ECP, with additional training, attending calls for conditions felt to benefit from ECP management	$n = 233$ ECP attended cases and $n = 772$ pre-intervention attended cases	% not conveyed following initial ECP attendance % avoiding admissions during pre-intervention period	Intervention arm: $n = 171/233$ (73%) Pre-intervention arm: $n = 369/772$ (48%)
					% ED attendance or hospital admission in 72 hours	Intervention arm: $n = 21$ further hospital attendances in 72 hours
					% ED attendance or hospital admission in 28 days	Intervention arm: $n = 131$ remained at home at 28 days (44% ED attendance/admission rate)
						Authors describe: ECPs reduce admissions by 17% at 28 days relative to the ED 'initial contact' figure (from 52% to 44%, $n = 1005$, $df = 1$, $p = 0.05$)
^a Kue et al., 2009 (USA) ⁹²	Cohort study to describe the experience of implementing three referral programmes and to report frequencies of referrals made	Adults aged ≥ 60 years who called the emergency services in the city EMS area and were not conveyed	Paramedic in-service training; paramedic referrals to social services and retrospective EMS physician referral on chart review	$n = 721$	Number of referrals for fall-related complaints Number of acceptance of referral	$n = 32$ $n = 24$

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Logan <i>et al.</i> , 2010 (UK) ⁶¹	RCT to evaluate whether or not a service to prevent falls in the community would reduce the rate of falls in older people who called the emergency services and were not taken to hospital	Adults aged ≥ 60 years living at home or in residential care in one of the PCT areas who were not taken to hospital	Retrospective identification of potential participants from ambulance service records. Use of a randomisation sequence, individualised multifactorial intervention programme following UK clinical fall guidelines	$n = 204$ ($n = 102$ in each group)	Died by 12-month follow-up At least one falls-related fracture admitted to hospital Number of emergency ambulance calls received for falls over 12 months Number of days in hospital per year Total number of hospital admissions Median Falls Efficacy Scale (fear of falling) Median Nottingham Extended Activities of Daily Living score Median Barthel Index of Activities of Daily Living score Number with at least one fall during follow-up Median days to first fall incidence rate ratio, over 12 months (primary outcome)	Control, $n = 16$; intervention, $n = 14$ ($p = 0.74$) Control, $n = 6$; intervention, $n = 3$ ($p = 0.35$) Control, $n = 365$; intervention, $n = 245$ ($p = 0.018$) Control, $n = 1141$; intervention, $n = 1257$ ($p = 0.70$) Control, $n = 99$; intervention, $n = 97$ ($p = 0.93$) Control, $n = 76$; intervention, $n = 57$ ($p < 0.001$) Control, $n = 6$; intervention, $n = 8$ ($p < 0.001$) Control, $n = 15$; intervention, $n = 15$ ($p = 0.021$) Control, $n = 96$; intervention, $n = 81$; incidence rate ratio = 0.86 ($p < 0.001$) Overall incidence of falls intervention group = 3.46 per person and control group 7.68; incidence rate ratio 0.45 (95% CI 0.35 to 0.58; $p < 0.001$)

continued

TABLE 38 Data extraction table (continued)

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Shah <i>et al.</i> , 2010 (USA) ⁸³	Cohort study to describe the development, implementation and, experience of a novel, community-based programme to identify rural-dwelling older adults with unmet medical and social needs	Community-dwelling older adults, aged 60 years and above, who accessed EMS for emergency care	EMS staff provided with training specific to geriatrics. The screening question looked at whether or not the patient had fallen in the previous year or if the EMS provider identified any falls risks in the home. Case managers intercepted the findings and contacted patients after 2 weeks to arrange a home visit	n = 814 (adults screened by EMS for falls)	Number of patients screened for falls Number patients referred due to falls Number patients at risk of falls	n = 814 n = 124 n = 552
Halter <i>et al.</i> , 2011 (UK) ⁸⁴	Qualitative study to understand the decision-making processes of emergency ambulance staff dealing with older people who have fallen	12 ambulance staff trained in, and used the, clinical assessment tool	Paramedics received 2-hour training session in the cause of falls, assessment of older person who has fallen and research methods. Paramedics completed clinical assessment tool for those patients who had a fall	12 paramedics	Paramedics' self-reported process of decision-making following 6 months use of the clinical assessment tool	<p>Paramedics described the decision-making process as follows:</p> <ul style="list-style-type: none"> • Pre-arrival: forming an early opinion from information from the emergency call. Initial contact: assessing the need for any immediate action and establishing a rapport • Continuing assessment: gathering and assimilating medical and social information • Making a conveyance decision: negotiation, referral and professional defence, using professional experience and instinct <p>The completion of the clinical assessment tool was low</p>

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Hoyle <i>et al.</i> , 2012 (New Zealand) ⁹⁵	To evaluate the first 1000 patients seen by emergency care paramedics	Average patient age of 62 years	Assessment by individual dispatchers to decide whether or not to send emergency care paramedic to patient. Emergency care paramedics followed clinical protocols that were specific to patient condition	n = 131 (number of falls out of the 1000 presentations attended by emergency care paramedics)	Patient disposition including ED and treated in community Reduced unnecessary transportation	87 (66%) patients were treated in the community and 44 (34%) patients transported to ED 40% of patients transported; 74% of patients seen by standard ECPs were transported
Sach <i>et al.</i> , 2012 (UK) ⁶²	Cost-effectiveness study to estimate the cost-effectiveness of a community falls prevention service compared with usual care, over the 12-month trial period	Patients aged ≥ 60 years, attended for a fall at home in four Nottinghamshire PCTs and were not transported to hospital	Participants in the intervention group were referred to a community falls prevention programme. Control group underwent normal care	n = 157 (82 intervention; 75 control)	Cost utility (QALY) Cost-effectiveness (falls per patient)	Mean difference in QALYs of 0.070, 95% CI -0.010 to 0.150; $p = 0.086$ Difference of mean number of falls per patient during 12 month follow-up period: -5.34, 95% CI -7.06 to -3.62; $p < 0.001$ Mean difference: -£1551, 95% CI -£5932.00 to £2829.00
Studneck <i>et al.</i> , 2012 (USA) ⁹⁶	Cohort study to describe the experience of an EMS agency using the MPDS low-acuity omega protocol (omega = calls classified as not requiring immediate ambulance response or transport and appropriate transfer to an advice-line nurse)	Emergency service callers classified as low acuity by MPDS	Introduction of MPDS Omega responses assigned to patients deemed to be low acuity by MPDS including falls. Phase 1: triaged and ambulance dispatched. Phase 2: patients offered to speak to an advice-line nurse for secondary triage, with ambulance responses continued per standard protocol	Phase 1: fall, n = 160; phase 2 fall, n = 101	% admitted to hospital % admitted to ICU	Phase 1, n = 56/160 (35%) hospital admission; phase 2, n = 66/101 (25.7%) hospital admission Phase 1, n = 3/160 (5.4%) ICU admission; phase 2, n = 0/101 ICU admission

continued

TABLE 38 Data extraction table (*continued*)

Study (country)	Study design and purpose	Population characteristics	Intervention	Sample size	Outcome measure	Results
Comans <i>et al.</i> , 2013 (Australia) ⁹⁷	Cohort study to discuss the potential barriers and enablers of a collaborative approach between two health services and to compare referral pathways	Patients older than 65 years, living in the community; and had had a recent fall attended by a paramedic trained in the referral technique	Referral pathways were direct from a paramedic or self-referral after information provided by a paramedic. Participants received a falls assessment from the rehabilitation service with referral to an 8-week falls programme	<i>n</i> = 638	Characteristics of referred patients (Baseline Frenchay Activities Index, Abbreviated Mental Test Score, EQ-5D and median falls in past 6 months showed)	No between-group descriptive differences
					Strategies to improve referral rates	Repeated education sessions with officers; contact from rehabilitation service to confirm referral; rehabilitation observation days
					Number of participants referred	<i>n</i> = 21
					Number of participants consented to referral	<i>n</i> = 17
					Patients consented to study and initial assessment	<i>n</i> = 13
					Patients enrolled onto programme	<i>n</i> = 8
					Patients completed programme	<i>n</i> = 5

df, degrees of freedom; ICU, intensive care unit; MPDS, Medical Priority Dispatch System; PCP, primary care physician.

a Kue *et al.*⁹² notes that it was unclear if sample size included all possible complaints or just falls. The number of accepted referrals was not differentiated by paramedic or physician. Only 7 of the 32 referrals were made by paramedics.

TABLE 39 Quality assessment table (cohort studies)

Quality assessment	1.1: clearly focused question	1.2: comparable source populations	1.3: participation rates stated	1.4: outcome at time of enrolment taken into account	1.5: percentage of participants dropping out	1.6: comparison made between participants and those lost to follow-up	1.7: outcomes clearly defined	1.8: blinding outcome	1.9: if blinding not possible, recognition that exposure could influence outcome	1.10: measure of exposure assessment is reliable	1.11: other sources cited to demonstrate outcome assessment is valid	1.12: exposure level assessed more than once	1.13: confounders identified and taken into account	1.14: CIs provided	2.1: how well study done to minimise risk of bias?	2.2: how strong is association between exposure and outcome?
^a Metcalfe, 2006 ⁵⁸	No	n/a	n/a	No	0%	n/a	Yes	n/a	No	Yes	n/a	No	No	No	Low	Weak
^b Shah <i>et al.</i> , 2006 ³⁰	Yes	Yes	Yes	n/a	63%	No	Yes	No	Yes	Yes	n/a	n/a	No	Yes	Acceptable	Moderate
^c Shandro <i>et al.</i> , 2007 ³¹	Yes	n/a	n/a	No	35%	n/a	Yes	n/a	n/a	Yes	n/a	No	No	No	Low	Moderate
^d Gray and Walker, 2008 ⁸⁰	Yes	Cannot say	No	No	n/a	n/a	Yes	n/a	No	Yes	n/a	n/a	No	No	Acceptable	Moderate
^e Kue <i>et al.</i> , 2009 ³²	Yes	n/a	n/a	Cannot say	25%	n/a	No	n/a	n/a	Cannot say	n/a	n/a	Cannot say	Yes	Low	Weak
^f Shah <i>et al.</i> , 2010 ³³	No	n/a	n/a	No	Cannot say	n/a	Yes	n/a	n/a	Yes	n/a	No	No	No	Low	Strong
^g Hoyle <i>et al.</i> , 2012 ³⁵	Yes	n/a	n/a	n/a	n/a	n/a	Yes	n/a	n/a	Yes	n/a	n/a	Yes	No	Low	Moderate
^h Studnek <i>et al.</i> , 2012 ³⁶	Yes	n/a	n/a	n/a	n/a	n/a	Yes	n/a	Yes	Yes	n/a	No	No	No	Low	Weak
ⁱ Comans <i>et al.</i> , 2013 ³⁷	Yes	Yes	Yes	No	75%	No	Yes	n/a	n/a	Yes	n/a	No	No	No	Low	Weak

n/a, not applicable.

a Poorly reported. The methods are poorly stated; the authors' conclusions do not match the results.

b EMS screening rates were high in the study; this is a feasible intervention for standard practice. Educational materials given out by EMS were not recalled and are not recommended.

c An intensive intervention is needed to see an impact of EMS screening.

d This study is very limited in its presentation of EMS-specific data and outcomes of care. Article does not state how many patients were exposed.

e Different data were collected for the two groups, limiting comparison.

f Data are not presented clearly. They are limited by small study numbers and being in one site.

g Low-quality study: no comparison arm or follow-up of patient outcomes.

h Subjective assessment and possibly bias in dispatch, disparity in transportation rates and could not check if patients attended at another hospital.

i This study includes 'fall' as a category to which the 'delayed dispatch' is applied. It is very limited in terms of outcomes measured. The study changes sites in the two phases and does not have any comparator/baseline data.

i Limited data to use.

TABLE 40 Quality assessment table (RCT)

Quality assessment	1.1: the study addresses an appropriate and clearly focused question	1.2: the assignment of subjects to treatment groups is randomised	1.3: adequate concealment method used	1.4: subjects and investigators are kept 'blind' to treatment allocation	1.5: treatment and control groups similar at the start of the trial	1.6: only difference between groups is treatment under investigation	1.7: all relevant outcomes measured in a standard, valid and reliable way	1.8: what percentage of the individuals or clusters recruited into each treatment arm of the study dropped	1.9: all the subjects analysed in the groups to which they were randomly allocated	1.10: where the study is carried out at more than one site, results are comparable for all sites	2.1: how well was the study done to minimise the risk of bias or confounding	2.2: is the overall effect is due to the intervention
^a Logan <i>et al.</i> , 2010 ⁶¹	Yes	Yes	Yes	No (subjects were not blinded, but investigators were)	Yes	Yes	Yes	6%	Yes	n/a	High	Yes
Mason <i>et al.</i> , 2007 ⁶⁶	Yes	Yes (by week)	Yes (forward roster concealed)	n/a	Yes	Yes	Yes	27%	Yes	n/a	High quality	Yes

n/a, not applicable.

^a Participants were not blinded. This should have been a cluster trial.

TABLE 41 Quality assessment table (cost-effectiveness)

Quality assessment	1.1: the study addresses an appropriate and clearly focused question	1.2: the economic importance of the question is clear	1.3: the choice of study design is justified	1.4: all costs that are relevant from the viewpoint of the study are included and are measured and valued appropriately	1.5: the outcome measures used to answer the question are relevant to that purpose and are measured and valued appropriately	1.6: if discounting of future costs and outcomes is necessary, it's been performed correctly	1.7: assumptions are explicit, sensitivity analysis performed	1.8: the decision rule explicit and comparisons made on the basis of incremental costs and outcomes	1.9: the results provide information of relevance to policy makers	2.1: how well was the study conducted	Are the results of this study directly applicable to the review question?
^a Sach <i>et al.</i> , 2012 ⁶²	Yes	Yes	Yes	Yes	Yes	n/a	Yes	Yes	Yes	High quality	Yes

n/a, not applicable.

^a The community falls prevention service delivered in this trial was cost-effective with little decision uncertainty. This study further justifies the development of clinical pathways linking the emergency ambulance services to community therapy services. Some measures in this study are self-reported and the participants were not blinded; this may have influenced study outcomes.

TABLE 42 Quality assessment table (qualitative)

Quality assessment	Clear statement of, and rationale for, research questions/ aims/purposes	Study thoroughly contextualised by existing literature	Method/design apparent, and consistent with research intent	Data collection strategy appropriate	Sample and sampling method appropriate	Analytical approach appropriate	Context described and taken into account of in interpretation		Data used to support interpretation	Researcher reflexivity demonstrated	Demonstration of sensitivity to ethical concerns	Relevance and transferability evident
							Clear audit trail given	Yes				
Halter <i>et al.</i> , 2011 ³⁴	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	No	Yes

Appendix 3 Excluded studies

The total number of excluded studies was 128; this excludes the three duplicate articles.

Limits (n = 12)

Not English (n = 10)

Asakawa Y, Takahashi R, Kagawa J. [Falling accidents among metropolitan elderly resulting in emergency ambulance transfer.] *Nihon Ronen Igakkai Zasshi* 2001;**38**:534–9.

Bauer C, Rietsch C, Groger I, Gassmann KG. [Mobility and safety for elderly (MoSi), a new intervention to improve mobility and gait in elderly people.] *Z Gerontol Geriatr* 2009;**42**:360–4.

Casas Herrero A, Martinez Velilla N, Alonso Renedo FJ. [Cognitive impairment and the risk of falling in the elderly.] *Rev Esp Geriatr Gerontol* 2011;**46**:311–18.

Casas-Herrero A, Izquierdo M. [Physical exercise as an efficient intervention in frail elderly persons.] *An Sist Sanit Navar* 2012;**35**:69–85.

Mamerow R. [Risk of accidental falls 2: and when it happens anyway.] *Pflege Z* 2003;**56**:268–9.

Niki R. [Critical appraisal of preventive care – from the perspective of health economics and policy.] *Nihon Ronen Igakkai Zasshi* 2012;**49**:54–7.

Pruckner S, Luiz T, Steinbach-Nordmann S, Nehmer J, Danner K, Madler C. [Emergency medicine – medicine for an ageing society. A contribution to the context of emergency missions for elderly people.] *Anaesthesist* 2008;**57**:391–6.

Rose DJ. [Preventing falls among older adults: no ‘one size suits all’ intervention strategy.] *Rehabil Med* 2011;**15**:37–49.

Szczerbinska K, Topor-Madry R. [The characteristics of falls based on the prospective registration in nursing home.] *Przegl Lek* 2011;**68**:576–84.

Yoshimoto Y. [The association between standardized incidence ratio of fall accidents with ambulance responses of elderly people and socio-economic status.] *Nihon Koshu Eisei Zasshi* 2011;**58**:183–9.

Abstract only (n = 1)

Snooks H, Cheung WY, Gwini S, Humphreys I, Sánchez A, Siriwardena N. Can older people who fall be identified in the ambulance call centre to enable alternative responses or care pathways? *EMJ* 2011;**28**:e1.

Review article (n = 1)

Mikolaizak AS, Simpson PM, Tiedemann A, Lord SR, Close JC. Systematic review of non-transportation rates and outcomes for older people who have fallen after ambulance service call-out. *Australas J Ageing* 2013;**32**:147–57.

Context (not emergency medical services) (n = 65)

Albert SM. Behavioral interventions to reduce risk of falls in nursing home residents. *J Am Med Dir Assoc* 2009;**10**:593–4.

Beswick A, Goberman-Hill R, Smith A, Wylde V, Ebrahim S. Maintaining independence in older people. *Rev Clin Gerontol* 2010;**20**:128–53.

Bleijlevens MH, Hendriks MR, van Haastregt JC, van Rossum E, Kempen GI, Diederiks JP, *et al*. Process factors explaining the ineffectiveness of a multidisciplinary fall prevention programme: a process evaluation. *BMC Public Health* 2008;**8**:332.

Bouman A, van Rossum E, Evers S, Ambergen T, Kempen G, Knipschild P. Effects on health care use and associated cost of a home visiting program for older people with poor health status: a randomized clinical trial in the Netherlands. *J Gerontol A Biol Sci Med Sci* 2008;**63**:291–7.

Bradley SM. Falls in older adults. *Mt Sinai J Med* 2011;**78**:590–5.

Campbell AJ, Robertson MC. Reply. *Age Ageing* 2008;**37**:353.

Carpenter CR, Platts-Mills TF. Evolving prehospital, emergency department, and 'inpatient' management models for geriatric emergencies. *Clin Geriatr Med* 2013;**29**:31–47.

Carpenter CR, Shah MN, Hustey FM, Heard K, Gerson LW, Miller DK. High yield research opportunities in geriatric emergency medicine: prehospital care, delirium, adverse drug events, and falls. *J Gerontol A Biol Sci Med Sci* 2011;**66**:775–83.

Catarina LN, Pereira V, Peter V, Fátima B. Role of physical activity in the prevention of falls and their consequences in the elderly. *Eur Rev Aging Phys Act* 2008;**5**:51–8.

Choi M, Hector M. Effectiveness of intervention programs in preventing falls: a systematic review of recent 10 years & meta-analysis. *J Am Med Dir Assoc* 2012;**13**:188(e13–21).

Cryer C. *What Works to Prevent Accidental Injury Amongst Older People. Report to the Health Development Agency*. Canterbury: University of Kent; 2001.

Davis JC, Robertson MC, Ashe MC, Liu-Ambrose T, Khan KM, Marra CA. Does a home-based strength and balance programme in people aged 80 years provide the best value for money to prevent falls? A systematic review of economic evaluations of falls prevention interventions. *Br J Sports Med* 2010;**44**:80–9.

Davison J, Marrinan S. Falls. *Rev Clin Gerontol* 2007;**17**:93–107.

Deasy C, Ryan D, O'Donnell C, Cusack S. The impact of a pre-hospital medical response unit on patient care & emergency department attendances. *Irish Med J* 2008;**101**:44–6.

Fairhall N, Sherrington C, Clemson L, Cameron ID. Do exercise interventions designed to prevent falls affect participation in life roles? A systematic review and meta-analysis. *Age Ageing* 2011;**40**:666–74.

Fitzharris MP, Day L, Lord SR, Gordon I, Fildes B. The Whitehorse No Falls trial: effects on fall rates and injurious fall rates. *Age Ageing* 2010;**39**:728–33.

Freiberger E, Haberle L, Spirduso WW, Zijlstra GA. Long-term effects of three multicomponent exercise interventions on physical performance and fall-related psychological outcomes in community-dwelling older adults: a randomized controlled trial. *J Am Geriatr Soc* 2012;**60**:437–46.

- Frith J, Davison J. Falls. *Rev Clin Gerontol* 2013;**23**:101–17.
- Ganz DA, Alkema GE, Wu S. It takes a village to prevent falls: reconceptualizing fall prevention and management for older adults. *Inj Prev* 2008;**14**:266–71.
- Gates S, Fisher JD, Cooke M, Carter Y, Lamb SE. Multifactorial assessment and targeted intervention for preventing falls and injuries among older people in community and emergency care settings: systematic review and meta-analysis. *BMJ* 2008;**336**:130–3.
- Gilbert T, Szczerbińska K, Ziere G, Ait S, Bonnefoy M, van der Velde N, *et al.* Management of falls: France, the Netherlands, Poland. *Eur Geriatr Med* 2012;**3**:246–53.
- Gillespie LD, Robertson MC, Gillespie WJ, Lamb SE, Gates S, Cumming RG, *et al.* Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2009;**2**:CD007146.
- Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, *et al.* Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2012;**9**:CD007146.
- Gunn H, Freeman J. Repeated falls: a key outcome or an adverse event? *Physiother Res Int* 2012;**17**:1–3.
- Hendriks MR, Evers SM, Bleijlevens MH, van Haastregt JC, Crebolder HF, van Eijk JT. Cost-effectiveness of a multidisciplinary fall prevention program in community-dwelling elderly people: a randomized controlled trial. *Int J Technol Assess Health Care* 2008;**24**:193–202.
- Holmes J, Bethel J. A different kind of care. *Emerg Nurse* 2007;**15**:8–12.
- Holroyd-Leduc JM. Review: exercise interventions reduce falls in elderly people living in the community. *Evid Based Med* 2009;**14**:176.
- Jenkyn KB, Hoch JS, Speechley M. How much are we willing to pay to prevent a fall? Cost-effectiveness of a multifactorial falls prevention program for community-dwelling older adults. *Can J Aging* 2012;**31**:121–37.
- Kannus P, Sievanen H, Palvanen M, Jarvinen T, Parkkari J. Prevention of falls and consequent injuries in elderly people. *Lancet* 2005;**366**:1885–93.
- Kurrle SE. Improving acute care services for older people. *Med J Aust* 2006;**184**:427–8.
- Lamb SE. Multidisciplinary assessment of elderly people with a history of multiple falls reduces the risk of further falls. *Aust J Physiother* 2009;**55**:139.
- Mayer D. Guidelines for care of the elderly. *Ann Emerg Med* 1998;**32**:115.
- McClure RJ, Hughes K, Ren C, McKenzie K, Dietrich U, Vardon P, *et al.* The population approach to falls injury prevention in older people: findings of a two community trial. *BMC Public Health* 2010;**10**:79.
- McKay C, Anderson KE. How to manage falls in community dwelling older adults: a review of the evidence. *Postgrad Med J* 2010;**86**:299–306.
- Meier DE, Beresford L. Palliative care's challenge: facilitating transitions of care. *J Palliat Med* 2008;**11**:416–21.
- Meirink K. Guidelines for care of the elderly. *Ann Emerg Med* 1998;**32**:114–15.

- Merrett A, Thomas P, Stephens A, Moghabghab R, Gruneir M. A collaborative approach to fall prevention. *Can Nurse* 2011;**107**:24–9.
- Michael YL, Whitlock EP, Lin JS, Fu R, O'Connor EA, Gold R. Primary care-relevant interventions to prevent falling in older adults: a systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2010;**153**:815–25.
- Miller ET. Safe after 60. *Rehabil Nurs* 2011;**36**:2.
- Murphy TE, Allore HG, Leo-Summers L, Carlin BP. Bayesian hierarchical modeling for a non-randomized, longitudinal fall prevention trial with spatially correlated observations. *Stat Med* 2011;**30**:522–30.
- Norton R, Kobusingye O. Global health injuries. *N Engl J Med* 2013;**368**:1723–30.
- Nowak A, Hubbard RE. Falls and frailty: lessons from complex systems. *J R Soc Med* 2009;**102**:98–102.
- Odom JV, Odom CV, Leys MJ. Does improving vision reduce the risk of falls? A review. *Insight* 2011;**4**:92–9.
- Oh DH, Park JE, Lee ES, Oh SW, Cho SI, Jang SN, *et al*. Intensive exercise reduces the fear of additional falls in elderly people: findings from the Korea falls prevention study. *Korean J Intern Med* 2012;**27**:417–25.
- Oliver D. Development of services for older patients with falls and fractures in England: successes, failures, lessons and controversies. *Arch Gerontol Geriatr* 2009;**49**(Suppl. 2):S7–12.
- Pan J-I, Yung C-J, Liang C-C, Lai L-F. *An Intelligent Homecare Emergency Service System for Elder Falling. World Congress on Medical Physics and Biomedical Engineering 2006*. New York, NY: Springer; 2007.
- Peeters GM, Heymans MW, de Vries OJ, Bouter LM, Lips P, van Tulder MW. Multifactorial evaluation and treatment of persons with a high risk of recurrent falling was not cost-effective. *Osteoporos Int* 2011;**22**:2187–96.
- Petridou ET, Manti EG, Ntinapogias AG, Negri E, Szczerbinska K. What works better for community-dwelling older people at risk to fall?: a meta-analysis of multifactorial versus physical exercise-alone interventions. *J Aging Health* 2009;**21**:713–29.
- Quigley P. Prevention of fall-related injuries: a clinical research agenda 2009–2014. *J Rehabil Res Dev* 2009;**46**:vii–xii.
- Rand D, Miller WC, Yiu J, Eng JJ. Interventions for addressing low balance confidence in older adults: a systematic review and meta-analysis. *Age Ageing* 2011;**40**:297–306.
- Rose DJ, Hernandez D. The role of exercise in fall prevention for older adults. *Clin Geriatr Med* 2010;**26**:607–31.
- Rose DJ. Preventing falls among older adults: no 'one size suits all' intervention strategy. *J Rehabil Res Dev* 2008;**45**:1153–66.
- Salminen M, Vahlberg T, Kivela SL. The long-term effect of a multifactorial fall prevention programme on the incidence of falls requiring medical treatment. *Public Health* 2009;**123**:809–13.
- Scott V, Votova K, Scanlan A, Close J. Multifactorial and functional mobility assessment tools for fall risk among older adults in community, home-support, long-term and acute care settings. *Age Ageing* 2007;**36**:130–9.

Shah MN, Rajasekaran K, Sheahan WD, Wimbush T, Karuza J. The effect of the geriatrics education for emergency medical services training program in a rural community. *J Am Geriatr Soc* 2008;**56**:1134–9.

Shyu YI, Tsai WC, Chen MC, Liang J, Cheng HS, Wu CC, *et al.* Two-year effects of an interdisciplinary intervention on recovery following hip fracture in older Taiwanese with cognitive impairment. *Int J Geriatr Psychiatry* 2012;**27**:529–38.

Skelton D. Exercise and falls prevention in older people. *CME Geriatric Medicine* 2007;**9**:16–21.

Smulders E, Enkelaar L, Schoon Y, Geurts AC, van Schrojenstein Lantman-de Valk H, Weerdesteyn V. Falls prevention in persons with intellectual disabilities: development, implementation, and process evaluation of a tailored multifactorial fall risk assessment and intervention strategy. *Res Dev Disabil* 2013;**34**:2788–98.

Soriano G, Roman E, Cordoba J, Torrens M, Poca M, Torras X, *et al.* Cognitive dysfunction in cirrhosis is associated with falls: a prospective study. *Hepatology* 2012;**55**:1922–30.

Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and non-fatal falls among older adults. *Injury Prevention* 2006;**12**:290–5.

Stewart J, McVittie C. Living with falls: house-bound older people's experiences of health and community care. *Eur J Ageing* 2011;**8**:271–9.

Umegaki H, Suzuki Y, Yanagawa M, Nonogaki Z, Endo H. Falls in elderly at high risk of requiring care. *Geriatr Gerontol Int* 2012;**12**:147–8.

Vellas B, Villars H, Abellan G, Soto ME, Rolland Y, Guigoz Y, *et al.* Overview of the MNA – Its history and challenges. *J Nutr Health Aging* 2006;**10**:456–65.

Webster J, Courtney M, O'Rourke P, Marsh N, Gale C, Abbott B, *et al.* Should elderly patients be screened for their 'falls risk'? Validity of the STRATIFY falls screening tool and predictors of falls in a large acute hospital. *Age Ageing* 2008;**37**:702–6.

Wu S, Keeler EB, Rubenstein LZ, Maglione MA, Shekelle PG. A cost-effectiveness analysis of a proposed national falls prevention program. *Clin Geriatr Med* 2010;**26**:751–66.

Population (clinical presentation not falls) (n = 6)

Augustine J. 'He keeps falling out' in the bathroom. *Emerg Med Serv* 2005;**34**:32.

Charney P, Sieber C, Cederholm T, Vellas B, Jensen G, Chumlea C, *et al.* Discussion. *J Nutr Health Aging* 2006;**10**:492–4.

Cooke M. Emergency care practitioners: a new safe effective role? *Qual Saf Health Care* 2006;**15**:387.

Rezende FM, Gaujac C, Rocha AC, Peres MP. A prospective study of dentoalveolar trauma at the Hospital das Clinicas, São Paulo University Medical School. *Clinics* 2007;**62**:133–8.

Shah MN, Lerner EB, Chiumento S, Davis EA. An evaluation of paramedics' ability to screen older adults during emergency responses. *Prehosp Emerg Care* 2004;**8**:298–303.

Swain AH, Hoyle SR, Long AW. The changing face of prehospital care in New Zealand: the role of extended care paramedics. *NZ Med J* 2010;**123**:11–14.

Study design and reporting format (none or entirely insufficient empirical data) (n = 24)

Augustine JJ. No history, no meds. The challenges from an elderly woman's fall just start with the broken hip. *EMS Mag* 2009;**38**:18–19.

Beghe C. Review: evidence from single studies shows that a few fall risk assessment tools can predict falls in elderly people. *Evid Based Med* 2007;**12**:186.

Brown K. Identifying patients at risk for falls. Intervention can improve outcomes and reduce repeat calls to 911. *EMS Insider* 2011;**38**:3–5.

Chase CA, Mann K, Wasek S, Arbesman M. Systematic review of the effect of home modification and fall prevention programs on falls and the performance of community-dwelling older adults. *Am J Occup Ther* 2012;**66**:284–91.

Clemson L. Prevention of falls in the community. *BMJ* 2010;**340**:c2244.

El Sanadi N. Geriatric injury prevention: a new role for EMS personnel? *Emerg Med Serv* 1996;**25**:51–3.

Emergency Department Management. Leverage front-line expertise to maximize trauma prevention efforts. *ED Manag* 2012;**24**:65–6.

Flavell E, Boyle M. Falls in the prehospital environment. *J Paramedic Practice* 2011;**3**:238–43.

Independent Nurse. Rapid community referral cuts falls. *Independent Nurse* 2010:4.

Lindstrom A. King County prevents falls. *JEMS* 2007;**32**:106.

Robitaille Y, Gauvin L. Fall prevention in older adults: towards an integrated population-based perspective. *Inj Prev* 2008;**14**:147–8.

Sadowski CA. Prevention of falls in older adults. *Can Pharm J* 2011;**144**:S17–18.

Scott T. Preventing emergency department admissions for older people. *Br J Community Nurs* 2011;**16**:392.

Shah MN, Rajasekaran K, Sheahan WD, Wimbush T, Karuza J. The effect of the geriatrics education for emergency medical services training program in a rural community. *J Am Geriatr Soc* 2008;**56**:1134–9.

Skelton DA, Hauer K, Lamb S. Re: 'Falls definition validation'. *Age Ageing* 2007;**36**:111–12.

Snooks H, Cheung WY, Close J, Dale J, Gaze S, Humphreys I, *et al*. Support and Assessment for Fall Emergency Referrals (SAFER 1) trial protocol. Computerised on-scene decision support for emergency ambulance staff to assess and plan care for older people who have fallen: evaluation of costs and benefits using a pragmatic cluster randomised trial. *BMC Emerg Med* 2010;**10**:2.

Snooks H, Anthony R, Chatters R, Cheung W-Y, Dale J, Donohoe R, *et al*. Support and assessment for fall emergency referrals (SAFER 2) research protocol: cluster randomised trial of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community-based care. *BMJ Open* 2012;**2**:6.

Snooks HA, Kingston MR, Anthony RE, Russell IT. New models of emergency prehospital care that avoid unnecessary conveyance to emergency department: translation of research evidence into practice? *Sci World J* 2013;182102.

Speechley M. Unintentional falls in older adults: a methodological historical review. *Can J Aging* 2011;**30**:21–32.

Spivack BS. Preventing falls: Direct intervention is needed. *BMJ* 2008;**336**:343.

Straus S. A 16-week tai chi programme prevented falls in healthy older adults. *Evid Based Med* 2008;**13**:54.

Udell J, Drahota A, Dean T, Sander R, Mackenzie H. Interventions for preventing falls in older people: an overview of Cochrane Reviews (Protocol). *Cochrane Database Syst Rev* 2011;**4**:CD009074.

van Hateren KJ, Kamper AM, Bilo HJ. Lack of evaluation of the effects of single preventive measures for falling. *Arch Intern Med* 2011;**171**:181–2.

Velhal G. Domestic accidents among the elderly. *Ann Trop Med Public Health* 2012;**5**:61–2.

Intervention (no intervention) (n = 14)

Close JC, Halter M, Elrick A, Brain G, Swift CG. Falls in the older population: a pilot study to assess those attended by London ambulance service but not taken to A&E. *Age Ageing* 2002;**31**:488–9.

Cone DC, Ahern J, Lee CH, Baker D, Murphy T, Bogucki S. A descriptive study of the 'lift-assist' call. *Prehosp Emerg Care* 2013;**17**:51–6.

Darnell G, Mason SM, Snooks H. Elderly falls: a national survey of UK ambulance services. *Emerg Med J* 2012;**29**:1009–10.

Dick T. Are you hip to this? Slip-and-fall injuries revisited. *Emerg Med Serv* 2003;**32**:32.

Halter M, Snooks H, Close J, Cheung WY, Moore F, Roberts SE. Emergency Care of Older People who fall: a missed opportunity. *Emerg Med J* 2006;**23**:321.

Newton JL, Kyle P, Liversidge P, Robinson G, Wilton K, Reeve P. The costs of falls in the community to the North East Ambulance Service. *Emerg Med J* 2006;**23**:479–81.

Page D. Is the lift worth it? New study concluded many lift-assist calls will result in callbacks. *JEMS* 2013;**38**:24–5.

Porter A, Snooks H, Youren A, Gaze S, Whitfield R, Rapport F, *et al.* 'Covering our backs': ambulance crews' attitudes towards clinical documentation when emergency (999) patients are not conveyed to hospital. *Emerg Med J* 2008;**25**:292–5.

Snooks HA, Halter M, Close JC, Cheung WY, Moore F, Roberts SE. Emergency care of older people who fall: a missed opportunity. *Qual Saf Health Care* 2006;**15**:390–2.

Snooks H, Cheung WY, Gwini SM, Humphreys I, Sanchez A, Siriwardena N. Can older people who fall be identified in the ambulance call centre to enable alternative responses or care pathways? *Emerg Med J* 2011;**28**:e1.

Spaite DW, Criss EA, Valenzuela TD, Meislin HW, Ross J. Geriatric injury: an analysis of prehospital demographics, mechanisms & patterns. *Ann Emerg Med* 1990;**19**:1418–21.

Thomas SL, Muscatello DJ, Middleton PM, Zheng W. Characteristics of fall-related injuries attended by an ambulance in Sydney, Australia: a surveillance summary. *NSW Public Health Bull* 2011;**22**:49–54.

Tiedemann A, Mikolaizak AS, Sherrington C, Segin K, Lord SR, Close JC. Older fallers attended by an ambulance but not transported to hospital: a vulnerable population at high risk of future falls. *Aust NZ J Public Health* 2013;**37**:179–85.

Weiss SJ, Chong R, Ong M, Ernst AA, Balash M. Emergency medical services screening of elderly falls in the home. *Prehosp Emerg Care* 2003;**7**:79–84.

Outcomes (outcomes for older people who fall not differentiated from those of other conditions) (n = 7)

Dale J, Williams S, Foster T, Higgins J, Snooks H, Crouch R, *et al.* Safety of telephone consultation for 'non-serious' emergency ambulance service patients. *Qual Saf Health Care* 2004;**13**:363–73.

Larsson B, Sixt E. Via Senetutis. Proceedings of the 7th Congress of the EUGMS, Malaga, Spain. 28–30 September 2011. *Eur Geriatr Med* 2011;**2**:144–5.

Mason S, O'Keeffe C, Knowles E, Bradburn M, Campbell M, Coleman P, *et al.* A pragmatic quasi-experimental multi-site community intervention trial evaluating the impact of Emergency Care Practitioners in different UK health settings on patient pathways (NEECaP Trial). *Emerg Med J* 2012;**29**:47–53.


Mason S, Wardrope J, Perrin J. Developing a community paramedic practitioner intermediate care support scheme for older people with minor conditions. *Emerg Med J* 2003;**20**:196–8.

O'Hara R, O'Keeffe C, Mason S, Coster JE, Hutchinson A. Quality and safety of care provided by emergency care practitioners. *Emerg Med J* 2012;**29**:327–32.

Pointer JE, Levitt MA, Young JC, Promes SB, Messana BJ, Ader ME. Can paramedics using guidelines accurately triage patients? *Ann Emerg Med* 2001;**38**:268–77.


Schmidt T, Neely KW, Adams AL, Newgard CD, Wittwer L, Muhr M, *et al.* Is it possible to safely triage callers to EMS dispatch centers to alternative resources? *Prehosp Emerg Care* 2003;**7**:368–74.

Appendix 4 Support and Assessment for Fall Emergency Referrals 2 organisational flow chart




Appendix 5 Station location maps

Welsh Ambulance Services NHS Trust Ambulance Stations and Acute Trusts




Map data ©2014 Google

London Ambulance Service Study Ambulance Stations and Acute Trusts



East Midlands Ambulance Service Study Ambulance Stations and Acute Trusts



Map data ©2014 Google

Appendix 6 Support and Assessment for Fall Emergency Referrals 2 postal consent pack

Safer 2 Participant Invitation Letter

Version 5

May 2011

**Ymddiriedolaeth GIG Gwasanaethau
Ambiwlans Cymru**

**Welsh Ambulance Services NHS
Trust**



Date

Dear

We are contacting you following your recent 999 call. The Welsh Ambulance Service is currently carrying out a study with Swansea University called SAFER 2, which is looking at how to improve the emergency care of older people.

As part of the study, we would like to follow up any 999 calls and unplanned A&E or hospital visits you may make over the next six months. You would not need to do anything for this to happen, as the information can be collected routinely. Please can you tick the box on the Consent Form enclosed if you are happy for us to see your information.

We have also included a questionnaire about your health and well-being for you (or someone who cares for you) to complete.

The information sheet provided gives further details about the study. If we haven't heard back from you after two weeks, we will telephone you to check whether you have received this letter and if you would like to talk to someone about it.

As a thank you for your time considering of these documents, we have enclosed a £5 voucher and a pen.

Please note that participation is voluntary and your care will not be affected in any way.

Yours sincerely,

Practice Research & Development Manager
Welsh Ambulance Services NHS Trust

Contact Details for Study

Helen Snooks
Professor of Health Services Research
SAFER 2 lead
Swansea University.

**Ymddiriedolaeth GIG Gwasanaethau
Ambiwlans Cymru**

Welsh Ambulance Services NHS Trust



**Swansea University
Prifysgol Abertawe**

SAFER 2: Participant Information Sheet

What is the study about?

The research study is helping to develop new ways for the ambulance service to work. Currently they either take people to the Accident and Emergency (A&E) department or leave them at home. Previous studies suggest these may not always be the best options for patients. This research looks at a new way of providing care in which paramedics can leave people at home when they do not need to go to hospital, but still make sure they receive the help they need.

Why have I been chosen?

We are writing to people aged 65 or over who have recently been attended by one of the paramedics who are taking part in the study.

Do I have to take part?

No. Participation is voluntary and there will be no impact on the care and treatment you receive, whether you take part or not.

What will happen if I do take part?

i) We would like to follow-up any further 999 calls or unplanned hospital attendances you may make over the next six months. To do this, we need permission to collect your information from ambulance service and hospital records. Once this has been done, all identifiable information will be removed.

ii) A questionnaire has been sent with this pack. If you are willing, we would like you to complete this and send it back in the freepost envelope provided.

The questionnaire asks about your general health and well-being following your 999 call and the type and quality of care you received. We have tried to keep the number of pages to the minimum possible.

If there is someone who is caring for you (a relative or friend), we are interested in their views as well, and they could complete the questionnaire on your behalf if you prefer this.

You will be sent a similar questionnaire again in six months time, which we would like you to complete and return to us. You can let us know if you are happy for this to happen by ticking the box at the end of the questionnaire.

Taking Part

A consent form is included with this letter. We would like you to read all the information carefully before deciding whether you wish to take part or not. If you have someone who helps look after you, please show them this information.

We would like you to tick one box, sign and date the form and return it in the FREEPOST envelope provided.

If you do not understand anything, would like some help or would like to talk to someone further - please call Moira Morgan on XXX, who will take your details. The best person to help you will then call you back as soon as possible.

Who will see my information?

All information is treated as confidential. Your personal details will only be seen by the clinical care team within the ambulance service until you have agreed to take part. Your contact details will then be passed onto the research team, who will ensure they are securely stored.

What happens with my information?

If you agree, a copy of the information will be given to the research team by the Ambulance Service.

Can I see what information of mine you are using?

Of course, under the Data Protection Act you have a right to see the information. Should you require this please contact Moira Morgan (study administrator) at Swansea University for a copy of the information. Contact details are at the end of this letter.

What if there is a problem?

We do not believe there will be any problems arising from your taking part in this trial. However, if there is anything you are not happy with, please, in the first instance, contact the study manager (details below). If you remain unhappy and wish to complain formally, please do so through the NHS complaints procedure.

What if I change my mind?

If you change your mind and do not wish to continue your participation in the study, you are free to withdraw at any time. Please contact Moira Morgan on the number given in this sheet. Your care will remain unaffected.

Who is organising and funding the research?

The study is organised by researchers in the School of Medicine at Swansea University, in collaboration with the Welsh Ambulance Services NHS Trust (WAST). It is funded by the Health Technology Assessment programme at the Department of Health.

Who has reviewed the study?

The study has been reviewed by Wales Research Ethics Committee and by the Health Technology Assessment programme.

Contact Details

Study Manager College of Medicine,
 Swansea University
 Tel: XXX

Study Administrator College of Medicine,
 Swansea University

Tel: XXX

*Safer 2 Participation Form**Version 4**May 2011*

**Ymddiriedolaeth GIG Gwasanaethau
Ambiwlans Cymru**

Welsh Ambulance Services NHS Trust



SAFER 2 Consent Form

Please tick one of the boxes and sign the form below.

(The care and treatment you receive will not be affected in any way by your response.)

I give consent for the study team to follow up any 999 calls, A&E attendances or hospital visits I may make for 6 months.

I do not wish to participate in the SAFER 2 study.

Signed _____ Date _____

If someone else has filled this form in for you:

Form completed by _____ Relationship _____

Signed _____ Date _____

Please return the completed form in the FREEPOST envelope provided

Name _____

Study ID

--	--	--	--	--

Appendix 7 Support and Assessment for Fall Emergency Referrals 2 1-month questionnaire



SAFER 2 one month questionnaire v4 May 2011

Study Number:



SAFER 2

ONE MONTH QUESTIONNAIRE

CONFIDENTIAL

Date of questionnaire completion $\overset{d}{\square} \overset{d}{\square} / \overset{m}{\square} \overset{m}{\square} / \overset{y}{\square} \overset{y}{\square} \overset{y}{\square} \overset{y}{\square}$
 / / 20

Is someone completing this survey on your behalf YES NO

Please let us know their relationship to you _____

If you would like help with this questionnaire,
please telephone XXX

If there is no answer, please leave your details
and a member of the SAFER 2 team will return
your call as soon as possible.

Please return the completed questionnaire in
the FREEPOST envelope provided.

THANK YOU



52519

Study Number

--	--	--	--	--

PLEASE READ THESE INSTRUCTIONS CAREFULLY

- ▶ Please use a blue or black pen, not a pencil
- ▶ Please mark your answers with an X clearly inside the box
- ▶ Please answer every question
- ▶ If you find it difficult to answer a question, do the best you can

SECTION A: 999 Care

d d m m y y y y

This section asks about the care you received from the ambulance service on / /

A1. Overall, how would you rate your general health before this call?

Excellent Good Fair Poor Very Poor

A2. Do you feel the medical condition that you called 999 for was...

Extremely Serious Very Serious Moderately Serious Slightly Serious Not Serious

Very Dissatisfied Dissatisfied Neutral Satisfied Very Satisfied

A3. How do you rate the following:

a) Waiting time for the ambulance to arrive?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b) Amount of time ambulance person spent with you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c) Decisions made by the person who attended you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d) The ambulance persons' concern about you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e) Receiving satisfactory answers to your questions from the ambulance person?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f) Explanation of any further referrals made by the ambulance person?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



52519

Study Number:

--	--	--	--	--	--

	Very Dissatisfied	Dissatisfied	Neutral	Satisfied	Very Satisfied
g) The thoroughness of the care you received?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h) The overall quality of the care you received from the ambulance service?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i) The outcome of your 999 call?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

SECTION B: Care and help needed

The next section asks questions about the care and help you may have needed following your 999 call

B1. **Please circle** how many times have you had a fall during the past month?

0 1 2 3 4 5 6 7 8 9 10

B2. **Please circle** the number of times you have had contact with each service related to your fall and 999 call. Please mark all the answers.

a) GP telephone advice	0	1	2	3	4	5	6	7	8	9	10
b) GP surgery visit	0	1	2	3	4	5	6	7	8	9	10
c) GP home visit	0	1	2	3	4	5	6	7	8	9	10
d) NHS Direct	0	1	2	3	4	5	6	7	8	9	10
e) Community nurse visit	0	1	2	3	4	5	6	7	8	9	10
f) Out-Patient attendance	0	1	2	3	4	5	6	7	8	9	10
g) Other Social Service provision (eg home help)	0	1	2	3	4	5	6	7	8	9	10

Please state _____



Study Number:

The next section asks questions about the care and help you may have needed following your fall and how this has affected you and those who care for you.

B3. a. Please let us know your CURRENT place of residence (please tick)

Own home

Staying with relatives

Residential Home

Hospital in-patient

Other

Please state _____

b. Is this different from your NORMAL place of residence YES NO

c. If yes, please indicate your normal place of residence

B4. Please let us know what (if any) special equipment or furniture Social Services have installed in your home to help you since your fall?



Study Number

--	--	--	--	--

SECTION C: Your health now

For this section exploring health-related quality of life we used version 2 of the Short Form questionnaire-12 items (SF-12).¹⁰⁵



52519

Study Number

--	--	--	--	--

SECTION D: Fear of falling

This section is about how confident you are about being able to do things without falling.

Please circle your answer for each of the activities below, with 0 meaning "not confident at all", 5 meaning "fairly confident" and 10 meaning "completely confident"

How confident are you that you can	Not confident at all					Fairly confident					Completely confident				
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
D1. Get dressed and undressed	0	1	2	3	4	5	6	7	8	9	10				
D2. Prepare a simple meal	0	1	2	3	4	5	6	7	8	9	10				
D3. Take a bath or a shower	0	1	2	3	4	5	6	7	8	9	10				
D4. Get in/out of a chair	0	1	2	3	4	5	6	7	8	9	10				
D5. Get in/out of bed	0	1	2	3	4	5	6	7	8	9	10				
D6. Answer the door or telephone	0	1	2	3	4	5	6	7	8	9	10				
D7. Walk around the inside of your house	0	1	2	3	4	5	6	7	8	9	10				
D8. Reach into cupboards or wardrobes	0	1	2	3	4	5	6	7	8	9	10				
D9. Do light housekeeping	0	1	2	3	4	5	6	7	8	9	10				
D10. Do simple shopping	0	1	2	3	4	5	6	7	8	9	10				
D11. Use public transport	0	1	2	3	4	5	6	7	8	9	10				
D12. Cross roads	0	1	2	3	4	5	6	7	8	9	10				
D13. Do light gardening or hang out the washing (please rate whichever you do most frequently)	0	1	2	3	4	5	6	7	8	9	10				
D14. Using front or rear steps at home	0	1	2	3	4	5	6	7	8	9	10				



Study Number

--	--	--	--	--

SAFER 2

*Please note - these details will be kept separately
from the questionnaire*

I am happy to be sent a questionnaire in 6 months

YES

NO

We would like to contact a small number of people to talk to them face to face about their experiences. If you do not mind being contacted by a member of the research team, please fill in your details below.

Name _____

Address _____

Phone No. _____

*Thank you very much for your time and effort
in completing this questionnaire*

Appendix 8 Final definition of the intervention

TABLE 43 Final definition of the intervention

Criterion	Core essential	To be agreed locally	East Midlands local agreement	London local agreement	Wales local agreement	Source
Training	<p>Training to take the form of half- to 1-day face-to-face content followed up with home study DVD and competency assessment. Sheffield clinicians to train the trainers. Local ambulance service trainers to cascade training to paramedics. The core of the intervention is the paramedic protocol. Training will therefore be concerned with tailoring existing skills rather than providing education and training for new skills</p>	<p>Precise length and arrangements for face-to-face training and competency assessment. Processes for cascade training</p>	<p>No cascade training. Training form is variable, from group sessions with the Sheffield trainers to one-to-one sessions with the study research champion and research associate</p>	<p>Sheffield team will deliver training to assigned training officers and RSO at intervention stations. Training officers to deliver training to study paramedics across three sessions with one-to-one sessions if necessary</p>	<p>The RSO attended a train the trainer day with the Sheffield Training Team. Training was then cascaded through the RSO to groups of paramedics, with one-to-one sessions as necessary</p>	<p>As specified in original proposal. Discussed and modified at definitions meeting</p>
Assessment tool	<p>Two elements:</p> <ol style="list-style-type: none"> assessment of safety for non-conveyance: red flag approach to patient assessment to ensure that factors that would indicate transfer to A&E are identified assessment of risk of further falls: basic falls risk assessment (based on FRAT tool) 	<p>To be used as an aide memoire or a specific additional form to PRF. Consent process for patient agreement to be referred to falls service to be confirmed</p>	<p>The assessment tool is to be used as an aide memoire. Paramedics have also been given a list of GP surgeries that patients have to be registered to in order to be eligible for referral to falls service. Paramedics receive verbal consent from the patient for referral to the falls service</p>	<p>The assessment tool is to be used as an aide memoire with an accompanying Older Person who has Fallen Assessment Form which will log the outcome of the assessment. If the decision is made to refer the patient to the falls team, the patient will be informed; if referral is refused, this will be recorded on the PCR</p>	<p>The assessment tool is to be used as an aide memoire with an accompanying Older Person who has Fallen Assessment Form which will log the outcome of the assessment. If the decision is made to refer the patient to the falls team, the patient will be informed; if referral is refused, this will be recorded on the PCR</p>	<p>As specified in original proposal. Discussed and modified at definitions meeting</p>
Referral tool	<p>Basic demographic, clinical and contact details to be completed in order to make referral to falls service</p>	<p>Specific tool to be agreed locally. Based on Welsh Ambulance Services NHS Trust form</p>	<p>Referral document produced in conjunction with EMAS, the falls services and the research team</p>	<p>Referral document produced in conjunction with the LAS EBS, the falls services and the research team</p>	<p>Referral document agreed locally with Welsh Ambulance Services NHS Trust, falls service and control room representatives</p>	<p>As specified in original proposal. Discussed and modified at definitions meeting</p>

Criterion	Core essential	To be agreed locally	East Midlands local agreement	London local agreement	Wales local agreement	Source
Process of referral	It is essential that there is some system in place to allow auditing of receipt of referral forms by falls service	The process and medium of referral from ambulance service to falls service to be agreed locally (referral via ambulance control, direct from scene or from station between jobs, by fax, or other route)	Referral made via control room via a specific telephone number (the 'Safeguarding number', as this line is also used for social services referrals). Referral taken by an emergency medical dispatcher (24/7). Project support officer collects referral forms from control room and faxes the referral to the appropriate falls service	Referral will be made through a single-point-of-access within LAS (the EBS desk). Access will be via assigned radio channel. EBS will fill out a referral proforma based on the information provided by the referring paramedic. EBS will call the appropriate falls service during office hours and will confirm receipt of referral and transfer proforma via fax	Control will be informed of the referral by the paramedic who will inform the falls service by faxing the locally agreed referral form to a single point-of-contact number. The referral form will record date and time referral received and the RSO will be informed by both control and the falls team	Discussed at final definitions meeting
Falls service response	Falls service to follow NICE guidance in terms of minimum follow-up of patient. Non-clinical assessment telephone contact with patient initially. Call to be made by a professional with the ability to recognise cases where more urgent input is needed and refer if necessary. Initial assessment of patient by an appropriately skilled and experienced individual health-care professional to take place following initial telephone contact. The timescale should fit with NICE guidance. Patient to be fed into an appropriate multifactorial intervention process following this assessment (based on NICE guidance)	Period of follow-up to be defined and agreed locally but should adhere to NICE guidance	Follow-up period variable depends on clinical need (which is recorded on the referral form). In all four of the locally participating falls services, the patient is called to book a home visit, the home visit is under taken, and then the patient is fed into the appropriate pathway. The patient is rung to assess clinical need and to arrange a home visit	Timescales agreed according to NICE guidance	Following referral, telephone contact is to be made by a district nurse within 72 hours and an in-home assessment will take place within 1 week	As specified in original proposal. Discussed and modified at definitions meeting

continued

TABLE 43 Final definition of the intervention (*continued*)

Criterion	Core essential	To be agreed locally	East Midlands local agreement	London local agreement	Wales local agreement	Source
Falls service make-up	Appropriate skill set based on NICE criteria	Specific details of service and patient intervention to be recorded locally	The home visit is undertaken to assess which 'treatment stream' the patient is placed into; subsequent treatment is provided by occupational therapists, physiotherapists, and nurses. Some people may be referred to doctors, social workers, dieticians and psychologists	Specific details of service and patient intervention to be recorded by relevant falls service. Subsequent treatment is provided by occupational health, physiotherapists and rehabilitation	Within Wales, the Cwm Taf District Nursing Teams will be responsible for receiving the referral and making the initial phone call and assessment. Depending on assessment outcome, further services will be available to the patient. Further services include the intermediate care and rehabilitation service, The National Exercise Referral Scheme Falls Prevention Programme or GP	As specified in original proposal. Discussed and modified at definitions meeting

Criterion	To be agreed locally	East Midlands local agreement	London local agreement	Wales local agreement	Source
Clinical support/ feedback loop	<p>Clinical support debriefs with A&E every 2–4 weeks. Feedback loop between falls services and paramedics to report outcomes for individuals referred essential</p> <p>Specific feedback mechanisms to be agreed locally</p>	<p>Paramedics able to call/ e-mail research team to ask questions. Clinical or operational questions to be forwarded to an EMAS clinical tutor or research champion for answering. PRFs audited once a month to see if the correct path was followed. The Falls Services will complete a 'data collection sheet' for every patient referred – these are to be collected from the falls service, and 'feedback information' transferred to a 'paramedic feedback sheet' (i.e. referral date, date assessed and outcome of assessment). These details will be e-mailed to the referring paramedic</p>	<p>Details on referral rates and paramedic queries will be fed back to the study team at regular intervals and used for clinical feedback to participating paramedics. Where possible falls services to provide both short- and long-term patient outcomes for feedback. On-station clinical and operational support delivered through RSO (LAS paramedic on secondment to study)</p>	<p>A data collection and paramedic feedback form has been agreed locally which records the main outcomes of the initial assessment. Once the patient has been assessed the district nurse will send the feedback to each paramedic. To support paramedics clinically and operationally a support network has been put in place. This involved the main study team for general inquiries, the RSO for training and ongoing use of the decision tool and referral pathways and a clinical on calls system which is available to paramedics to address clinical queries. Paramedics use of the decision tool and referral pathways will be monitored by the research team</p> <p>paramedics contacted to discuss issues</p>	<p>As specified in original proposal. To be discussed and agreed</p>

DVD, digital versatile disc; EBS, Emergency Bed System; EMAS, East Midlands Ambulance Service; FRAT, Falls Risk Assessment Tool; LAS, London Ambulance Service; PCR, Patient Clinical Record; RSO, research support officer.

Appendix 9 Support and Assessment for Fall Emergency Referrals 2 paramedic training session programme



SAFER 2 Support and Assessment for Fall Emergency Referrals Paramedic Training Session Programme

11:00 a.m. – 11:15 a.m.	Welcome and introduction <i>Aims, scope, time-line, paramedic involvement.</i>
11:15 a.m. – 11:30 a.m.	Teaching of clinical assessment tools <i>Watching of SAFER 2 clinical assessment DVD with discussion. Includes social and falls history and clinical examination to rule out the need for referral to the Emergency Department.</i>
11:30 a.m. – 12:00 p.m.	Falls service overview and protocol for referring to falls service <i>Falls service make-up, falls prevention interventions, referral protocol.</i>
12:00 p.m. – 12:15 p.m.	Clinical decision making and use of the paramedic clinical decision flow chart
12:15 p.m. – 12:45 p.m.	Lunch
12:45 p.m. – 1:15 p.m.	SAFER 2 patient scenarios <i>Teaching of clinical assessment tools through likely scenarios.</i>
1:15 p.m. – 2:00 p.m.	Assessment of competence <i>Assessed clinical scenario.</i>
2:00 p.m. – 2:15 p.m.	Question and answer session. Summary and close. <i>Including details of clinical support available during the trial.</i>

Appendix 10 Support and Assessment for Fall Emergency Referrals 2 6-month questionnaire



60784

SAFER 2 six month questionnaire V1 Sep 2011

Study Number:

--	--	--	--	--



SAFER 2

SIX MONTHS QUESTIONNAIRE

CONFIDENTIAL

Date of questionnaire completion $\begin{matrix} d & d \\ \square & \square \end{matrix}$ / $\begin{matrix} m & m \\ \square & \square \end{matrix}$ / $\begin{matrix} y & y & y & y \\ 2 & 0 & \square & \square \end{matrix}$

Is someone completing this survey on your behalf YES NO

Please let us know their relationship to you _____

If you would like help with this questionnaire,
please telephone XXX

If there is no answer, please leave your details
and a member of the SAFER 2 team will return
your call as soon as possible.

Please return the completed questionnaire in
the FREEPOST envelope provided.

THANK YOU



60784

Study Number

--	--	--	--	--

PLEASE READ THESE INSTRUCTIONS CAREFULLY

- ▶ Please use a blue or black pen, not a pencil
- ▶ Please mark your answers with an X clearly inside the box
- ▶ Please answer every question
- ▶ If you find it difficult to answer a question, do the best you can

SECTION A: 999 Care

This section asks about the care you received from the ambulance service on

d	d	m	m	y	y	y	y	y	y

A1. Overall, how would you rate your general health before this call?

Excellent Good Fair Poor Very Poor

A2. Do you feel the medical condition that you called 999 for was...

Extremely Serious Very Serious Moderately Serious Slightly Serious Not Serious

	Very Dissatisfied	Dissatisfied	Neutral	Satisfied	Very Satisfied
A3. How do you rate the following:					
a) Waiting time for the ambulance to arrive?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b) Amount of time ambulance person spent with you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c) Decisions made by the person who attended you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d) The ambulance persons' concern about you?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e) Receiving satisfactory answers to your questions from the ambulance person?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f) Explanation of any further referrals made by the ambulance person?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



60784

Study Number:

--	--	--	--	--

	Very Dissatisfied	Dissatisfied	Neutral	Satisfied	Very Satisfied
g) The thoroughness of the care you received?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h) The overall quality of the care you received from the ambulance service?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i) The outcome of your 999 call?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

SECTION B: Care and help needed

The next section asks questions about the care and help you may have needed after your 999 call and over the following six months.

B1. Thinking back to the time after your 999 call, please circle the number of contacts you had with each service **because** of the condition you called for. Please answer all the questions.

a) GP telephone advice	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
b) GP surgery visit	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
c) GP home visit	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
d) NHS Direct	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
e) Community nurse visit	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
f) Out-Patient attendance	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		
<hr/>												
g) Other Social Service provision (eg home help)	0	1	2	3	4	5	6	7	8	9	10	11
	12	13	14	15	16	17	18	19	20	More than 20		

Please state: _____



Study Number:

B2. a. Please let us know your CURRENT place of residence (please tick)

Own home

Staying with relatives

Residential Home

Hospital in-patient

Other

Please state _____

b. How long have you been resident here? _____

c. Is this different from your NORMAL place of residence YES NO

d. If yes, please indicate your normal place of residence _____

B3. Please let us know what (if any) special equipment or furniture Social Services have installed in your home to help you since your fall?

B4. Please circle how many times in total you have had a fall during the past **SIX** months?

0 1 2 3 4 5 6 7 8 9 10 11

12 13 14 15 16 17 18 19 20 more than 20



60784

Study Number

--	--	--	--	--

For this section exploring health-related quality of life we used version 2 of the Short Form questionnaire-12 items (SF-12).¹⁰⁵



60784

Study Number

--	--	--	--	--

SECTION D: Fear of falling

This section is about how confident you are **now** about being able to do things without falling.

Please circle your answer for each of the activities below, with 0 meaning "not confident at all", 5 meaning "fairly confident" and 10 meaning "completely confident"

How confident are you that you can	Not confident at all					Fairly confident					Completely confident				
D1. Get dressed and undressed	0	1	2	3	4	5	6	7	8	9	10				
D2. Prepare a simple meal	0	1	2	3	4	5	6	7	8	9	10				
D3. Take a bath or a shower	0	1	2	3	4	5	6	7	8	9	10				
D4. Get in/out of a chair	0	1	2	3	4	5	6	7	8	9	10				
D5. Get in/out of bed	0	1	2	3	4	5	6	7	8	9	10				
D6. Answer the door or telephone	0	1	2	3	4	5	6	7	8	9	10				
D7. Walk around the inside of your house	0	1	2	3	4	5	6	7	8	9	10				
D8. Reach into cupboards or wardrobes	0	1	2	3	4	5	6	7	8	9	10				
D9. Do light housekeeping	0	1	2	3	4	5	6	7	8	9	10				
D10. Do simple shopping	0	1	2	3	4	5	6	7	8	9	10				
D11. Use public transport	0	1	2	3	4	5	6	7	8	9	10				
D12. Cross roads	0	1	2	3	4	5	6	7	8	9	10				
D13. Do light gardening or hang out the washing (please rate most frequent)	0	1	2	3	4	5	6	7	8	9	10				
D14. Using front or rear steps at home	0	1	2	3	4	5	6	7	8	9	10				

THANK YOU

Appendix 11 Support and Assessment for Fall Emergency Referrals 2 data analysis plan



V0.9 (18-11-2013)

SAFER 2 DATA ANALYSIS PLAN

Trial Title

SAFER 2: Support and Assessment for Fall Emergency Referrals 2

Trial Summary

Care of older people who fall: evaluation of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community based care

Trial Details

Trial Registration: ISRCTN 60481756
Chief Investigator: H.A. Snooks, Swansea University

Revision History

Revision Date	Release	Summary of Changes	Changes Made by
09/2009	1.0	Initial version	WYC
03/2010	1.1	Revised to include economic analysis	WYC
04/2010	1.2	Revised to include qualitative analysis	WYC
05/2010	1.3	Revised with protocol	HS
07/2010	1.4	Revised randomisation, PGI, non-consenter & non-respondent analyses	WYC
09/2010	0.4	Version numbers revised by DMEC 20/07/10	JP
02/2012	0.5	Revised with protocol	AJW
03/2012	0.6	Revised with protocol	SG
05/2012	0.7	Revised economic & qualitative analyses	BA
07/2012	0.8	Revised for DMEC meeting on 24/07/12	AJW & ITR
05/2013	0.9	Further revisions	SG, IH, AJW

Approvals

This document requires the following approvals. A signed copy should be placed in the project files.

Name	Signature	Title	Date of Issue	Version

Distribution

This document has been distributed to:

Name	Title	Date	Version
SAFER 2 DMEC		20-07-2010	1.4
SAFER 2 DMEC		24-07-2012	0.8
SAFER 2 DMEC/TSC		28-11-2013	0.9

Section 1: SUMMARY AND OVERVIEW

SAFER 2 is a pragmatic cluster randomised controlled trial (CRT) with a qualitative component, involving 25 ambulance stations from three UK centres (namely, the East Midlands, London, and Wales)^[1]. 223 paramedics taking part in the study have recruited approximately 6000 patients who call 999 following a fall. Half of the participating stations have been randomly allocated to an intervention group, and paramedics based at these stations deliver care to older people who have fallen according to new protocols which allow emergency ambulance paramedics to assess and refer patients to appropriate community based care. The control group comprises the remaining stations, and paramedics based at these deliver normal care.

Randomisation of stations to groups was carried out in accordance with the principles outlined in WWORTH SOP24 Randomisation^[2]; specifically, randomisation was undertaken *after* paramedics in study stations had volunteered to participate (thus avoiding possible selection bias), and used centre and the number of eligible calls attended (based on data from 2009) as stratification variables. Clinical relevance is related to, but not determined by, the criterion of one event avoided (or incurred) in ten, which was the clinically significant effect size used in the sample size calculation. The calculated sample size for binary outcomes (whether an event occurred or not) is also adequate for a statistical assessment of the observed differences in times to the first event in the intervention and control group.

This analysis plan for SAFER 2 has been developed using appropriate CONSORT (www.consort-statement.org) statements and checklists, including those relating to cluster trials and patient-reported outcomes.

Section 2: TRIAL OBJECTIVES

The overall aim of this trial is to assess the benefits and costs to patients and the NHS of a complex intervention comprising education, clinical protocols and pathways enabling paramedics to assess older people who have fallen and refer them to community-based falls services when appropriate. Specific objectives are to:

- 1 Compare outcomes, processes and costs of care between intervention and control groups:
 - a patient outcomes: rate and pattern of subsequent emergency healthcare contacts or deaths, for any reason and for falls; health related quality of life (HRQoL); psychological status (especially fear of falling); and change in place of residence
 - b processes of care: pathway of care at index fall; subsequent healthcare contacts; ambulance service operational indicators and protocol compliance including clinical documentation
 - c costs of care: provided by NHS and personal social services; incurred by patients and carers in seeking care;
- 2 Estimate wider system effects of the introduction of the intervention on ambulance service performance and costs;
- 3 Understand how patients experience the new health technology;
- 4 Identify factors which facilitate or hinder the use of the intervention;
- 5 Inform the development of methods for falls research, especially outcome measures recommended for trials of interventions for older people who fall^[3]

Section 3: DATA COLLECTION & HANDLING

3.1 Data Sources & Collection

Our primary outcome is “further emergency healthcare contacts (999 call or Emergency Department (ED) attendance or hospital admission) or death for any reason and for falls per recruited faller, and time to first contact or death” for six months following the index call. This CRT does not approach participants at the point of treatment, because they may be in distress and unable to give informed consent, but seeks instead retrospective consent to follow up through routine medical records and by postal questionnaire. We planned to follow-up all our patients anonymously in order to achieve the primary outcome for our target sample size. In Wales, arrangements for this process are already in place through the existing SAIL (Secure Anonymised Information Linkage) Database, and our application for permission through the Information Governance Review Panel has been granted. We applied to the NIGB Ethics and Confidentiality Committee for permission to follow-up our patients anonymously through the Information Centre in England and permission has been granted for this process. Conditional to this permission, people who have clearly dissented to identifiable follow-up must be excluded from the study. For clarity, if a patient has at any time declined to be part of the study (either in returning a consent form or orally on the telephone) they will be removed from the study.

3.2 Sample Size and Statistical Power

We made the conservative estimate that trial patients have about 50% chance of making another emergency contact within six months; in the absence of intra-cluster correlation (ICC), a sample of 4190 evaluable participants yields 90% power, when using two-sided 5% significance level, of detecting a change in this chance from 50% to <45% or to >55%. The analogous outcome in the SAFER 1^[4] trial indicated an ICC of zero in clusters of participants seen by paramedics from the same ambulance station. To be conservative, we therefore allowed for an ICC of 0.002, and sought 251.6 (calculated from $0.998 \cdot 4190 / \{25 - 0.002 \cdot 4190\}$) participants per ambulance station, making a total sample size target of 6290.

However, actual recruitment is slightly lower, at 5939 participants, and we will lose approximately 1000 patients that decline to be part of the study. Furthermore, previous experience suggests that we shall also lose about 10% of the remaining patients from analysis because we shall not be able to match them with their routine electronic data on contacts with the NHS. Hence we expect the recruited sample of 5939 participants to yield only 4400 suitable for accurate information linkage. With an ICC of 0.002 and equal cluster sizes, this expected evaluable sample size corresponds to 3260 independent participants, which, in turn, equates to approximately 80% statistical power. In short, the cumulative loss in recruitment, and from dissenting and un-matched patients will reduce the statistical power of SAFER 2 from a conservative 90% to the traditional 80%, even allowing for more intra-cluster correlation than observed in the SAFER 1^[4] trial.

3.3 Missing Data

General Principles

As per WWORTH SOP28 Statistics^[5], we shall adopt a consistent approach to missing data relating to both effectiveness and cost-effectiveness except where individual outcome measures require some variation in that approach. We shall exclude participants without follow-up data, and, for each variable, summarise the frequency of missing data, which directly influences the sample size in, and hence the statistical power of, some analyses. If there is reason to suspect that data are not Missing Completely At Random (MCAR), the trial statistician and chief investigator will discuss the findings. If there is no reason to suspect that data are not MCAR, we shall use appropriate imputation methods to mitigate the problem of missing data.

Internal imputation of HRQoL data at a particular data collection point

None of the questionnaires (SF-12, derived SF-6D, or Modified Falls Efficacy Scale (mFES)^[6]) has an official algorithm for imputing individually missing answers. Some patients may have particular reasons for missing some of the questionnaire items, and it is unlikely that a plausible model for conditional Missing At Random can be established even on a case-by-case basis. Nevertheless, to minimise missing values and use available information, any such missing values within patient interviews will be completed by imputation^[7] within the reduced dataset of individual responses to questions in the three measures, using the regression or expectation maximization (EM) algorithms of the Missing Value Analysis module within SPSS. Scale scores will then be calculated according to the relevant instructions for the measure. The use of multiple imputation methods will be considered if the incidence of missing data is high.

External imputation of HRQoL data scores

If a participant is dead at the data collection point, the SF-6D score will be taken as zero. To avoid outliers, SF-12 and mFES scores will be taken as the minimum value observed for that measure in the relevant treatment group

Otherwise, missing summary scale and subscale scores will be imputed by regression from all available values of that score at other data points and the allocated treatment group. Further predictors (eg gender, age, current hospitalisation) may be included, unless they are already used as covariates in the main analysis

If any potential predictor other than study group has an F value of less than 1 (that is, it increases the standard error of the prediction) it will be removed from the list of predictors and the prediction recalculated.

Missing Health Economic data

Addressing the problem of missing data may involve employing Mean Imputation (that is, assigning a mean value to the respondents with missing data) or Regression Imputation (that is, using regression models to provide estimates of missing data from complete data, when the missing data is part of a multivariate data set). The usual method for dealing with censored data relating to costs will be to employ the weighted cost method with known cost histories^[8].

3.4 Withdrawals

Patients can withdraw from the study whenever they wish, and do not have to give a reason, although any reasons given must be documented. Their subsequent treatment will not be affected in any way. Patients may also withdraw from the questionnaire element of the study but be retained for other follow up. Any patients lost must be traced and documented whenever possible.

Wherever possible, data missing because of withdrawal will be imputed from available data.

Section 4: LINKS BETWEEN OBJECTIVES & OUTCOMES

4.1 Trial Objective 1 'Compare costs, processes and outcomes of care between experimental and control groups'

Objective 1a Patient Outcomes

- | | | |
|---|---|--------------------------|
| 1 | Number of further reported contacts or deaths | [primary outcome] |
| 2 | Time to the first reported contact or death | [primary outcome] |

- Primary outcomes will be analysed using the hierarchy of outcomes (comprising 999 calls, ED attendances, emergency admissions, and death) established in SAFER1^[4].
- 3 Injuries related to further falls **[secondary outcome]**
This will be summarised by cross-tabulation of the numbers and proportions in each study group, split by types of injuries.
 - 4 Health-related Quality of Life **[secondary outcome]**
This is measured by SF-12 and derived SF-6D, and will be assessed and summarised at one month and six months.
 - 5 Satisfaction **[secondary outcome]**
This is measured by the Quality of Care Monitor, and will be assessed and summarised at one month.
 - 6 Psychological status **[secondary outcome]**
This is measured by the mFES, and will be assessed and summarised at one month and six months.
 - 7 Change in place of residence at six months **[secondary outcome]**

Objective 1b Processes of Care

- 1 Pathway of care **[secondary outcome]**
This is measured by health and social care contacts associated with the index fall and during follow-up period, split by types of contacts and adjusted by covariates.
- 2 Ambulance service operational indicators **[secondary outcomes]**
These comprise episode of care and job cycle times, adjusted by covariates.
- 3 Protocol compliance **[secondary outcome]**
This will be measured by a series of data items related to protocol compliance (such as the completion of clinical documentation; referral processes).

Objective 1c Cost of Care **[secondary outcome]**

This objective is addressed by estimating and comparing the resources utilised in both groups, adopting a 'bottom-up approach' from the perspective of the NHS, personal social services, and patients and their families. Specifically:

- 1 Estimation of costs of providing the intervention
Data relating to direct costs to the NHS will be assessed using data logged as a part of routine practice and from resource utilisation recording sheets, together with reference to patient records and discussions with relevant finance staff. Resources used will be translated into costs using relevant published unit costs.
The number of non-conveyances will be logged, and potential impact on costs to the ambulance service and hospital emergency departments estimated.
Further, the additional costs to community services will be documented and costs using relevant published unit costs.
- 2 Costs to patients and families
Patients and their families will complete a self-administered questionnaire specifically designed for this study but based on other instruments catalogued within the Database of Instruments for Resource Use Management (www.dirum.org).

Further, data on the use of the health service and social services resources will be collected for each patient using a combination of paramedic records and routine hospital records (for hospital events). Data relating to social services costs will be derived from discussion with relevant social services departments

4.2 Trial Objective 2 'Estimate wider system effects of the introduction of the intervention on ambulance service performance and costs'

We consider this objective two parts.

Objective 2a Wider effect on operational indicators

Ambulance service operational indicators **[secondary outcomes]**

Response times across the study catchment and surrounding areas during the trial period will be compared to pre-trial response times and 'rest of service' response times (adjusted for pre-trial differences) in order to identify any knock on or halo effects.

Objective 2b Wider effect on Costs of Care [secondary outcome]

Consequences of the scheme for the wider NHS and social services (eg ED attendances, inpatient admissions, GP consultations, out-of-hours GP contacts, NHS Direct contacts, social services utilisation) will be estimated using a combination of routine data sources, discussions with staff from relevant departments (eg social services) and from responses to patient completed questionnaires that highlight service contacts and utilisation..

Further, the time spent by ambulance staff with patients and the nature of their involvement will be documented so as to determine the direct cost of patient care, cost of patient transportation and other costs – reflecting travel, time spent at ED discharging patients and so on.

4.3 Trial Objective 3 'Gain an in-depth understanding of how the intervention is experienced by patients'

This objective is addressed using face-to-face semi-structured interviews conducted in co-operation with a carer or older person present at time of fall. Qualitative interviews to be undertaken with patients who: experienced a fall; were recruited to the trial; seen by an intervention paramedic and agreed to the interview. This will include patients: who were taken to ED; referred to a falls service; and patients who were neither taken to the ED nor referred to a falls service.

The aims of the interviews are to explore:

- Patient experience of ambulance service
- Patient experience of those seen by a falls service
- Patient health since fall
- Patient satisfaction with treatment

4.4 Trial Objective 4 'Understand how the intervention is delivered in practice, identifying factors which enable or hinder its use'

This objective is addressed using focus groups carried out with intervention paramedics 'pre' and 'post' trial. Focus groups will be carried out by two researchers, one to lead discussions and one to take notes and enable the linkage of texts to speakers as well as concerning other details, such as points of consensus or disagreement, issues that drew strong emotional responses such as anger, fear or anxiety.

The aims of the focus groups are to explore:

- Paramedics views and attitudes toward the new intervention
- Any preconceptions about the new way of working

- Factors which enable the use of the new referral pathway
- Factors which hinder the use of the new referral pathway

4.5 Trial Objective 5 'Inform the development of methods for falls research in relation to outcomes assessment'

We shall compare mFES scores with SF-12 component scores and derived SF-6D scores as recommended by Prevention of Falls Network Europe (ProFaNE)^[9] group to establish construct validity. We shall assess predictive validity by comparing mFES scores with the mean number of further falls and time to first subsequent fall.

Section 5: MAIN ANALYSES

5.1 Statistical Analysis

This will conform to principles outlined in WORTH SOP28 Statistics^[5]; specifically, the primary analysis will be by 'treatment allocated', reflecting the pragmatic nature of the trial design where patients are cared for by paramedics based at ambulance stations randomly allocated to deliver intervention or control care.

Primary outcomes comprise a hierarchy relating to further emergency healthcare contacts for falls, as measured by contacts with the 999 service, Emergency Department (ED) or emergency admission to hospital for a further fall, or deaths within six months of the index fall. Important predictors that may affect triage decisions and outcomes, including the distance between incidents and EDs at index fall, patient age and gender, patterns of presentation (eg, whether out-of-hours or not) will be considered as covariates, and included in statistical modelling. We shall also take account of potential confounders such as the sequence of base stations (and their trial allocation) from which paramedics attend patients who fall more than once. The principles used in analysing primary outcomes also apply to quantitative secondary outcomes.

Required analyses include: logistic regression (for binary outcomes); cross-tabulations and risk ratios (for categorical outcomes); and survival analysis, including Cox's proportional hazards models^[10], (for measurement outcomes, such as times to events). Primary outcomes will also be jointly analysed using methods developed for recurrent event analysis^[11-14]. Repeated observations will be analysed in two ways: as 'repeated measures'; and summarised by "area under curve" of HRQoL measures per patient.

Where appropriate, multilevel modelling will be used to estimate clustering effects for stations or centres. The exact number of levels in models will be determined using statistically significant changes in likelihood ratio tests according to the principle of parsimonious parameterisation. Potential covariates to be included in models will be tested; those with an F value of less than 1 (that is, they increase the standard error of the estimate) will be excluded and the analysis recalculated. Binary covariates where almost all cases (>90%) are in one category will also be excluded.

Residual diagnostics will be used where analyses assume Normality; if the distributions of residuals are markedly non-normal, data transformation or bootstrapping will be considered. Residual analysis will be used to identify outliers; identified outliers will be excluded and the analysis updated. Wherever possible, outcome descriptions, summaries and comparisons will be expressed in accordance with appropriate CONSORT (www.consort-statement.org) guidelines, including estimates with 95% confidence intervals (allowing two-tailed tests at the 5% significance level).

5.2 Health Economic Analysis

This will estimate costs of providing the new intervention, and consequences for the NHS (for instance, ED attendances, inpatient admissions) and personal social services. Data on the use of the health service resources will be collected for each patient using routine ambulance service and hospital records. Costs will then be calculated using unit-costs estimated through a micro-costing study within the trial. Direct costs to the NHS will be assessed using data logged as a part of routine practice and from resource utilisation recording sheets, together with reference to patient records and discussions with relevant finance staff.

SF6D scores (derived from SF12 scores) will also be used to estimate the quality adjusted life years (QALYs) gained from the intervention, and an incremental cost-per-QALY calculated. These ratios will be presented along with their associated Cost-effectiveness Acceptability Curves (CEAC). Sensitivity analyses will be undertaken to assess the robustness of results to changes in the configuration of the intervention and other health service costs. We aim to publish economic results alongside clinical results where possible (for instance, in the BMJ format of pairs of companion papers).

The Senior Trial Economist is responsible for the health economic analysis, which will generally adopt approaches employed in the main analysis of clinical data. Specifically: .

- Analysis will be by 'treatment allocated' when generating baseline findings, with inclusive analysis by treatment received or exclusive analysis of treatment per-protocol being used for sensitivity analysis when also undertaken in reporting clinical findings
- Addressing uncertainty by applying bootstrapping for CEACs and confidence intervals^[15-17].
- Using a range of time-horizons to estimate the effectiveness and cost-effectiveness of the intervention.
- Applying appropriate discount rates to costs and benefits, as required by NICE; applying appropriate threshold Incremental Cost-effectiveness Ratios (ICER), such as the £20,000 and £30,000 thresholds used by NICE, in cost-per-QALY calculations.

5.3 Qualitative Analysis

All interviews and focus groups, with the permission of participants, will be recorded and transcribed. As this is a team-based project involving collection of data in multiple sites, maintaining consistency in data collection and transcription is crucial. The qualitative analysis co-ordinator will set up a data management system consisting of instructions on converting raw data to computer files (in the form of a transcription protocol), organising data storage, data archiving steps and a data management checklist. The transcription protocol will ensure that standard conventions are adopted throughout the transcription process, and that a standard presentation format is used. Each transcript will be quality assured by the relevant researcher.

Following transcription, participants in the paramedic and service provider focus groups will be given the opportunity to review the anonymised transcripts and state whether they agree with the contents or wish to amend their quotes.

The data will be analysed thematically using the framework analysis approach for applied policy research^[18]. This is a systematic, dynamic and transparent method of analysis, which generates themes from the original accounts of participants^[19]. In the analysis the data is sifted, charted and sorted in accordance with key issues and themes using five steps: familiarisation; identifying a thematic framework; indexing, charting; and mapping and interpretation.

All data analysis will be carried out independently by two researchers, who will then meet to discuss and agree final coding and interpretation. Analysis will be conducted using NVivo. When reporting the results, we will ensure that the anonymity of responders is preserved and that no quote can be attributed to a particular individual.

Section 6: INTERIM & FURTHER ANALYSES

Patients will be recruited over a 15 month period and followed up for six months. The trial will end and be analysed after the last six month follow-up contact with any patient in the trial. No formal interim analyses of the trial outcomes, and no subgroup analyses, are planned within the trial.

6.1 Non-Consenter Analysis

Study patients will be compared with eligible patients who have not consented to the study in terms of age, gender, date and time of index call, on-scene time, main condition code and disposition by appropriate significance tests.

6.2 Non-Respondent Analysis

Study patients who have returned the study questionnaire will be compared with those who have not return the questionnaire by the same parameters of the non-consenter analysis. In addition, those responding to the questionnaire will also be compared to those not returning the questionnaire in terms of the primary outcomes.

6.3 Sensitivity Analyses

Subject to resources, we will carry out further appropriate sensitivity analyses, some identified as the main analyses proceed. These will include a comparison between identifiable data retrieved from hospitals and those retrieved through the anonymous data sources (where experience with SAFER1^[4] indicates that a good or perfect match between anonymised records and study participants should be possible in 90% of cases); consideration of patterns of referral to falls services; and analyses based on further refined definitions of the intervention group – for instance, excluding participants ineligible for falls service referral.

Section 7: REPORTING & RELATED ISSUES

The SAFER 2 DMEC will receive immediate notification of all SUSARs, and summaries of current trial data. This will allow the DMEC to monitor trial progress, and request unblinded comparisons where they have cause for concern. The DMEC may then make a final decision to modify or terminate the trial, as specified in its terms of reference.

REFERENCES

- [1] Snooks, H, Anthony, R, Chatters, R, Cheung, WY, Dale, J, Donohoe, R, Gaze, S, Halter, M, Konitou, M, Logan, P, Lyons, R, Mason, M, Nicholl, J, Phillips, C, Phillips, J, Russell, I, Siriwardena, AN, Wani, M, Watkins, A, Whitfield, R, Wilson, L. Support And Assessment For Fall Emergency Referrals (SAFER 2) Research Protocol, *BMJ Open*, 2012, 2:e002169.
- [2] WWORTH SOP24 Randomisation, WWORTH, Swansea University, 2012.
- [3] Ruta DA, Garatt AM, Leng M, Russell IT, Macdonald L. A New Approach to the Measurement of Quality of Life: The Patient Generated Index. *Medical Care*, 1994; 32, 1109-1126.

- [4] SAFER1 (Support And Assessment For Fall Emergency Referrals) Final Report, Swansea University, 2011.
- [5] WWORTH SOP28 Statistics, WWORTH, Swansea University, 2012.
- [6] Hill KD, Schwarz JA, Kalogeropoulos AJ, Gibson, SJ. Fear of falling revisited. *Archives of Physical Medicine and Rehabilitation*, 1996; 77, 1025-1029.
- [7] Brazier J, Ratcliffe J, Salomon JA, Tsuchiya A. *Measuring and valuing health benefits for economic evaluation*, 2007, Oxford: Oxford University Press.
- [8] Young TA. Estimating mean total costs in the presence of censoring: a comparative assessment of methods, *Pharmacoeconomics*, 2005; 23, 1229-1242.
- [9] Lamb SE, Jorstad-Stein EC, Hauer K, Becker C. Development of a Common Outcome Data Set for Fall Injury Prevention Trials: The Prevention of Falls Network Europe Consensus. *Journal of the American Geriatric Society*, 2005; 53:1618-1622
- [10] Cox, DR. Regression Models and Life Tables, *Journal of the Royal Statistical Society: Series B*, 1972; 34, 86-94.
- [11] Andersen, PK, Gill, RD. Cox's Regression Model for Counting Processes: A Large Sample Study, *Annals of Statistics*, 1982; 10, 1100-1120.
- [12] Box-Steffensmeier, JM, de Boef, S. Repeated Events Survival Models: The Conditional Frailty Model, *Statistics in Medicine*, 2006; 25, 3518-3533.
- [13] Cook, RJ, Lawless, JF. *The Statistical Analysis Of Recurrent Events*, 2007. New York, CRC.
- [14] Cook RJ, Andersen, PK. Recurrent Event Analyses [Editorial introduction to a Special Issue on Recurrent Event Analysis], *Lifetime Data Analysis*, 2010; 16, 449-450.
- [15] Glick HA, Doshi JA, Sonnad SS, Polsky D. *Economic Evaluation in Clinical Trials*, 2007. Oxford: Oxford University Press.
- [16] Groot Koerkamp B, Hunkink M, Stijnen T, Mannitt JK, Kuntz KM, Weinstein C. Limitations of acceptability curves for presenting uncertainty in cost-effectiveness analysis. *Medical Decision Making*, 2007; 27, 101-111.
- [17] National Institute for Health and Clinical Excellence (NICE), Review of the guide to the methods of technology appraisal (consultation document), 2007.
- [18] Ritchie J, Spencer L. Qualitative data analysis for applied policy research. Chapter 12 in Huberman AM, Miles MB (Eds): *The Qualitative Researcher's Companion*, 2002. Sage Publications.
- [19] Srivastava A, Thomson SB. Framework Analysis: A Qualitative Methodology for Applied Policy Research. *Journal of Administration & Governance*, 2009; 4, 72-79.

Appendix 12 Patient semistructured interviews schedule

Patient experience (of ambulance service)

1. Can you tell me what happened when you fell and an ambulance was called?

Prompt:

Where did you fall?

How did you fall?

Were you hurt?

2. What did the ambulance crew do when they arrived?

3. Were you informed of what would happen next (e.g. referral to a falls service or taken to hospital)?

Prompt:

Did you understand what this would mean?

4. What did you feel about what they did?

Prompt:

At the time of fall and on reflection.

5. Have you fallen before and were you seen by an ambulance crew? (If answer is yes go to question 5a)

5a. In what ways was the assessment and treatment you received different from previously?

6. How did you feel about this decision?

Prompt:

Was it right for you and were you part of the decision?

Patient experience (of those seen by a falls service)

1. How long was it before someone from the falls service contacted you?

2. How did you feel about the amount of time it took before you were contacted?

3. What did the Falls Service offer you?

Prompt:

How did you feel about what they did?

Prompts:

Exercise Programme.

Prompt any of the following:

- Home exercise programme to improve strength and balance.
- One to one strength, balance and mobility training.
- Referral to a balance group or other group.
- Floor transfer practice.

Falls Prevention programme

Prompt:

Advice and education regarding how to reduce risk of falls, including pacing activity, sitting to put on trousers and shoes and appropriate footwear, etc.

Environmental adaptations or assistive devices

Prompt any of the following:

- Walking aid provision such as walking frame, walking stick, wheelchair, etc. or any other devices such as pendant alarm, grab rails or anything else.
- Advice on environmental adaptations and onward referrals to services to address issues such as loose rugs, wires, etc.

Occupational therapist

Prompt:

Occupational Therapy assessment of home environment, bed and toilet transfers and wheelchair suitability/ or of ability to perform personal care (including dressing) and domestic tasks.

Physiotherapist

Prompt:

Physiotherapy assessment of strength, range of movement, balance and transfers.

Disability counsellor

Prompt:

To talk about feelings and problems in a supportive way.

Social services assessment

Prompt:

For a review of the adequacy of the care package or anything else.

Anything else

4. Could the service be improved in any way?

Prompt:

Were you happy with what they offered you and was there anything you were unhappy with?

Patient health (post fall)

1. Have you had a fall since?
2. In terms of your health, how is it different since your fall?
3. Since the fall have any areas of your life been affected?

Prompt:

Have any areas of your life improved?

Prompt:

Have any areas of your life got worse?

Patient satisfaction

1. This project is looking at helping ambulance paramedics decide the best treatment for you if you fall.
At the time of your fall did you know they were following a new process for assessing elderly fallers and what do you think about this?
2. What did you think of the overall process?
3. How would you feel about being treated by the ambulance crew in this way in the future?
4. Is there anything else you would like to tell us relating to your fall and the attention you received?

Appendix 13 Support and Assessment for Fall Emergency Referrals 2 paramedic pretrial focus group interview schedule

Support and Assessment for Fall Emergency Referrals 2 paramedic pretrial focus group interview schedule

Thank you for taking part today. This session is to find out a bit more about your current practice and to get your thoughts and views on the new referral pathways proposed for the SAFER 2 trial.

Check that we have got the consent and demographic info forms completed.

Questions

What motivated you to get involved in the study?

Have there been any other 'pathways of care' or new ways of working introduced while you've been a paramedic?

Prompts:

- How have new practices been introduced? What's worked, what hasn't?
- Issues around changing practice – are there any pressures?
- What are the attitudes towards new developments in the health service.
- Differences between attitudes of paramedics and their managers.
- Barriers and motivations to new developments?
- Issues around using new developments.
- Support for using new pathways/ways of working.
- Operational impacts.

Do new pathways of care sometimes bring with them additional decision-making for paramedics?

Prompts:

- Adapting to this.
- Resistance?
- Time to adopt new processes?
- Levels/amount of training.

How do you deal with calls from older fallers at the moment?

Prompts:

- Options available to the paramedic.
- Links to other services to support older fallers.
- Explore issues around deciding to leave at home or convey.
- Any frustrations with the current system.
- Risks with the current system – to patient.
- Risks with the current system – to paramedic (confidence to leave patients at home).

Do you think that having access to a falls pathway will make a difference?

Prompts:

- To the patient.
- To you.
- To the ambulance service.
- Suitability of this patient group for the new care pathway.

And similarly, what are your expectations for this new pathway?

We asked at the beginning about your motivations for getting involved. We wondered what you thought your colleagues would make of you taking part in the study.

Prompts:

- What could encourage more people to become involved?

Appendix 14 Support and Assessment for Fall Emergency Referrals 2 paramedic trial-end interview schedule

Thank you for taking part today. This session is to find out more about your involvement in the SAFER 2 study and experience of using the new protocol and to understand how you made your decisions on whether to convey patients to hospital, refer to a falls service or to leave the patient at home.

Your views, which will be kept confidential, will enable us to assess the benefit to patients of the 'intervention' and pathways to community-based falls services. We simply need to understand if it works and if it can improve patient outcomes in relation to injury, death and quality of life.

1. How did you find using the SAFER 2 trial flow chart in practice?

Prompts:

- Which parts were useful and which were not, i.e. the 'get up and go test'?
- Did you need to make any additional decisions because patients did not fit into the protocol?
- Did you keep a copy of the flow chart with you?
- Did it make you feel more or less confident in your decision-making?

2. What were your expectations of the new way of working and were they met?

Prompts:

- Increase your job satisfaction and/or confidence.
- Aid your decision-making and assessments.

3. The idea of the flow chart was to make it safe to leave a patient at home with a referral to a falls service. We found this didn't always happen and some patients were left at home without a referral – we'd like to find out what influenced your decision?

Prompts:

- Meeting response times and/or last job or on-scene time.
- Decision-making/confidence.
- Patient preference.
- Clinical judgement.
- Familiarity with process.
- Advising patients they did not need to go to hospital.
- Patient already had a care package in place/or similar.
- Something else.

4. How satisfied were you with the amount training you received?

Prompts:

- Would you have liked to have received more or less training?
- Was it informative enough – did you feel ready and confident to use the flow chart?
- Did it enhance your clinical knowledge?

5. Was there any support available to you while you were taking part in the study?

Prompts:

- Did anyone contact you by telephone or e-mail?
- Did you feel well supported by ambulance service management and trainers?

6. What would you think if the initiative was introduced across the service?

Prompts:

- Do you feel it would change your role at all or affect the way you work?
- What factors would make it difficult and what could help the process?

7. When comparing the SAFER 2 trial to other similar studies, involving older fallers, it seems a higher proportion of patients were conveyed to hospital – we'd like to find out if the SAFER 2 trial changed the way you normally work?

Prompts

- Did the intervention affect you decision-making?
- Difficult/easier to assess this group of patients?
- Did it make you convey patients more often or less often?

Appendix 15 Stakeholder trial-end interview schedules

Support and Assessment for Fall Emergency Referrals 2 ambulance personnel trial-end interview schedule

1. At the beginning of the study, what were your expectations of the SAFER 2 trial intervention and the new way of working?

Prompts:

- Increase job satisfaction and/or confidence.
- Aid decision-making and assessments.
- Better patient care.
- Increase workload/increased demand on service.
- Increased efficiency/turnaround time.
- How aware were they of the SAFER 2 trial and what it was meant to achieve?

2. In what ways did your involvement in the SAFER 2 trial impact on your role?

Prompts:

- Time.
- Advantages/Improvements.
- Disadvantages/difficulties.

3. In what way do you think the SAFER 2 intervention has changed the way that colleagues within the organisation (paramedics, dispatch personnel) conduct their work?

Prompts:

- Advantages.
- Disadvantages.
- More confident or less confident in decisions.
- Make decisions in a different way.
- Provides more protection.
- Undermines/supports autonomy and clinical judgement.

4. What factors do you think had an effect on the extent to which the SAFER 2 trial was adopted within your organisation?

Prompts:

Factors within the organisation:

- Time pressures.
- Dispatch coding.
- Beliefs.
- Work habits.

Factors outside the organisation:

- Other pathways or developments.
- Patient choice.
- Family of patients, etc.
- Contact between roles (university, ambulance, falls).
- Any other barriers.

5. We have found that some patients seen by an intervention paramedic were left at home without a referral. Why do you think this might have been?

Prompts:

- Patient already had a care package in place/or similar.
- Paramedics meeting response times and/or last job or on-scene time.
- Paramedic decision-making/confidence.
- Patient preference.
- Clinical judgement.
- Familiarity with process.
- Advising patients they did not need to go to hospital.
- Something else.

6. What would you think if the SAFER 2 trial was rolled out across a wider service area?

Prompts:

- Good thing or not?
 - Advantages to patients/paramedics/organisation.
 - What factors would make it difficult and what could help the process?
 - Differences between a small scale trial and a big service change.
 - Boundary and equity issues i.e. thought it wasn't fair that people living in some areas got the service and others didn't.
 - Risk of duplication of initiatives.
 - Changes in work practices such as following new protocols/flow charts are an inevitable process.
7. Do you have any recommendations for how the process could be improved for future use?
8. Describe the SAFER 2 trial intervention using only one word?
9. Is there anything else about the study that you would like to say?

Support and Assessment for Fall Emergency Referrals 2 wider stakeholder trial-end focus group schedule

1. At the beginning of the study, what were your expectations of the SAFER 2 trial intervention and the new way of working?

Prompts:

- Increase job satisfaction and/or confidence.
- Aid decision-making and assessments.
- Better patient care.
- Increased workload/increased demand on services.
- Increased efficiency/turnaround time.
- Were your expectations met?

2. In what ways did your involvement in the SAFER 2 trial impact on your role?

Prompts:

- Time?
- Improvements/difficulties?

3. What factors do you think had an effect on the use of the SAFER 2 trial intervention within your organisation?

Prompts:

- Internal communications/support.
- Time pressures.
- Problems referring patients onto further specialist assessment.
- Beliefs/attitudes.
- Expectations about role, clear or unclear.
- Processes to be followed, clear or unclear.

4. What factors do you think had an effect on the use of the SAFER 2 trial intervention outside your organisation?

Prompts:

- Other pathways or developments.
- Other organisations/support.
- Patient choice.
- Family of patients, etc.
- Contact between roles (university, ambulance, falls).

5. Have you had any feedback from service users about the SAFER 2 trial intervention?

Prompts:

- Positive feedback.
- Negative feedback.

6. What would you think if the new way of working was introduced across the service?

Prompts:

- Good thing or not?
- Advantages to patients/paramedics/organisation.
- What factors would make it difficult and what could help the process?
- Differences between a small scale trial and a big service change.
- Boundary and equity issues i.e. thought it wasn't fair that people living in some areas got the service and others didn't.
- Risk of duplication of initiatives.
- Changes in work practices such as following new protocols/flow charts are an inevitable process.

7. Do you have any recommendations for how the process could be improved for future use?

8. Describe the SAFER 2 trial using only one word?

9. Is there anything else about the study that you would like to say?

Appendix 16 Support and Assessment for Fall Emergency Referrals 2 framework for analysis of interviews and focus groups

Support and Assessment for Fall Emergency Referrals 2 framework or analysis of interviews and focus groups with staff: including paramedics and staff of other relevant service areas

This analysis is of 'before' and 'after' data together.

1. Core coding.
 - i. Data source (code entire transcript to either subcode below).
 - Interview.
 - Focus group.
 - ii. Site (code entire transcript to one of the subcodes below).
 - London Ambulance Service.
 - East Midlands Ambulance Service .
 - Welsh Ambulance Services NHS Trust.
 - iii. Role of speaker (each extract coded should also be crosscoded to one of the subcodes below).
 - Paramedic.
 - Other ambulance service staff.
 - Falls service.
 - Other.
2. Expectations of the SAFER 2 trial intervention (before).
 - i. Rate of referrals.
 - ii. Monitoring/performance management function.
 - iii. Better patient care.
 - iv. Anticipated impact on process (paperwork/time at scene).
 - v. Anticipated impact on other services (e.g. falls service).
 - vi. Unclear/uncertain expectations.
 - vii. Other.
3. Implementation process for the SAFER 2 trial intervention (after).
 - i. Impact on workload and role – ambulance service.
 - ii. Training and support.
 - iii. Teamwork/networking/communication.
 - iv. Other changes taking place at the same time.
 - v. Wider impact within ambulance service (e.g. raising awareness of falls as an issue).
 - vi. Impact on workload and role – falls service/other (including comments on appropriateness of referrals).

- vii. Overall comments on the SAFER 2 trial intervention.
 - Negative.
 - Positive.
4. Referral practice post-implementation.
- i. Reported reasons for not referring.
 - Negative feedback.
 - Taking easier option/embedded habits
 - Difficulties with patient assessment.
 - Time restrictions/end of shift.
 - ii. Confidence in/anxiety about pathways and outcomes.
 - iii. Suitability of patients for referral to falls service.
 - iv. Referral to other services (including GPs).
 - v. Role of patient and family in decision-making.
 - vi. Advice from/discussion with colleagues.
 - vii. Other.
5. Tension between research and real world practice.
- i. Motivation for involvement in research study or new innovation.
 - Persuasion by an individual.
 - Desire to improve service quality.
 - Personal development and skills.
 - ii. Factors discouraging involvement in research study.
 - iii. Perspective on study as a contained piece of work (bounded area, end date, etc. plus comments about any problems with timetable).
 - iv. Drive to recruit patients to target numbers.
 - v. Confusion/ambiguity around status and nature of research study (including references to other research studies in progress).
 - vi. Future rollout.
 - vii. Perceived benefits of study (e.g. evidence for commissioners).
 - viii. Other.
6. Suggestions.
- i. Follow-up/feedback on individual patients.
 - ii. Adapting model (e.g. case finding of patients at lower risk levels, scaling up falls service, more links to other services).
 - iii. Adapting the SAFER 2 trial process (e.g. putting a prompt on ePRF, devising flow chart, better communication).
 - iv. Adapting implementation of trial (e.g. better communication about progress).
 - v. Other.
7. Existing practice re older fallers.
- i. Decision-making.
 - ii. Outcomes.

- iii. Perceived problems with existing practice.
 - iv. Other.
8. Communication.
 9. Previous experience of pathways and protocols.
 10. Concerns, issues around pay, status, role, hours, breaks, etc.

Support and Assessment for Fall Emergency Referrals 2: framework for analysis of interviews with patients

1. Case coding for patient (note that the entire transcript will be coded to each of the nodes which start with 1).
 - i. Outcome of contact.
 - Patient attended hospital as a result of index fall.
 - Patient did not attend hospital as a result of index fall, referred to falls service.
 - Patient did not attend hospital as a result of index fall, **not** referred to falls service.
 - ii. Gender.
 - Female.
 - Male.
 - iii. Site.
 - London Ambulance Service.
 - East Midlands Ambulance Service.
 - Welsh Ambulance Services NHS Trust.
2. Other overall comments (these should be coded to specific points in the transcript).
 - i. Any evidence of/report of memory problems or confusion.
 - ii. Nature/cause of fall.
 - iii. Repeat falls.
 - Previous fall(s) reported.
 - Subsequent fall(s) reported.
 - iv. Other medical conditions.
 - v. Description and nature of injury.
3. Patient perspective on ambulance service based on their contact at time of index fall.
 - i. Perspective on paramedics/ambulance technicians.
 - Overall evaluation.
 - Personal qualities – kindness/calm, etc.
 - Impact on patient – maintaining dignity/minimising embarrassment/reassuring/asking rather than telling/explaining what's going on, etc.

- ii. Perspective on other aspects of the ambulance service.
 - Service overall – including waiting time.
 - Despatch.
4. Process – when ambulance service attended patient for index fall.
 - i. Physical checks/examination.
 - ii. Questions.
 - iii. Lifting.
 - iv. Social aspects – ‘having a laugh’/cup of tea/making lunch, etc.
 - v. Presence/role of family members/carers/others.
 - vi. Patient’s reflection/comparison with any previous experience.
 - vii. Paperwork/notes/records/communication.
5. Decision-making about transfer to hospital – when ambulance service attended patient for index fall.
 - i. Negotiation/paramedics offering choice.
 - ii. Patients’ expressed will/autonomy.
 - iii. Influences on patient’s decision-making.
6. Hospital attendance.
 - i. Experience of/perspective on attending hospital for index fall.
 - ii. Experience of/perspective on past hospital attendance.
7. Experience of falls service.
 - i. Patient evaluation.
 - ii. Confusion/ambiguity about role and contact.
8. Experience of other relevant health or support services.
 - i. Social care services (including voluntary sector).
 - ii. Other health services.
 - iii. Other or unclear.
 - iv. Private health care/own resources.
9. Consequences of fall.
 - i. Overall impact on quality of life.
 - ii. Impact on confidence/fear of falling.
 - iii. Pain/other health consequences.

Appendix 17 Support and Assessment for Fall Emergency Referrals 2 Involving People case study

Involving older people in a multi-centre randomised controlled trial of a complex intervention in pre-hospital emergency care

This case study has been developed by **Alun Toghill**, Involving People Network Member, and **Marina Koniotou**, Swansea University. It describes Alun's experience of public involvement in the SAFER 2 study.

The SAFER 2 trial will measure the costs and benefits of a protocol for use by emergency ambulance paramedics in the care of older people who have fallen, allowing the paramedic to assess and refer appropriate patients to a community based falls service.

The randomised controlled trial involves ambulance stations in three participating services (London, Wales, East Midlands) which are randomly allocated to 1) implement the new protocol (intervention group) or 2) continue to provide care according to their standard practice (control group). www.saferproject.org

My SAFER 2 journey began when I answered a request from Involving People to be involved in a research project.

Starting out

I was immediately drawn towards the project because it was about a falls prevention programme and it ignited my passion to help others.

It wasn't long before the ball was firmly rolling, when the trial researcher contacted me to explain what my role would involve. I vividly recall being warmly welcomed by the group as I explained to them my background and what skills I possessed; allowing them to decide where I would best be placed to help.

My background is diverse; I've worked in hospitals and in the elderly care private sector as a nurse, and I've also had wide experience in the community and voluntary sector, including setting up projects for the care and support of the elderly and at risk.

These experiences led me to be involved in a number of tasks, which would include working with the study team to assist in designing the project methodology i.e. developing the patient questionnaire.

Attending Meetings

I attended Task and Finish Group and Task Management Group meetings in Swansea and Bristol, by teleconference and in person. When I was unable to attend meetings, the trial researcher would arrange meetings at my home, as this was better for me. The flexibility of this approach was an example to all and is a model to be replicated.

As the study progressed I was able to participate fully in meetings, to monitor and review progress not only in Wales, but also at both London and East Midlands participating sites. Attendance at joint meetings and through teleconferencing meant that I was fully integrated within the research network and was included in writing days to help input into the final report.

Making a difference

Together with other members of the research team I took part in evaluating patient transcripts to ensure that any identifiable information was omitted appropriately. My work helped to ensure there was no breach of confidentiality, for example, I identified one lady from one of the trial sites because the transcript noted

her first name and the road of the accident.

I also helped with the design and formatting of the patient questionnaire. From a lay person's point of view, I could ensure ease of completion by looking at the questionnaire's clarity, use of English and the order of questions.

I advised on how to improve the low completion rate of the patient questionnaire and consent form by suggesting changing the colour of the forms from white to cream. This was based on previous projects I worked on where it was identified that changing the colour of the paper helps it not get lost in an avalanche of other papers. It is more likely to be completed and returned if it stands out from the crowd!

Role-play

Along with the Welsh-based researcher, I undertook a role-play activity as part of a process to refine the patient interview schedule. I played the patient, whilst they posed the interview questions.

This novel way of working allowed me to refine how the interview flowed and offer advice on where the schedule failed to articulate itself in a language and order the patient would understand and be comfortable with.

I was able to identify any duplication in the questions and advise how to pitch questions to this vulnerable group of patients. It is important that the patients in the target group have enough time to answer; young people are quicker at responding.

I emphasised the importance of building a relationship of mutual trust with the patient interviewee. When you ask for their name and, for arguments sake, they say Elizabeth Rose Taylor, you should ideally respond by asking them what they would like to be called. If, for example, this person is known as Lizzie, addressing them in a familiar way rather than a formal way helps put them at ease. If the interviewer cares about the interviewee it can show that they are serious about hearing their views and a more open response is likely.

Reflections

My involvement with the project has been an enjoyable and valuable learning experience. I have found that my varied life experiences have been used by the project to enhance the final outcome. I do feel proud that my contribution has had a marked effect on this activity and that shared participation has been of value to all parties.

I was always listened to and, as a result, I gave everything I could and I got a lot out of it. I felt the other researchers had a similar attitude; they gave what they could and were open, honest and friendly. There were no barriers and this encouraged trust and mutual respect between everyone.

I brought a different insight to the study and my drive and passion grew when I could see an end product that was going to make a difference. I know I have made a difference as my suggestions have more than been acknowledged; many have actually been adopted.


I've talked to other studies about the SAFER 2 model; too many service users feel let down by the process as they have not been given a defined role and they have not been appropriately involved enough.

A final message


This has been a positive experience of feeling as though I am working together with the team. I hope my positive experience will have a ripple effect, as I will continue to promote this excellent model of service user involvement.

Appendix 18 Site Consolidated Standards of Reporting Trials flow charts


Site 1



Site 2



Site 3



Appendix 19 Further clinical effectiveness results

Participant numbers and subgroups flow

As outlined in *Chapter 4*, a total of 5914 eligible patients were attended by 215 paramedics based at 31 ambulance stations across the three study sites. After dissenting patients were excluded, 80% ($n = 4704$) were available for follow-up: 2420 in the intervention group and 2284 in the control group. Various analyses in *Chapter 4* exclude $n = 49$ participants not matched anonymously, as data on key components in the primary outcome were unavailable. A further subgroup of participants in site 1 were registered at a GP practice unable to refer patients to the falls referral service. In this appendix, we present the main results tables for the subgroup which also excludes these participants ineligible for a falls referral.

TABLE 44 Primary outcomes at 1 month, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral

Primary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			ICC	
		Intervention	Control	Estimate, p-value	95% CI	Estimate	95% CI	
Proportion of patients dying (any cause) ^c	All	135/2161 (6.2%)	119/1982 (6.1%)	OR = 1.011; p = 0.931	0.783 to 1.307	0	n/a	
	Site 1	74/1081 (6.8%)	62/1055 (5.9%)					
	Site 2	31/540 (5.7%)	26/433 (6.0%)					
	Site 3	30/540 (5.6%)	31/494 (6.3%)					
Proportion of patients with further emergency admission to hospital, or death ^d	All	561/2161 (26.0%)	507/1982 (25.6%)	OR = 0.941; p = 0.430	0.808 to 1.095	0	n/a	
	Site 1	278/1081 (25.7%)	265/1055 (25.1%)					
	Site 2	121/540 (22.4%)	108/433 (24.9%)					
	Site 3	162/540 (30.0%)	134/494 (27.1%)					
Proportion of patients with further emergency event (ED attendance or emergency admission to hospital) or death ^e	All	673/2161 (31.1%)	610/1982 (30.8%)	OR = 0.956; p = 0.534	0.828 to 1.103	< 0.0001	0 to 0.0037	
	Site 1	326/1081 (30.2%)	320/1055 (30.3%)					
	Site 2	157/540 (29.1%)	135/433 (31.2%)					
	Site 3	190/540 (35.2%)	155/494 (31.4%)					
Proportion of patients with further emergency event (emergency service call, ED attendance, emergency admission to hospital) or death ^f	All	793/2161 (36.7%)	743/1982 (37.5%)	OR = 0.895; p = 0.118	0.780 to 1.028	0.0021	0.0002 to 0.0071	
	Site 1	404/1081 (37.4%)	409/1055 (38.8%)					
	Site 2	172/540 (31.9%)	152/433 (35.1%)					
	Site 3	217/540 (40.2%)	182/494 (36.8%)					

Primary outcomes	Raw data		Adjusted comparison ^{a,b}		ICC		
	Site	Intervention	Control	Estimate, p-value	95% CI	Estimate	95% CI
Further emergency events per patient (emergency service call, ED attendance, admission to hospital) per day at risk, ^g mean (SD) [n]	All	0.0780 (0.2254) [1979]	0.0762 (0.2215) [1836]	$\Delta = 0.0032$; $p = 0.650$	-0.0107 to 0.0172	0.0023	0.0003 to 0.0070
	Site 1	0.0673 (0.1841) [1012]	0.0688 (0.1959) [1001]				
	Site 2	0.0755 (0.2066) [466]	0.0675 (0.2021) [384]		$\Delta_1 = -0.0055$; $p = 0.943$	0.0016	0.0001 to 0.0061
	Site 3	0.1018 (0.3033) [501]	0.1003 (0.2817) [451]				
Mean time to first event (days, truncated at 30), ^h mean (proportion truncated)	All	22.48 (1368/2161)	22.40 (1239/1982)	$HR = 0.970$; $p = 0.557$	0.878 to 1.073	0.0026	0.0004 to 0.0076
	Site 1	22.57 (677/1081)	22.22 (646/1055)				
	Site 2	23.22 (368/540)	22.74 (281/433)				
	Site 3	21.54 (323/540)	22.49 (312/494)				

HR, hazard ratio; n/a, not applicable.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a Hazard Ratio from Cox regression models for survival analysis; an additive group effect [Δ_1 , in the same units as the dependent variable; Δ_1 refers to log-transformed data, using $\ln(x + 0.001)$ in place of x] from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Gender ($p < 0.001$); age ($p < 0.001$); seasonality ($p = 0.038$).

d Gender ($p = 0.002$); interaction between site 3 and group ($p = 0.011$); out of hours ($p = 0.036$).

e Gender ($p = 0.010$); seasonality ($p = 0.030$); interaction between site 3 and group ($p = 0.039$).

f Site 1 ($p = 0.034$); interaction between site 3 and group ($p = 0.010$); gender ($p = 0.004$); seasonality ($p = 0.013$); out of hours ($p = 0.011$).

g For Δ_1 : site 3 ($p < 0.001$); age ($p = 0.028$). For Δ_2 : site 3 ($p = 0.011$); out of hours ($p = 0.002$).

h Gender ($p < 0.001$); seasonality ($p = 0.006$); out of hours ($p = 0.019$).

TABLE 45 Components of the primary outcome at 1 month, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral

Primary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			ICC	
		Intervention	Control	Estimate, p-value	95% CI	Estimate	95% CI	
Proportion of patients dying (any cause)	All	135/2161 (6.2%)	119/1982 (6.0%)	OR = 1.011; p = 0.931	0.783 to 1.307	0	n/a	
	Site 1	74/1081 (6.8%)	62/1055 (5.9%)					
	Site 2	31/540 (5.7%)	26/433 (6.0%)					
Proportion of patients with further emergency admission to hospital ^c	Site 3	30/540 (5.6%)	31/494 (6.3%)					
	All	472/2161 (21.8%)	425/1982 (21.4%)	OR = 1.024; p = 0.754	0.883 to 1.188	0.0012	0 to 0.0052	
	Site 1	226/1081 (20.9%)	222/1055 (21.0%)					
Proportion of patients with further ED attendance ^d	Site 2	104/540 (19.3%)	88/433 (20.3%)					
	Site 3	142/540 (26.3%)	115/494 (23.3%)					
	All	419/2161 (19.4%)	367/1982 (18.5%)	OR = 1.067; p = 0.420	0.912 to 1.248	0.0075	0.0024 to 0.0160	
Further ED attendances per patient, ^e mean (SD) [n]	Site 1	212/1081 (19.6%)	193/1055 (18.3%)					
	Site 2	79/540 (14.6%)	63/433 (14.5%)					
	Site 3	128/540 (23.7%)	111/494 (22.5%)					
Further ED attendances per patient, ^e mean (SD) [n]	All	0.2683 (0.6322) [1979]	0.2609 (0.8087) [1836]	Δ = 0.839; p = 0.057	0.700 to 1.006	0.0052	0.0014 to 0.0120	
	Site 1	0.2747 (0.6945) [1012]	0.2268 (0.4995) [1001]					
	Site 2	0.2017 (0.5017) [466]	0.1901 (0.4824) [384]					
Further ED attendances per patient per day at risk, ^f mean (SD) [n]	Site 3	0.3174 (0.6042) [501]	0.3969 (1.3743) [451]					
	All	0.0242 (0.1060) [1979]	0.0219 (0.0816) [1836]	Δ = 0.0022; p = 0.483	-0.0039 to 0.0082	0.0054	0.0013 to 0.0131	
	Site 1	0.0225 (0.0821) [1012]	0.0188 (0.0694) [1001]					
Further ED attendances per patient per day at risk, ^f mean (SD) [n]	Site 2	0.0172 (0.0611) [466]	0.0138 (0.0661) [384]	Δ = 0.0486; p = 0.390	-0.0623 to 0.1596	0.0099	0.0034 to 0.0204	
	Site 3	0.0341 (0.1648) [501]	0.0357 (0.1115) [451]					

TABLE 45 Components of the primary outcome at 1 month, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral (*continued*)

Primary outcomes	Raw data		Adjusted comparison ^{a,b}		ICC		
	Site	Intervention	Control	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Proportion of patients with further emergency service call ^g	All	401/2161 (18.6%)	430/1982 (21.7%)	OR = 0.817; <i>p</i> = 0.010	0.700 to 0.953	0.0066	0.0021 to 0.0139
	Site 1	216/1081 (20.0%)	257/1055 (24.4%)				
	Site 2	78/540 (14.4%)	68/433 (15.7%)				
	Site 3	107/540 (19.8%)	105/494 (21.3%)				
Further emergency service calls per patient, ^h mean (SD) [<i>n</i>]	All	0.3032 (0.7931) [1979]	0.3328 (0.7314) [1836]	Λ = 0.907; <i>p</i> = 0.143	0.795 to 1.034	0.0041	0.0011 to 0.0097
	Site 1	0.3261 (0.8500) [1012]	0.3856 (0.8168) [1001]				
	Site 2	0.2318 (0.6508) [466]	0.2292 (0.5686) [384]				
	Site 3	0.3234 (0.7920) [501]	0.3038 (0.6384) [451]				
Further emergency service calls per patient per day at risk, ⁱ mean (SD) [<i>n</i>]	All	0.0206 (0.0639) [1979]	0.0245 (0.0823) [1836]	Δ = -0.0038; <i>p</i> = 0.100	-0.0084 to 0.0007	0.0045	0.0011 to 0.0106
	Site 1	0.0198 (0.0580) [1012]	0.0247 (0.0723) [1001]				
	Site 2	0.0191 (0.0642) [466]	0.0174 (0.0733) [384]	Δ_L = -0.1261; <i>p</i> = 0.029	-0.2391 to -0.0131	0.0051	0.0014 to 0.0116
	Site 3	0.0238 (0.0743) [501]	0.0301 (0.1066) [451]				

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a multiplicative event rate ratio (Λ , scalar) from negative binomial regression models for count data; an additive group effect (Δ , in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(x+0.001)$ in place of x) from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Site 3 (*p* = 0.002); out of hours (*p* = 0.025).

d Site 1 (*p* = 0.003); site 3 (*p* < 0.001); age (*p* = 0.021); age2 (*p* = 0.009); seasonality (*p* = 0.003).

e Site 3 (*p* < 0.001); interaction between site 1 and group (*p* < 0.001); age (*p* < 0.001); age2 (*p* = 0.002); seasonality (*p* = 0.001); days at risk (*p* < 0.001).

f For Δ : site 3 (*p* < 0.001). For Δ_L : site 1 (*p* = 0.018); site 3 (*p* < 0.001); age2 (*p* = 0.025); seasonality (*p* = 0.003).

g Site 1 (*p* < 0.001); site 3 (*p* = 0.002); age (*p* < 0.001); seasonality (*p* = 0.010); out of hours (*p* < 0.001); recruitment point (*p* = 0.001).

h Site 1 (*p* < 0.001); site 3 (*p* = 0.002); age (*p* = 0.003); recruitment point (*p* = 0.001); out of hours (*p* < 0.001).

i For Δ : age (*p* < 0.001); site 3 (*p* = 0.005); out of hours (*p* = 0.038). For Δ_L : site 1 (*p* = 0.001); site 3 (*p* = 0.008); age (*p* < 0.001); out of hours (*p* = 0.001); recruitment point (*p* = 0.001); seasonality (*p* = 0.016).

TABLE 46 Primary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral

Primary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			ICC
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	95% CI	
Proportion of patients dying (any cause) ^c	All	420/2161 (19.4%)	362/1982 (18.3%)	OR = 1.199; <i>p</i> = 0.083	0.977 to 1.472	0	n/a
	Site 1	211/1081 (19.5%)	213/1055 (20.2%)				
	Site 2	116/540 (21.5%)	75/433 (17.3%)				
	Site 3	93/540 (17.2%)	74/494 (15.0%)				
Proportion of patients with further emergency admission to hospital, or death ^d	All	1212/2161 (56.1%)	1107/1982 (55.9%)	OR = 1.120; <i>p</i> = 0.147	0.961 to 1.305	0.0010	0 to 0.0050
	Site 1	576/1081 (53.3%)	580/1055 (55.0%)				
	Site 2	302/540 (55.9%)	238/433 (55.0%)				
	Site 3	334/540 (61.9%)	289/494 (58.5%)				
Proportion of patients with further emergency event (ED attendance or emergency admission to hospital), or death ^e	All	1388/2161 (64.2%)	1250/1982 (63.1%)	OR = 0.957; <i>p</i> = 0.532	0.834 to 1.098	0.0002	0 to 0.0038
	Site 1	664/1081 (61.4%)	655/1055 (62.1%)				
	Site 2	349/540 (64.6%)	273/433 (63.0%)				
	Site 3	375/540 (69.4%)	322/494 (65.2%)				
Proportion of patients with further emergency event (emergency service call, ED attendance, emergency admission to hospital) or death ^f	All	1532/2161 (70.9%)	1395/1982 (70.4%)	OR = 0.999; <i>p</i> = 0.990	0.872 to 1.144	0	n/a
	Site 1	764/1081 (70.7%)	751/1055 (71.2%)				
	Site 2	372/540 (68.9%)	289/433 (66.7%)				
	Site 3	396/540 (73.3%)	355/494 (71.9%)				

TABLE 46 Primary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral (*continued*)

Primary outcomes	Raw data		Adjusted comparison ^{a,b}		ICC		
	Site	Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Further emergency events per patient (emergency service call, ED attendance, admission to hospital) per day at risk, ^g mean (SD) [<i>n</i>]	All	0.0477 (0.1552) [2150]	0.0454 (0.1445) [1975]	$\Delta = 0.0024$; <i>p</i> = 0.614	-0.0068 to 0.0115	0.0016	0.0001 to 0.0057
	Site 1	0.0419 (0.1373) [1080]	0.0398 (0.1025) [1054]				
	Site 2	0.0464 (0.1577) [535]	0.0438 (0.1806) [429]	$\Delta_L = -0.0265$; <i>p</i> = 0.647	-0.1399 to 0.0869	0.0023	0.0002 to 0.0074
	Site 3	0.0607 (0.1838) [535]	0.0589 (0.1808) [492]				
Mean time to first event (days, truncated at 180), ^h mean (proportion truncated)	All	84.36 (629/2161)	84.54 (587/1982)	HR = 0.995; <i>p</i> = 0.903	0.926 to 1.071	0.0028	0.0004 to 0.0080
	Site 1	84.28 (317/1081)	82.26 (304/1055)				
	Site 2	90.69 (168/540)	90.87 (144/433)				
	Site 3	78.17 (144/540)	83.88 (139/494)				

HR, hazard ratio; *n/a*, not applicable.

^a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

^b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a Hazard Ratio (HR) from Cox regression models for survival analysis; an additive group effect [Δ], in the same units as the dependent variable; Δ_L refers to log-transformed data, using $\ln(x + 0.001)$ from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

^c Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* = 0.004); site 3 (*p* = 0.003); interaction between site 1 and group (*p* = 0.046); distance to ED (*p* = 0.039).

^d Age (*p* < 0.001); gender (*p* < 0.001); distance to ED (*p* = 0.005); interaction between site 1 and group (*p* = 0.010); out of hours (*p* = 0.019); recruitment point (*p* = 0.036).

^e Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* = 0.002); interaction between site 1 and group (*p* = 0.005); seasonality (*p* = 0.027); recruitment point (*p* = 0.010).

^f Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* < 0.001); recruitment point (*p* = 0.004); seasonality (*p* = 0.025).

^g For Δ : seasonality (*p* = 0.010); recruitment point (*p* = 0.020); site 3 (*p* = 0.013) For Δ_L : site 3 (*p* = 0.030); gender (*p* = 0.001); age (*p* < 0.001); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); distance to ED (*p* = 0.022).

^h Age (*p* < 0.001); gender (*p* < 0.001); out of hours (*p* < 0.001); recruitment point (*p* = 0.017); seasonality (*p* = 0.014); site 1 (*p* = 0.033); site 3 (*p* = 0.008).

TABLE 47 Components of the primary outcome at 6 months, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral

Primary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			ICC	
		Intervention	Control	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI	
Proportion of patients dying (any cause)	All	420/2161 (19.4%)	362/1982 (18.3%)	OR = 1.199; <i>p</i> = 0.083	0.977 to 1.472	0	n/a	
	Site 1	211/1081 (19.5%)	213/1055 (20.2%)					
	Site 2	116/540 (21.5%)	75/433 (17.3%)					
Proportion of patients with further emergency admission to hospital ^c	Site 3	93/540 (17.2%)	74/494 (15.0%)					
	All	1039/2161 (48.1%)	967/1982 (48.8%)	OR = 1.092; <i>p</i> = 0.252	0.940 to 1.269	0.0048	0.0013 to 0.0111	
	Site 1	483/1081 (44.7%)	503/1055 (47.7%)					
Proportion of patients with further ED attendance ^d	Site 2	258/540 (47.8%)	206/433 (47.6%)					
	Site 3	298/540 (55.2%)	258/494 (52.2%)					
	All	961/2161 (44.5%)	899/1982 (45.4%)	OR = 0.961; <i>p</i> = 0.535	0.849 to 1.089	0.0220	0.0099 to 0.0391	
Further ED attendances per patient, ^e mean (SD) [n]	Site 1	474/1081 (43.8%)	462/1055 (43.8%)					
	Site 2	196/540 (36.3%)	167/433 (38.6%)					
	Site 3	291/540 (53.9%)	270/494 (54.7%)					
Further ED attendances per patient, ^e mean (SD) [n]	All	0.843 (1.416) [2150]	0.939 (2.886) [1975]	Λ = 0.809; <i>p</i> < 0.001	0.720 to 0.910	0.0133	0.0057 to 0.0246	
	Site 1	0.804 (1.404) [1080]	0.806 (1.375) [1054]					
	Site 2	0.561 (0.912) [535]	0.678 (1.083) [429]					
Further ED attendances per patient per day at risk, ^f mean (SD) [n]	Site 3	1.204 (1.743) [535]	1.453 (5.296) [492]					
	All	0.0178 (0.0951) [2150]	0.0142 (0.0673) [1975]	Δ = 0.0037; <i>p</i> = 0.157	-0.0014 to 0.0087	0.0039	0.0009 to 0.0098	
	Site 1	0.0168 (0.0940) [1080]	0.0106 (0.0330) [1054]					
Further ED attendances per patient per day at risk, ^f mean (SD) [n]	Site 2	0.0145 (0.1033) [535]	0.0153 (0.1117) [429]	Δ_L = -0.0354; <i>p</i> = 0.450	-0.1272 to 0.0564	0.0277	0.0134 to 0.0475	
	Site 3	0.0231 (0.0886) [535]	0.0209 (0.0701) [492]					

TABLE 47 Components of the primary outcome at 6 months, analysis by treatment allocated, adjusted for significant covariates for anonymised linked data for matched participants eligible for a falls referral (*continued*)

Primary outcomes	Raw data		Adjusted comparison ^{a,b}		ICC		
	Site	Intervention	Control	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Proportion of patients with further emergency service call ^g	All	940/2161 (43.5%)	910/1982 (45.9%)	OR = 0.898; <i>p</i> = 0.092	0.792 to 1.018	0.0025	0.0004 to 0.0074
	Site 1	484/1081 (44.8%)	499/1055 (47.3%)				
	Site 2	233/540 (43.1%)	169/433 (39.0%)				
	Site 3	223/540 (41.3%)	242/494 (49.0%)				
Further emergency service calls per patient, ^h mean (SD) [<i>n</i>]	All	1.129 (2.560) [2150]	1.250 (2.685) [1975]	Λ = 0.921; <i>p</i> = 0.055	0.846 to 1.002	0.0054	0.0017 to 0.0115
	Site 1	1.272 (3.086) [1080]	1.453 (3.299) [1054]				
	Site 2	0.849 (1.497) [535]	0.839 (1.456) [429]				
	Site 3	1.120 (2.183) [535]	1.173 (1.881) [492]				
Further emergency service calls per patient per day at risk, ⁱ mean (SD) [<i>n</i>]	All	0.0128 (0.0377) [2150]	0.0166 (0.0541) [1975]	Δ = -0.0036 <i>p</i> = 0.010	-0.0064 to -0.0009	0.0019	0.0005 to 0.0055
	Site 1	0.0122 (0.0328) [1080]	0.0168 (0.0494) [1054]				
	Site 2	0.0125 (0.0376) [535]	0.0139 (0.0620) [429]	Δ_1 = -0.120; <i>p</i> = 0.014	-0.2156 to -0.0246	0.0036	0.0008 to 0.0088
	Site 3	0.0144 (0.0460) [535]	0.0184 (0.0565) [492]				

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; a multiplicative Event Rate Ratio (Λ , scalar) from negative binomial regression models for count data; an additive group effect Δ , in the same units as the dependent variable; Δ_1 refers to log-transformed data, using $\ln(x+0.001)$ in place of x from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Interaction between site 1 and group (*p* = 0.007); gender (*p* = 0.010); age (*p* = 0.031); recruitment point (*p* = 0.026); distance to ED (*p* = 0.003).

d Site 1 (*p* = 0.044); site 3 (*p* < 0.001); out of hours (*p* = 0.012); distance to ED (*p* = 0.008); age (*p* = 0.001); seasonality (*p* = 0.032).

e Site 3 (*p* < 0.001); interaction between site 1 and group (*p* = 0.001); gender (*p* < 0.001); seasonality (*p* < 0.001); recruitment point (*p* = 0.013); distance to ED (*p* < 0.001); days at risk (*p* < 0.001).

f For Δ : seasonality (*p* = 0.008); recruitment point (*p* = 0.012). For Δ_1 : site 3 (*p* < 0.001); age (*p* < 0.001); distance to ED (*p* < 0.001); seasonality (*p* = 0.004); out of hours (*p* = 0.016).

g Age (*p* < 0.001); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); distance to ED (*p* = 0.009).

h Site 1 (*p* < 0.001); site 3 (*p* = 0.027); recruitment point (*p* < 0.001); out of hours (*p* < 0.001); distance to ED (*p* = 0.005).

i For Δ : out of hours (*p* < 0.001); seasonality (*p* = 0.003); age (*p* = 0.011). For Δ_1 : out of hours (*p* < 0.001); recruitment point (*p* < 0.001); age (*p* < 0.001); distance to ED (*p* = 0.001).

TABLE 48 Secondary outcomes at 1 month, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral

Secondary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			ICC	
		Intervention (A)	Control (B)	Estimate, p-value	95% CI	Estimate	95% CI	
Duration of subsequent inpatient episodes: nights in hospital, truncated at 30 days, ^c mean (SD) [n]	All	2.26 (6.14) [2161]	2.16 (6.13) [1982]	$\Delta = 0.432; p = 0.061$	-0.019 to 0.884	0.0003	0 to 0.0031	
	Site 1	1.95 (5.53) [1081]	2.05 (5.81) [1055]					
	Site 2	2.51 (6.72) [540]	2.42 (6.77) [433]					
	Site 3	2.64 (6.67) [540]	2.17 (6.20) [494]					
Proportion with further reported injuries (fractures) ^d	All	93/2161 (4.3%)	81/1982 (4.1%)	OR = 1.043; $p = 0.790$	0.763 to 1.426	0.0048	0.0012 to 0.0113	
	Site 1	33/1081 (3.1%)	32/1055 (3.0%)					
	Site 2	41/540 (7.6%)	27/433 (6.2%)					
	Site 3	19/540 (3.5%)	22/494 (4.5%)					
Quality of life								
SF-12 MCS, ^e mean (SD) [n]	All	39.65 (12.46) [415]	38.66 (12.04) [368]	$\Delta = 1.071; p = 0.220$	-0.642 to 2.784	0.0063	0.0002 to 0.0286	
	Site 1	40.48 (12.92) [182]	40.15 (13.04) [175]					
	Site 2	37.76 (11.59) [126]	35.63 (10.69) [104]					
	Site 3	40.44 (12.53) [107]	39.26 (10.88) [89]					
SF-12 PCS, ^f mean (SD) [n]	All	29.14 (9.99) [415]	29.41 (10.08) [368]	$\Delta = -0.424; p = 0.553$	-1.829 to 0.980	0	n/a	
	Site 1	29.06 (9.85) [182]	29.70 (11.36) [175]					
	Site 2	28.13 (9.66) [126]	28.39 (7.99) [104]					
	Site 3	30.47 (10.55) [107]	30.03 (9.58) [89]					
Derived SF-6D, ^g mean (SD) [n]	All	0.5555 (0.1271) [444]	0.5527 (0.1264) [406]	$\Delta = -0.0001; p = 0.994$	-0.0170 to 0.0168	0.0099	0.0011 to 0.0306	
	Site 1	0.5597 (0.1260) [196]	0.5658 (0.1338) [195]					
	Site 2	0.5353 (0.1103) [134]	0.5226 (0.1089) [114]					
	Site 3	0.5719 (0.1443) [114]	0.5620 (0.1254) [97]					

Secondary outcomes	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Patient satisfaction							
QCM Technical, ^b mean (SD) [n]	All	62.66 (8.10) [519]	63.07 (8.37) [495]	$\Delta = -0.293$; $p = 0.572$	-1.310 to 0.725	0.0092	0.0013 to 0.0270
	Site 1	62.85 (7.63) [230]	63.62 (8.38) [239]				
	Site 2	61.37 (9.78) [163]	61.64 (8.63) [137]				
	Site 3	64.01 (6.07) [126]	63.58 (7.91) [119]				
QCM Interpersonal, ^c mean (SD) [n]	All	68.98 (8.78) [519]	68.07 (9.22) [495]	$\Delta = 3.105$; $p < 0.001$	1.522 to 4.689	0.0134	0.0020 to 0.0379
	Site 1	67.47 (10.03) [230]	67.97 (10.29) [239]				
	Site 2	71.18 (7.42) [163]	68.25 (8.51) [137]				
	Site 3	68.88 (7.28) [126]	68.06 (7.68) [119]				
Fall-related self efficacy (fear of falling), ^c mean (SD) [n]	All	3.676 (3.005) [585]	3.734 (3.059) [535]	$\Delta = -0.010$; $p = 0.956$	-0.353 to 0.334	0.0081	0.0009 to 0.0254
	Site 1	3.772 (3.039) [253]	3.942 (3.207) [256]				
	Site 2	3.148 (2.866) [182]	3.194 (2.666) [151]				
	Site 3	4.156 (3.035) [150]	3.955 (3.133) [128]				

TABLE 48 Secondary outcomes at 1 month, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral (continued)

Secondary outcomes	Site	Raw data		Adjusted comparison ^{a,b}		ICC	
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Proportion of patients who reported one or more further falls ^c	All	382/576 (66.3%)	373/528 (70.6%)	OR = 0.682; <i>p</i> = 0.010	0.510 to 0.914	0.0141	0.0029 to 0.0356
	Site 1	180/251 (71.7%)	179/253 (70.8%)				
	Site 2	112/178 (62.9%)	105/149 (70.5%)				
	Site 3	90/147 (61.2%)	89/126 (70.6%)				

QCM, Quality of Care Monitor.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality; indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; an additive group effect (Δ), in the same units as the dependent variable; Δ_k refers to observed data from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Age (*p* = 0.012); interaction between site 1 and group (*p* = 0.015).

d Site 1 (*p* = 0.001); distance to ED (*p* = 0.001); gender (*p* = 0.007); age (*p* = 0.015).

e Site 1 (*p* = 0.001); site 3 (*p* = 0.009).

f Age² (*p* = 0.001); out of hours (*p* = 0.015).

g Site 1 (*p* = 0.001); site 3 (*p* = 0.003); out of hours (*p* = 0.006); gender (*p* = 0.003).

h Distance to ED (*p* = 0.003).

i Interaction between site 1 and group (*p* < 0.001); interaction between site 3 and group (*p* = 0.031).

j Age (*p* < 0.001); out of hours (*p* = 0.001); site 1 (*p* < 0.001); site 3 (*p* = 0.019); recruitment point (*p* = 0.047).

k Interaction between site 1 and group (*p* = 0.016).

TABLE 49 Secondary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral

Secondary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			
		Intervention (A)	Control (B)	Estimate, p-value	95% CI	Estimate	95% CI
Duration of subsequent inpatient episodes: nights in hospital, truncated at 180 days, ^c mean (SD) [n]	All	11.18 (23.05) [2161]	12.08 (23.52) [1982]	$\Delta = 1.334; p = 0.132$	-0.403 to 3.070	0.0061	0.0018 to 0.0135
	Site 1	9.11 (19.07) [1081]	10.35 (20.41) [1055]				
	Site 2	14.48 (29.56) [540]	14.75 (30.01) [433]				
	Site 3	12.03 (22.52) [540]	13.42 (25.53) [494]				
Proportion with further reported subsequent injuries (fractures) ^d	All	213/2161 (9.9%)	195/1982 (9.8%)	OR = 1.200; $p = 0.106$	0.962 to 1.497	0.0166	0.0069 to 0.0309
	Site 1	78/1081 (7.2%)	80/1055 (7.6%)				
	Site 2	92/540 (17.0%)	59/433 (13.6%)				
	Site 3	43/540 (8.0%)	56/494 (11.3%)				
Quality of life							
SF-12 MCS, ^e mean (SD) [n]	All	42.93 (12.44) [238]	42.90 (12.02) [218]	$\Delta = 0.043; p = 0.970$	-2.203 to 2.290	0	n/a
	Site 1	43.52 (12.23) [107]	43.71 (12.12) [121]				
	Site 2	41.84 (12.42) [53]	38.95 (12.77) [43]				
	Site 3	42.84 (12.84) [78]	44.24 (10.67) [54]				
SF-12 PCS, ^f mean (SD) [n]	All	30.38 (11.31) [238]	31.31 (11.24) [218]	$\Delta = -0.93; p = 0.368$	-2.961 to 1.099	0	n/a
	Site 1	29.57 (11.22) [107]	31.42 (10.99) [121]				
	Site 2	31.61 (12.89) [53]	30.77 (12.79) [43]				
	Site 3	30.64 (10.31) [78]	31.49 (10.70) [54]				
Derived SF-6D, ^g mean (SD) [n]	All	0.5865 (0.1368) [260]	0.5933 (0.1349) [232]	$\Delta = -0.0069; p = 0.572$	-0.0308 to 0.0170	0	n/a
	Site 1	0.5842 (0.1395) [119]	0.6055 (0.1387) [128]				
	Site 2	0.5841 (0.1439) [57]	0.5619 (0.1352) [47]				
	Site 3	0.5913 (0.1292) [84]	0.5916 (0.1236) [57]				

continued

TABLE 49 Secondary outcomes at 6 months, analysis by treatment allocated, adjusted for significant covariates for matched participants eligible for a falls referral (continued)

Secondary outcomes	Site	Raw data		Adjusted comparison ^{a,b}			
		Intervention (A)	Control (B)	Estimate, <i>p</i> -value	95% CI	Estimate	95% CI
Fall-related self efficacy (Fear of falling) ^h mean (SD) [<i>n</i>]	All	4.474 (3.295) [313]	4.712 (3.345) [280]	$\Delta = -0.230$; $p = 0.328$	-0.776 to 0.260	0	n/a
	Site 1	4.497 (3.349) [139]	4.999 (3.416) [146]				
	Site 2	4.217 (3.254) [72]	4.010 (3.423) [60]				
	Site 3	4.624 (3.273) [102]	4.716 (3.090) [74]				
Proportion with change in place of residence at 6 months ⁱ	All	29/293 (9.9%)	15/252 (6.0%)	OR = 1.719; $p = 0.102$	0.897 to 3.292	0.0102	0.0007 to 0.0368
	Site 1	12/130 (9.2%)	5/135 (3.7%)				
	Site 2	5/73 (6.8%)	4/50 (8.0%)				
Proportion of patients who reported one or more further fall ^j	Site 3	12/90 (13.3%)	6/67 (9.0%)				
	All	209/304 (68.8%)	174/267 (65.2%)	OR = 0.418; $p = 0.009$	0.217 to 0.807	0.0171	0.0030 to 0.0457
	Site 1	91/134 (67.9%)	87/143 (60.8%)				
	Site 2	44/74 (59.5%)	43/56 (76.8%)				
	Site 3	74/96 (77.1%)	44/68 (64.7%)				

n/a, not applicable.

a As well as indicators for group, site, and their interaction, 'core' covariates considered are age (in years) and its square; distance to ED (in miles); recruitment point (based on days since start of study); seasonality, indicators of gender and whether or not the index call was made during out of [GP] hours.

b The comparison between groups reflects the variable under consideration; specifically, we report an OR from logistic regression models for binary variables; an additive group effect (Δ), from linear models for measurement variables. Statistically significant covariates are listed in further footnotes.

c Site 1 ($p < 0.001$); age ($p = 0.002$); age² ($p = 0.023$).

d Gender ($p < 0.001$); site 1 ($p < 0.001$); interaction between site 3 and group ($p < 0.001$).

e Age ($p = 0.031$).

f Age ($p < 0.001$); gender ($p = 0.043$).

g Age ($p = 0.003$).

h Age² ($p < 0.001$); out of hours ($p = 0.032$).

i Age² ($p = 0.042$).

j Interaction between site 3 and group ($p < 0.001$); interaction between site 1 and group ($p = 0.005$); age ($p = 0.022$); site 1 ($p = 0.015$); recruitment point ($p = 0.013$); distance to ED ($p = 0.010$).

A decorative graphic consisting of numerous thin, parallel green lines that curve from the left side of the page towards the right, creating a sense of movement and depth.

**EME
HS&DR
HTA
PGfAR
PHR**

Part of the NIHR Journals Library
www.journalslibrary.nihr.ac.uk

This report presents independent research funded by the National Institute for Health Research (NIHR). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health

Published by the NIHR Journals Library