

This is a repository copy of *Nanosegregation and Structuring in the Bulk and at the Surface of Ionic-Liquid Mixtures*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/115928/>

Version: Accepted Version

Article:

Slattery, John Martin orcid.org/0000-0001-6491-8302, Bruce, Duncan Watson orcid.org/0000-0002-1365-2222, Cabry, Christopher Patrick et al. (12 more authors) (2017) Nanosegregation and Structuring in the Bulk and at the Surface of Ionic-Liquid Mixtures. *Journal of Physical Chemistry B*. pp. 6002-6020. ISSN 1520-5207

<https://doi.org/10.1021/acs.jpcb.7b01654>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Nano-Segregation and Structuring in the Bulk and at the Surface of Ionic-Liquid Mixtures

Journal:	<i>The Journal of Physical Chemistry</i>
Manuscript ID	jp-2017-016542.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Bruce, Duncan; University of York, Chemistry Cabry, Christopher; University of York, Chemistry Nuno Canongia Lopes, Jose; Universidade de Lisboa Instituto Superior Tecnico Costen, Matthew; Heriot-Watt University, Institute of Chemical Sciences D'Andrea, Lucia; University of York, Chemistry Grillo, Isabelle; Institut Laue-Langevin Marshall, Brooks; Montana State University, Department of Chemistry and Biochemistry McKendrick, Kenneth; Heriot-Watt University, School of Engineering and Physical Sciences Minton, Timothy; Montana State University, Department of Chemistry and Biochemistry; Purcell, Simon; Heriot-Watt University, Institute of Chemical Sciences Rogers, Sarah; ISIS-STFC Neutron Scattering Facility, Harwell Science and Innovation Campus Slattery, John; University of York, Chemistry Shimizu, Karina; Instituto Superior Técnico, Smoll, Eric; Montana State University Bozeman Tesa-Serrate, Maria; Heriot-Watt University, Institute of Chemical Sciences</p>
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p>	
<p>toc_v3_small.tif</p>	

SCHOLARONE™
Manuscripts

Nano-Segregation and Structuring in the Bulk and at the Surface of Ionic-Liquid Mixtures

Duncan W. Bruce^{1,*}, Christopher P. Cabry¹, José N. Canongia Lopes^{4,5,*}, Matthew L. Costen^{2,*}, Lucía D'Andrea¹, Isabelle Grillo⁷, Brooks C. Marshall³, Kenneth G. McKendrick^{2,*}, Timothy K. Minton^{3,*}, Simon M. Purcell², Sarah Rogers⁶, John M. Slattery^{1,*}, Karina Shimizu^{4,5}, Eric Smoll³, María A. Tesa-Serrate²

¹*Department of Chemistry, University of York, Heslington, York YO10 5DD, UK*

²*Institute of Chemical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK*

³*Department of Chemistry and Biochemistry, Montana State University, Montana 59717, USA*

⁴*Centro de Química Estrutural, IST, Universidade de Lisboa, 1049-001 Lisboa, Portugal*

⁵*ITQB, Universidade Nova de Lisboa, Avenida República, 2780-157 Oeiras, Portugal*

⁶*ISIS, Science & Technology Facilities Council, Rutherford Appleton Laboratory, Chilton, UK*

⁷*ILL, 71 Avenue des Martyrs, CS 20156-38042, Grenoble Cedex 9, France*

Abstract

Ionic-liquid (IL) mixtures hold great promise, as they allow liquids with a wide range of properties to be formed by mixing two common components, rather than by synthesizing a large array of pure ILs with different chemical structures. In addition, these mixtures can exhibit a range of properties and structural organization that depend on their composition, which opens up new possibilities for the composition-dependent control of IL properties for particular applications. However, the fundamental properties, structure and dynamics of IL mixtures are currently poorly understood, which limits their more widespread application. This paper presents the first comprehensive investigation into the bulk and surface properties of IL mixtures formed from two commonly encountered ILs: 1-ethyl-3-methylimidazolium and 1-dodecyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide

1
2
3 ([C₂mim][Tf₂N] and [C₁₂mim][Tf₂N]). Physical property measurements (viscosity, conductivity and
4 density) find that these IL mixtures are not well described by simple mixing laws, suggesting that
5 their structure and dynamics are strongly composition-dependent. Small-angle X-ray and neutron
6 scattering (SAXS and SANS) measurements, alongside molecular dynamics (MD) simulations, show
7 that at low mole fractions of [C₁₂mim][Tf₂N], the bulk of the IL is composed of small aggregates of
8 [C₁₂mim]⁺ ions in a [C₂mim][Tf₂N] matrix, which is driven by nano-segregation of the long alkyl
9 chains and the polar parts of the IL. As the proportion of [C₁₂mim][Tf₂N] in the mixtures increases,
10 the size and number of aggregates increases until the C12 alkyl chains percolate through the system
11 and a bicontinuous network of polar and non-polar domains is formed. Reactive atom scattering-laser-
12 induced fluorescence (RAS-LIF) experiments, also supported by MD simulations, have been used to
13 probe the surface structure of these mixtures. It is found that the vacuum-IL interface is enriched
14 significantly in C12 alkyl chains, even in mixtures low in the long-chain component. These data
15 show, contrary to previous suggestions, that the [C₁₂mim]⁺ ion is surface active in this binary IL
16 mixture. However, the surface does not become saturated in C12 chains as its proportion in the
17 mixtures increases and remains unsaturated in pure [C₁₂mim][Tf₂N].
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 Introduction

37 Ionic liquids (ILs) are molten salts that are liquid at relatively low temperatures. They are often
38 defined as salts with melting temperatures below 100 °C, but a great many are liquid at room
39 temperature and below. There has been much interest in ILs in recent years and their unusual
40 combination of properties, such as very low vapor pressures, wide electrochemical windows, good
41 thermal stabilities, and ability to dissolve a wide range of solutes, has led to their use in a range of
42 areas. These include as reaction media for sustainable chemistry and catalysis;¹⁻¹⁰ in the
43 electrochemical deposition of metals and semiconductors;^{1, 11-16} nanoscience;¹⁷⁻²³ biomass
44 processing;²⁴ carbon capture and storage (CCS);²⁵⁻²⁶ analytical chemistry²⁷⁻³² and many more.
45
46
47
48
49
50
51
52
53

54 One of the advantages of ILs is that their properties can be altered substantially by changing
55 the anion, cation or both. This offers significant potential to tune the properties of an IL for a
56 particular application; for this reason, ILs are often described as 'designer solvents'. However, the
57
58
59
60

1
2
3 downside of this flexibility is that there are a huge number of potential combinations of ions that will
4 form ILs and it would be impossible to synthesize and test all of these for every application. Recent
5 studies have shown that it is possible to generate a library of liquids with a range of properties by
6 simply mixing two or more ILs.³³⁻³⁴ As such, IL mixtures offer the attractive prospect of being able to
7 access a range of properties from a relatively small number of common ILs by mixing, rather than
8 having to synthesize and store large numbers of different ILs. However, while significant progress has
9 been made in recent years in our understanding of pure ILs,³⁵⁻³⁶ comparatively little is known about
10 the fundamental properties of IL mixtures and how these relate to their composition.
11
12
13
14
15
16
17
18

19 The structure and dynamics of ILs, both in the bulk and at the gas-liquid interface, are critical
20 to many of their applications. In the bulk, the amphiphilic nature of many IL ions results in nano-
21 segregation of the liquid into polar and non-polar domains,³⁷ leading to the development of
22 nanostructure that has been studied using a range of experimental and computational approaches.³⁵
23 Considerably less is known about the effects of mixing ILs on their nanostructure, or indeed their
24 fundamental physical properties compared to pure ILs. Some reports of physicochemical
25 investigations of IL mixtures have found that they exhibit properties that are quite well described by
26 simple mixing laws,³⁸⁻⁵³ although this is somewhat surprising given that the chemical structures of
27 some of the ions involved are significantly different. However, non-ideal behavior has also been
28 observed in other studies (*e.g.* of conductivity,^{40-41, 54-56} density,^{46, 50, 57-59} viscosity^{48, 52, 60-61} and phase
29 behavior⁶²) and, interestingly, some IL mixtures that display ideal mixing behavior in some of their
30 physical properties exhibit significantly non-ideal behavior in others. This has led to the suggestion
31 that IL mixtures are best considered as 'double salt ionic liquids' (DSILs)³⁴ whose properties are
32 defined by their ultimate ionic composition, rather than by the properties of the pure ILs from which
33 they are derived.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 In most cases the origins of non-ideal behavior in IL mixtures are not well understood, but
50 factors such as size differences between the cations and/or the different hydrogen-bonding abilities of
51 ions in a mixture have been proposed as important.^{34, 48} As is so often the case with ILs, general
52 explanations for these phenomena that apply to all of the diverse range of ILs that are known are
53 unlikely to be found. Instead, detailed experimental and theoretical investigation of a range of
54
55
56
57
58
59
60

1
2
3 different types of IL mixtures is important to understand their behavior.^{42, 63-69} One area that has not
4 received significant attention, but is likely to play a key role in dictating the properties of IL mixtures,
5 is the nanostructure that results from aggregation of one IL in another. A small number of studies
6 have investigated the self-organization of ionic amphiphiles (*e.g.* long-chain imidazolium, pyridinium
7 and piperidinium halide or $[\text{BF}_4]^-$ salts) in both protic ($[\text{EtNH}_3][\text{NO}_3]$)⁷⁰⁻⁷⁶ and aprotic ($[\text{C}_4\text{mim}][\text{Tf}_2\text{N}]$
8 and $[\text{C}_n\text{mim}][\text{FeCl}_4]$ ($n = 2$ and 4)⁷⁷⁻⁷⁹ ILs. It is also well known that ILs can be used as solvents for
9 the self-organization of non-ionic surfactants.⁸⁰⁻⁸¹ What is currently unclear is the extent to which self-
10 organization plays a role in the observed properties of IL mixtures containing ions that are more
11 similar in their chemical structure. To the best of our knowledge there have been no reports of self-
12 organization (*i.e.* aggregation/micelle formation) in binary IL mixtures (mixtures containing a
13 common anion and two different cations or *vice versa*).

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The structures of IL surfaces have also been studied in some detail and are of particular interest in applications where the gas–liquid interface is important *e.g.* CCS, multi-phase catalysis and gas separation. A range of approaches have been used to study IL surfaces and some general features and design rules are beginning to emerge, although there is still much to learn in this area.^{36, 82-86} IL surface structure is influenced by variations of both the anion and cation and, within a series of ILs, increasing the alkyl chain length on the cation leads to a surface that is less polar and more saturated with alkyl chains.⁸⁷ Moving from large anions (*e.g.* $[\text{Tf}_2\text{N}]^-$) to smaller anions (*e.g.* $[\text{BF}_4]^-$) for a given chain length increases the density of alkyl chains at the surface, due to more efficient packing of the ions.⁸³ However, there have been relatively few investigations into the surface structure of IL mixtures.^{67, 84, 88-89} Information of the surface structure can be obtained from indirect classical measurements such as surface tension measurements, which we do not consider further here. More direct experimental techniques used include angle-resolved X-ray photoelectron spectroscopy (ARXPS),⁹⁰ Rutherford backscattering spectroscopy (RBS),^{86, 91} low-energy ion scattering (LEIS)⁸² and time-of-flight secondary ion mass spectrometry (ToF-SIMS),^{86, 92} and the technique which we have pioneered and use here, reactive-atom scattering with laser-induced fluorescence detection (RAS-LIF).^{83, 93-94} They have also been investigated by MD simulations.⁹⁵⁻⁹⁶ There are large discrepancies in the degree of surface enrichment reported, ranging from stoichiometric surface

compositions (detected by ARXPS)⁹⁰ to the complete absence of some constituting ions at the surface (reported in an LEIS study).⁸² It should be noted that these studies were concerned with different IL combinations, which might have distinct mixing behaviors. However, the same IL mixture investigated using two techniques (RBS and ToF-SIMS) showed different surface compositions.⁸⁶




Figure 1: The chemical structures of the IL ions used here.

This work provides a detailed picture of the physical properties and structure (in the bulk and at the gas-liquid interface) of binary IL mixtures that display properties that deviate significantly from ideal mixing behavior. The mixtures of interest are formed from two imidazolium ILs: [C₂mim][Tf₂N] and [C₁₂mim][Tf₂N] (figure 1) in various proportions (these mixtures will be denoted [C₂mim]_{1-x}[C₁₂mim]_x[Tf₂N], where $x = 0$ to 1).^{33, 53} By mixing an IL with no nanostructure, [C₂mim][Tf₂N] (figure 2, left), with one containing the amphiphilic cation, [C₁₂mim][Tf₂N] (figure 2, right), which is known to form a nano-segregated structure in the bulk and display significant surface ordering, interesting bulk and surface phenomena are observed.^{87, 93-94} This is the first time that many of these features have been reported, and that a unified discussion of both bulk and surface properties has been presented for binary IL mixtures.




Figure 2: MD simulation snapshots illustrating the bulk-phase structure of $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ (left) and $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ (right).⁹⁷ Atoms associated with the $[\text{Tf}_2\text{N}]^-$ ion are colored red. For $[\text{C}_2\text{mim}]^+$ and $[\text{C}_{12}\text{mim}]^+$ ions, the methyl substituent, imidazolium ring and the first methylene group of the alkyl substituent are shown in blue. All other atoms of the $[\text{C}_{12}\text{mim}]^+$ and $[\text{C}_2\text{mim}]^+$ cation alkyl substituents are shown in grey. Nano-segregation between the polar network (red/blue mesh) and non-polar domains (grey and green beads) can be seen in $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$, but no nano-segregation is seen in $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$.

Results

The bulk properties of $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures ($x = 0$ to 1) were investigated by measuring physical properties (viscosity, conductivity and density), as well as with SANS and SAXS experiments, and MD simulations. The structure of the vacuum-liquid interface was investigated using reactive-atom scattering probed *via* laser-induced-fluorescence (RAS-LIF) and MD simulations. As such, this is a most extensive characterization of bulk and surface properties of these mixtures.

Physical property measurements

The viscosity and conductivity of ILs are known to be quite sensitive to the nature of both the anion and cation. However, in previous work on IL mixtures, composition-dependent viscosities and

1
2
3 conductivities have often been found to be quite well described by simple mixing laws. In particular,
4
5 the Arrhenius or Grunberg-Nissan mixing law:⁹⁸⁻¹⁰⁰
6

$$\log \eta = x \log \eta_1 + (1 - x) \log \eta_2 \quad (1)$$

7
8
9 (where x is the mole fraction of one component of a binary mixture and η , η_1 and η_2 are the viscosity
10
11 of the mixture, component 1 and component 2, respectively) or the Katti and Chaudhri mixing law:¹⁰¹
12

$$\log \eta V_m = x \log \eta_1 V_{m,1} + (1 - x) \log \eta_2 V_{m,2} + \frac{\Delta G^E}{RT} \quad (2)$$

13
14
15 (where V_m , $V_{m,1}$ and $V_{m,2}$ are the molar volumes of the mixture and the two components, respectively,
16
17 and ΔG^E is the excess molar Gibbs energy of activation for flow).
18
19

20
21 Temperature-dependent viscosities and ionic conductivities were measured for at least twelve
22
23 compositions of $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ from 293 to 333 K (see ESI for details). The temperature
24
25 dependence of both the viscosity and conductivity data fitted well to the Vogel-Fulcher-Tammann
26
27 (VFT) equation for glass-forming systems (full experimental data and fitting is given in the ESI) and
28
29 all ILs behaved as Newtonian fluids under the shear-rate range used. The viscosities of mixtures in the
30
31 $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ system are poorly described by a linear mixing law:
32

$$\eta = x \eta_1 + (1 - x) \eta_2 \quad (3)$$

33
34 as expected for ILs (see ESI for details). However, large deviations from ideal mixing behavior are
35
36 also seen when using the Grunberg-Nissan mixing law (figure 3). Attempted fitting of the data to the
37
38 Katti and Chaudhri mixing law showed similar behavior, as the molar volumes of mixtures in this
39
40 system show only small deviations from ideal mixing behavior with changing composition (*vide*
41
42 *infra*).
43
44

45
46 Figure 3 (black squares) shows that according to the Grunberg-Nissan mixing law, at low
47
48 $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ loadings (up to 16 mol%) the viscosity of the mixtures behaves in essentially an ideal
49
50 way (within the error of the experiment), but from the 24 mol% point onwards there is a significant
51
52 deviation, which appears greatest at around 50 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. In this region, the IL mixtures
53
54 show a positive deviation and are more viscous than would be expected based on the Grunberg-Nissan
55
56 law.
57
58
59
60




Figure 3: Percentage deviation of viscosity and conductivity (at 293 K) from the Grunberg-Nissan mixing law for $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ mixtures as a function of composition.

Conductivity measurements on the $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ mixtures give a similar picture (figure 3, red circles), although the conductivity data deviate less significantly from the Grunberg-Nissan mixing law compared to the viscosity data. Again, there is a significant (within experimental error) difference between the observed and predicted conductivity from 24 mol% $[C_{12}mim][Tf_2N]$ onwards, being greatest at around 50 mol%. In the region where there is a significant difference between the observed and predicted conductivities, the mixtures have lower conductivities than expected from Grunberg-Nissan mixing behavior. This is consistent with the viscosity data, as conductivities are usually found to be approximately inversely proportional to viscosities in ILs.^{1, 102-}

104

The density (figure 4, black squares) and molar volumes of the $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ mixture system exhibit smaller deviations from ideal mixing behavior (using a linear mixing law as shown in equation 3) than the viscosity and conductivity. The observed densities are lower than expected based on a linear mixing law and the maximum deviation occurs at around $x = 0.5$. Excess

molar volumes calculated from the density data are somewhat scattered (figure 4, red circles), probably because they represent very small differences and any small errors in the measured densities are magnified, and do not show a particularly clear trend across the composition range. However, there is a tendency for small, positive excess molar volumes in these mixtures.




Figure 4: Percentage deviation of density and molar volume (at 293 K) from ideal behavior modelled using a linear mixing law for $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures as a function of composition. Note that error bars are included but are not visible as they are very small.

Small-angle X-ray scattering measurements

SAXS measurements have been shown in a range of studies to give good insight into the nanostructure present in pure ILs.¹⁰⁵⁻¹¹⁹ In order to further probe the bulk structure of $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures, SAXS measurements were performed on 16 compositions from $x = 0$ to $x = 1$ at 301 K (figure 5).




Figure 5: Selected SAXS data at 301 K for $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures as a function of composition. Data have been smoothed and stacked using an arbitrary offset on the y-axis to improve legibility and the PNPP, COP and CP peaks are labelled (see main text for descriptions). Data including all compositions can be found in the ESI.

Three peaks are often seen in SAXS data for ILs. These can be assigned to i) the characteristic distances between ions of different charge at close-contact (contact peak, CP) at high q ($\sim 1.3 \text{ \AA}^{-1}$); ii) characteristic distances related to the alternation of ions within the polar network of the IL (charge-ordering peak, COP) ($\sim 0.8\text{-}0.9 \text{ \AA}^{-1}$); and iii) characteristic separations caused by the segregation of polar and non-polar domains within the IL (polar/non-polar peak, PNPP) at low q ($\sim 0.2\text{-}0.25 \text{ \AA}^{-1}$).¹²⁰⁻¹²¹ In pure $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ only the COP is clearly visible, although there is a suggestion of a broad CP, which has been identified by others for this IL.^{113, 120} This is consistent with the fact that $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ does not form a nanostructured liquid. In contrast, in pure

1
2
3 [C₁₂mim][Tf₂N] the PNPP is clearly visible and this is indicative of nanoscale organization within the
4 liquid, with the PNPP indicating the formation, and length scale, of local bilayer structure formed by
5 the separation of the polar parts of the IL by the non-polar C12 chains on the cation. In the case of
6 [C₁₂mim][Tf₂N], MD simulations suggest that this nanostructure can be identified as a bicontinuous
7 network of polar and non-polar domains that percolates through the IL.¹²⁰

8
9
10
11
12
13 What has been discovered here is that in [C₂mim]_{1-x}[C₁₂mim]_x[Tf₂N] mixtures the PNPP
14 begins to appear as an incipient peak at 24 mol% [C₁₂mim][Tf₂N] and, as the concentration of
15 [C₁₂mim][Tf₂N] increases it gains in intensity, sharpens and moves to higher q (from ~ 0.20 to 0.24 \AA^{-1}
16 ¹, representing length scales of ~ 31 to 26 \AA , respectively). The COP changes little in intensity with
17 composition, but its position shifts slightly, representing an increase of around 5.5%, from 7.1 \AA in
18 pure [C₂mim][Tf₂N] to 7.5 \AA in pure [C₁₂mim][Tf₂N]. Finally, the CP becomes more visible as the
19 concentration of [C₁₂mim][Tf₂N] increases, although its position appears to vary little. These changes
20 will be discussed further below in the context of our complementary neutron scattering results and
21 MD simulations, and compared to data obtained for a series of pure [C_{*n*}mim][Tf₂N] ILs for $2 \leq n \leq$
22 10 .¹²⁰

23 24 25 26 27 28 29 30 31 32 33 34 35 36 **Small-angle neutron scattering measurements**

37 In order to gain additional insight into the structure of the bulk phase, SANS measurements were
38 performed on the [C₂mim-*d*₁₁]_{1-x}[C₁₂mim]_x[Tf₂N] system; data for seven compositions ($x = 0.04, 0.24,$
39 $0.32, 0.52, 0.74, 0.87$ and 1) are presented here. Data were collected on the D22 small-angle
40 diffractometer at the Institut Laue-Langevin (ILL) in Grenoble. Perdeuterated [C₂mim-*d*₁₁][Tf₂N] was
41 used to give contrast between the [C₂mim]⁺ and [C₁₂mim]⁺ ions in the mixtures, as protonated and
42 deuterated species typically have very different neutron scattering length densities (SLDs). Data were
43 collected over a q -range of *ca.* 0.01 to 0.9 \AA^{-1} , which corresponds to length scales of *ca.* 630 to 7 \AA .
44 Full details of the synthetic procedure used to prepare [C₂mim-*d*₁₁][Tf₂N] and SANS experimental
45 details are given in the ESI.




Figure 6: SANS data for $[\text{C}_2\text{mim-}d_{11}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures as a function of composition. Data are normalized to the background values determined during data fitting.

The SANS data in figure 6 show clear differences in neutron scattering as a function of composition. At lower loadings of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ in $[\text{C}_2\text{mim-}d_{11}][\text{Tf}_2\text{N}]$ (4 and 24 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$) the scattering intensity decreases relatively smoothly with increasing q . However, the data for the mixtures containing 52, 74 and 87 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ are dominated by a large peak between *ca.* 0.2 and 0.25 \AA^{-1} , which is also visible in the data at 32 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. This peak corresponds to the PNPP peak seen in the SAXS data.

The SANS data suggest a picture that is qualitatively similar to that obtained from the SAXS data, in that the PNPP appears in the data between 24 and 32 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. However, a more detailed picture can be obtained from the SANS data by fitting to scattering models; two different models have been employed. An ellipsoidal model describes a system composed of ellipsoidal scattering aggregates with a particular SLD (in this case the approximate SLD of non-

deuterated alkyl chains, $-3.7 \times 10^{-7} \text{ \AA}^{-2}$) dispersed within a solvent with a particular SLD (in this case the SLD of the $[\text{C}_2\text{mim}-d_{11}][\text{Tf}_2\text{N}]$ and cationic head groups of the $[\text{C}_{12}\text{mim}]^+$ ions, which depends on the composition of the mixture). The ellipsoids are characterized by axial and equatorial radii, and the volume fraction of the system that they occupy. At low loadings of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ (4 and 24 mol%) the ellipsoidal model fits the data quite well (see ESI for fits), although the fit is poorer at 24 mol%. This is potentially because an incipient PNPP at this composition changes the form of the scattering data without being obvious as a peak, due to the low q scattering that is present. The ellipsoidal model does not fit the data at $x \geq 0.32$.

The Teubner-Strey model, which has the following form, can be used to fit the data across all compositions (see ESI).¹²²⁻¹²³

$$I(q) = \frac{1}{a+c_1q^2+c_2q^4} + \text{background} \quad (4)$$

where a , c_1 and c_2 are empirical fitting parameters that can be related to three quantities of interest by the following relationships:

$$d\text{-spacing} = \frac{2\pi}{\sqrt{\frac{1}{2}\sqrt{\frac{a}{c_2} + \frac{1c_1}{4c_2}}}} \quad (5)$$

$$\text{correlation length} = \frac{1}{\sqrt{\frac{1}{2}\sqrt{\frac{a}{c_2} + \frac{1c_1}{4c_2}}}} \quad (6)$$

$$\gamma(\text{amphiphile strength}) = \frac{c_1}{\sqrt{4ac_2}} \quad (7)$$

The Teubner-Strey model is often used to describe systems that undergo a transition between discrete aggregates (or micelles) and a bicontinuous network as a function of composition (*e.g.* microemulsions).¹²²⁻¹²³ The model fits the data well at low $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ loadings, also reproducing the key features of the data at higher mole fractions of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ in the mixtures. These two models give complementary information about the system and the fitting parameters that were used are summarized in table 1.

Table 1: Fitting parameters for an ellipsoid model of the SANS data for [C₂mim-*d*₁₁]_{1-x}[C₁₂mim]_x[Tf₂N] mixtures (*x* = 0.04 and 0.24) and derived parameters for the Teubner-Strey model of the SANS data for all [C₂mim-*d*₁₁]_{1-x}[C₁₂mim]_x[Tf₂N] mixtures.

IL system	Ellipsoid model				Teubner-Strey model		
	Axial radius (Å)	Equatorial radius (Å)	Volume fraction occupied by ellipsoidal scatterers	Theoretical volume fraction occupied by C12 chains	<i>d</i> -spacing (Å)	Correlation length (Å)	Amphiphile strength, γ
<i>x</i> = 0.04	12(1)	65(1)	0.01(1)	0.03	42	5	0.3
<i>x</i> = 0.24	15(1)	6(1)	0.16(1)	0.14	40	6	0.1
<i>x</i> = 0.32	-	-	-	0.18	37	7	-0.1
<i>x</i> = 0.52	-	-	-	0.26	30	9	-0.5
<i>x</i> = 0.74	-	-	-	0.34	27	12	-0.8
<i>x</i> = 0.87	-	-	-	0.38	26	13	-0.8
<i>x</i> = 1	-	-	-	0.41	25	16	-0.9

At 4 and 24 mol% [C₁₂mim][Tf₂N], when the ellipsoid model can be used to fit the data, the scattering ellipsoids are found to be prolate and their dimensions correspond to approximately two or three C12 chains (an extended C12 chain is ~ 16 Å long) that interdigitate slightly. As expected, the volume fraction of the system occupied by the aggregated C12 alkyl chains increases with mole fraction of [C₁₂mim][Tf₂N] in the mixture. Those in the fitting model can be compared with a theoretical value calculated from the approximate volumes of the non-polar parts of the C12 alkyl chains and the polar parts of the system (the cationic head groups, including alkyl carbon atoms connected directly to nitrogen and the anions) assessed using a group-contribution model described previously.¹²⁴⁻¹²⁶ There is good agreement between the volume fractions used in the fitting model at 24 mol% [C₁₂mim][Tf₂N], but the fitted value is somewhat low at 4 mol% [C₁₂mim][Tf₂N], which may be due to the presence of isolated C12 chains in this mixture that may not contribute significantly to the observed neutron scattering.

The Teubner-Strey model allows the extraction of the 'amphiphile strength' parameter (γ): typically for $\gamma > 1$ the system is regarded as disordered, aggregate/micelle formation is seen for $0 < \gamma < 1$, bicontinuous structures are found where $-1 < \gamma < 0$ and lamellar structures are seen when $\gamma < -1$. As the SANS data for the [C₂mim-*d*₁₁]_{1-x}[C₁₂mim]_x[Tf₂N] mixtures are quite well described by the Teubner-Strey model (see ESI for fits), the γ parameter for these mixtures can be interpreted in the same way to give a picture of the structure in the bulk. For mixtures containing 4 and 24 mol%


1
2
3 [C₁₂mim][Tf₂N], $0 < \gamma < 1$ suggesting that isolated aggregates of C12 chains from the [C₁₂mim]⁺ ion
4 exist. This is consistent with the fact that the SANS data for these compositions can also be fitted to a
5 model based on ellipsoidal scattering objects. By 32 mol% [C₁₂mim][Tf₂N], γ has become just
6 negative and decreases in value steadily as the amount of [C₁₂mim][Tf₂N] in the mixture increases,
7 ultimately reaching a value of -0.9 for pure [C₁₂mim][Tf₂N]. This suggests a change in structure from
8 isolated aggregates of C12 chains to a bicontinuous network of non-polar and polar domains within
9 the ILs as a function of composition.
10
11
12
13
14
15
16

17
18 At compositions where $x < 0.32$, the d -spacing (which can be interpreted as the average
19 distance between scattering objects) decreases with increasing amounts of [C₁₂mim][Tf₂N] in the
20 mixture, as one would expect with an increasing concentration of long alkyl chains and thus an
21 increase in the aggregate density. As the amount of [C₁₂mim][Tf₂N] in the mixture increases, and the
22 structure switches from isolated aggregates to form a continuous non-polar sub-phase, there is a large
23 decrease in d -spacing (from 40 to 30 Å between 24 and 52 mol% [C₁₂mim][Tf₂N]). The d -spacing
24 then decreases steadily with increasing mole fraction of [C₁₂mim][Tf₂N] in the mixture to 25 Å in the
25 pure IL. Within this regime, the d -spacing is derived from the position of the PNPP, as in the SAXS
26 data, and can be interpreted in the same way *i.e.* as the length scale of local bilayer structures formed
27 from the C12 alkyl chains.
28
29
30
31
32
33
34
35
36
37
38
39

40 **Molecular dynamics simulations of the bulk**

41
42 In order to better understand the bulk structure of this system and to aid interpretation of the
43 SANS/SAXS data, molecular dynamics (MD) simulations were performed on [C₂mim]_{1-x}
44 [C₁₂mim]_x[Tf₂N] mixtures, where $x = 0, 0.04, 0.24, 0.32, 0.52, 0.74, 0.87$ and 1, matching
45 compositions studied *in vitro*. Condensed-phase MD simulations were carried out using the DLPOLY
46 package.¹²⁷ All molecules were modeled using the CL&P atomistic force field,¹²⁸⁻¹²⁹ which is based on
47 the OPLS-AA framework¹³⁰ but was to a large extent developed specifically for encompassing entire
48 homologous series of ionic liquids. Due to the slow dynamics of this type of system, special care was
49 taken to ensure the attainment of equilibrium conditions: (i) equilibrations started from initial low
50
51
52
53
54
55
56
57
58
59
60

1
2
3 density configurations, with ions placed randomly in periodic cubic boxes; (ii) typical equilibrations
4 were implemented for more than 1 ns; (iii) multiple re-equilibrations through the use of temperature
5 annealing and switching off and on of the Coulomb interactions were performed; (iv) further
6 simulation runs were used to produce equilibrated systems at the studied temperatures. The
7 equilibrated systems were used in production runs of at least 4 ns at 500 K (see ESI for details). The
8 simulations were performed at 500 K to improve the dynamics and equilibration of the simulated
9 systems. Nonetheless, comparisons between the main structural features of systems simulated at 500K
10 and 300K are possible, as proven by the green and grey lines in figures 6(a) and 6(b), corresponding
11 to pure $[C_2mim][Tf_2N]$ and $[C_{12}mim][Tf_2N]$ at the two temperatures (*vide infra*). The simulations
12 allow structure factor functions, $S(q)$, to be calculated for each IL (shown in figure 7). The structural
13 analysis has focused on the low- q region of the $S(q)$ functions ($0.2 < q/\text{\AA}^{-1} < 1.8$). In that range most
14 ionic liquids feature three peaks, the CP, COP and PNPP, as described above. These can be compared
15 to the SAXS data presented above, with which they are in excellent agreement, giving confidence in
16 the robustness of the computational methodology. In addition, the MD trajectories can be analyzed to
17 provide a more detailed understanding of the aggregation and structural characteristics of the ILs
18 under study (discussed in detail below).



1
2
3 Figure 7: (a) Experimental SAXS data for [C₂mim][Tf₂N], [C₁₂mim][Tf₂N] and selected [C₂mim]_{1-x}
4 [C₁₂mim]_x[Tf₂N] mixtures. (b) Calculated total structure factor functions, $S(q)$ for the same ILs and
5 IL mixtures at 500 K. (c) Calculated $S(q)$ functions of six pure [C_{*n*}mim][Tf₂N] ionic liquids at 300
6
7
8
9 K.^{119, 126} All $S(q)$ functions were shifted, in proportion to the volume fraction of the liquid occupied by
10
11 the non-polar alkyl chains, in the y-axis direction in order to improve legibility. Shifting the data in
12
13 this way also allows the IL mixtures to be compared to the pure ILs (see text for details and also note
14
15 the position of the colored markers on the right hand side of the middle graph).
16
17
18
19
20

21
22 Figure 7b shows the structure factor functions, $S(q)$, for the IL mixtures in this study and 6c
23 shows $S(q)$ functions of six pure ionic liquids (all even-numbered chains from [C₂mim][Tf₂N] to
24 [C₁₂mim][Tf₂N]) at 300 K.^{97, 120} Also shown for comparison (figure 7a) are experimental SAXS data
25
26 collected as part of this study (*vide supra*). A comparison of the experimental SAXS data and
27
28 calculated $S(q)$ functions for the IL mixtures shows good agreement between experiment and theory.
29
30 In both sets of data, the PNPP begins to appear as an incipient peak at $x = 0.32$, which grows in
31
32 intensity and shifts to higher q with increasing mole fraction of [C₁₂mim][Tf₂N]. In the case of the
33
34 MD simulations, the incipient PNPPs in the 24 to 32 mol% regions are difficult to ascertain due to the
35
36 impossibility of calculating data below q -values of 0.2 \AA^{-1} in the simulations, due to finite-size effects
37
38 of the simulation boxes. In addition, the COP shifts steadily to lower q with increasing mole fraction
39
40 of [C₁₂mim][Tf₂N]. A comparison of the [C₂mim][Tf₂N] and [C₁₂mim][Tf₂N] pure ionic liquids at 300
41
42 and 500 K (green and grey lines in figures 6b and 6c) shows that the $S(q)$ differences between the two
43
44 simulation temperatures are small (COPs and CPs become less pronounced at higher temperatures).
45
46 Thus, it is possible to make a comparison between the IL mixtures and pure ILs simulated at the two
47
48 temperatures.
49
50

51
52 It was shown in previous work^{57, 131} that the volumetric behavior of IL mixtures is additive
53
54 and almost ideal: that means that at a given temperature, the volume occupied by an ion in an ionic
55
56 liquid is almost constant, regardless of the counter-ion. Indeed, the fact that densities only deviate
57
58 very slightly from ideal mixing behavior in the system under study here (figure 4) suggests that this is
59
60

1
2
3 true in these IL mixtures. This notion supports the use of group-contribution rules to estimate the
4 molar volume of ionic liquids. Likewise, the contribution of the alkyl side chains to the molar volume
5 of an ion is also additive and constant: each methylene $-CH_2-$ group added to a chain at 298 K
6 increases the molar volume of the ion by around $17 \text{ cm}^3 \text{ mol}^{-1}$. This suggests that a comparison can be
7 made between different ILs (pure or mixtures) based on the volumes occupied by the polar network
8 and the non-polar domains. For instance, an equimolar ($[C_2\text{mim}][\text{Tf}_2\text{N}] + [C_{12}\text{mim}][\text{Tf}_2\text{N}]$) mixture
9 has around the same non-polar volume fraction as pure $[C_7\text{mim}][\text{Tf}_2\text{N}]$, 4 mol% $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ has a
10 non-polar volume fraction in-between that of $[C_2\text{mim}][\text{Tf}_2\text{N}]$ and $[C_3\text{mim}][\text{Tf}_2\text{N}]$ and so on. A
11 comparison of the mixtures and pure ILs shows the following revealing similarities and differences:

21 i) the PNPP and CP intensity trends in the mixtures are similar to those observed for the pure
22 ionic liquids, *i.e.*, both increase for systems with long-chain cations. The intensities of the PNPPs are
23 higher and are found at a lower q -value (see point ii) in the mixtures than in pure ionic liquids with
24 similar non-polar volume fractions. As in the SAXS data, PNPPs start to emerge as an incipient peak
25 in the 24 mol% $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ mixture and between the pure ILs $[C_4\text{mim}][\text{Tf}_2\text{N}]$ and
26 $[C_6\text{mim}][\text{Tf}_2\text{N}]$, and grow in intensity either with increasing mole fraction of $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ in the
27 mixture or with increasing chain length in the pure ILs. Thus, there is some correlation between the
28 appearance, and growth in intensity, of the PNPP and the non-polar volume-fraction of the liquid,
29 regardless of whether or not this was achieved through mixing or by increasing chain length on the
30 cation. This correlation is striking for the $[C_2\text{mim}]_{1-x}[C_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures investigated here, but
31 more work would be needed on related systems to establish whether it is a generally causal effect.

32 ii) the PNPP peak positions shift substantially to lower q with increasing chain length in the
33 pure ILs, corresponding to an increase in length scale of the non-polar network with increasing chain
34 length. However, the PNPP position in the mixtures is much less sensitive to composition, and shifts
35 in the opposite direction, to higher q with increasing mole fraction of $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ (from ~ 32 to 26
36 \AA). This origin of this shift is discussed below. Thus, the q -values of the PNPPs are controlled
37 predominantly by the longest chains present in the systems, since these dictate the length scale of the
38 non-polar network. Thus, in structural terms, the mixtures do not behave as though they were single
39 components with a chain length that is an average determined by composition.

iii) the COP intensities (relative to the CP intensities) are depressed in the case of the mixtures. Moreover, the COP intensity increases slightly in mixtures containing a larger proportion of [C₁₂mim][Tf₂N], whereas it decreases for pure ILs with increasing cation chain length. The less structured polar network of the mixtures may be related to the existence of isolated C2 moieties that perturb its organization.

iv) in both the SAXS data and MD simulations, the COP positions in the $S(q)$ functions shift monotonically to lower q -values both in the mixtures and in the pure ILs. It was found previously,^{97, 120-121, 132-134} that such a trend is related to the stretching of the polar network as it has to accommodate larger non-polar domains in its midst.

The MD trajectories were analyzed carefully to establish the different aggregation patterns that are possible within the non-polar domains that constitute the different systems.¹²⁰⁻¹²¹ Figure 8 shows the tail-tail aggregation distributions in the mixtures. In this figure, the probability of finding a chain belonging to an aggregate of a particular size is given on the y-axis, and the aggregate size is given on the x-axis.




Figure 8: Discrete probability distribution function, $P(n_a)$, of a given chain belonging to a non-polar aggregate of size n_a . Aggregates of only C12 chains are shown in the top panels (a, c, e, g, i and k) and aggregates containing both C12 and C2 chains are shown in the bottom panels (b, d, f, h, j and l). All y-axes are on the same (probability) scale, but different x-axes (and selected inserts) are used to highlight the key information. ' n_{12} ' and ' n_T ' represent the total number of C12 and (C2 + C12) chains

1
2
3 present in the mixtures, respectively. The numbers and arrows in the text correspond to out-of-the-
4
5 scale values, e.g. in 7a $P(n_a=1) = 68\%$ i.e. 68% of the C12 chains are isolated; $P(n_a=2) = 21\%$ i.e. 21%
6
7 of the C12 chains form “C12 dimers”; $P(n_a=3) = 8\%$ i.e. 8% of the C12 chains form “C12 trimers”.
8
9

10
11 The first composition (4 mol% [C₁₂mim][Tf₂N]) of figure 8a and 8b shows that the aggregates
12
13 formed by only dodecyl or both chains are always quite small (aggregates of ethyl chains from
14
15 [C₂mim]⁺ are small at all compositions, see ESI for details). On moving to the second composition (24
16
17 mol% [C₁₂mim][Tf₂N]) the mixture exhibits more interconnected, non-polar domains and the light
18
19 orange bar distribution in figure 8c shows the existence of large aggregates of C12 chains, which
20
21 nevertheless never contain all (350) of the dodecyl chains present in the system. Due to their length,
22
23 the majority of dodecyl chains are in contact with other dodecyl chains (only about 10% are isolated).
24
25 However, ethyl chains from the [C₂mim]⁺ ion essentially do not aggregate, as would be expected from
26
27 this essentially non-amphiphilic ion and around 67% of the ethyl chains are completely isolated (see
28
29 ESI). The next composition (32 mol% [C₁₂mim][Tf₂N]) is much closer to the percolation limit of the
30
31 alkyl side chains contained in the box. In fact, figures 7e and 7f show that the probability of C12
32
33 chains belonging to very large aggregates is already greater than that corresponding to the inclusion in
34
35 smaller aggregates (a bump starts to develop on the right-hand side of the histograms). The other
36
37 mixtures (figures 7g to 7l) show non-polar domains formed by large, percolating clusters of the long
38
39 chains, which can be described as being 'decorated' by ethyl chains from [C₂mim]⁺ ions. In these last
40
41 mixtures, the ethyl chains that are not connected to the dodecyl moieties become more and more
42
43 isolated from each other (ethyl-only clusters become scarcer, see ESI).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 9: Aggregate volume-to-box volume ratio, $R_V(n_a)$, as a function of maximum aggregate length-to-box-diagonal ratio, $R_d(n_a)$ for all chains aggregates in the mixture series. The black guidelines represent the volume-to-maximum-length ratios for spheres, oblate ellipsoids and prolate ellipsoids in order to guide the eye.

Figure 9 shows the relationship between the length and volume of the tail aggregates – a measure of their shapes – as their sizes become progressively larger (from the lower left to the upper right corners of the graphs). It shows that the shape of the aggregate is more or less independent of the type of chains it contains, but strongly dependent on its size: small aggregates have a prolate shape that evolves gradually as their size increases to more globular structures, perhaps best viewed as fragments of a non-polar network, which eventually coalesce and percolate throughout the system. This is consistent with the SANS modelling, where prolate ellipsoidal scattering objects fit the data reasonably well at low mole fractions of $[C_{12}mim][Tf_2N]$ in the mixtures and the system is described as a bicontinuous network of polar and non-polar domains when more $[C_{12}mim][Tf_2N]$ is present.




Figure 10: (a – c) Average size of contact neighbors, N_i , in the chain aggregates (filled color circles) as a function of the non-polar volume fraction of the liquids plotted as the x -axis. (a) ethyl aggregates; (b) dodecyl aggregates; (c) all non-polar aggregates.

The aggregation analysis can also calculate the average number of direct contact neighbors (other chains) that a given side chain may have within the non-polar domains (figure 10). The number of neighbors depends strongly on the size of the chains, since longer chains have more potential contact points, and their degree of aggregation. Figures 9b and 9c show that as the non-polar volume fraction of the liquid increases (with increasing $[C_{12}mim][Tf_2N]$ mole fraction) the number of nearest neighbors increases, which is similar to the trends observed for pure ILs with increasing alkyl chain length.¹³⁴ The ethyl chains show different behavior (figure 10a). These chains have very low average neighbor numbers, observed values below 1 denote a large proportion of isolated ethyl chains, and the number of neighbors decreases with increasing non-polar volume fraction. Thus, as in the pure IL $[C_2mim][Tf_2N]$ (figure 2, left), the ethyl chains of the $[C_2mim]^+$ ion in these mixtures are not strongly organized and are predominantly found as isolated ethyl chains distributed throughout the IL.

RAS-LIF experiments

As in our previous work on IL surface structure,^{83, 87, 93-94, 135-136} RAS-LIF experiments were used to probe the vacuum-liquid interface in $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ mixtures, where $x = 0, 0.01, 0.04,$

1
2
3 0.08, 0.16, 0.24, 0.52, 0.74, 0.87 and 1 (see ESI for experimental details). RAS-LIF is a very surface-
4 sensitive technique that aims to detect selectively the abstractable hydrogen atoms exposed at the
5 surface of an IL by reaction with gas-phase O(³P) atoms.^{83, 137} This produces OH radicals that scatter
6 back into the gas phase, where they are detected by laser-induced fluorescence (LIF). The
7 translational energy distribution of the incident O(³P) atoms is such that a significant fraction of the
8 O(³P) atoms are capable of abstracting an H atom from weaker, 2° C-H bonds, but not from stronger
9 C-H bonds.^{83, 136} Consequently, the detection of CH₂ groups in the alkyl chains is strongly favored
10 over other types of H, including those in the CH₃ groups or attached to the imidazolium ring. Thus,
11 interfacial reactivity observed in a RAS-LIF experiment is a quantitative measure of the accessibility
12 of these CH₂ groups at the surface.
13
14
15
16
17
18
19
20
21
22

23 The rotational distributions of the OH products were characterized by recording LIF
24 excitation spectra at a fixed delay between the photolysis and probe laser pulses in order to confirm
25 that the distributions did not vary materially across the range of mixtures. This was indeed the case as
26 described in detail in the ESI. The LIF intensity of a single transition could, therefore, be taken to be
27 representative of the total number density of OH scattered from the surface.
28
29
30
31
32

33 Appearance profiles (the LIF intensity as a function of delay between photolysis and probe
34 lasers) were measured on the most intense line in the spectrum corresponding to the Q₁(1) transition.
35 Details of the procedure and the correction for minor background signals are given in the ESI. The
36 resulting profiles for the pure ILs, selected mixtures and the squalane reference are presented in figure
37 11. Consistent with our previous results,^{83 87, 93-94} the OH yield from [C₂mim][Tf₂N] is almost below
38 the measurable level while the [C₁₂mim][Tf₂N] surface has slightly more than half the reactivity of
39 squalane. The yield of OH increases rapidly with *x*, showing significant reactivity for *x* of just 0.01.
40 At *x* ≥ 0.24, the profiles start to cluster together as they converge towards pure [C₁₂mim][Tf₂N].
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 11: Selected OH appearance profiles for the reference compound squalane (black squares), against which all signals are normalised, $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ (grey diamonds), $[C_2\text{mim}][\text{Tf}_2\text{N}]$ (green diamonds) and representative $[C_2\text{mim}]_{1-x}[C_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures (circles, color code in figure). Each profile is an average of three independent sets of ten profiles each as described in the text. Error bars are 1σ standard error of the mean.

Relative values of reactivity were obtained from the appearance profiles by performing a density-to-flux transformation and integrating the resulting flux profiles between 0 and $30\ \mu\text{s}$, as described previously.⁸³ The resulting values are tabulated in the ESI (Table S5) and are plotted against mole fraction in figure 12a, where they are expressed as a fraction of the yield from pure $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$. To allow a straightforward visual assessment of the departure from stoichiometry, a diagonal line has been included connecting the yields from pure $[C_2\text{mim}][\text{Tf}_2\text{N}]$ and $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$, respectively (i.e. between the points with $x = 0$ and 1). To the extent that mole fractions represent the proportions of the surface occupied by different ions correctly, this straight line corresponds to stoichiometric behavior. It is clear that on this basis the RAS-LIF experiments find that the interfacial

1
2
3 reactivity is higher than the stoichiometric result across almost the full range. However, some care is
4
5 needed here; it is questionable whether mole fraction is an unbiased independent variable against
6
7 which to measure stoichiometric surface occupancy, simply because it does not recognize the
8
9 different sizes of the two cation types. This point is explored quantitatively in the ESI, where two
10
11 alternative versions of figure 12 are provided. These are constructed using as x -axis either the volume
12
13 fraction (figure S50), calculated from the known relative molar volumes, V_m , of [C₂mim][Tf₂N] and
14
15 [C₁₂mim][Tf₂N], or the area fraction (figure S51, taken to be proportional to $V_m^{2/3}$ (*i.e.* the cross-
16
17 sectional area if the molecules were spherical). The positive deviations from stoichiometry remain
18
19 clear in both of these plots. The choice of construction does not therefore change the qualitative
20
21 conclusions, of which the principal one is that there is a clear, non-stoichiometric preference for the
22
23 longer-chain [C₁₂mim]⁺ cation to be exposed to attack at the surface at all concentrations except for
24
25 the very richest mixtures with x close to 1. This is emphasized in figure 12b, which shows, on the same
26
27 vertical and horizontal scales, the difference between the measured and stoichiometric values. The
28
29 maximum deviation occurs in the vicinity of a mole fraction $x = 0.24$ (which, for comparison, is
30
31 equivalent to a volume fraction of 0.34).
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 12: (a) Measured relative OH flux for [C₂mim]_{1-x}[C₁₂mim]_x[Tf₂N] mixtures plotted against mole fraction of [C₁₂mim]⁺. The diagonal line represents hypothetical stoichiometric behavior if the surface occupancy were simply proportional to mole fraction. (b) The difference between the experimental data points and the stoichiometric line.

Molecular dynamics simulations of the vacuum-IL interface

RAS-LIF vacuum-liquid scattering is ultimately controlled by the many-atom dynamics during and after gaseous projectile collision with the vacuum-liquid interface. However, previous studies demonstrate that liquid-dependent trends in RAS-LIF product flux can be qualitatively reproduced with an analysis of static vacuum-liquid interface composition.^{87, 135-136} To complement experiment, an additional set of MD simulations were performed to model the vacuum-liquid interface of $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures (where $x = 0, 0.04, 0.08, 0.16, 0.24, 0.52, 0.87$ and 1) using the same Lopes/Pádua all-atom, OPLS-type force field.^{129, 138-142} Vacuum-liquid-vacuum slab structures were prepared and subjected to 5 sequential equilibration cycles where each equilibration cycle consists of a phase of elevated temperature (500 K) propagation followed by a phase of room-temperature (298 K) propagation. Data analysis was only performed on the final phase of room-temperature propagation. Full simulation details are given in the ESI.




Figure 13: Top-down view of the vacuum-liquid interface of eight $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures where $x=0, 0.04, 0.08, 0.16, 0.24, 0.52, 0.87$, and 1. One periodic boundary condition unit cell, referred to in the text as “unit cell”, is shown for each liquid. Atoms associated with the $[\text{Tf}_2\text{N}]^-$ ion are colored red. The methyl group that terminates the ethyl chain in $[\text{C}_2\text{mim}]^+$ is colored green.

1
2
3 For $[\text{C}_2\text{mim}]^+$ and $[\text{C}_{12}\text{mim}]^+$ ions, the methyl substituent (4 atoms), imidazolium ring (8 atoms) and
4
5 the first methylene group of the alkyl substituent (3 atoms) are shown in blue. All other atoms of the
6
7 $[\text{C}_{12}\text{mim}]^+$ cation alkyl substituent are shown in grey.
8
9

10
11 Figure 13 is a compound image that displays the final trajectory frame of eight different $[\text{C}_2\text{mim}]_1$ -
12
13 $_x[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures. Each frame presents a top-down view of the vacuum-liquid interface for a
14
15 single unit cell and liquid molecules are rendered as a collection of opaque, van der Waals spheres.
16
17 Accessible surface area (ASA) calculations were used to identify interfacial atoms and quantify cation
18
19 surface density (Figure 14a) and surface 2° hydrogen density (Figure 14b) as a function of
20
21 $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ mole fraction. Interfacial atoms are defined as all liquid atoms near the vacuum-
22
23 liquid interface that can accommodate contact with a 1.5 \AA radius spherical probe.^{83, 136} In figure 14c,
24
25 the excess surface 2° hydrogen density computed by ASA analysis of molecular dynamics data is
26
27 compared to the excess OH yield from RAS-LIF experiments. As described earlier, these excess
28
29 properties are derived by computing the deviation of the observed signal from hypothetical linear
30
31 behavior between pure $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ and pure $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ with x as a dependent variable. Note
32
33 that the excess properties in Figure 14c are computed from data normalized to the signal of pure
34
35 $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$.
36

37
38 Since the nonpolar methyl group of $[\text{C}_2\text{mim}]^+$ (green in figure 13) is associated with 2 ethyl-
39
40 group methylene hydrogen atoms and the nonpolar undecyl group of $[\text{C}_{12}\text{mim}]^+$ (grey in figure 13) is
41
42 associated with 22 dodecyl-group methylene hydrogen atoms, figure 13 suggests that the total
43
44 methylene hydrogen atom density at the vacuum-liquid interface monotonically increases with x . This
45
46 is consistent with the experimental RAS-LIF data in figure 12a and the computed surface 2° hydrogen
47
48 density in figure 14b. The MD simulations show that this increase in 2° hydrogen atoms at the
49
50 vacuum-liquid interface is dominated by the $[\text{C}_{12}\text{mim}]^+$ dodecyl group at all mixture concentrations
51
52 studied. Due to the length of the dodecyl group, every surface-active $[\text{C}_{12}\text{mim}]^+$ cation adds a large
53
54 number of vacuum-accessible secondary hydrogen atoms. This increase in surface 2° hydrogen atoms
55
56 is most dramatic for mixtures with a low mole fraction of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$.
57
58
59
60

1
2
3 For pure [C₂mim][Tf₂N], the vacuum-liquid interface shows a strict pattern of alternating
4 charged ion packing. As the mole fraction of [C₁₂mim][Tf₂N] increases from 0, the number density of
5 [C₂mim]⁺ at the vacuum-liquid interface decreases and the number density of [C₁₂mim]⁺ at the
6 vacuum-liquid interface increases (figure 14a). Figure 14a also demonstrates that the surface C12
7 mole fraction approaches 0.5 when the bulk mole fraction (approximated from the total system ion
8 count) is ~0.16. Therefore, the C12 cation has a higher preference for the vacuum-liquid interface than
9 the C2 cation. This observation is consistent with other studies of ionic liquid mixtures where, in
10 general, larger ions are more surface active.^{67, 86, 91-92} Comparing the excess OH yield from RAS-LIF
11 experiments with the excess surface 2° hydrogen density from ASA analysis of MD simulations,
12 excellent agreement is observed in the shape of the excess curve, the peak excess location, and the
13 peak excess value (see figure 14c). Small modifications to the ASA parameters used to identify
14 vacuum-accessible atoms do not change the peak excess position and only introduce small changes to
15 the shape of the excess curve and the peak excess value. There are a number of possible technical
16 reasons for the remaining small deviation between experiment including the simulation size and
17 various limitations associated with the forcefield description of the liquid. However, the level of
18 agreement is particularly striking because we reiterate that the correlation being drawn is between the
19 RAS-LIF OH yield and the static vacuum-liquid interface composition as measured by a probe
20 particle in the ASA analysis. This particle is not constrained to approach the surface only along
21 trajectories that would lead to reaction of the corresponding O(³P) atoms. The correspondence
22 between experiment and simulation supports a correlation between RAS-LIF OH yield and the
23 [C₁₂mim]⁺ number density at the vacuum-liquid interface and provides validation of the vacuum-
24 liquid interface predicted by classical molecular dynamics.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 14: Surface structure and composition of $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ mixtures, as analyzed by accessible surface area (ASA) calculations. Surface area is defined by the x and y dimensions of the unit cell and does not account for corrugation of the liquid surface induced by capillary waves. The dashed diagonal line in panel b represents hypothetical linear or stoichiometric behavior between pure $[C_2mim][Tf_2N]$ and pure $[C_{12}mim][Tf_2N]$ as a function of total system mole fraction (see the discussion in the text relating to the construction of figure 12). In panel c, both experiment and simulation are normalized to the signal from pure $[C_{12}mim][Tf_2N]$ before computing the deviation from stoichiometric behavior. Error bars are $\pm 1\sigma$ standard error of the mean.

Discussion

A detailed molecular insight into the bulk-phase structure of the $[C_2mim]_{1-x}[C_{12}mim]_x[Tf_2N]$ system has been gained using a range of complementary techniques. This understanding allows a proposal to be made as to why the physical properties of this system change as they do with changing composition, and importantly to speculate on why some properties in this system deviate more strongly from ideal behavior than others.

Four key observations on the bulk properties in particular require explanation and they are considered in the discussion that follows:

- 1) The onset of significant deviation from ideal bulk physical property behavior occurs at around 24 mol% $[C_{12}mim][Tf_2N]$.
- 2) Viscosity data deviate more significantly from ideal mixing behavior than conductivity data, whereas densities show only small deviations from ideal behavior.

- 1
2
3 3) Observed viscosities are higher, and conductivities lower, in these mixtures than would be
4 expected based on simple mixing laws.
5
6
7 4) Densities are slightly lower than ideal, and there is a slight tendency for small, positive excess
8 molar volumes, in these mixtures.
9
10

11 The SAXS, SANS and MD simulations can now be brought together to give a consistent
12 picture of the bulk-phase structure of the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures. At low mole fractions
13 of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ (e.g. 4 mol%) only small aggregates (2-3 alkyl chains) of C12 chains are present in
14 the liquid. The appearance of an incipient PNPP from 24 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ reflects the presence
15 of some aggregates of C12 alkyl chains with a bilayer structure of sufficient size and coherence length
16 to produce a PNPP. MD simulations suggest that although small aggregates are more common,
17 aggregates with a range of sizes form at this composition, the largest containing ~17% of the C12
18 chains (see figure 8 for quantitative data regarding the number of alkyl chains found in aggregates at
19 each composition). SANS fitting gives a slightly positive γ value, which agrees with the simulations
20 that this structure is best described as isolated aggregates rather than bicontinuous network at this
21 composition. These aggregates subsequently coalesce with increasing $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ concentration,
22 initially to form globular fragments of a non-polar network, and ultimately a continuous, non-polar
23 sub-phase. This coalescence appears to be all but complete by 52 mol% $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ according to
24 negative γ values in the SANS fitting and aggregate analyses from the MD simulations. The observed
25 shift in position of the PNPP to higher q (and associated reduction in d -spacing) as the concentration
26 of C12 chains increases can be linked to the length scale of the bilayer structures present in the liquids
27 (i.e. the separation of the polar network by the alkyl chains of the non-polar sub-phase). Analysis of
28 the bulk-phase MD trajectories suggests that there is no significant change in C12 alkyl chain
29 interdigitation with increasing mole fraction of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ (see ESI for details). Thus, this small
30 change in length scale appears not to be related to the degree of interdigitation. Rather, there seems to
31 be a general contraction in the non-polar network with increasing mole fraction of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$
32 that maybe caused by small rearrangements in the relative morphologies of the two sub-phases (e.g.
33 loss of branched connectivity within the polar network or fewer “decorating” $[\text{C}_2\text{mim}]^+$ cations).
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 These observations can be compared to the behavior of pure $[C_n\text{mim}][\text{Tf}_2\text{N}]$ ILs with
4 increasing alkyl chain length. The PNPP appears and grows in intensity in a similar way in the
5 mixtures and pure ILs. In addition, SANS modelling and MD aggregate analysis of the mixtures and
6 pure ILs¹²⁰ suggests that larger aggregates of alkyl chains begin to appear around $[C_4\text{mim}][\text{Tf}_2\text{N}]$ and
7 $[C_2\text{mim}]_{0.76}[C_{12}\text{mim}]_{0.24}[\text{Tf}_2\text{N}]$, which have similar non-polar volume fractions, with percolation of the
8 non-polar network occurring by around $[C_6\text{mim}][\text{Tf}_2\text{N}]$ and by 52 mol% $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ in the
9 mixtures. SANS fitting and MD simulations suggest that percolation occurs at slightly different
10 compositions, both are based on models with some degree of error so this can be expected, but both
11 agree that percolation is complete at 52 mol% $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$. This suggests that the formation of
12 non-polar aggregates and their subsequent coalescence to form a bicontinuous network of polar and
13 non-polar domains in these ILs is driven primarily by the relative volume fractions of the non-polar
14 and polar parts of the system. However, the length scales of the non-polar domains are different in the
15 mixtures and pure ILs, being dictated by the length of the longest alkyl chain. Thus, it is possible to
16 relate some of the features of $[C_2\text{mim}]_{1-x}[C_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures to pure ILs with which they are
17 'equivalent' in terms of non-polar volume fraction, but not all. The IL mixtures are structurally
18 distinct from the pure ILs.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 This picture of the self-organized structure of the $[C_2\text{mim}]_{1-x}[C_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures can be
37 summarized visually in the four selected MD simulation snapshots presented in figure 15. The
38 percolation of the non-polar sub-phase for the two mixtures richer in $[C_{12}\text{mim}][\text{Tf}_2\text{N}]$ is obvious in
39 figures 14c and 14d. The large non-polar domains are dominated by interconnected C12 chains (grey
40 beads) 'decorated' at their periphery by a few C2 chains (light green beads). Some of the latter can still
41 be found in the midst of the polar network (red/blue mesh), separated from the non-polar sub-phase.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 15: Four MD simulation snapshots illustrating the nano-segregation between the polar network (red/blue mesh) and non-polar domains (grey and green beads) in $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ (a) $x = 0.04$, (b) $x = 0.24$, (c) $x = 0.52$ and (d) $x = 0.87$. Atoms associated with the $[\text{Tf}_2\text{N}]^-$ ion are colored red. For $[\text{C}_2\text{mim}]^+$ and $[\text{C}_{12}\text{mim}]^+$ ions, the methyl substituent, imidazolium ring and the first methylene group of the alkyl substituent are shown in blue. All other atoms of the $[\text{C}_{12}\text{mim}]^+$ cation alkyl substituent are shown in grey.

Armed with this understanding of the bulk structure of the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures, it is possible to propose why the physical properties of this system behave as they do with changing composition. In particular, the origins of the four key physical property observations that are highlighted above can be rationalized. Significant deviations from ideal behavior for the viscosities and conductivities of these mixtures become apparent at $x = 0.24$, with both sets of data suggesting

1
2
3 that ion mobility is lower than would be expected based on ideal mixing behavior. As discussed
4
5 above, this composition is composed mainly of small aggregates of C12 alkyl chains, but where larger
6
7 aggregates (up to around 60 chains, 17% of all C12 chains) can also be seen. It is proposed that it is
8
9 the presence of these larger aggregates, containing more connected C12 chains (which represent
10
11 fragments of a nascent non-polar network) that lead, through enhanced chain-chain interactions, to
12
13 lower ion mobility and the observed deviation from ideal behavior for the observed viscosities and
14
15 conductivities.

16
17 The fact that viscosity is the property that deviates most from ideal behavior in this system
18
19 could be linked to the reorganization of the longer-range ordering of the non-polar sub-phase that is
20
21 required in response to a shearing force. This may provide an additional resistance to flow alongside
22
23 local interionic interactions in the polar network. Recent work has suggested that network structures,
24
25 albeit based on rather different intermolecular interactions, formed in molten-salt mixtures also
26
27 contribute to higher than expected viscosities.¹⁴³⁻¹⁴⁴ In these cases MD simulations using polarizable
28
29 force fields, which capture the dynamics of the systems more accurately, have shown that viscous
30
31 relaxation can be dominated by reorganization of network structures, which form as a function of their
32
33 composition, within the liquids. In the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures, the fact that the
34
35 conductivity deviates less significantly from ideal behavior than viscosity presumably indicates that a
36
37 subtly different mechanism for ion mobility operates in the presence of an applied electric field.

38
39 The observation that density/excess molar volume data for the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$
40
41 mixtures deviate relatively little from ideal behavior compared with viscosity and conductivity data
42
43 suggests that overall, the sum of all interionic interactions changes relatively little when $[\text{C}_2\text{mim}]^+$
44
45 ions are substituted for $[\text{C}_{12}\text{mim}]^+$ ions. However, the small deviations from ideality that are observed
46
47 (lower densities and small, positive excess molar volumes) are consistent with slightly weakened
48
49 overall ion-ion interactions, consistent with the shift in COP position to lower q in SAXS experiments
50
51 (slightly longer cation···cation/anion···anion separations), which may result from the disruption of the
52
53 polar network by the formation of the non-polar sub-phase.
54
55
56
57
58
59
60

1
2
3 The corresponding changes taking place at the surface of the mixtures as the composition is
4 varied are now considered. Contrary to previous reports based on ARXPS measurements,⁹⁰ significant
5 surface enrichment of C12 alkyl chains was seen in RAS-LIF experiments and MD simulations. The
6
7 corresponding key features of the surface properties are: i) the surface deviates substantially from
8
9 stoichiometry in terms of excess C12 chains, from the lowest values of x ; ii) the deviation from
10
11 stoichiometry is at a maximum in the region of $x = 0.24$; iii) there is a region at very high x where the
12
13 behavior returns to the stoichiometric line; iv) even in pure $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ (*i.e.* at $x = 1$), the surface
14
15 is not fully saturated with C12 chains.
16
17
18

19 The picture that has emerged from the RAS-LIF measurements, supported by the MD
20
21 simulations, can be summarized in figure 16, which shows approximately two unit cells from the MD
22
23 simulations of each mixture slab, viewed from the side (camera perpendicular to surface normal).
24
25 Neat $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ shows no ordering at the vacuum liquid-interface. Figure 16 provides visual
26
27 confirmation of $[\text{C}_{12}\text{mim}]^+$ surface enrichment at all mixture mole fractions of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. At
28
29 very low mol% of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$, there is a preference for the dodecyl chain of surface active
30
31 $[\text{C}_{12}\text{mim}]^+$ ions to lay perpendicular to the interface-normal. However, in mixtures containing larger
32
33 amounts of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ the C12 chains are observed to assume a disordered but somewhat more
34
35 upright position.
36
37

38 The propensity for C12 chains to occupy the surface at low x is clearly an expression of their
39
40 relative energetic preference for the surface over the bulk in a dilute mixture in $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$
41
42 compared to the same competition between surface and bulk in pure $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. This can be
43
44 understood by a simple analogy to an amphiphilic surfactant in a polar solvent (here $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$)
45
46 where the amphiphile (in this case the $[\text{C}_{12}\text{mim}]^+$ ion) will occupy the surface preferentially to avoid
47
48 unfavorable hydrophobic interactions. Data from SANS and MD simulations show that by $x = 0.24$,
49
50 the $[\text{C}_{12}\text{mim}]^+$ ions form aggregates in the bulk and there is evidence that some of these aggregates are
51
52 quite large, which is interpreted as arising from a hydrophobic effect. As such, these aggregates
53
54 provide an alternative location for $[\text{C}_{12}\text{mim}]^+$ ions as x increases. The combination of surface and bulk
55
56 analysis then suggests that by $x = 0.24$ the two environments compete quite well as locations for
57
58 $[\text{C}_{12}\text{mim}]^+$ ions, so that the degree of surface enrichment increases more slowly. However, a thorough
59
60

1
2
3 analysis would also have to take into account cooperative behavior between the dodecyl chains at the
4 surface. Visual inspection of the simulations in figures 13 (in particular) and 16 shows that there
5
6 clearly is such cooperative behavior in the surface occupation by C12 chains, which tend to aggregate
7
8 even at low values of x . This clustering is still apparent at higher mole fractions, including in neat
9
10 $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$. As noted earlier, the surface of pure $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ does not appear entirely
11
12 saturated with alkyl chains. Indeed, the relatively large size of the $[\text{Tf}_2\text{N}]^-$ anion and the proximity of
13
14 cation and anion near the surface means that 'true' saturation will not be achieved – *i.e.* the situation as
15
16 visualized in Figure 13 is as saturated as this system can become, because the cations cannot pack
17
18 more closely together than this. In conventional surfactant chemistry, addition of surfactant to water
19
20 leads to a rapid reduction in the surface tension as the surfactants assemble at the surface, with a sharp
21
22 change in slope when the critical micelle concentration (cmc) is reached. However, careful analysis
23
24 and consideration of these IL mixtures, for which the C12 component is much more modestly surface
25
26 active than a typical surfactant in aqueous solution, shows that the surface only becomes saturated
27
28 relatively slowly and continues to change with $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ mole fraction all the way to $x = 1$.
29
30 A cmc is not evident in this system and it is clear that surface saturation *per se* does not drive
31
32 aggregation. Given that formation of aggregates in the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures is
33
34 observed by $x = 0.24$, at which composition the surface is clearly not saturated, it is clear that there is
35
36 little real analogy between this IL mixture system and conventional aqueous solutions of surfactants.
37
38 If the system were to be considered as a $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$ surfactant dissolved in $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$, the
39
40 data suggest that its surface activity is relatively weak and so it is clear that bulk aggregation in this
41
42 system is not driven by the absence of available surface sites.
43
44

45
46 Finally, it is observed (figure 12) that there is an initial, near stoichiometric, reduction in C12
47
48 surface coverage with the addition of a small proportion of $[\text{C}_2\text{mim}][\text{Tf}_2\text{N}]$ to neat $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$.
49
50 This is consistent with a primarily entropic, colligative effect, where the C2 chains distribute
51
52 statistically amongst highly aggregated C12 chains at both the surface and in the bulk.
53
54
55
56
57
58
59
60




Figure 16: Side view of the vacuum-liquid surface of $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures. One unit cell is shown for each liquid. For clarity, only the non-polar parts of the cation alkyl groups are shown: $[\text{C}_2\text{mim}]^+$ methyl group on the ethyl chain is shown in green and $[\text{C}_{12}\text{mim}]^+$ final eleven carbon atoms of the dodecyl chain are shown in white.

Conclusions

In a comprehensive study, a range of approaches have been used to gain unprecedented insight into the bulk and surface structure of the binary IL mixtures $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ and have related this understanding to key physical properties of these liquids. Unlike some IL mixtures, the structure and properties of the $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ system are strongly dependent on composition. The IL surface structure is influenced substantially by composition, and, while previous studies have suggested otherwise,⁹⁰ clear evidence of surface enrichment of $[\text{C}_{12}\text{mim}]^+$ ions is found, with this effect being most pronounced around $x = 0.24$. At low mole fractions of $[\text{C}_{12}\text{mim}][\text{Tf}_2\text{N}]$, the bulk physical properties (viscosities and conductivities) show no significant deviation from ideal behavior. However, from $x = 0.24$ onwards these properties behave in a non-ideal way. At this composition,

1
2
3 larger aggregates of C12 alkyl chains are present in the liquid, which coalesce with increasing x to
4 form a continuous non-polar sub-phase by $x = 0.52$ and it is proposed that increased chain-chain
5 interactions in these larger aggregates are linked to the observed deviations in viscosities and
6 conductivities. The length scales measured by scattering studies demonstrate that these IL mixtures
7 are structurally distinct from pure ILs with which they share similar non-polar volume fractions. In
8 the systems studied, the length scale of local bilayer structure in the liquids is clearly related to the
9 length of the longest alkyl chain in the IL. However, the appearance of the PNPP in SAXS and SANS
10 studies and the progression from small non-polar alkyl-chain aggregates to a continuous non-polar
11 sub-phase in MD simulations do show some correlation with the non-polar volume fraction of both
12 the pure ILs and mixtures. To sum up, there are some similarities and some differences between the
13 IL mixtures investigated here and related pure ILs, but it is clear that they are a distinct system. The
14 observations reported here are exciting, as they show that a range of structural and physical properties,
15 both in the bulk and at the surface, can be controlled simply by the composition of an IL mixture,
16 rather than by synthesizing a large number of pure ILs. The detailed understanding that this work
17 brings promises to allow the rational selection of specific surface or bulk structure/properties in
18 $[\text{C}_2\text{mim}]_{1-x}[\text{C}_{12}\text{mim}]_x[\text{Tf}_2\text{N}]$ mixtures when they are being considered for particular applications.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Supporting Information.**

39 Full experimental and computational details are given in the supplementary information.
40
41
42
43
44
45

46 **Acknowledgements**

47 The authors are grateful to the NSF and EPSRC (grant numbers NSF-CHE-1266032 and
48 EP/K032062/1), the University of York, Heriot-Watt University, Montana State University, the
49 Technical University of Lisbon and the Royal Society (International Exchanges Grants) for funding.
50 Gratitude is also expressed to the Science and Technology Facilities Council for neutron scattering
51 beam time at ISIS and the ILL (SANS Xpress and ILL experiment 6-03-435) and to Professor George
52
53
54
55
56
57
58
59
60

Schatz, Northwestern University, for helpful discussions and other input into the project. All data created during this research are available by request from the University of York Data Catalogue.

References

1. Wasserscheid, P.; Welton, T., *Ionic Liquids in Synthesis*, 2nd ed.; Wiley-VCH: Weinheim, 2008.
2. Zhao, D. B.; Wu, M.; Kou, Y.; Min, E., Ionic Liquids: Applications in Catalysis. *Catal. Today* **2002**, *74*, 157-189.
3. Welton, T., Ionic Liquids in Catalysis. *Coord. Chem. Rev.* **2004**, *248*, 2459-2477.
4. Giernoth, R., Homogeneous Catalysis in Ionic Liquids. In *In Situ NMR Methods in Catalysis*, Bargon, J.; Kuhn, L. T., Eds. 2007; Vol. 276, pp 1-23.
5. Parvulescu, V. I.; Hardacre, C., Catalysis in Ionic Liquids. *Chem. Rev.* **2007**, *107*, 2615-2665.
6. Betz, D.; Altmann, P.; Cokoja, M.; Herrmann, W. A.; Kuehn, F. E., Recent Advances in Oxidation Catalysis Using Ionic Liquids as Solvents. *Coord. Chem. Rev.* **2011**, *255*, 1518-1540.
7. Zhang, Q.; Zhang, S.; Deng, Y., Recent Advances in Ionic Liquid Catalysis. *Green Chem.* **2011**, *13*, 2619-2637.
8. Welton, T., Room-Temperature Ionic Liquids. Solvents for Synthesis and Catalysis. *Chem. Rev.* **1999**, *99*, 2071-2083.
9. Sheldon, R. A.; Lau, R. M.; Sorgedraeger, M. J.; van Rantwijk, F.; Seddon, K. R., Biocatalysis in Ionic Liquids. *Green Chem.* **2002**, *4*, 147-151.
10. Hallett, J. P.; Welton, T., Room-Temperature Ionic Liquids: Solvents for Synthesis and Catalysis. 2. *Chem. Rev.* **2011**, *111*, 3508-3576.
11. Endres, F.; MacFarlane, D.; Abbott, A., *Electrodeposition from Ionic Liquids*; Wiley-VCH: Weinheim, 2008.
12. Ohno, H., *Electrochemical Aspects of Ionic Liquids*; Wiley-Interscience: New Jersey, 2005.
13. Liu, F.; Deng, Y. D.; Han, X. P.; Hu, W. B.; Zhong, C., Electrodeposition of Metals and Alloys from Ionic Liquids. *J. Alloys Compd.* **2016**, *654*, 163-170.
14. Endres, F.; Bukowski, M.; Hempelmann, R.; Natter, H., Electrodeposition of Nanocrystalline Metals and Alloys from Ionic Liquids. *Angew. Chem., Int. Ed.* **2003**, *42*, 3428-3430.
15. Abbott, A. P.; McKenzie, K. J., Application of Ionic Liquids to the Electrodeposition of Metals. *Phys. Chem. Chem. Phys.* **2006**, *8*, 4265-4279.
16. Endres, F., Ionic Liquids: Solvents for the Electrodeposition of Metals and Semiconductors. *ChemPhysChem* **2002**, *3*, 144-154.
17. Dupont, J.; Scholten, J. D., On the Structural and Surface Properties of Transition-Metal Nanoparticles in Ionic Liquids. *Chem. Soc. Rev.* **2010**, *39*, 1780-1804.
18. Luska, K. L.; Migowski, P.; Leitner, W., Ionic Liquid-Stabilized Nanoparticles as Catalysts for the Conversion of Biomass. *Green Chem.* **2015**, *17*, 3195-3206.
19. Scholten, J. D.; Leal, B. C.; Dupont, J., Transition Metal Nanoparticle Catalysis in Ionic Liquids. *ACS Catal.* **2012**, *2*, 184-200.
20. Vollmer, C.; Janiak, C., Naked Metal Nanoparticles from Metal Carbonyls in Ionic Liquids: Easy Synthesis and Stabilization. *Coord. Chem. Rev.* **2011**, *255*, 2039-2057.
21. Precht, M. H. G.; Campbell, P. S., Metal Oxide and Bimetallic Nanoparticles in Ionic Liquids: Synthesis and Application in Multiphase Catalysis. *Nanotech. Rev.* **2013**, *2*, 577-595.
22. Antonietti, M.; Kuang, D. B.; Smarsly, B.; Yong, Z., Ionic Liquids for the Convenient Synthesis of Functional Nanoparticles and Other Inorganic Nanostructures. *Angew. Chem., Int. Ed.* **2004**, *43*, 4988-4992.
23. Torimoto, T.; Tsuda, T.; Okazaki, K.; Kuwabata, S., New Frontiers in Materials Science Opened by Ionic Liquids. *Adv. Mater.* **2010**, *22*, 1196-1221.
24. Brandt, A.; Grasvik, J.; Hallett, J. P.; Welton, T., Deconstruction of Lignocellulosic Biomass with Ionic Liquids. *Green Chem.* **2013**, *15*, 550-583.

- 1
2
3 25. Wang, C. M.; Luo, X. Y.; Zhu, X.; Cui, G. K.; Jiang, D. E.; Deng, D. S.; Li, H. R.; Dai, S.,
4 The Strategies for Improving Carbon Dioxide Chemisorption by Functionalized Ionic Liquids. *RSC*
5 *Adv.* **2013**, *3*, 15518-15527.
- 6 26. Ramdin, M.; de Loos, T. W.; Vlugt, T. J. H., State-of-the-Art of CO₂ Capture with Ionic
7 Liquids. *Ind. Eng. Chem. Res.* **2012**, *51*, 8149-8177.
- 8 27. Anderson, J. L.; Armstrong, D. W.; Wei, G. T., Ionic Liquids in Analytical Chemistry. *Anal.*
9 *Chem.* **2006**, *78*, 2892-2902.
- 10 28. Pandey, S., Analytical Applications of Room-Temperature Ionic Liquids: A Review of
11 Recent Efforts. *Anal. Chim. Acta* **2006**, *556*, 38-45.
- 12 29. Han, X.; Armstrong, D. W., Ionic Liquids in Separations. *Acc. Chem. Res.* **2007**, *40*, 1079-
13 1086.
- 14 30. Berthod, A.; Ruiz-Angel, M.; Carda-Broch, S., Ionic Liquids in Separation Techniques. *J*
15 *Chromatogr. A* **2008**, *1184*, 6-18.
- 16 31. Sun, P.; Armstrong, D. W., Ionic Liquids in Analytical Chemistry. *Anal. Chim. Acta* **2010**,
17 *661*, 1-16.
- 18 32. Ho, T. D.; Zhang, C.; Hantao, L. W.; Anderson, J. L., Ionic Liquids in Analytical Chemistry:
19 Fundamentals, Advances, and Perspectives. *Anal. Chem.* **2014**, *86*, 262-285.
- 20 33. Niedermeyer, H.; Hallett, J. P.; Villar-Garcia, I. J.; Hunt, P. A.; Welton, T., Mixtures of Ionic
21 Liquids. *Chem. Soc. Rev.* **2012**, *41*, 7780-7802.
- 22 34. Chatel, G.; Pereira, J. F. B.; Debbeti, V.; Wang, H.; Rogers, R. D., Mixing Ionic Liquids -
23 "Simple Mixtures" or "Double Salts"? *Green Chem.* **2014**, *16*, 2051-2083.
- 24 35. Hayes, R.; Warr, G. G.; Atkin, R., Structure and Nanostructure in Ionic Liquids. *Chem. Rev.*
25 **2015**, *115*, 6357-6426.
- 26 36. Lovelock, K. R. J., Influence of the Ionic Liquid/Gas Surface on Ionic Liquid Chemistry.
27 *Phys. Chem. Chem. Phys.* **2012**, *14*, 5071-5089.
- 28 37. Lopes, J. N. A. C.; Padua, A. A. H., Nanostructural Organization in Ionic Liquids. *J. Phys.*
29 *Chem. B* **2006**, *110*, 3330-3335.
- 30 38. Egashira, M.; Okada, S.; Yamaki, J., The Effect of the Coexistence of Anion Species in
31 Imidazolium Cation-Based Molten Salt Systems. *Solid State Ionics* **2002**, *148*, 457-461.
- 32 39. Navia, P.; Troncoso, J.; Romani, L., Viscosities for Ionic Liquid Binary Mixtures with a
33 Common Ion. *J. Solution Chem.* **2008**, *37*, 677-688.
- 34 40. Stoppa, A.; Buchner, R.; Hefter, G., How Ideal Are Binary Mixtures of Room-Temperature
35 Ionic Liquids? *J. Mol. Liq.* **2010**, *153*, 46-51.
- 36 41. Annat, G.; Forsyth, M.; MacFarlane, D. R., Ionic Liquid Mixtures-Variations in Physical
37 Properties and Their Origins in Molecular Structure. *J. Phys. Chem. B* **2012**, *116*, 8251-8258.
- 38 42. Aparicio, S.; Atilhan, M., Mixed Ionic Liquids: The Case of Pyridinium-Based Fluids. *J.*
39 *Phys. Chem. B* **2012**, *116*, 2526-2537.
- 40 43. Kakibe, T.; Hishii, J. Y.; Yoshimoto, N.; Egashira, M.; Morita, M., Binary Ionic Liquid
41 Electrolytes Containing Organo-Magnesium Complex for Rechargeable Magnesium Batteries. *J.*
42 *Power Sources* **2012**, *203*, 195-200.
- 43 44. Larriba, M.; Garcia, S.; Navarro, P.; Garcia, J.; Rodriguez, F., Physical Properties of N-
44 Butylpyridinium Tetrafluoroborate and N-Butylpyridinium Bis(Trifluoromethylsulfonyl)Imide Binary
45 Ionic Liquid Mixtures. *J. Chem. Eng. Data* **2012**, *57*, 1318-1325.
- 46 45. Ning, H.; Hou, M. Q.; Mei, Q. Q.; Liu, Y. H.; Yang, D. Z.; Han, B. X., The Physicochemical
47 Properties of Some Imidazolium-Based Ionic Liquids and Their Binary Mixtures. *Sci. China. Chem.*
48 **2012**, *55*, 1509-1518.
- 49 46. Song, D. Y.; Chen, J., Densities and Viscosities for Ionic Liquids Mixtures Containing
50 Eohmim Bf₄, Bmim Bf₄ and Bpy Bf₄. *J. Chem. Thermodyn.* **2014**, *77*, 137-143.
- 51 47. Song, D. Y.; Chen, J., Density and Viscosity Data for Mixtures of Ionic Liquids with a
52 Common Anion. *J. Chem. Eng. Data* **2014**, *59*, 257-262.
- 53 48. Larriba, M.; Navarro, P.; Beigbeder, J. B.; Garcia, J.; Rodriguez, F., Mixing and
54 Decomposition Behavior of { 4bmpy Tf₂N + emim EtSO₄ } and { 4bmpy Tf₂N + emim TfES } Ionic
55 Liquid Mixtures. *J. Chem. Thermodyn.* **2015**, *82*, 58-75.
- 56 49. Stolarska, O.; Soto, A.; Rodriguez, H.; Smiglak, M., Properties Modification by Eutectic
57 Formation in Mixtures of Ionic Liquids. *RSC Adv.* **2015**, *5*, 22178-22187.
- 58
59
60

- 1
2
3 50. Pinto, A. M.; Rodriguez, H.; Colon, Y. J.; Arce, A.; Arce, A.; Soto, A., Absorption of Carbon
4 Dioxide in Two Binary Mixtures of Ionic Liquids. *Ind. Eng. Chem. Res.* **2013**, *52*, 5975-5984.
- 5 51. Fox, E. T.; Weaver, J. E. F.; Henderson, W. A., Tuning Binary Ionic Liquid Mixtures:
6 Linking Alkyl Chain Length to Phase Behavior and Ionic Conductivity. *J. Phys. Chem. C* **2012**, *116*,
7 5270-5274.
- 8 52. Castiglione, F.; Raos, G.; Battista Appetecchi, G.; Montanino, M.; Passerini, S.; Moreno, M.;
9 Famulari, A.; Mele, A., Blending Ionic Liquids: How Physico-Chemical Properties Change. *Phys.*
10 *Chem. Chem. Phys.* **2010**, *12*, 1784-1792.
- 11 53. Clough, M. T.; Crick, C. R.; Gräsvik, J.; Hunt, P. A.; Niedermeyer, H.; Welton, T.;
12 Whitaker, O. P., A Physicochemical Investigation of Ionic Liquid Mixtures. *Chem. Sci.* **2015**, *215*,
13 1101-1114.
- 14 54. Every, H.; Bishop, A. G.; Forsyth, M.; MacFarlane, D. R., Ion Diffusion in Molten Salt
15 Mixtures. *Electrochim. Acta* **2000**, *45*, 1279-1284.
- 16 55. Bayley, P. M.; Best, A. S.; MacFarlane, D. R.; Forsyth, M., Transport Properties and Phase
17 Behaviour in Binary and Ternary Ionic Liquid Electrolyte Systems of Interest in Lithium Batteries.
18 *ChemPhysChem* **2011**, *12*, 823-827.
- 19 56. Taige, M. A.; Hilbert, D.; Schubert, T. J. S., Mixtures of Ionic Liquids as Possible
20 Electrolytes for Lithium Ion Batteries. *Z. Phys. Chem.* **2012**, *226*, 129-139.
- 21 57. Lopes, J. N. C.; Cordeiro, T. C.; Esperanca, J. M. S. S.; Guedes, H. J. R.; Huq, S.; Rebelo, L.
22 P. N.; Seddon, K. R., Deviations from Ideality in Mixtures of Two Ionic Liquids Containing a
23 Common Ion. *J. Phys. Chem. B* **2005**, *109*, 3519-3525.
- 24 58. Navia, P.; Troncoso, J.; Romani, L., Excess Magnitudes for Ionic Liquid Binary Mixtures
25 with a Common Ion. *J. Chem. Eng. Data* **2007**, *52*, 1369-1374.
- 26 59. Timperman, L.; Vigeant, A.; Anouti, M., Eutectic Mixture of Protic Ionic Liquids as an
27 Electrolyte for Activated Carbon-Based Supercapacitors. *Electrochim. Acta* **2015**, *155*, 164-173.
- 28 60. Khupse, N. D.; Kurolikar, S. R.; Kumar, A., Temperature Dependent Viscosity of Mixtures of
29 Ionic Liquids at Different Compositions. *Indian J Chem A* **2010**, *49*, 727-730.
- 30 61. Fumino, K.; Bonga, A. M.; Golub, B.; Paschek, D.; Ludwig, R., Non-Ideal Mixing Behaviour
31 of Hydrogen Bonding in Mixtures of Protic Ionic Liquids. *ChemPhysChem* **2015**, *16*, 299-304.
- 32 62. Maximo, G. J.; Santos, R.; Brandao, P.; Esperanca, J.; Costa, M. C.; Meirelles, A. J. A.;
33 Freire, M. G.; Coutinho, J. A. P., Generating Ionic Liquids from Ionic Solids: An Investigation of the
34 Melting Behavior of Binary Mixtures of Ionic Liquids. *Cryst. Growth Des.* **2014**, *14*, 4270-4277.
- 35 63. Shimizu, K.; Tariq, M.; Rebelo, L. P. N.; Lopes, J. N. C., Binary Mixtures of Ionic Liquids
36 with a Common Ion Revisited: A Molecular Dynamics Simulation Study. *J. Mol. Liq.* **2010**, *153*, 52-
37 56.
- 38 64. Llovel, F.; Valente, E.; Vilaseca, O.; Vega, L. F., Modeling Complex Associating Mixtures
39 with C-N-Mim Tf₂N Ionic Liquids: Predictions from the Soft-Saft Equation. *J. Phys. Chem. B* **2011**,
40 *115*, 4387-4398.
- 41 65. Brussel, M.; Brehm, M.; Pensado, A. S.; Malberg, F.; Ramzan, M.; Stark, A.; Kirchner, B.,
42 On the Ideality of Binary Mixtures of Ionic Liquids. *Phys. Chem. Chem. Phys.* **2012**, *14*, 13204-
43 13215.
- 44 66. Omar, S.; Lemus, J.; Ruiz, E.; Ferro, V. R.; Ortega, J.; Palomar, J., Ionic Liquid Mixtures-an
45 Analysis of Their Mutual Miscibility. *J. Phys. Chem. B* **2014**, *118*, 2442-2450.
- 46 67. Palchowdhury, S.; Bhargava, B. L., Segregation of Ions at the Interface: Molecular Dynamics
47 Studies of the Bulk and Liquid-Vapor Interface Structure of Equimolar Binary Mixtures of Ionic
48 Liquids. *Phys. Chem. Chem. Phys.* **2015**, *17*, 19919-19928.
- 49 68. Paschek, D.; Golub, B.; Ludwig, R., Hydrogen Bonding in a Mixture of Protic Ionic Liquids:
50 A Molecular Dynamics Simulation Study. *Phys. Chem. Chem. Phys.* **2015**, *17*, 8431-8440.
- 51 69. Zahn, S.; Stark, A., Order in the Chaos: The Secret of the Large Negative Entropy of
52 Dissolving 1-Alkyl-3-Methylimidazolium Chloride in Trihexyltetradecylphosphonium Chloride.
53 *Phys. Chem. Chem. Phys.* **2015**, *17*, 4034-4037.
- 54 70. Thomaier, S.; Kunz, W., Aggregates in Mixtures of Ionic Liquids. *J. Mol. Liq.* **2007**, *130*,
55 104-107.
56
57
58
59
60

- 1
2
3 71. Zhao, Y. R.; Chen, X.; Wang, X. D., Liquid Crystalline Phases Self-Organized from a
4 Surfactant-Like Ionic Liquid C(16)MIMCl in Ethylammonium Nitrate. *J. Phys. Chem. B* **2009**, *113*,
5 2024-2030.
- 6 72. Heintz, A.; Lehmann, J. K.; Kozlova, S. A.; Balantseva, E. V.; Bazyleva, A. B.; Ondo, D.,
7 Micelle Formation of Alkylimidazolium Ionic Liquids in Water and in Ethylammonium Nitrate Ionic
8 Liquid: A Calorimetric Study. *Fluid Phase Equilib.* **2010**, *294*, 187-196.
- 9 73. Kang, W. P.; Dong, B.; Gao, Y. N.; Zheng, L. Q., Aggregation Behavior of Long-Chain
10 Imidazolium Ionic Liquids in Ethylammonium Nitrate. *Colloid Polym. Sci.* **2010**, *288*, 1225-1232.
- 11 74. Shi, L.; Zhao, M.; Zheng, L., Micelle Formation by N-Alkyl-N-Methylpyrrolidinium Bromide
12 in Ethylammonium Nitrate. *Colloids Surf. A* **2011**, *392*, 305-312.
- 13 75. Shi, L. J.; Zheng, L. Q., Aggregation Behavior of Surface Active Imidazolium Ionic Liquids
14 in Ethylammonium Nitrate: Effect of Alkyl Chain Length, Cations, and Counterions. *J. Phys. Chem.*
15 *B* **2012**, *116*, 2162-2172.
- 16 76. Dai, C.; Du, M. Y.; Liu, Y. F.; Wang, S. L.; Zhao, J. H.; Chen, A.; Peng, D. X.; Zhao, M. W.,
17 Aggregation Behavior of Long-Chain Piperidinium Ionic Liquids in Ethylammonium Nitrate.
18 *Molecules* **2014**, *19*, 20157-20169.
- 19 77. Li, N.; Zhang, S. H.; Zheng, L. Q.; Dong, B.; Li, X. W.; Yu, L., Aggregation Behavior of
20 Long-Chain Ionic Liquids in an Ionic Liquid. *Phys. Chem. Chem. Phys.* **2008**, *10*, 4375-4377.
- 21 78. Shi, L. J.; Li, N.; Zheng, L. Q., Aggregation Behavior of Long-Chain N-Aryl Imidazolium
22 Bromide in a Room Temperature Ionic Liquid. *J. Phys. Chem. C* **2011**, *115*, 18295-18301.
- 23 79. Klee, A.; Prevost, S.; Gradzielski, M., Self-Assembly of Imidazolium-Based Surfactants in
24 Magnetic Room-Temperature Ionic Liquids: Binary Mixtures. *ChemPhysChem* **2014**, *15*, 4032-4041.
- 25 80. Greaves, T. L.; Drummond, C. J., Ionic Liquids as Amphiphile Self-Assembly Media. *Chem.*
26 *Soc. Rev.* **2008**, *37*, 1709-1726.
- 27 81. Greaves, T. L.; Drummond, C. J., Solvent Nanostructure, the Solvophobic Effect and
28 Amphiphile Self-Assembly in Ionic Liquids. *Chem. Soc. Rev.* **2013**, *42*, 1096-1120.
- 29 82. Villar-Garcia, I. J.; Fearn, S.; De Gregorio, G. F.; Ismail, N. L.; Gschwend, F. J. V.;
30 McIntosh, A. J. S.; Lovelock, K. R. J., The Ionic Liquid-Vacuum Outer Atomic Surface: A Low-
31 Energy Ion Scattering Study. *Chem. Sci.* **2014**, *5*, 4404-4418.
- 32 83. Tesa-Serrate, M. A.; Marshall, B. C.; Smoll, E. J.; Purcell, S. M.; Costen, M. L.; Slattery, J.
33 M.; Minton, T. K.; McKendrick, K. G., Ionic Liquid-Vacuum Interfaces Probed by Reactive Atom
34 Scattering: Influence of Alkyl Chain Length and Anion Volume. *J. Phys. Chem. C* **2015**, *119*, 5491-
35 5505.
- 36 84. Villar-Garcia, I. J.; Fearn, S.; Ismail, N. L.; McIntosh, A. J. S.; Lovelock, K. R. J., Fine
37 Tuning the Ionic Liquid-Vacuum Outer Atomic Surface Using Ion Mixtures. *Chem. Commun.* **2015**,
38 *51*, 5367-5370.
- 39 85. Ziemkiewich, M. P.; Zutz, A.; Nesbitt, D. J., Inelastic Scattering of Radicals at the Gas-Ionic
40 Liquid Interface: Probing Surface Dynamics of BMIM-Cl, BMIM-BF₄, and BMIM-Tf₂N by
41 Rovibronic Scattering of No [(2)Pi(1/2)(0.5)]. *J. Phys. Chem. C* **2012**, *116*, 14284-14294.
- 42 86. Nakajima, K.; Miyashita, M.; Suzuki, M.; Kimura, K., Surface Structures of Binary Mixtures
43 of Imidazolium-Based Ionic Liquids Using High-Resolution Rutherford Backscattering Spectroscopy
44 and Time of Flight Secondary Ion Mass Spectroscopy. *J. Chem. Phys.* **2013**, *139*, 224701.
- 45 87. Wu, B. H.; Zhang, J. M.; Minton, T. K.; McKendrick, K. G.; Slattery, J. M.; Yockel, S.;
46 Schatz, G. C., Scattering Dynamics of Hyperthermal Oxygen Atoms on Ionic Liquid Surfaces:
47 [EMIM][NTf₂] and [C₁₂MIM][NTf₂]. *J. Phys. Chem. C* **2010**, *114*, 4015-4027.
- 48 88. Oliveira, M. B.; Dominguez-Perez, M.; Freire, M. G.; Llovel, F.; Cabeza, O.; Lopes-da-
49 Silva, J. A.; Vega, L. F.; Coutinho, J. A. P., Surface Tension of Binary Mixtures of 1-Alkyl-3-
50 Methylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquids: Experimental Measurements
51 and Soft-Saft Modeling. *J. Phys. Chem. B* **2012**, *116*, 12133-12141.
- 52 89. Oliveira, M. B.; Dominguez-Perez, M.; Cabeza, O.; Lopes-da-Silva, J. A.; Freire, M. G.;
53 Coutinho, J. A. P., Surface Tensions of Binary Mixtures of Ionic Liquids with
54 Bis(trifluoromethylsulfonyl)imide as the Common Anion. *J. Chem. Thermodyn.* **2013**, *64*, 22-27.
- 55 90. Maier, F.; Cremer, T.; Kolbeck, C.; Lovelock, K. R. J.; Paape, N.; Schulz, P. S.;
56 Wasserscheid, P.; Steinrueck, H. P., Insights into the Surface Composition and Enrichment Effects of
57 Ionic Liquids and Ionic Liquid Mixtures. *Phys. Chem. Chem. Phys.* **2010**, *12*, 1905-1915.
- 58
59
60

- 1
2
3 91. Nakajima, K.; Oshima, S.; Suzuki, M.; Kimura, K., Surface Structures of Equimolar Mixtures
4 of Imidazolium-Based Ionic Liquids Using High-Resolution Rutherford Backscattering Spectroscopy.
5 *Surf. Sci.* **2012**, *606*, 1693-1699.
- 6 92. Souda, R., Surface Segregation in Binary Mixtures of Imidazolium-Based Ionic Liquids. *Surf.*
7 *Sci.* **2010**, *604*, 1694-1697.
- 8 93. Waring, C.; Bagot, P. A. J.; Slattery, J. M.; Costen, M. L.; McKendrick, K. G., O(3P) Atoms
9 as a Chemical Probe of Surface Ordering in Ionic Liquids. *J. Phys. Chem. A* **2010**, *114*, 4896-4904.
- 10 94. Waring, C.; Bagot, P. A. J.; Slattery, J. M.; Costen, M. L.; McKendrick, K. G., O(3P) Atoms
11 as a Probe of Surface Ordering in 1-Alkyl-3-Methylimidazolium-Based Ionic Liquids. *J. Phys. Chem.*
12 *Lett.* **2010**, *1*, 429-433.
- 13 95. Bhargava, B. L.; Balasubramanian, S., Layering at an Ionic Liquid-Vapor Interface: A
14 Molecular Dynamics Simulation Study of [BMIM][PF₆]. *J. Am. Chem. Soc.* **2006**, *128*, 10073-10078.
- 15 96. Garcia, G.; Atilhan, M.; Aparicio, S., Interfacial Properties of Double Salt Ionic Liquids: A
16 Molecular Dynamics Study. *J. Phys. Chem. C* **2015**, *119*, 28405-28416.
- 17 97. Shimizu, K.; Lopes, J. N. C., Probing the Structural Features of the 1-Alkyl-3-
18 Methylimidazolium Hexafluorophosphate Ionic Liquid Series Using Molecular Dynamics
19 Simulations. *J. Mol. Liq.* **2015**, *210*, 257-263.
- 20 98. Arrhenius, S., Origin of Grunberg-Nissan Mixing Law. *Z. Phys. Chem.* **1887**, *1*, 285.
- 21 99. Powell, R. E.; Roseveare, W. E.; Eyring, H., Diffusion, Thermal Conductivity, and Viscous
22 Flow of Liquids. *Ind. Eng. Chem.* **1941**, *33*, 430-435.
- 23 100. Grunberg, L.; Nissan, A. H., Mixture Law for Viscosity. *Nature* **1949**, *164*, 799-800.
- 24 101. Katti, P. K.; Chaudhri, M. M., Viscosities of Binary Mixtures of Benzyl Acetate with
25 Dioxane, Aniline, and M-Cresol. *J. Chem. Eng. Data* **1964**, *9*, 442-443.
- 26 102. MacFarlane, D. R.; Forsyth, M.; Izgorodina, E. I.; Abbott, A. P.; Annat, G.; Fraser, K., On the
27 Concept of Ionicity in Ionic Liquids. *Phys. Chem. Chem. Phys.* **2009**, *11*, 4962-4967.
- 28 103. Pinkert, A.; Ang, K. L.; Marsh, K. N.; Pang, S. S., Density, Viscosity and Electrical
29 Conductivity of Protic Alkanolammonium Ionic Liquids. *Phys. Chem. Chem. Phys.* **2011**, *13*, 5136-
30 5143.
- 31 104. Schreiner, C.; Zugmann, S.; Hartl, R.; Gores, H. J., Fractional Walden Rule for Ionic Liquids:
32 Examples from Recent Measurements and a Critique of the So-Called Ideal Kcl Line for the Walden
33 Plot. *J. Chem. Eng. Data* **2010**, *55*, 1784-1788.
- 34 105. Annapureddy, H. V. R.; Kashyap, H. K.; De Biase, P. M.; Margulis, C. J., What Is the Origin
35 of the Prepeak in the X-Ray Scattering of Imidazolium-Based Room-Temperature Ionic Liquids? *J.*
36 *Phys. Chem. B* **2010**, *114*, 16838-16846.
- 37 106. Gontrani, L.; Russina, O.; Lo Celso, F.; Caminiti, R.; Annat, G.; Triolo, A., Liquid Structure
38 of Trihexyltetradecylphosphonium Chloride at Ambient Temperature: An X-Ray Scattering and
39 Simulation Study. *J. Phys. Chem. B* **2009**, *113*, 9235-9240.
- 40 107. Hettige, J. J.; Kashyap, H. K.; Annapureddy, H. V. R.; Margulis, C. J., Anions, the Reporters
41 of Structure in Ionic Liquids. *J. Phys. Chem. Lett.* **2013**, *4*, 105-110.
- 42 108. Kashyap, H. K.; Hettige, J. J.; Annapureddy, H. V. R.; Margulis, C. J., Saxs Anti-Peaks
43 Reveal the Length-Scales of Dual Positive-Negative and Polar-Apolar Ordering in Room-
44 Temperature Ionic Liquids. *Chem. Commun.* **2012**, *48*, 5103-5105.
- 45 109. Rocha, M. A. A.; Neves, C. M. S. S.; Freire, M. G.; Russina, O.; Triolo, A.; Coutinho, J. A.
46 P.; Santos, L. M. N. B. F., Alkylimidazolium Based Ionic Liquids: Impact of Cation Symmetry on
47 Their Nanoscale Structural Organization. *J. Phys. Chem. B* **2013**, *117*, 10889-10897.
- 48 110. Russina, O.; Gontrani, L.; Fazio, B.; Lombardo, D.; Triolo, A.; Caminiti, R., Selected
49 Chemical-Physical Properties and Structural Heterogeneities in 1-Ethyl-3-Methylimidazolium Alkyl-
50 Sulfate Room Temperature Ionic Liquids. *Chem. Phys. Lett.* **2010**, *493*, 259-262.
- 51 111. Russina, O.; Triolo, A., New Experimental Evidence Supporting the Mesoscopic Segregation
52 Model in Room Temperature Ionic Liquids. *Faraday Discuss.* **2012**, *154*, 97-109.
- 53 112. Russina, O.; Triolo, A.; Gontrani, L.; Caminiti, R., Mesoscopic Structural Heterogeneities in
54 Room-Temperature Ionic Liquids. *J. Phys. Chem. Lett.* **2012**, *3*, 27-33.
- 55 113. Russina, O.; Triolo, A.; Gontrani, L.; Caminiti, R.; Xiao, D.; Hines, L. G.; Bartsch, R. A.;
56 Quitevis, E. L.; Plechkova, N.; Seddon, K. R., Morphology and Intermolecular Dynamics of 1-Alkyl-
57
58
59
60

- 3-Methylimidazolium Bis{(Trifluoromethane)Sulfonyl}Amide Ionic Liquids: Structural and Dynamic Evidence of Nanoscale Segregation. *J. Phys.: Condens. Mat.* **2009**, *21*.
114. Triolo, A.; Russina, O.; Bleif, H. J.; Di Cola, E., Nanoscale Segregation in Room Temperature Ionic Liquids. *J. Phys. Chem. B* **2007**, *111*, 4641-4644.
115. Triolo, A.; Russina, O.; Caminiti, R.; Shirota, H.; Lee, H. Y.; Santos, C. S.; Murthy, N. S.; Castner, E. W., Comparing Intermediate Range Order for Alkyl- Vs. Ether-Substituted Cations in Ionic Liquids. *Chem. Commun.* **2012**, *48*, 4959-4961.
116. Triolo, A.; Russina, O.; Fazio, B.; Appetecchi, G. B.; Carewska, M.; Passerini, S., Nanoscale Organization in Piperidinium-Based Room Temperature Ionic Liquids. *J. Chem. Phys.* **2009**, *130*.
117. Triolo, A.; Russina, O.; Fazio, B.; Triolo, R.; Di Cola, E., Morphology of 1-Alkyl-3-Methylimidazolium Hexafluorophosphate Room Temperature Ionic Liquids. *Chem. Phys. Lett.* **2008**, *457*, 362-365.
118. Xiao, D.; Hines, L. G.; Li, S. F.; Bartsch, R. A.; Quitevis, E. L.; Russina, O.; Triolo, A., Effect of Cation Symmetry and Alkyl Chain Length on the Structure and Intermolecular Dynamics of 1,3-Dialkylimidazolium Bis(Trifluoromethanesulfonyl)Amide Ionic Liquids. *J. Phys. Chem. B* **2009**, *113*, 6426-6433.
119. Zheng, W.; Mohammed, A.; Hines, L. G.; Xiao, D.; Martinez, O. J.; Bartsch, R. A.; Simon, S. L.; Russina, O.; Triolo, A.; Quitevis, E. L., Effect of Cation Symmetry on the Morphology and Physicochemical Properties of Imidazolium Ionic Liquids. *J. Phys. Chem. B* **2011**, *115*, 6572-6584.
120. Shimizu, K.; Bernardes, C. E. S.; Lopes, J. N. C., Structure and Aggregation in the 1-Alkyl-3-Methylimidazolium Bis(Trifluoromethylsulfonyl)Imide Ionic Liquid Homologous Series. *J. Phys. Chem. B* **2014**, *118*, 567-576.
121. Bernardes, C. E. S.; Shimizu, K.; Ferreira, A. I. M. C. L.; Santos, L. M. N. B. F.; Lopes, J. N. C., Structure and Aggregation in the 1,3-Dialkyl-Imidazolium Bis(Trifluoromethylsulfonyl)Imide Ionic Liquid Family: 2. From Single to Double Long Alkyl Side Chains. *J. Phys. Chem. B* **2014**, *118*, 6885-6895.
122. Schubert, K. V.; Strey, R.; Kline, S. R.; Kaler, E. W., Small-Angle Neutron-Scattering near Lifshitz Lines - Transition from Weakly Structured Mixtures to Microemulsions. *J. Chem. Phys.* **1994**, *101*, 5343-5355.
123. Teubner, M.; Strey, R., Origin of the Scattering Peak in Microemulsions. *J. Chem. Phys.* **1987**, *87*, 3195-3200.
124. Krossing, I.; Slattery, J. M.; Daguene, C.; Dyson, P. J.; Oleinikova, A.; Weingartner, H., Why Are Ionic Liquids Liquid? A Simple Explanation Based on Lattice and Solvation Energies. *J. Am. Chem. Soc.* **2006**, *128*, 13427-13434.
125. Slattery, J. M.; Daguene, C.; Dyson, P. J.; Schubert, T. J. S.; Krossing, I., How to Predict the Physical Properties of Ionic Liquids: A Volume-Based Approach. *Angew. Chem. Int. Ed.* **2007**, *46*, 5384-5388.
126. Bulut, S.; Eiden, P.; Beichel, W.; Slattery, J. M.; Beyersdorff, T. F.; Schubert, T. J. S.; Krossing, I., Temperature Dependence of the Viscosity and Conductivity of Mildly Functionalized and Non-Functionalized [Tf₂N] Ionic Liquids. *ChemPhysChem* **2011**, *12*, 2296-2310.
127. Smith, W.; Forester, T. R. *The DL_Poly Package of Molecular Simulation Routines (V.2.2)*, The Council for The Central Laboratory of Research Councils.: Warrington: Daresbury Laboratory, 2006.
128. Lopes, J. N. C.; Deschamps, J.; Padua, A. A. H., Modeling Ionic Liquids Using a Systematic All-Atom Force Field. *J. Phys. Chem. B* **2004**, *108*, 2038-2047.
129. Lopes, J. N. C.; Padua, A. A. H., Molecular Force Field for Ionic Liquids Composed of Triflate or Bistriflylimide Anions. *J. Phys. Chem. B* **2004**, *108*, 16893-16898.
130. Jorgensen, W. L.; Maxwell, D. S.; TiradoRives, J., Development and Testing of the Opls All-Atom Force Field on Conformational Energetics and Properties of Organic Liquids. *J. Am. Chem. Soc.* **1996**, *118*, 11225-11236.
131. Rebelo, L. P. N.; Lopes, J. N. C.; Esperanca, J. M. S. S.; Guedes, H. J. R.; Lachwa, J.; Najdanovic-Visak, V.; Visak, Z. P., Accounting for the Unique, Doubly Dual Nature of Ionic Liquids from a Molecular Thermodynamic, and Modeling Standpoint. *Accounts of Chemical Research* **2007**, *40*, 1114-1121.

- 1
2
3 132. Freitas, A. A.; Shimizu, K.; Lopes, J. N. C., Complex Structure of Ionic Liquids. Molecular
4 Dynamics Studies with Different Cation-Anion Combinations. *J. Chem. Eng. Data* **2014**, *59*, 3120-
5 3129.
- 6 133. Shimizu, K.; Tariq, M.; Freitas, A. A.; Padua, A. A. H.; Lopes, J. N. C., Self-Organization in
7 Ionic Liquids: From Bulk to Interfaces and Films. *J. Braz. Chem. Soc.* **2016**, *27*, 349-362.
- 8 134. Shimizu, K.; Lopes, J. N. C., Comparing the Structure of Different Ionic Liquid Series:
9 Bistriflamide V. Hexafluorophosphate; Pure V. Equimolar Mixtures. *Fluid Phase Equilib.* **2016**, *418*,
10 181-191.
- 11 135. Purcell, S. M.; Tesa-Serrate, M. A.; Marshall, B. C.; Bruce, D. W.; D'Andrea, L.; Costen, M.
12 L.; Slattery, J. M.; Smoll, E. J.; Minton, T. K.; McKendrick, K. G., Reactive-Atom Scattering from
13 Liquid Crystals at the Liquid-Vacuum Interface: [C₁₂MIM][BF₄] and 4-Cyano-4'-Octylbiphenyl
14 (8cb). *Langmuir* **2016**, *32*, 9938-9949.
- 15 136. Tesa-Serrate, M. A.; Smoll, E. J.; D'Andrea, L.; Purcell, S. M.; Costen, M. L.; Bruce, D. W.;
16 Slattery, J. M.; Minton, T. K.; McKendrick, K. G., Hiding the Headgroup? Remarkable Similarity in
17 Alkyl Coverage of the Surfaces of Pyrrolidinium- and Imidazolium-Based Ionic Liquids. *J. Phys.*
18 *Chem. C* **2016**, *120*, 27369-27379.
- 19 137. Tesa-Serrate, M. A.; Smoll, E. J.; Minton, T. K.; McKendrick, K. G., Atomic and Molecular
20 Collisions at Liquid Surfaces. *Annual Review of Physical Chemistry, Vol 67* **2016**, *67*, 515-540.
- 21 138. Lopes, J. N. A. C.; Padua, A. A. H.; Deschamps, J., Modeling Ionic Liquids Using an
22 Extended Opls-Aa Force Field. *Abstr. Pap. Am. Chem. Soc.* **2003**, *226*, U618-U618.
- 23 139. Lopes, J. N. C.; Deschamps, J.; Padua, A. A. H., Modeling Ionic Liquids Using a Systematic
24 All-Atom Force Field. *J. Phys. Chem. B* **2004**, *108*, 2038-2047.
- 25 140. Lopes, J. N. C.; Padua, A. A. H., Molecular Force Field for Ionic Liquids Iii: Imidazolium,
26 Pyridinium, and Phosphonium Cations; Chloride, Bromide, and Dicyanamide Anions. *J. Phys. Chem.*
27 *B* **2006**, *110*, 19586-19592.
- 28 141. Lopes, J. N. C.; Padua, A. A. H.; Shimizu, K., Molecular Force Field for Ionic Liquids Iv:
29 Trialkylimidazolium and Alkoxy-carbonyl-Imidazolium Cations; Alkylsulfonate and Alkylsulfate
30 Anions. *J. Phys. Chem. B* **2008**, *112*, 5039-5046.
- 31 142. Shimizu, K.; Almantariotis, D.; Gomes, M. F. C.; Padua, A. A. H.; Lopes, J. N. C., Molecular
32 Force Field for Ionic Liquids V: Hydroxyethylimidazolium, Dimethoxy-2-Methylimidazolium, and
33 Fluoroalkylimidazolium Cations and Bis(Fluorosulfonyl)Amide, Perfluoroalkanesulfonylamide, and
34 Fluoroalkylfluorophosphate Anions. *J. Phys. Chem. B* **2010**, *114*, 3592-3600.
- 35 143. Salanne, M.; Simon, C.; Turq, P.; Madden, P. A., Conductivity-Viscosity-Structure:
36 Unpicking the Relationship in an Ionic Liquid. *J. Phys. Chem. B* **2007**, *111*, 4678-4684.
- 37 144. Corradini, D.; Madden, P. A.; Salanne, M., Coordination Numbers and Physical Properties in
38 Molten Salts and Their Mixtures. *Faraday Discuss.* **2016**, *190*, 471-486.
- 39
40
41
42

TOC Graphic

