



UNIVERSITY OF LEEDS

This is a repository copy of *Crustal structure of western Hispaniola (Haiti) from a teleseismic receiver function study*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/116077/>

Version: Accepted Version

Article:

Corbeau, J, Rolandone, F, Leroy, S et al. (13 more authors) (2017) Crustal structure of western Hispaniola (Haiti) from a teleseismic receiver function study. *Tectonophysics*, 709. pp. 9-19. ISSN 0040-1951

<https://doi.org/10.1016/j.tecto.2017.04.029>

© 2017 Elsevier B.V. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.




eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Accepted Manuscript

Crustal structure of western Hispaniola (Haiti) from a teleseismic receiver function study

J. Corbeau, F. Rolandone, S. Leroy, K. Guerrier, D. Keir, G. Stuart, V. Clouard, R. Gallacher, S. Ulysse, D. Boisson, R. Bien-aimé Momplaisir, F. Saint Preux, C. Prépetit, J.-M. Saurel, B. Mercier de Lépinay, B. Meyer



PII: S0040-1951(17)30179-8
DOI: doi: [10.1016/j.tecto.2017.04.029](https://doi.org/10.1016/j.tecto.2017.04.029)
Reference: TECTO 127474
To appear in: *Tectonophysics*
Received date: 12 December 2016
Revised date: 25 April 2017
Accepted date: 26 April 2017

Please cite this article as: J. Corbeau, F. Rolandone, S. Leroy, K. Guerrier, D. Keir, G. Stuart, V. Clouard, R. Gallacher, S. Ulysse, D. Boisson, R. Bien-aimé Momplaisir, F. Saint Preux, C. Prépetit, J.-M. Saurel, B. Mercier de Lépinay, B. Meyer, Crustal structure of western Hispaniola (Haiti) from a teleseismic receiver function study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Tecto(2017), doi: [10.1016/j.tecto.2017.04.029](https://doi.org/10.1016/j.tecto.2017.04.029)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Crustal structure of western Hispaniola (Haiti) from a teleseismic receiver function study

J. Corbeau^{1,2}, F. Rolandone¹, S. Leroy¹, K. Guerrier³, D. Keir^{4,5}, G. Stuart⁶, V. Clouard²,
R. Gallacher⁴, S. Ulysse³, D. Boisson³, R. Bien-aimé Momplaisir³, F. Saint Preux⁷, C.
Prépetit⁷, J-M. Saurel², B. Mercier de Lépinay⁸ and B. Meyer¹

¹Sorbonne Universités, UPMC Univ Paris 06, CNRS, Institut des Sciences de la Terre de Paris (ISTeP), 4 place Jussieu, 75005 Paris, France

²Now at Observatoire Volcanologique et Sismologique de Martinique (OVSM), Institut de Physique du Globe de Paris (IPGP), UMR 7154, CNRS Sorbonne Paris Cité, Paris, France

³URGeo, Faculté des Sciences, Université d'Etat d'Haiti, Port-au-Prince, Haiti

⁴Ocean and Earth Science, National Oceanography Center Southampton, University of Southampton, Southampton, SO14 3ZH, UK

⁵Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Florence 50121, Italy

⁶Institute of Geophysics and Tectonics, School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

⁷Bureau des Mines et de l'Énergie, Unité Technique de Sismologie, Port-au-Prince, Haiti

⁸Géoazur, CNRS, Univ. Sophia-Antipolis, Valbonne, France

Corresponding author: Jordane Corbeau, jordane.corbeau@gmail.com

Abstract

Haiti, located at the northern Caribbean plate boundary, records a geological history of terrane accretion from Cretaceous island arc formations to the Eocene to Recent oblique collision with the Bahamas platform. Little is presently known about the underlying crustal structure of the island. We analyze P-waveforms arriving at 27 temporary broadband seismic stations deployed over a distance of 200 km across the major terrane boundaries in Haiti to determine the crustal structure of western Hispaniola. We compute teleseismic receiver functions using the Extended-Time Multi-Taper method and determine crustal thickness and bulk composition (V_p/V_s) using the H-k stacking method. Three distinctive and fault-bounded crustal domains, defined by their characteristic Moho depth distributions and bulk crustal V_p/V_s , are imaged across Haiti. We relate these domains to three crustal terranes that have been accreted along the plate boundary during the northeastwards displacement of the Caribbean plate and are presently being deformed in a localized fold and thrust belt. In the northern domain, made up of volcanic arc facies, the crust has a thickness of ~23 km and V_p/V_s of 1.75 +/- 0.1 typical of average continental crust. The crust in the southern domain is part of the Caribbean Large Igneous Province (Caribbean LIP), and is ~22 km thick with V_p/V_s of 1.80 +/- 0.03 consistent with plume-related rocks of late Cretaceous age. Significantly thicker, the crust in central Haiti has values of Moho depths averaging ~41 km and with V_p/V_s of 1.80 +/- 0.05. We propose that the central domain is likely constructed of an island arc upper crust with fragments of dense material originating from mafic lavas or LIP material. We produce a crustal profile along a N-S transect across Haiti accounting for the surface geology, shallow structural history, and new seismological constraints provided by variations of crustal thickness and bulk composition.

Keywords: Receivers Function, Haiti, crustal structure

1- Introduction

1.1- Overview

One of the keys to understanding the transpressive northern Caribbean plate boundary and its geodynamic evolution is constraining crustal thickness and composition. Several geological field studies have led to a description of the shallow structure and stratigraphy of the island of Hispaniola (e.g., Mann et al., 1995; Pubellier et al., 2000). The M_w 7.0 2010 Haiti earthquake prompted several geological and geophysical studies to constrain the fault geometry and the crustal structure in the area of the mainshock (e.g., Douilly et al., 2013). However, our knowledge of the mid- and lower-crustal tectonics remains very limited in the absence of whole crustal geophysical studies in Haiti. In Dominican Republic, the eastern part of Hispaniola Island, interpretation of crustal thickness from gravity data show thick crust below the southern flank of the Dominican Central Cordillera (Bowin, 1976). From April 2013 for nearly 14 months, a temporary seismic network consisting of 27 stations was deployed in Haiti (Trans-Haiti project) to determine its crustal thickness and bulk composition using teleseismic P-wave data. In addition, we analyze seismograms from 3 permanent stations of the Canadian National network (2010-2015) operated by the Bureau des Mines et de l'Energie and UTS. This paper is the first one about works on temporary deployment of seismometers cross-cutting the whole of the island.

The purpose of the present study is to determine crustal structure across Haiti from receiver function (RF) analysis below Haiti so as to better constrain plate scale tectonic evolution of the region responsible for the development of the current fold-thrust belt (Mann et al., 1995; Pubellier et al., 2000; Hernaiz Huerta et al., 2007). The receiver function analysis technique is an appropriate approach to image major discontinuities within the crust and upper mantle from incoming teleseismic P-waveforms (e.g. Ammon, 1991).

1.2- Geological and tectonic setting of Haiti

Haiti, the western part of the Hispaniola island, is located on the Northern Caribbean plate boundary, which separates the Caribbean plate from the North American plate (Fig. 1). Currently, the Caribbean plate moves in a east-northeast direction at about 20 mm/yr relative to the North American plate (Symithe et al., 2015). As the plate boundary is oriented E-W and the displacement vector of the Caribbean plate is about $N70^\circ$, the area undergoes oblique collision and transpression at a large restraining bend in the strike-slip plate boundary. The deformation of the Northern Caribbean plate boundary in Haiti is partitioned and two major E-W left-lateral strike-slip faults, the Septentrional-Oriente Fault Zone (SOFZ) in the North and the Enriquillo-Plantain-Garden Fault Zone (EPGFZ) in the south with both faults accommodating the strike-slip component of the displacement (Fig. 1). GPS geodesy shows that compression is mainly accommodated by Miocene to recent folding and thrusting in the North-Haitian thrust fault and the Massif de la Selle in southern Haiti (Symithe and Calais, 2016). The Trans-Haitian fold-and-thrust belt in central Haiti (Fig. 1) was active until the late Neogene (Mann et al., 1995) but does not appear to accommodate significant shortening today.

The geological and geodynamical history of Haiti is complex, however, two distinct domains have been identified. Haiti is part of the Cretaceous volcanic island arc constituted at the boundary of the Pacific realm (Pindell et al., 2006) called the Greater Antilles arc. The Greater Antilles volcanic arc was initiated by an eastward dipping subduction in Central America (Pindell et al., 2012; Van der Lelij, 2013; Hastie *et al.*, 2013). The Greater Antilles arc now consists of a part of Cuba, Hispaniola and Puerto Rico islands (Mann et al., 1995). This island arc constitutes two thirds of Hispaniola Island, and is mainly made up of arc magmatic facies (Boisson, 1987; Escuder Viruete et al., 2006). The southern part of Haiti has been interpreted to be part of the Caribbean Large Igneous Province (LIP), formed during the Cretaceous on the Pacific Farallon plate, over the Galapagos hotspot (Duncan and Hargraves, 1984). The LIP outcrops as a tholeiitic substratum associated with Upper Cretaceous sediments in the Southern Peninsula of Haiti (Calmus, 1983), and has been imaged south and west of Haiti with refraction and reflection data (Leroy et al., 2000; Mauffret et al., 2001; Corbeau et al., 2016a). The volcanic island arc and the LIP subsequently moved north- and eastwards from their Pacific position between the North and South American plates, thus partitioning the current Caribbean plate (Pindell, 2012). The Greater Antilles arc became an inactive intra-oceanic arc at the end of the Upper Cretaceous when it collided with the Bahamas carbonate platform (Leroy et al., 2000; Cruz-Orosa et al., 2012; Iturralde-Vinent et al., 2006). Between the island arc and the LIP lies the Quaternary Cul-de-Sac sedimentary basin (Fig. 1), which is bounded to the North by the thrusts of the Trans-Haitian belt, also called Haiti fold and thrust belt. This belt is formed of NW-SE thrusts having propagated towards the SW since the Lower Miocene (Pubellier et al., 2000).

1.3- Previous geophysical work

Previous geophysical studies have placed crude constraints on crustal structure in the south and in the vicinity of Haiti that help place our results in context and aid interpretation.

A compilation of seismic refraction data acquired in the Caribbean plate shows that the thickness of the crust is not uniform (Diebold et al., 1981; Mauffret and Leroy, 1997; Mauffret et al., 2001). The Caribbean oceanic crust is 5-km thick in Haitian sub-basin, Colombia and Venezuela basins (Fig. 1). In the middle of the Caribbean plate, the original oceanic crust is underplated by ultra-mafic material (Leroy et al., 2000), forming a LIP of ~10 to 15 km in thickness. The Beata Ridge (Fig. 1), 20-30 km thick, is composed of oceanic crust underplated by a ~15-20 km higher velocity (6 to 8 km/s) ultra-mafic magmatic material and covered by a thin volcanic layer of ~2 km (basaltic with a P-wave velocity of 4.5 to 6 km/s). Submersible sampling along the Beata Ridge to the South of Haiti (Mauffret et al., 2001) confirmed the deep origin of the underplated materials that are ultra-mafic rocks such as picrites or komatiites (Revillon et al., 2000).

In addition, Douilly et al. (2013) provide a velocity model for the crust in the southern part of Haiti, and show that the mean V_p/V_s in this area is 1.80, which is typical for oceanic crust mafic rocks (Christensen, 1996). Velocity model and receiver functions studies made by Moreno et al. (2002) and Gonzalez et al. (2012) show that the depth of the Moho is approximately 20 km in the south of Cuba. In eastern Hispaniola, a seismic refraction study shows that the Moho is at approximately 24 km depth, and reaches roughly 30 km deep in the central part of Hispaniola (Nuñez et al., 2015). The S- and Lg-wave study

of McNamara et al. (2012) implies that the Moho in the southwestern part of Hispaniola is 20.6 km depth.

2- Data and Method

2.1- Data

We use data from the 27 broadband stations of the temporary Trans-Haiti network deployment (April 2013 - June 2014), spanning Haiti from North to South (Fig. 2). This unique data set used SEIS-UK and iSTeP UPMC instruments that included 15 CMG-40TD and 12 CMG-6TD seismometers (30s natural period, 100 sps sampling rate). Additionally, we use data from 3 broadband permanent stations of the Canadian National Seismic network installed in Haiti after the 2010 Mw 7.0 earthquake and co-operated by the Bureau des Mines et de l'Energie - UTS (Fig. 2).

To construct receiver functions we use seismograms from teleseismic earthquakes of magnitudes $M_w > 5.0$, with epicentral distances between 30° and 90° . The initial data set includes a total of 580 events recorded by the Trans-Haiti stations between April 2013 and June 2014, and 2473 events for the Canadian stations between February 2010 and August 2015. The data are generally noisy, due to cultural noise sources, and a second order Butterworth filter with corner frequencies of 0.03 and 2 Hz is applied to the data prior to analysis. A visual quality control is subsequently performed on all the data such that only seismograms that have clear direct-P arrivals are used to produce receiver functions. This results in a reduced data set of 149 event records, with a range of between 2 and 13 per station. A map showing the epicentral distribution of the analyzed earthquakes is given in Fig. 3. Note that the majority of the events arrive from the south of the seismic network.

2.2- Receiver function technique

To image the crustal structure beneath Haiti we use the receiver function (RF) method (e.g. Ammon, 1991), which is a deconvolution technique for isolating the P-to-S converted phases (Ps) and associated reverberations (PpPs and PsPs + PpSs) of an incoming P-waveform beneath a seismic station. The incident P-wave on the vertical component is deconvolved from the radial and tangential components to remove source complexity and leave a series of pulses that represent the P-to-S phase conversions (the so-called receiver function).

Several deconvolution methods have been developed with different noise stabilization processes: deconvolution in the frequency domain (Langston, 1979; Owens et al., 1983; Ammon, 1991); deconvolution in the time domain by least squares estimation (Abers et al., 1995); iterative deconvolution in the time domain (Gurrola et al., 1995; Ligorria and Ammon, 1999) and multi-taper frequency domain cross-correlation (MTRF) (Park and Levin, 2000; Helffrich, 2006). The advantage of the last method is that it works on the correlated signal and mitigates against noise on the RFs. It has been recommended in the case of oceanic island studies, such as the Cap Verde islands (Helffrich et al., 2010), the Seychelles islands (Hammond et al., 2013), and the Canary islands (Lodge et al., 2012; Martinez-Arevalo et al., 2013), and in our case we choose it for its capacity to remove anthropic noise.

To estimate the Moho depth below the island of Haiti, we use the Extended-Time Multitaper Frequency domain Cross-Correlation Receiver-Function (ETMTRF) technique of Helffrich (2006), which is based on the MTRF method of Park and Levin (2000). The ETMTRF

method uses a series of short and overlapping multiple tapers which window the time series across its full length, and sum the individual Fourier transformed signals to produce a RF estimate.

2.3- Moho depth and Vp/Vs estimation

We use a receiver function stacking technique (H- κ stacking) developed by Zhu and Kanamori (2000) to estimate the Moho depth (H) and the bulk crust ratio V_p/V_s (κ). This method uses the following equations (1), (2), (3), and (4) for stacking the RFs, and gives the results on a 10,000 point grid which shows the plausible range of values for H on increments of 0.4 km between 10 and 50 km and κ on increments of 0.004 between 1.6 and 2.0 (Christensen, 1996):

$$s(H, \kappa) = \sum_{j=1}^N w_1 r_j(t_1) + w_2 r_j(t_2) - w_3 r_j(t_3), \quad (1)$$

$$t_1 = H \left[\sqrt{\frac{1}{V_s^2} - p^2} - \sqrt{\frac{1}{V_p^2} - p^2} \right], \quad (2)$$

$$t_2 = H \left[\sqrt{\frac{1}{V_s^2} - p^2} + \sqrt{\frac{1}{V_p^2} - p^2} \right], \quad (3)$$

$$t_3 = 2H \sqrt{\frac{1}{V_s^2} - p^2}, \quad (4)$$

where w_1 , w_2 , w_3 are weights, $r_j(t_i)$ are the amplitudes at the arrival times for each of the raypaths evaluated, N is the number of receiver functions, and p is the ray parameter determined from the IASPEI travel time tables (Kennet, 1991). The weights were chosen as $w_1 = 0.5$, $w_2 = 0.3$ and $w_3 = 0.2$ as suggested by Zhu and Kanamori (2000). The

errors in the results were calculated by taking the maximum axes of the 95% confidence interval of the grid plotted using the Zhu and Kanamori (2000) stacking method. The H- κ stack was performed assuming a mean crustal P-wave velocity (V_p) of 6.6 km/s. This mean crustal P-wave velocity comes from a study of the Pg seismic phase over Hispaniola (McNamara et al., 2012). Crustal thickness, H, and average Vp/Vs determination are not very sensitive to variations in the average Vp, as pointed out by Zhu and Kanamori (2000), and testing Vp within the range 6.4-6.7 km/s on increments of 0.1 km/s provides very similar results of H and Vp/Vs ratio.

Our final analysis uses the RFs computed from 149 earthquakes, from which one or more reverberant Moho phases (PpPs or PpSs + PsPs) were clearly identifiable.

3- Results

Of the 30 stations, 27 yielded receiver functions with coherent P-to-S arrivals. Of these, 19 displayed PpPs and PpSs + PsPs arrivals as well as Ps, enabling H- κ analysis (Table 1). Examples of stacked receiver functions and H- κ plots for stations PIGN, LGNH, and MGOA are shown in Figure 4. For the 8 stations at which no clear reverberant phases were found automatically by the program, RFs were combined to form single station stacks and enhance the signal of the reverberations. At these stations, a clear Ps arrival is identifiable and can be picked manually, but the H- κ stacking technique cannot be performed. We can still however estimate the Moho depth using equation (2) and by assuming a Vp/Vs ratio after several tests. Final Vp/Vs ratios chosen are the values providing arrival times for the PpPs and PpSS + PsPs reverberations that match the signal stacks, and which are coherent with the Vp/Vs of the surrounding stations (Table 2).

The results show large but systematic variations in Moho depth and V_p/V_s ratio throughout Haiti (Figures 5 and 6). The Moho depths can be split into three main groups shown in Fig. 5 and based on location and values (in latitudinal order GGIL, MEND, STRA, PIGN and CAYH in the northern part of Haiti; BOIS, DROU, GADC, THOM, CANG, DUFF, MIRB, ARCH, GIMB, BMCA, CBOQ, BMES, PAPH and LGNH in the central part, and MGOA, PETG, GRBO, JAKH, JACM, FERM, FURC, and MARG in the southern part of Haiti).

3.1- Moho depths in Haiti

Results from the H-K stacking method in the northern domain of Haiti are well resolved, with the stations STRA, PIGN and CAYH particularly well constrained. Results are also consistent across this domain giving Moho depths of between 20.1 km and 28.2 km (Table 1, Fig. 5).

Results in the southern part of Haiti are also well resolved, with the stations MGOA, JACM and PETG particularly well constrained. Results across the domain give Moho depths ranging between 16.1 km and 29.8 km, increasing in the center of the area at the stations GRBO, MARG, FURC (Table 1, Fig. 5). The stations MARG and JAKH have clearly identifiable Ps arrivals, but due to the lack of observable multiples the H-K stacking technique cannot be performed. However, we can still estimate the depth of the Moho by using equation (2) (Table 2, Fig. 5).

Our results show that the central part of Haiti exhibits greater Moho depths than in the northern and southern parts, with a Moho discontinuity always deeper than 30 km. The RFs of the stations located in this part of Haiti are more complicated due to intracrustal reverberations, probably due to sediments layers. However, consistent values are found for

the stations THOM, CANG, DUFF, BMCA, CBOQ, DROU, LGNH and BOIS with Moho depths ranging between 34.6 km and 45.2 km (Table 1, Fig. 5). We estimate the Moho depth for the stations GADC, MIRB, GIMB, BMES, ARCH, and PAPH by assuming a V_p/V_s ratio from the closest stations (Table 2, Fig. 5). Transect A in Fig. 7 shows the abrupt change in Moho depth of ~10 km between the southern part and the central part of Haiti.

3.2- V_p/V_s ratio

The average crustal V_p/V_s ratio is sensitive to the bulk composition of the crust (Zandt and Ammon, 1995; Chevrot and Van der Hilst, 2000, Stuart et al., 2006). Mineralogy is an important factor influencing V_p/V_s ratio (Christensen, 1996): For felsic quartz-rich rocks, such as granite, V_p/V_s ratio is ~1.71; for intermediate rocks, such as diorite, V_p/V_s ratio is typically ~1.78; and for mafic rocks, such as gabbro V_p/V_s ratio is ~1.87. Higher values of V_p/V_s ratio are associated with the presence of partial melting or fluids (Watanabe, 1993; Thompson et al., 2010).

The V_p/V_s ratio over Haiti ranges from 1.64 to 1.99 (Fig. 6) suggesting a wide range of mineral composition lateral variation. In particular, the stations MEND, CANG, BMCA and LGNH show particularly high V_p/V_s ratios of 1.94, 1.99, 1.98 and 1.96 respectively (Fig. 6). We observe that these stations are located on or near major tectonic features (Fig. 2), and therefore fluid migration along active faults could feasibly locally increase the V_p/V_s ratio. In order to interpret regional variations in crustal structure and since the interstation variability is relatively high, we calculate the mean V_p/V_s ratio for each of the 3 domains identified in Fig. 5, excluding the four very high values. Regional averages of V_p/V_s seem to provide more insights than single-station values, excepting for particularly high values that

could show fluid migration (Rossi et al., 2006). The mean V_p/V_s ratio is 1.75 ± 0.10 in the northern part of Haiti, 1.80 ± 0.05 in the central part, and 1.80 ± 0.03 in the southern part.

3.3- Shallow discontinuity

In addition to P to S Moho converted phases, one station, CAYH, exhibits another clear shallow discontinuity. The RF plot for station CAYH versus back-azimuth (top of Fig. 8) shows a shallow discontinuity in the first 2 s of the signal, prior to the P_s arrival from the Moho. Clear Moho P_s arrivals are obvious just after 2 s, and are easily identified in the upper RFs (group a, Fig. 8), coming from the south (back-azimuths between 176° and 178° , Fig. 8). The H-K stack method for this group gives a Moho depth of 22.1 km, which is consistent with the closest station PIGN (Fig. 5 and Table 1), and a V_p/V_s ratio of 1.64. We estimate the depth of the shallow discontinuity easily visible in the RFs of the group b assuming that the upper crust has an average V_p of 5.8 km/s and an average V_p/V_s ratio of 1.77. These average properties are derived from the P velocity model in Haiti of Douilly et al. (2013). This discontinuity is thus estimated at ~ 9.7 km depth.

4- Discussion

4.1- Identification of 3 distinct crustal domains

The depth distribution of the Moho in Haiti shows 3 distinct regions that we relate to 3 distinct geological domains (Fig. 9).

The Southern Peninsula of Hispaniola (Fig. 9) is known to be a part of the Caribbean LIP, and to be composed of Cretaceous tholeiitic material (Calmus, 1983; Bien-Aimé Momplaisir, 1987). In this part of the island, the Moho depths from our study vary from 16.1

km to 29.8 km along a W-E profile (Fig. 7E) and from 20.5 km to 27.5 km along a N-S profile (Fig. 7A), giving an average crustal thickness of 22 ± 5 km. The variability of the Moho depths we image in the Southern Peninsula may reflect the large differences in the LIP thickness identified in previous studies between the oceanic crust and the areas with more or less of underplated material (Diebold et al., 1981; Mauffret and Leroy, 1997; Leroy et al., 2000; Mauffret et al., 2001). In our results the mean V_p/V_s ratio is 1.80 ± 0.03 , consistent with previous studies (Douilly et al., 2013) and the geology of the Southern Peninsula (Calmus, 1983; Bien-Aimé Momplaisir, 1987). The northern limit of the LIP crustal domain is given by the Moho depth calculated at the station LGNH (34.6 km; Fig. 5) and could correspond to the trace of the EPGFZ (Fig. 2) in this area (south of LGNH), as suggested by previous studies (Saint-Fleur et al., 2015).

The northern part of Hispaniola (Fig 9) is known to be a part of the Great Arc of the Caribbean (Burke, 1988), an inactive intra-oceanic island arc formed in the Cretaceous at the boundary of the Pacific domain and the proto-Caribbean oceanic crust (Pindell et al., 2006). The Moho depth values we compute are ranging between 20.1 km and 28.2 km, providing an average crustal thickness of 23 ± 3 km in agreement with the values found in southern Cuba (approximately 20 km, Moreno et al., 2002; Gonzalez et al., 2012) and eastern Hispaniola (about 24 km, Nuñez et al., 2015). Furthermore, our results delineate the southwestern limit of the island arc domain. The shift of the Moho depths, from 22.1 km (CAYH) to 36.4 km (GADC), occurs across the trace of a major thrust evidenced by geological studies (Figs 5 and 9). Both this major thrust and the sharp offset of the Moho depths outline a distinct crustal domain to the west and south of this feature.

In addition to the southern and northern domains discussed above, we identify a

central domain distinct from both the LIP in the South and the island arc in the North (Fig. 9), which is characterized by a thicker crust ranging from 32.3 km to 45.4 km, with an average of 41 ± 4 km. The relatively deep Moho is well constrained by 13 stations. In addition, the preliminary results of a recent seismic refraction study by Nuñez et al. (2015) indicate that the thickness of the crust increases from eastern Hispaniola towards the west reaching a value of ~40 km at the boundary between Haiti and the Dominican Republic near our profile. The Cul-de-Sac sedimentary basin (Fig. 1) and the Trans-Haitian belt belong to the same thick domain, where the nature of the crust below these superficial structures has still to be determined. We further discuss the different hypothesis regarding the nature of the crust of this domain.

4.2 Vp/Vs ratio

We find very high Vp/Vs ratios, between 1.94 and 1.99 at four stations, located near major faults: LGNH near the strike-slip EPGFZ and MEND, CANG and BMCA near the main thrust faults (Fig. 2 and 6). High Vp/Vs ratios are often associated with partial melt, basaltic intrusions or the presence of fluids (Stuart et al, 2006; Hammond et al., 2011). We favor an explanation whereby the high Vp/Vs ratios associated with major crustal faults are related to the presence of fluids within the fault zones (Thurber et al., 2003; Zhao et al., 1996). Nevertheless, all the stations located near major faults do not present high Vp/Vs values (e.g. CAYH, fig. 2).

The mean Vp/Vs ratio (1.80 ± 0.03) found in the southern domain is in agreement with its late Cretaceous oceanic plateau origin (Christensen, 1996; Douilly et al., 2013). In the northern domain composed of a late Cretaceous-Eocene, intra-oceanic volcanic arc and

forearc-accretionary prism, the V_p/V_s ratio of 1.75 ± 0.10 is close to that expected for the average continental crust, 1.768 (Christensen, 1996). The central part of Haiti is characterized by V_p/V_s ratio of 1.80 ± 0.05 , indicating that mafic rocks may be part of the crustal lithology of this domain as in the southern domain.

4.3- Implications for the tectonic history of Hispaniola

We identify three distinct tectonic domains with different crustal thickness and bulk composition. Based on our computed Moho depths and V_p/V_s ratios, and with additional constraints from the major thrusts identified at the surface and the geodynamic reconstructions (Leroy et al., 2000; Pubellier et al., 2000; Calais et al., 2016), we propose a geological model along a N-S profile across Haiti (Fig. 10). This transect shows the LIP in the South, the Cretaceous volcanic arc in the North, and a central thicker domain.

The central domain is characterized by relatively thick crust of ~41 km, between two domains with significantly thinner crusts of 20-30 km. Surprisingly, the thick crust of the central domain is not associated with major relief (~1 km). Based on field studies and geodynamic reconstructions proposed for this area (e.g. Stéphan et al., 1990; Meschede and Frisch, 1998; Mann et al 1991; Calais et al., 2016), the nature of the crust of the central domain is likely to be island arc and fore-arc type, considered as either back-arc (Mann and Lawrence, 1991) or a Cretaceous-Eocene remnant arc (Heubeck et al., 1991). Other stratigraphic and tectonic studies (e.g. Pubellier et al., 2000) suggest that the fold-and-thrust belt of this crustal domain may be rooted on the continuation of the rifted crust of the eastern Cayman continental passive margin (Fig. 9). However, V_p/V_s ratio of 1.80, typical for mafic rocks (average of values for this domain; Tables 1 & 2), may indicate that

the central crustal domain could not be composed only of island arc crust, and may imply the presence of dense material. This dense material could be either related to Pliocene-Quaternary mafic center of volcanism coming from an ancient continental mantle lithosphere fragment (Kamenov et al., 2011) or associated to LIP material fragment coming from the south and both trapped early in central Haiti. Geochemical studies of the LIP south of Hispaniola from submersible dive samples (Revillon et al., 2000; Mauffret et al., 2001) show indeed that the lower part of the LIP is made of ultra-mafic rocks and geophysical studies (Mauffret and Leroy, 1997; 1999) evidence a high velocity layer composed of dense material (velocity 7.5 to 7.9 km/s; density of 3 to 3.1; Herzberg et al., 1983). The mafic Quaternary rocks are also MgO-rich (Kamenov et al 2011) and give high V_p/V_s ratio.

On the transect in Fig. 10 we also draw the intracrustal discontinuity determined from our receiver functions at the station CAYH. The estimated depth of this intracrustal discontinuity (~10 km) may provide information on the possible depth for the Late-Cretaceous to Pleistocene flysch of the Peralta belt (Witschard and Dolan, 1990; Dolan et al., 1991), which outcrops at the front of the Cretaceous volcanic arc (Fig. 10). At the surface, three major thrusts are geologically mapped (the Montagnes Noires, the Chaîne des Matheux and the Gonâve Island; Fig. 1 and 2; Mann et al., 1995; Pubellier et al., 2000; Hernaiz Huerta et al., 2007). In our transect, we propose to extend these thrusts to greater depth along three large crustal slices (Fig. 10). The three major thrusts of the Trans-Haitian belt show small cumulative throws at outcrop and little evidence of strong compressional tectonics (Mann et al 1995; Pubellier et al 2000, Corbeau et al., 2016b). SW-vergent fold and thrust belt related to the Hispaniola restraining bend and back arc basin collapse

(Heubeck et al., 1991) could have contributed to the overall thickening.

5- Conclusions

A receiver function study of 23 stations is used to determine the crustal structure across Haiti, in order to place better constraints on the history and geometry of late Cretaceous to recent terrane accretion. In the northern part of Haiti, consisting of the Cretaceous volcanic island arc and forearc of the Greater Antilles, the Moho discontinuity is imaged at a depth of ~23 km with V_p/V_s ratio of 1.75. In the southern part of Haiti, made of the folded and thrust Cretaceous oceanic Large Igneous Province (LIP), the Moho depth is imaged at ~22 km, and V_p/V_s ratio is 1.80. In the central part of Haiti, encompassing the Haitian fold-and-thrust belt, the Moho discontinuity is deeper with a mean depth of ~41 km. We propose that this unexpected thick central domain with V_p/V_s ratio of 1.80 corresponds to a distinct crustal domain. This domain is likely to consist of island-arc type crust related to the Great Arc of the Caribbean, or the proximal part of the Cayman Trough eastern passive margin, associated with mafic material coming from fragments of lithospheric mantle from the north or mafic LIP material from the south.

Acknowledgements

The seismological equipment was loaned from the SEIS-UK equipment pool and from the ITeP facilities. The facilities of SEIS-UK are supported by the Natural Environment Research Council (NERC) under Agreement R8/H10/64. DK time on the research is funded

by NERC grant NE/L013932/1. This work could not have been done without Gérard Laborde and DIGICEL, and without our collaboration with the Bureau des Mines et de l'Energie and the Université d'Etat d'Haiti in Haiti. The paper benefits from fruitful discussions with Nicolas Bellahsen and Claudio Rosenberg. We thank the Editor, the reviewer David Schlaphorst and an anonymous reviewer for their constructive comments that helped improved the manuscript.

References

- Abers, G. A., X. Hu and L. R. Sykes, 1995. Source scaling of earthquakes in the Shumagin region, Alaska : time-domain inversions of regional waveforms. *Geophysical Journal International*, vol. 123, no 1, p. 41-58.
- Ammon, C. J., 1991. The isolation of receiver effects from teleseismic P waveforms. *Bull. Seismol. Soc. Am.*, vol. 81, no 6, p. 2504-2510.
- Bien-Aimé Momplaisir, R., 1986. Contribution à l'étude géologique de la partie orientale du Massif de la Hotte (Presqu'île du Sud d'Haïti) : Synthèse structurale des marges de la presqu'île à partir de données sismiques, thèse de doctorat, Ph. D. thesis, 210 pp., Univ. Pierre-et-Marie-Curie (Paris VI), Paris.
- Boisson, D., 1987. Etude géologique du massif du nord d'Haïti (Hispaniola-grandes Antilles), thèse de doctorat, Ph. D. thesis, Univ. Pierre et Marie Curie (Paris VI), Paris.
- Bowin, C.O, 1976. Caribbean Gravity Field and Plate Tectonics, Geological Society of America Special Paper, vol. 169.
- Burke, K., 1988. Tectonic evolution of the Caribbean, Annual Review of Earth and Planetary Sciences, vol. 16, p. 201-230.
- Calais, E. and B. Mercier de Lépinay, 1991. From transtension to transpression along the northern Caribbean plate boundary off Cuba : Implications for the recent motion of the Caribbean plate, *Tectonophysics*, vol. 186, no 3, p. 329-350.
- Calais, E., S. Smithe, B. Mercier de Lépinay and C. Prépetit, 2016. Plate boundary segmentation in the northeastern Caribbean from geodetic measurements and Neogene geological observations, *Comptes Rendus Geoscience*, 348(1), 42-51.
- Calmus, T., 1983. Contribution à l'étude géologique du massif de Macaya(sud-ouest d'Haïti, Grandes Antilles) : sa place dans l'évolution de l'orogène Nord-Caraïbe, thèse de doctorat, Univ. Paris VI, Paris.
- Chevrot, S. and R. D. van der Hilst, 2000. The poisson ratio of the Australian crust : geological and geophysical implications, *Earth and Planetary Science Letters*, vol. 183, no 1, p. 121-132.
- Christensen, N. I., 1996. Poisson's ratio and crustal seismology, *Journal of Geophysical Research: Solid Earth (1978-2012)*, vol. 101, no B2, p. 3139-3156.
- Corbeau, J., F. Rolandone, S. Leroy, B. Mercier de Lépinay, B. Meyer, N. Ellouz-Zimmermann and R. Momplaisir, 2016a. The Northern Caribbean plate boundary in the Jamaica Passage: structure and seismic stratigraphy, *Tectonophysics*, doi:10.1016/j.tecto.2016.03.022.

Corbeau, J., F. Rolandone, S. Leroy, B. Meyer, B. Mercier de Lépinay, N. Ellouz-Zimmermann and R. Momplaisir, 2016b. How transpressive is the Northern Caribbean plate boundary ?, *Tectonics*, doi:10.1002/2015TC003996.

Cruz-Orosa, I., F. Sàbat, E. Ramos, L. Rivero and Y. M. Vázquez-Taset, 2012. Structural evolution of the La Trocha fault zone : Oblique collision and strike-slip basins in the Cuban Orogen, *Tectonics*, vol. 31, no 5.

Diebold, J., P. Stoffa, P. Buhl and M. Truchan, 1981. Venezuela Basin crustal structure, *Journal of Geophysical Research : Solid Earth (1978-2012)*, vol. 86, no B9, p. 7901-7923.

Dolan, J., P. Mann, R. de Zoeten, C. Heubeck, J. Shiroma and S. Monechi, 1991. Sedimentologic, stratigraphic, and tectonic synthesis of Eocene-Miocene sedimentary basins, Hispaniola and Puerto Rico, *Geological Society of America Special Papers*, vol. 262, p.217-264.

Douilly, R., J. S. Haase, W. L. Ellsworth, M.-P. Bouin, E. Calais, S. J. Symithe, J. G. Armbruster, B. Mercier de Lépinay, A. Deschamps, S.-L. Mildor and collab., 2013. Crustal Structure and Fault Geometry of the 2010 Haiti Earthquake from Temporary Seismometer Deployments, *Bulletin of the Seismological Society of America*, vol. 103, no 4, p. 2305-2325.

Duncan, R. and R. Hargraves, 1984. Plate tectonic evolution of the Caribbean region in the mantle reference frame, *Geological Society of America Memoirs*, vol. 162, p. 81-94.

Escuder Viruete, J., Contreras F., Stein, G., Urien, P., Joubert, M., Ullrich, T.D., Mortensen, J., Perez Estaun, A., 2006. Transpression and strike-slip partitioning in the Caribbean island arc: fabric development, kinematics and Ar-Ar ages of syntectonic emplacement of the Loma de Cabrera batholith, Dominican Republic, *Journal of Structural Geology*, 28, 1496-1519.

González, O., B. Moreno, F. Romanelli and G. F. Panza, 2012. Lithospheric structure below seismic stations in Cuba from the joint inversion of Rayleigh surface waves dispersion and receiver functions, *Geophysical Journal International*, vol. 189, p. 1047-1059.

Granja-Bruña, J., A. Carbó-Gorosabel, P. L. Estrada, A. Muñoz-Martín, U. ten Brink, M. G. Ballesteros, M. Druet and A. Pazos, 2014. Morphostructure at the junction between the Beata ridge and the Greater Antilles island arc (offshore Hispaniola southern slope), *Tectonophysics*, vol. 618, p. 138-163.

Gurrola, H., G. E. Baker and J. B. Minster, 1995. Simultaneous time-domain deconvolution with application to the computation of receiver functions, *Geophysical Journal International*, vol. 120, no 3, p. 537-543.

Hammond, J., J.-M. Kendall, G. Stuart, D. Keir, C. Ebinger, A. Ayele and M. Belachew, 2011. The nature of the crust beneath the Afar triple junction : Evidence from receiver functions, *Geochemistry, Geophysics, Geosystems*, vol. 12, no 12.

Hammond, J., J.-M. Kendall, G. Stuart, C. Ebinger, I. Bastow, D. Keir, A. Ayele, M. Belachew, B. Goitom, G. Ogubazghi and collab., 2013. Mantle upwelling and initiation of rift segmentation beneath the Afar Depression, *Geology*, vol. 41, no 6, p. 635-638.

Hastie, A. R., S. F. Mitchell, P. J. Treloar, A. C. Kerr, I. Neill and D. N. Barfod, 2013. Geochemical components in a Cretaceous island arc : The Th/La-(Ce/Ce*)Nd diagram and implications for subduction initiation in the inter-American region, *Lithos*, vol. 162, p. 57-69.

Helfrich, G., 2006. Extended-time multitaper frequency domain cross-correlation receiver-function estimation, *Bulletin of the Seismological Society of America*, vol. 96, no 1, p. 344-347.

Helfrich, G., B. Faria, J. F. Fonseca, A. Lodge and S. Kaneshima, 2010. Transition zone structure under a stationary hot spot: Cape Verde, *Earth and Planetary Science Letters*, vol. 289, no 1, p. 156-161.

Hernaiz Huerta, P.P., Diaz de Neira, J.A., Garcia Senz, J. Deschamps, I., Genna, A., Nicole, N., Lopera, E., Escuder Virruete, J., Ardevol Oro, Ll., Perez Estaun, A., 2007. La estructura del suroeste de la Republica Dominicana: un ejemplo de deformacion en regimen transpresivo, *Boletin Geologico y Minero*, 118(2) : 337-358.

Herzberg, C., W. Fyfe and M. Carr, 1983. Density constraints on the formation of the continental Moho and crust, *Contributions to Mineralogy and Petrology*, vol. 84, no 1, p. 1-5.

Heubeck, C., P. Mann, J. Dolan and S. Monechi, 1991. Diachronous uplift and recycling of sedimentary basins during cenozoic tectonic transpression, northeastern caribbean plate margin, *Sedimentary geology*, vol. 70, no 1, p. 1-32.

Iturralde-Vinent, M. A., 2006. Meso-Cenozoic Caribbean paleogeography : implications for the historical biogeography of the region, *International Geology Review*, vol. 48, no 9, p. 791-827.

Kamenov, G. D. et al., 2011. Ancient lithospheric source for Quaternary lavas in Hispaniola. *Nature Geoscience*, vol. 4, p. 554-557.

Kennet, B., 1991. IASPEI 1991 seismological tables, *Terra Nova*, vol. 3, no 2, p. 122-122.

Langston, C. A., 1979. Structure under Mount Rainier, Washington, inferred from teleseismic body waves, *Journal of Geophysical Research : Solid Earth* (1978-2012), vol. 84, no B9, p. 4749-4762.

Leroy, S., B. Mercier de Lépinay, A. Mauffret and M. Pubellier, 1996. Structural and tectonic evolution of the eastern Cayman Trough (Caribbean Sea) from seismic reflection data, *AAPG bulletin*, vol. 80, no 2, p. 222-247.

Leroy, S., A. Mauffret, P. Patriat and B. Mercier de Lépinay, 2000. An alternative interpretation of the Cayman trough evolution from a reidentification of magnetic anomalies, *Geophysical Journal International*, vol. 141, no 3, p. 539-557.

Leroy, S., N. Ellouz-Zimmermann, J. Corbeau, F. Rolandone, B. Mercier de Lépinay, B. Meyer, R. Momplaisir, J. L. Granja Bruna, A. Battani, C. Baurion, E. Burov and collab., 2015. Segmentation and kinematics of the North America-Caribbean plate boundary offshore Hispaniola, *Terra Nova*.

Ligorria, J. P. and C. J. Ammon, 1999. Iterative deconvolution and receiver-function estimation, *Bulletin of the seismological Society of America*, vol. 89, no 5, p. 1395-1400.

Lodge, A., S. Nippress, A. Rietbrock, A. García-Yeguas and J. Ibáñez, 2012. Evidence for magmatic underplating and partial melt beneath the canary islands derived using teleseismic receiver functions, *Physics of the Earth and Planetary Interiors*, vol. 212, p. 44-54.

Mauffret, A. and S. Leroy, 1997. Seismic stratigraphy and structure of the Caribbean igneous province, *Tectonophysics*, vol. 283, no 1, p. 61-104.

Mauffret, A., S. Leroy, É. d'Acremont, A. Maillard, B. Mercier de Lépinay, A. T. Dos Reis, N. Miller, A. Nercessian, R. Pérez-Vega and D. Perez, 2001. Une coupe de la province volcanique Caraïbe : premiers résultats de la campagne sismique Casis 2, *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science*, vol. 333, no 10, p. 659-667.

Mann, P., G. Draper and J. F. Lewis, 1991. An overview of the geologic and tectonic development of Hispaniola, *Geological Society of America Special Papers*, vol. 262, p. 1-28.

Mann, P. and S. Lawrence, 1991. Petroleum potential of southern Hispaniola, *Journal of Petroleum Geology*, vol. 14, no 2, p. 291-308.

Mann, P., F. Taylor, R. L. Edwards and T.-L. Ku, 1995. Actively evolving microplate formation by oblique collision and sideways motion along strike-slip faults : An example from the northeastern Caribbean plate margin, *Tectonophysics*, vol. 246, no 1, p. 1-69.

Martinez-Arevalo, C., F. de Lis Mancilla, G. Helffrich and A. Garcia, 2013. Seismic evidence of a regional sublithospheric low velocity layer beneath the Canary islands, *Tectonophysics*, vol. 608, p. 586-599.

- McNamara, D., M. Meremonte, J. Maharrey, S.-L. Mildore, J. Altidore, D. Anglade, S. Hough, D. Given, H. Benz, L. Gee and collab., 2012. Frequency-dependent seismic attenuation within the Hispaniola Island region of the Caribbean Sea, *Bulletin of the Seismological Society of America*, vol. 102, no 2, p. 773-782.
- Meschede, M. and W. Frisch, 1998. A plate-tectonic model for the Mesozoic and Early Cenozoic history of the Caribbean plate, *Tectonophysics*, vol. 296, no 3, p. 269-291.
- Moreno, B., M. Grandison and K. Atakan, 2002. Crustal velocity model along the southern Cuban margin : implications for the tectonic regime at an active plate boundary, *Geophysical Journal International*, vol. 151, no 2, p. 632-645.
- Nuñez, D., D. Cordoba, F. J. Nuñez-Cornu and M. O. Cotilla, 2015. Lithosphere structure from Cordillera Central to Cordillera Oriental (Dominican Republic), *EGU General Assembly Conference Abstracts*, vol. 17, p. 14467.
- Owens, T., S. Taylor and G. Zandt, 1983. Isolation and enhancement of the response of local seismic structure from teleseismic P-waveforms, *cahier de recherche*, Lawrence Livermore National Lab., CA (USA).
- Park, J. and V. Levin, 2000. Receiver functions from multiple-taper spectral correlation estimates, *Bulletin of the Seismological Society of America*, vol. 90, no 6, p. 1507-1520.
- Pindell, J., L. Kennan, K. P. Stanek, W. Maresch and G. Draper, 2006. Foundations of Gulf of Mexico and Caribbean evolution: eight controversies resolved, *Geologica Acta*, vol. 4, no 1-2, p. 303.
- Pindell, J., W. V. Maresch, U. Martens and K. Stanek, 2012. The Greater Antillean Arc : Early Cretaceous origin and proposed relationship to Central American subduction mélanges : implications for models of Caribbean evolution, *International Geology Review*, vol. 54, no 2, p. 131-143.
- Pubellier, M., A. Mauffret, S. Leroy, J. M. Vila and H. Amilcar. 2000, «Plate boundary readjustment in oblique convergence : Example of the Neogene of Hispaniola, Greater Antilles», *Tectonics*, vol. 19, no 4, p. 630-648.
- Revillon, S., E. Hallot, N. Arndt, C. Chauvel and R. Duncan, 2000. A complex history for the Caribbean Plateau: petrology, geochemistry, and geochronology of the Beata Ridge, South Hispaniola, *The Journal of Geology*, vol. 108, no 6, p. 641-661.
- Rossi, G., Abers, G. A., Rondenay, S., & Christensen, D. H., 2006. Unusual mantle Poisson's ratio, subduction, and crustal structure in central Alaska, *Journal of Geophysical Research: Solid Earth*, vol. 111, no B9.

Saint Fleur, N., N. Feuillet, R. Grandin, E. Jacques, J. Weil-Accardo and Y. Klinger, 2015. Seismotectonics of southern Haiti: A new faulting model for the 12 January 2010 M7 earthquake, *Geophysical Research Letters*.

Symithe, S. and E. Calais, 2016. Present-day shortening in Southern Haiti from GPS measurements and implications for seismic hazard, *Tectonophysics*, 679, 117-124.

Symithe, S., E. Calais, J. B. Chabaliere, R. Robertson and M. Higgins, 2015. Current block motions and strain accumulation on active faults in the Caribbean, *Journal of Geophysical Research: Solid Earth*, 120(5), 3748-3774.

Stéphan, J.-F., B. Mercier de Lépinay, E. Calais and M. Tardy, 1990. Paleogeodynamic maps of the Caribbean: 14 steps from Lias to Present, *Bull. Soc. géol. Ft*, p. 0-6.

Stuart, G., I. Bastow and C. Ebinger, 2006. Crustal structure of the northern Main Ethiopian Rift from receiver function studies, *Geological Society*, London, Special Publications, vol. 259, no 1, p. 253-267.

Thompson, D., I. Bastow, G. Helffrich, J. Kendall, J. Wookey, D. Snyder, and D. Eaton, 2010. Precambrian crustal evolution: Seismic constraints from the Canadian Shield, *Earth Planet. Sci. Lett.*, 297, 655-666, doi:10.1016/j.epsl.2010.07.021.

Thurber, C., S. Roecker, K. Roberts, M. Gold, L. Powell and K. Rittger, 2003. Earthquake locations and three-dimensional fault zone structure along the creeping section of the San Andreas fault near Parkfield, CA : Preparing for SAFOD, *Geophysical Research Letters*, vol. 30, no 3.

VanDerLelij, R., 2013. Reconstructing north-western Gondwana with implications for the evolution of the Iapetus and Rheic Oceans: a geochronological, thermochronological and geochemical study, thèse de doctorat, University of Geneva.

Watanabe, T., 1993. Effects of water and melt on seismic velocities and their application to characterization of seismic reflectors, *Geophysical Research Letters*, vol. 20, no 24, p. 2933-2936.

Witschard, M. and J. F. Dolan, 1990. Contrasting structural styles in siliciclastic and carbonate rocks of an offscraped sequence: The Peralta accretionary prism, Hispaniola, *Geological Society of America Bulletin*, vol. 102, no 6, p. 792-806.

Zandt, G. and C. J. Ammon, 1995. Continental crust composition constrained by measurements of crustal Poisson's ratio, *Nature*, vol. 374, no 6518, p. 152-154.

Zhao, D., H. Kanamori, H. Negishi and D. Wiens, 1996. Tomography of the source area of the 1995 Kobe earthquake : evidence for fluids at the hypocenter ?, *Science*, vol. 274, no 5294, p. 1891-1894.

Zhu, L. and H. Kanamori, 2000. Moho depth variation in southern California from teleseismic receiver functions, *Journal of Geophysical Research B*, vol. 105, no B2, p. 2969-2980.

Figure 1: Tectonic map of the Northern Caribbean plate boundary. Faults are from previous studies (Calais and Mercier de Lépinay, 1991; Mann et al., 1995; Leroy et al., 1996; 2015; Granja Bruna et al., 2014). NHF: North-Hispaniola Fault; SOFZ: Septentrional-Oriente Fault Zone; EPGFZ: Enriquillo-Plantain-Garden Fault Zone.

Figure 2: Locations of the seismic stations used in this receiver function study in Haiti, superimposed on the topographic map and with the major tectonic features. Red circles are the stations of the Trans-Haiti network and blue circles are the 3 permanent stations of the Canadian National Network. Faults are from Pubellier et al. (2000). SOFZ: Septentrional-Oriente Fault Zone; EPGFZ: Enriquillo-Plantain-Garden Fault Zone. The red line indicates the localization of the transect proposed Fig. 10.

Figure 3: Global distribution of the earthquakes used when computing Receiver Functions. They are plotted with an azimuthal equidistant map projection around our network.

Figure 4: Examples of Moho depth (H) versus V_p/V_s plots (left) from the method of Zhu and Kanamori (2000) and receiver functions associated (right) for one station of each domain. On each receiver function the arrival time of the Moho phase P_s (t_1) and the reverberations P_pP_s (t_2) and $P_sP_s+P_pS_s$ (t_3) are marked based on the results of the H-K stack (Table 1) and the equations (2) to (4). The receiver functions are plotted as a function of the back-azimuth (number at the right of each trace). See Figs. 2 and 5 for the location of the stations and the different domains.

Figure 5: Variations in Moho depth (Tables 1 and 2) across Haiti determined from receiver function analysis. The blue lines A, B, C, D, and E show the orientation of transects in Figure 7. The black dashed lines delimit the boundary between 3 distinct domains of Moho depth values, numbered 1, 2 and 3 (see text for explanation).

Figure 6: Variations in V_p/V_s ratio (Tables 1 and 2) across Haiti determined from receiver function analysis. The orange lines A, B, C, D, and E show the orientation of transects in Figure 7.

Figure 7: Variations in elevation, Moho depth, and V_p/V_s ratio across Haiti. The elevation exaggeration is $\times 10$. (A) SW-NW across the central and southern part of Haiti, (B) and (C) SE-NW and SW-NE across the northern part of Haiti, (D) S-N across the West of the central and southern part, and (E) W-E across the southern part of Haiti. The orientations of the transects are shown on Figures 5 and 6.

Figure 8: Estimation of the depth of an intracrustal layer for the station CAYH. The Moho depth is estimated at 22.1 km with P_s arrival and P_pP_s and $P_sP_s+P_pS_s$ reverberations, using the H-K stack method and the RFs coming from the South (back-azimuth of about 170° , group a). A reverberation is imaged before the P_s arrival on RFs coming from the North-East (back-azimuth of about 55° , group b), and the corresponding estimated depth with P_s pickeds method ($V_p=5.8$ km/s and $K=1.77$) is 9.7 km.

Figure 9: Location of the proposed different crustal domains we identified in Hispaniola. (1) the Caribbean Large Igneous Province, (2) the Cretaceous island arc and (3) the fold and thrust belt. C: Cuba; J: Jamaica; H: Haiti; D.R: Dominican Republic. NHF: North Haitian Fault; SOFZ: Septentrional-Oriente Fault Zone; THB: Trans-Haitian Belt; EPGFZ: Enriquillo-Plantain-Garden Fault Zone; CT: Cayman Trough; LNR: Lower Nicaraguan Rise. East CMS: eastern Cayman Trough margin system; LIP: Large Igneous Province. The black line indicates the localization of the transect shown in Fig. 10. The thicknesses of the LIP at sea are from Mauffret and Leroy (1997). The boundary between the East CMS and the LIP at sea is from Corbeau et al. (2016a).

Figure 10: Summary conceptual geological transect across Haiti based on the Moho and intracrustal discontinuity depths. See Figs 2 and 9 for location of the profile. The seismologic stations are indicated. EPGFZ: Enriquillo-Plantain-Garden fault zone; SOFZ: Septentrional-Oriente fault zone. The log south of the transect is a model of the LIP crustal thickness based on a compilation of seismic refraction results: the layer 2V (original oceanic crust) is sandwiched between an upper basaltic layer (in black) and the layer 3V (original crust underplated); R is the top of a high velocity layer (Mauffret and Leroy, 1997).

Table 1: Moho depth (H) and Vp/Vs ratio (κ) estimated by the H-K stacking method.

Stations	N	H(km)	δH (km)	K	δK
BMCA (line A)	13	44.3	0.1	1.98	0.01
BOIS (line D)	2	41.5	0.1	1.75	0.01
CANG (line A)	4	41.5	0.4	1.99	0.01
CAYH (line B)	5	22.1	0.4	1.64	0.02
CBOQ (line A)	7	39.5	0.7	1.82	0.02
DROU (line D)	7	47.2	0.4	1.85	0.02
DUFF (line A)	3	45.2	0.4	1.73	0.01
FERM (line A)	2	20.5	0.4	1.79	0.03
FURC (line A)	3	26.6	0.3	1.83	0.02
GGIL (line C)	3	20.1	0.7	1.81	0.04
GRBO (line E)	3	29.8	0.3	1.83	0.02
JACM (line D)	3	20.9	0.4	1.80	0.03
LGNH (line D)	8	34.6	0.4	1.96	0.02
MEND (line C)	6	22.5	0.4	1.94	0.02
MGOA (line E)	10	16.1	0.5	1.74	0.04
PETG (line E)	3	19.7	0.4	1.83	0.03
PIGN (line B)	4	22.1	0.5	1.86	0.04
STRA (line B)	4	28.2	0.4	1.67	0.03
THOM (line A)	5	37.9	0.5	1.87	0.02

Table 2: Moho depth (H) estimated by Ps picked method.

Stations	N	H(km)	K
ARCH (line D)	5	32.3	1.80
BMES (line A)	5	41.9	1.80
GADC (line A)	4	36.4	1.85
GIMB (line A)	9	44.5	1.75
JAKH (line D)	9	19.0	1.80
MARG (line A)	3	25.5	1.80
MIRB (line A)	3	45.4	1.74
PAPH (line A)	9	42.3	1.85




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10

Highlights:

- We explore the crustal structure of Haiti along the northern Caribbean plate boundary
- We use receiver-function technique to determine crustal thickness and bulk composition
- Three distinct crustal domains are imaged on a 200-km-long transect across Haiti
- The central domain is surprisingly thick and may contain fragments of dense material