



UNIVERSITY OF LEEDS

This is a repository copy of *Patient characteristics predicting failure to receive indicated care for type 2 diabetes*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/116473/>

Version: Accepted Version

Article:

Mounce, LTA, Steel, N, Hardcastle, AC et al. (6 more authors) (2014) Patient characteristics predicting failure to receive indicated care for type 2 diabetes. *Diabetes Research and Clinical Practice*, 107 (2). pp. 247-258. ISSN 0168-8227

<https://doi.org/10.1016/j.diabres.2014.11.009>

© 2014, Elsevier Ireland Ltd. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Figure 1. Deriving the numerators and denominators of the glyated haemoglobin (HbA1c), proteinuria and foot examination quality indicators from ELSA questions.

