



UNIVERSITY OF LEEDS

This is a repository copy of *Defects in the Cell Signaling Mediator  $\beta$ -Catenin Cause the Retinal Vascular Condition FEVR*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/117491/>

Version: Supplemental Material

---

**Article:**

Panagiotou, ES, Sanjurjo Soriano, C, Poulter, JA [orcid.org/0000-0003-2048-5693](https://orcid.org/0000-0003-2048-5693) et al. (15 more authors) (2017) Defects in the Cell Signaling Mediator  $\beta$ -Catenin Cause the Retinal Vascular Condition FEVR. *The American Journal of Human Genetics*, 100 (6). pp. 960-968. ISSN 0002-9297

<https://doi.org/10.1016/j.ajhg.2017.05.001>

---

© 2017, American Society of Human Genetics. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

**Reuse**


Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



**Figure S1. Clinical appearance of the proband from family F1321 aged one month.** (A) Ultrasound biomicroscopy photograph of the left eye showing vitreous hemorrhage. (B) Ultrasound biomicroscopy photograph of the right eye showing a funnel-shaped retinal detachment and vitreous hemorrhage. (C) Computerized tomography scan of the brain showing hyper-dense left globe content and no acute intracranial hemorrhage.

		p.Arg710Cys			
		↓			
Human	700	----	IGAQGEPLG-YRQDDPSYRSFHSG----	722	
Chimp	700	----	IGAQGEPLG-YRQDDPSYRSFHSG----	722	
Monkey	700	----	IGAQGEPLG-YRQDDPSYRSFHSG----	722	
Dog	700	----	IGAQGEPLG-YRQDDPSYRSFHSG----	722	
Cow	700	----	IGAQGEPLG-YRQDDPSYRSFHSG----	722	
Mouse	700	----	IGAQGEALG-YRQDDPSYRSFHSG----	722	
Rat	700	----	IGAQGEALG-YRQDDPSYRSFHSG----	722	
Chicken	700	----	IGAQGEPLG-YRPDDPSYRSFHSG----	722	
Zebrafish	699	----	IGAQGEPLG-YRQDDPSYRSFHSG----	721	
Fruit fly	714		LQDMLGPPEEAYEGLYGQGPVHSSHGGRF	744	
Mosquito	712		LQDILSPEQAYEGLYGQGPASVHSSHGGRF	742	
Frog	700	----	IGAQGEPLG-YRQDDPSYRSFHAP----	722	

**Figure S2. Protein sequence alignment of human  $\beta$ -catenin with homologues.** Multiple protein

alignment was calculated using HomoloGene (<http://www.ncbi.nlm.nih.gov/homologene>) [Edgar, R.C.

(2004) Nucleic Acids Res. 32, 1792-7]. Thirty amino acid residues surrounding the p.Arg710Cys variant are


shown. Conserved amino acid residues are shaded. Accession numbers: Human NP\_001091679.1;

Chimpanzee XP\_001138023.1; Monkey NP\_001244847.1; Dog NP\_001131124.1; Cow NP\_001069609.1;


Mouse NP\_001159374.1; Rat NP\_445809.2; Chicken NP\_990412.1; Zebrafish NP\_571134.2; Fruit fly

(armadillo) NP\_996328.1; Mosquito (Arm\_Anoga) XP\_309245.5; Frog NP\_001016958.1.


A



B



**Figure S3. Published intragenic heterozygous *CTNNB1* mutations causing syndromic intellectual disability (ID) and autism spectrum disorder (ASD).** Schematic representation of the *CTNNB1* gene and protein showing the location of *CTNNB1* mutations identified in individuals with ID syndrome (blue) and ASD (green) (de Ligt et al. 2012; O’Roak et al. 2012; Tucci et al. 2014; Kuechler et al. 2015; Kharbanda et al. 2017) alongside the mutations identified in the present study (red). The mutation in purple was identified in a child with ID syndrome but with an eye phenotype consistent with a diagnosis of FEVR (Dixon et al 2016). The amino-terminal domain (NTD) spans amino acids 1-137, 12 armadillo repeats (ARM) span amino acids 138-664, the carboxy-terminal domain (CTD) spans 667-781 and the Helix-C domain spans amino acids 667-683. All mutations are written using the standard recommended guidelines (<http://varnomen.hgvs.org>). The frameshift mutations reported in Kharbanda et al. 2017 (c.1038\_1044delGCTATCTinsGCT; c.1038\_1044delGCTATCTinsGCT; c.799\_809delGAAGGAGCTAAinsGAA) have therefore been changed to fit these standards.



**Figure S4. Western blot analysis and quantitative densitometry to confirm the expression of the wildtype and mutant  $\beta$ -catenin constructs.** (A) Western blot of *CTNNB1* expression constructs. STF cells were transiently transfected with expression vectors for wild-type (WT) or mutant  $\beta$ -catenin (Arg710Cys or His720\*) or empty vector (pDEST40). Mock transfection (Mock) and untreated STF cells were additional controls. Forty-eight hours after transfection, whole cell lysates were prepared and subjected to western blot analysis with anti- $\beta$ -catenin polyclonal antibody (#9562 Cell Signalling Technology). Anti- $\beta$ -actin monoclonal antibody (A5441 Sigma Aldrich) was used as the loading control. (B) Western band intensities were measured using Image Lab 5.2.1 (Bio-Rad) and quantified relative to the cells transfected with empty vector. Results are shown as ratios to the loading control. Note that this experiment was only performed once and the samples are not normalized for transfection efficiency but to total number of cells.

**Table S1. *CTNNB1* primers for PCR/sequencing.**

<b>Exon</b>	<b>Forward primer (5'-3')</b>	<b>Reverse primer (5'-3')</b>	<b>Size (bp)</b>
2	CAGGTATCCCAGTGACTTAGGAG	GCAGAAAATGGAGCAAAGG	268
3&4	TGCTTTTCTTGGCTGTCTTTC	AGTTTTCAAGTACTGGTATTGGGT	913
5&6	TGAAGTAAATGCTCAAGGGGA	CCACAACCCATTCATGGAAA	758
7	GCAAGCTGGCTGAAATTCTT	TCAGTAGTTAAAGTTCTACCACCTTTT	351
8&9	CAGAAGGACACCTCCTAAGGC	CCCTATCGCAGCCATACTTC	761
10	GCGATAGGGGTAAGATTCTGAAAT	CTCTTCAGGAAGACGGATGG	411
11	TTACGGGGAACTTCGGGTAT	TCATAAAATTAAATGTTGGTAACCC	345
12&13	TGTGAATGCCTCTTGCACTC	CAATGCAAATGAATGTGTACTAAGTG	569
14	TTGTTCCTTTTGTAATCTGAAAGTATG	CTGCCAACACTGGTTTCCC	267
15	TTTGGATGCCCTAACCTCAG	TAGCCTAAACCACTCCCACC	419

<b>Mutation</b>	<b>PolyPhen2</b>	<b>MutationTaster</b>	<b>CADD*</b>
c.1434_1435insC p.Glu479Argfs*18	N/A	Disease causing (prediction probability 1)	Score 35
c.2128C>T p.Arg710Cys	Possibly damaging (score 0.54)	Disease causing (prediction probability 0.999)	Score 25.4
c.2142_2157 dup p.His720*	N/A	Disease causing (prediction probability 1)	Score 35

**Table S2. Summary of bioinformatics analyses undertaken to predict the pathogenic nature of the *CTNNB1* mutations.** URLs: PolyPhen2, <http://genetics.bwh.harvard.edu/pph2/> [Adzhubei, I.A. et al. (2010). Nat. Methods 7, 248-9]; Mutationtaster, <http://www.mutationtaster.org/> [Schwarz, J.M. et al. (2010). Nat. Methods 7, 575-6]; Scaled CADD (Combined Annotation Dependent Depletion) scores generated using version 1.3 <http://cadd.gs.washington.edu> [Kircher, M. et al. (2014). Nat. Genet. 46, 310-5]. \*Scaled CADD scores of 20 means that the variant is amongst the top 1% of deleterious variants in the human genome and a score of 30 means that the variant is in the top 0.1%.