



This is a repository copy of *Measurement of the B⁰ and B⁻ meson lifetimes*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/120152/>

Version: Submitted Version

Article:

BUSKULIC, D., DECAMP, D., GOY, C. et al. (403 more authors) (1993) Measurement of the B⁰ and B⁻ meson lifetimes. *Physics Letters B*, 307 (1-2). pp. 194-208. ISSN 0370-2693

[https://doi.org/10.1016/0370-2693\(93\)90211-Y](https://doi.org/10.1016/0370-2693(93)90211-Y)

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Measurement of B_s^0 and D_s^- meson lifetimes

The LHCb collaboration[†]

Abstract

We report on a measurement of the flavor-specific B_s^0 lifetime and of the D_s^- lifetime using proton-proton collisions at center-of-mass energies of 7 and 8 TeV, collected by the LHCb experiment and corresponding to 3.0 fb^{-1} of integrated luminosity. Approximately 407 000 $B_s^0 \rightarrow D_s^{(*)-} \mu^+ \nu_\mu$ decays are partially reconstructed in the $K^+ K^- \pi^- \mu^+$ final state. The B_s^0 and D_s^- natural widths are determined using, as a reference, kinematically similar $B^0 \rightarrow D^{(*)-} \mu^+ \nu_\mu$ decays reconstructed in the same final state. The resulting differences between widths of B_s^0 and B^0 mesons and of D_s^- and D^- mesons are $\Delta_\Gamma(B) = -0.0115 \pm 0.0053 \text{ (stat)} \pm 0.0041 \text{ (syst)} \text{ ps}^{-1}$ and $\Delta_\Gamma(D) = 1.0131 \pm 0.0117 \text{ (stat)} \pm 0.0065 \text{ (syst)} \text{ ps}^{-1}$, respectively. Combined with the known B^0 and D^- lifetimes, these yield the flavor-specific B_s^0 lifetime, $\tau_{B_s^0}^{\text{fs}} = 1.547 \pm 0.013 \text{ (stat)} \pm 0.010 \text{ (syst)} \pm 0.004 \text{ (}\tau_B\text{)} \text{ ps}$ and the D_s^- lifetime, $\tau_{D_s^-} = 0.5064 \pm 0.0030 \text{ (stat)} \pm 0.0017 \text{ (syst)} \pm 0.0017 \text{ (}\tau_D\text{)} \text{ ps}$. The last uncertainties originate from the limited knowledge of the B^0 and D^- lifetimes. The results improve upon current determinations.

Submitted to Phys. Rev. Lett.

[†]Authors are listed at the end of this paper.

Comparisons of precise measurements and predictions associated with quark-flavor dynamics probe the existence of unknown particles at energies much higher than those directly accessible at particle colliders. The precision of the predictions is often limited by the strong-interaction theory at low energies, where calculations are intractable. Predictive power is recovered by resorting to effective models such as heavy-quark effective theory [1–10], which rely on an expansion of the quantum chromodynamics corrections in powers of $1/m$, where m is the mass of the heavy quark in a bound system of a heavy quark and a light quark. These predictions are validated and refined using lifetime measurements of bottom and charm hadrons. Hence, improved lifetime measurements ultimately enhance the reach in searches for non-standard-model physics.

Measurements of the “flavor-specific” B_s^0 meson lifetime, $\tau_{B_s^0}^{\text{fs}}$, are particularly relevant [11]. This empirical quantity is a function of the natural widths of the two mass eigenstates resulting from B_s^0 – \bar{B}_s^0 oscillations; it is measured with a single-exponential fit to the distribution of decay time in final states to which only one of B_s^0 and \bar{B}_s^0 mesons can decay [12]. The current best determination, $\tau_{B_s^0}^{\text{fs}} = 1.535 \pm 0.015(\text{stat}) \pm 0.014(\text{syst})$ ps [13], obtained by the LHCb collaboration using hadronic $B_s^0 \rightarrow D_s^- \pi^+$ decays, has similar statistical and systematic uncertainties. Semileptonic B_s^0 decays, owing to larger signal yields than in hadronic decays, offer richer potential for precise $\tau_{B_s^0}^{\text{fs}}$ measurements. However, neutrinos and low-momentum neutral final-state particles prevent the full reconstruction of such decays. This introduces systematic limitations associated with poor knowledge of backgrounds and difficulties in obtaining the decay time from the observed decay-length distribution. Indeed, the result $\tau_{B_s^0}^{\text{fs}} = 1.479 \pm 0.010(\text{stat}) \pm 0.021(\text{syst})$ ps [14], based on a $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu X$ sample from the D0 collaboration, is limited by the systematic uncertainty. Throughout this Letter, the symbol X identifies any decay product, other than neutrinos, not included in the candidate reconstruction, and the inclusion of charge-conjugate processes is implied.

In this Letter, we use a novel approach that suppresses the above limitations and achieves a precise measurement of the flavor-specific B_s^0 meson lifetime. The lifetime is determined from the variation in the B_s^0 signal yield as a function of decay time, relative to that of B^0 decays that are reconstructed in the same final state and whose lifetime is precisely known. The use of kinematically similar B^0 decays as a reference allows the reduction of the uncertainties from partial reconstruction and lifetime-biasing selection criteria. The analysis also yields a significantly improved determination of the D_s^- lifetime over the current best result, $\tau_{D_s^-} = 0.5074 \pm 0.0055(\text{stat}) \pm 0.0051(\text{syst})$ ps, reported by the FOCUS collaboration [15].

We analyze proton-proton collisions at center-of-mass energies of 7 and 8 TeV collected by the LHCb experiment in 2011 and 2012 and corresponding to an integrated luminosity of 3.0 fb^{-1} . We use a sample of approximately 407 000 $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$ and $B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu$ “signal” decays, and a sample of approximately 108 000 $B^0 \rightarrow D^{*-} \mu^+ \nu_\mu$ and $B^0 \rightarrow D^- \mu^+ \nu_\mu$ “reference” decays. The D candidates are reconstructed as combinations of K^+ , K^- , and π^- candidates originating from a common vertex, displaced from any proton-proton interaction vertex. The $B_{(s)}^0$ candidates, $K^+ K^- \pi^- \mu^+$, are formed by D candidates associated with muon candidates originating from another common displaced vertex. We collectively refer to the signal and reference decays as $B_s^0 \rightarrow [K^+ K^- \pi^-]_{D_s^{(*)-}} \mu^+ \nu_\mu$ and $B^0 \rightarrow [K^+ K^- \pi^-]_{D^{(*)-}} \mu^+ \nu_\mu$, respectively. A fit to the ratio of event yields between the signal and reference decays as a function of $B_{(s)}^0$ decay time, t , determines $\Delta_\Gamma(B) \equiv$

$1/\tau_{B_s^0}^{\text{fs}} - \Gamma_d$, where Γ_d is the known natural width of the B^0 meson. A similar fit performed as a function of the $D_{(s)}^-$ decay time determines the decay-width difference between D_s^- and D^- mesons, $\Delta_\Gamma(D)$. Event yields are determined by fitting the “corrected-mass” distribution of the candidates, $m_{\text{corr}} = p_{\perp, D\mu} + \sqrt{m_{D\mu}^2 + p_{\perp, D\mu}^2}$ [16]. This is determined from the invariant mass of the $D_{(s)}^- \mu^+$ pair, $m_{D\mu}$, and the component of its momentum perpendicular to the $B_{(s)}^0$ flight direction, $p_{\perp, D\mu}$, to compensate for the average momentum of unreconstructed decay products. The flight direction is the line connecting the $B_{(s)}^0$ production and decay vertices; the decay time $t = m_B L k / p_{D\mu}$ uses the known $B_{(s)}^0$ mass, m_B [17], the measured $B_{(s)}^0$ decay length, L , and the momentum of the $D_{(s)}^- \mu^+$ pair, $p_{D\mu}$. The scale factor k corrects $p_{D\mu}$ for the average momentum fraction carried by decay products excluded from the reconstruction [18, 19]. The effects of decay-time acceptances and resolutions, determined from simulation, are included.

The LHCb detector is a single-arm forward spectrometer equipped with precise charged-particle vertexing and tracking detectors, hadron-identification detectors, calorimeters, and muon detectors, optimized for the study of bottom- and charm-hadron decays [20, 21]. Simulation [22, 23] is used to identify all relevant sources of bottom-hadron decays, model the mass distributions, and correct for the effects of incomplete kinematic reconstructions, relative decay-time acceptances, and decay-time resolutions. The unknown details of the B_s^0 decay dynamics are modeled in the simulation through empirical form-factor parameters [24], assuming values inspired by the known B^0 form factors [11]. We assess the impact of these assumptions on the systematic uncertainties.

The online selection requires a high-transverse-momentum muon candidate associated with one, two, or three charged particles, all with origins displaced from the proton-proton interaction points [25]. In the offline reconstruction, the muon is combined with charged particles consistent with the topology and kinematics of signal $B_s^0 \rightarrow [K^+ K^- \pi^-]_{D_s^{(*)-}} \mu^+ \nu_\mu$ and reference $B^0 \rightarrow [K^+ K^- \pi^-]_{D^{(*)-}} \mu^+ \nu_\mu$ decays. The range of $K^+ K^- \pi^-$ mass is restricted around the known values of the $D_{(s)}^-$ meson masses such that cross-contamination between signal and reference samples is smaller than 0.1%, as estimated from simulation. We also reconstruct “same-sign” $K^+ K^- \pi^- \mu^-$ candidates, formed by charm and muon candidates with same-sign charge, to model combinatorial background from accidental $D_{(s)}^- \mu^+$ associations. The event selection is optimized toward suppressing the background under the charm signals and making same-sign candidates a reliable model for the combinatorial background: track- and vertex-quality, vertex-displacement, transverse-momentum, and particle-identification criteria are chosen to minimize shape and yield differences between same-sign and signal candidates in the $m_{D\mu} > 5.5 \text{ GeV}/c^2$ region, where genuine bottom-hadron decays are kinematically excluded and combinatorial background dominates. Mass vetoes suppress background from misreconstructed decays such as $B_s^0 \rightarrow \psi^{(\prime)}(\rightarrow \mu^+ \mu^-) \phi(\rightarrow K^+ K^-)$ decays where a muon is misidentified as a pion, $\Lambda_b^0 \rightarrow \Lambda_c^+(\rightarrow p K^- \pi^+) \mu^- \bar{\nu}_\mu X$ decays where the proton is misidentified as a kaon or a pion, and $B_{(s)}^0 \rightarrow D_{(s)}^- \pi^-$ decays where the pion is misidentified as a muon. Significant contributions arise from decays of a bottom hadron into pairs of charm hadrons, one peaking at the $D_{(s)}^-$ mass and the other decaying semileptonically, or into single charm hadrons and other particles. Such decays include $B_{(s)}^0 \rightarrow D_{(s)}^{(*)-} D_{(s)}^+$, $B^+ \rightarrow \bar{D}^{*0} D^{*+}$, $B^+ \rightarrow D^- \mu^+ \nu_\mu X$, $B^+ \rightarrow D_s^{*-} K^+ \mu^+ \nu_\mu X$, $B^0 \rightarrow D_s^{*-} K^0 \mu^+ \nu_\mu X$, $B_s^0 \rightarrow D^0 D_s^- K^+$, $B_s^0 \rightarrow D^- D_s^+ K^0$, $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^{*-} X$, and $\Lambda_b^0 \rightarrow D_s^+ \Lambda \mu^- \bar{\nu}_\mu X$ decays. We suppress these

backgrounds with a threshold, linearly dependent on m_{corr} , applied to the $D_{(s)}^-$ momentum component perpendicular to the $B_{(s)}^0$ flight direction. Finally, a $t > 0.1$ ps requirement on the $D_{(s)}^-$ proper decay-time renders the signal- and reference-decay acceptances as functions of decay time more similar, with little penalty for signal.

A total of approximately 468 000 (141 000) signal (reference) candidates, formed by combining $K^+K^-\pi^-$ candidates in the D_s^- (D^-) signal range with μ^+ candidates, satisfy the selection. Figure 1 shows the relevant mass distributions. The enhancements of the signal and reference distributions over the corresponding same-sign distributions for $m_{D\mu} < 5.5$ GeV/ c^2 are due to bottom-hadron decays. The absence of candidates at $m_{D\mu} \approx 5.3$ GeV/ c^2 results from the $B_{(s)}^0 \rightarrow D_{(s)}^-\pi^-$ veto. The two peaks in the $K^+K^-\pi^-$ distributions of same-sign candidates are due to genuine charm decays accidentally combined with muon candidates. Along with $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{(*)-}}\mu^+\nu_\mu$ decays, many B_s^0 decays potentially useful for the lifetime measurement contribute signal candidates, including decays into $D_{(s)}^{**}(\rightarrow D_{(s)}^{(*)-}X)\mu^+\nu_\mu$, $D_s^-\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$, $D_s^{*-}(\rightarrow D_s^-X)\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$, and $D_s^{**}(\rightarrow D_s^{(*)-}X)\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$ final states.¹ Similarly, along with the $B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ decays, potential reference candidates come from B^0 decays into $D^{**}(\rightarrow D^{(*)-}X)\mu^+\nu_\mu$, $D^-\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$, $D^{*-}(\rightarrow D^-X)\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$, and $D^{**}(\rightarrow D^{(*)-}X)\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau$ final states. However, we restrict the signal (reference) decays solely to the $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{(*)-}}\mu^+\nu_\mu$ ($B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$) channels because they contribute 95% (91%) of the inclusive $K^+K^-\pi^-\mu^+$ yield from semileptonic B^0 (B_s^0) decays and require smaller and better-known k -factor corrections to relate the observed decay times to their true values.

A reliable understanding of the sample composition is essential for unbiased lifetime results. An unbiased determination from simulation of the acceptances and mass distributions as functions of decay time requires that the simulated sample mirrors the data composition. We therefore weight the composition of the simulated samples according to the results of a least-squares fit to the m_{corr} distributions in data (Fig. 2). In the B_s^0 sample, such a global composition-fit includes the two signal components, $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^-}\mu^+\nu_\mu$ and $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{*-}}\mu^+\nu_\mu$; a combinatorial component; and two physics backgrounds. The physics backgrounds are formed by grouping together contributions with similar corrected-mass distributions, determined from simulation. They are divided into contributions at lower values of corrected mass ($B^0 \rightarrow D^{(*)-}D_s^{(*)+}$, $B^+ \rightarrow \bar{D}^{(*)0}D_s^{(*)+}$, and $D^{**}(\rightarrow D_s^{(*)-}X)\mu^+\nu_\mu$) and at higher corrected-mass values ($B^+ \rightarrow D_s^{(*)-}K^+\mu^+\nu_\mu X$, $B^0 \rightarrow D_s^{(*)-}K^0\mu^+\nu_\mu X$, and $B_s^0 \rightarrow D_s^-\tau^+(\rightarrow \mu^+\nu_\mu\bar{\nu}_\tau)\nu_\tau X$). The distributions of all components are modeled empirically from simulation, except for the combinatorial component, which is modeled using same-sign data. Contributions expected to be smaller than 0.5% are neglected. The effect of this approximation and of possible variations of the relative proportions within each fit category are treated as contributions to the systematic uncertainties. The fit p -value is 62.1% and the fractions of each component are determined with absolute statistical uncertainties in the range 0.13%–0.91%. A simpler composition fit is used for the B^0 sample. Signal and combinatorial components are chosen similarly to the B_s^0 case; the contributions from $B^0 \rightarrow D^{**}(\rightarrow D^{(*)-}X)\mu^+\nu_\mu$ and $B^+ \rightarrow D^-\mu^+\nu_\mu X$ decays have sufficiently similar distributions to be merged into a single physics-background component. The results of the corrected-mass fit of the reference sample also offer a

¹Here and in the following, the symbol $D_{(s)}^{**}$ identifies collectively higher orbital excitations of $D_{(s)}^-$ mesons.




Figure 1: Distributions of $D\mu$ mass for (top panel) reference candidates, formed by combining $D^- \rightarrow K^+K^-\pi^-$ candidates with μ^+ candidates, and (bottom panel) signal candidates formed by $D_s^- \rightarrow K^+K^-\pi^-$ candidates combined with μ^+ candidates. The inset shows the $K^+K^-\pi^-$ -mass distribution with vertical lines enclosing the D^- (D_s^-) candidates used to form the reference (signal) candidates. The dark-filled histograms show same-sign candidate distributions.

validation of the approach, since the composition of this sample is known precisely from other experiments. The largest discrepancy observed among the individual fractional contributions is 1.3 statistical standard deviations.

The composition fit is sufficient for the determination of $\Delta_\Gamma(D)$, where no k -factor corrections are needed since the final state is fully reconstructed. We determine $\Delta_\Gamma(D)$ through a least-squares fit of the ratio of signal B_s^0 and reference B^0 yields as a function of the charm-meson decay time in the range 0.1–4.0 ps. The yields of signal $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{(*)-}}\mu^+\nu_\mu$ and reference $B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ decays are determined in each of 20 decay-time bins with a m_{corr} fit similar to the global composition-fit. The two signal and the two physics-background contributions are each merged into a single component according to the total proportions determined by the global fit and their decay-time dependence as determined from simulation. The fit includes the decay-time resolution and the ratio between signal and reference decay-time acceptances, which is determined from simulation to be uniform within 1%. The fit is shown in the top panel of Fig. 3; it has 34% p -value and determines $\Delta_\Gamma(D) = 1.0131 \pm 0.0117 \text{ ps}^{-1}$.

The measurement of $\Delta_\Gamma(B)$ requires an acceptance correction for the differences between signal and reference decays and the k -factor correction. The acceptance correction accounts for the difference in decay-time-dependent efficiency due to the combined effect of the difference between D^- and D_s^- lifetimes and the online requirements on the spatial separation between $D_{(s)}^-$ and $B_{(s)}^0$ decay vertices: we apply to the B_s^0 sample a per-candidate




Figure 2: Corrected-mass distributions for (top panel) reference $B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ and (bottom panel) signal $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{(*)-}}\mu^+\nu_\mu$ candidates satisfying the selection. Results of the global composition-fit are overlaid. In the signal sample the lower- and higher-mass backgrounds are independently fit as described in the text, but merged into a single “physics background” component in the above projection.

weight, based on the $\Delta_\Gamma(D)$ result, such that the D_s^- and D^- decay-time distributions become consistent. The k -factor correction is a candidate-specific correction, where the average missing momentum in a simulated sample is used to correct the reconstructed momentum in data. The k -factor dependence on the kinematic properties of each candidate is included through a dependence on $m_{D\mu}$, $k(m_{D\mu}) = \langle p_{D\mu}/p_{\text{true}} \rangle$, where p_{true} indicates the true momentum of the $B_{(s)}^0$ meson. The equalization of the compositions of simulated and experimental data samples ensures that the k -factor distribution specific to each of the four signal and reference decays is unbiased. We determine $\Delta_\Gamma(B)$ with the same fit of m_{corr} used to measure $\Delta_\Gamma(D)$ but where the ratios of signal and reference yields are determined as functions of the $B_{(s)}^0$ decay time. The decay-time smearing due to the k -factor spread is included in the fit. After the D_s^- lifetime weighting, the decay-time acceptances of simulated signal and reference modes are consistent, with a p -value of 83%, and are not included in the fit. The fit is shown in the middle panel of Fig. 3; the resulting width difference is $\Delta_\Gamma(B) = -0.0115 \pm 0.0053 \text{ ps}^{-1}$, with 91% p -value.

To check against biases due to differing acceptances and kinematic properties, the analysis is validated with a null test. We repeat the width-difference determination by using the same reference $B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ sample and replacing the signal decays with 2.1 million $B^0 \rightarrow [K^+\pi^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ decays, where the D^- is reconstructed in the $K^+\pi^-\pi^-$ final state (Fig. 3, bottom panel). Differing momentum and vertex-displacement selection criteria induce up to 10% differences between acceptances as a




Figure 3: Ratio between acceptance-corrected yields of signal $B_s^0 \rightarrow [K^+K^-\pi^-]_{D_s^{(*)-}}\mu^+\nu_\mu$ and reference $B^0 \rightarrow [K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ decay yields as a function of (top panel) charm-meson and (middle panel) bottom-meson decay time. The bottom panel shows the ratio between acceptance-corrected B^0 decay yields in the $[K^+\pi^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ and $[K^+K^-\pi^-]_{D^{(*)-}}\mu^+\nu_\mu$ channels as a function of B^0 decay time. Fit results are overlaid. Relevant for the results is only the slope of the ratios as a function of decay time; absolute ratios, which depend on the decay yields, weighting, and efficiencies, are irrelevant.

function of D^- decay time and up to 25% variations as a function of B^0 decay time. Acceptance ratios are therefore included in the fit. The p -values are 21% for the B^0 fit and 33% for the D^- fit. The resulting width differences, $\Delta_\Gamma(D) = (-19 \pm 10) \times 10^{-3} \text{ ps}^{-1}$ and $\Delta_\Gamma(B) = (-4.1 \pm 5.4) \times 10^{-3} \text{ ps}^{-1}$, are consistent with zero.

We assess independent systematic uncertainties due to (i) potential fit biases; (ii) assumptions on the components contributing to the sample and their mass distributions; (iii) assumptions on the signal decay model, e.g., choice of $B_s^0 \rightarrow D_s^{*-}$ form factors; (iv) uncertainties on the decay-time acceptances; (v) uncertainties on the decay-time resolution; (vi) contamination from B_s^0 candidates produced in B_c^+ decays; and (vii) mismodeling of transverse-momentum (p_T) differences between B^0 and B_s^0 mesons. We evaluate each contribution by including the relevant effect in the model and repeating the whole analysis on ensembles of simulated experiments that mirror the data. For the $\Delta_\Gamma(D)$ result, the systematic uncertainty is dominated by a 0.0049 ps^{-1} contribution due to the decay-time acceptance, and a 0.0039 ps^{-1} contribution due to the decay-time resolution. A smaller contribution of 0.0018 ps^{-1} arises from possible mismodeling of p_T differences in B^0 and B_s^0 production. For the $\Delta_\Gamma(B)$ result, a 0.0028 ps^{-1} uncertainty from mismodeling of p_T differences between B^0 and B_s^0 mesons and a 0.0025 ps^{-1} contribution from the B_s^0 decay model dominate. Smaller contributions arise from B_c^+ feed-down (0.0010 ps^{-1}), residual

fit biases (0.0009 ps^{-1}), sample composition (0.0005 ps^{-1}), and decay-time acceptance and resolution (0.0004 ps^{-1} each). The uncertainties associated with the limited size of simulated samples are included in the fit χ^2 and contribute up to 20% of the statistical uncertainties. The uncertainty in the decay length has negligible impact. Consistency checks based on repeating the measurement independently on subsamples chosen according to data-taking time, online-selection criteria, charged-particle and vertex multiplicities, momentum of the $K^+K^-\pi^-\mu^+$ system, and whether only the $D_s^-\mu^+\nu_\mu$ or the $D_s^{*-}\mu^+\nu_\mu$ channel is considered as signal, all yield results compatible with statistical fluctuations.

In summary, we report world-leading measurements of B_s^0 and D_s^- meson lifetimes using a novel method. We reconstruct $B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu$ and $B_s^0 \rightarrow D_s^-\mu^+\nu_\mu$ decays from proton-proton collision data collected by the LHCb experiment and corresponding to 3.0 fb^{-1} of integrated luminosity. We use $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ and $B^0 \rightarrow D^-\mu^+\nu_\mu$ decays reconstructed in the same final state as a reference to suppress systematic uncertainties. The resulting width differences are $\Delta_\Gamma(B) = -0.0115 \pm 0.0053 \text{ (stat)} \pm 0.0041 \text{ (syst)} \text{ ps}^{-1}$ and $\Delta_\Gamma(D) = 1.0131 \pm 0.0117 \text{ (stat)} \pm 0.0065 \text{ (syst)} \text{ ps}^{-1}$. Their correlation is negligible. Using the known values of the B^0 [17, 26] and D^- lifetimes [17, 27], we determine the flavor-specific B_s^0 lifetime, $\tau_{B_s^0}^{\text{fs}} = 1.547 \pm 0.013 \text{ (stat)} \pm 0.010 \text{ (syst)} \pm 0.004 \text{ (}\tau_B\text{)} \text{ ps}$, and the D_s^- lifetime, $\tau_{D_s^-} = 0.5064 \pm 0.0030 \text{ (stat)} \pm 0.0017 \text{ (syst)} \pm 0.0017 \text{ (}\tau_D\text{)} \text{ ps}$; the last uncertainties are due to the limited knowledge of the B^0 and D^- lifetime, respectively. The results are consistent with, and significantly more precise than the current values [13–15]. They might offer improved insight into the interplay between strong and weak interactions in the dynamics of heavy mesons and sharpen the reach of current and future indirect searches for non-standard-model physics.

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).

References

- [1] M. A. Shifman and M. B. Voloshin, *On annihilation of mesons built from heavy and light quark and B^0 - \bar{B}^0 oscillations*, Sov. J. Nucl. Phys. **45** (1987) 292.
- [2] M. A. Shifman and M. B. Voloshin, *On production of D and D^* mesons in B meson decays*, Sov. J. Nucl. Phys. **47** (1988) 511.
- [3] E. Eichten and B. R. Hill, *An effective field theory for the calculation of matrix elements involving heavy quarks*, Phys. Lett. **B234** (1990) 511.
- [4] B. Guberina, R. D. Peccei, and R. Rückl, *Weak decays of heavy quarks*, Phys. Lett. **B91** (1980) 116.
- [5] B. Guberina, R. D. Peccei, and R. Rückl, *Effects of QCD on the Cabibbo patterns in B meson decays*, Phys. Lett. **B90** (1980) 169.
- [6] N. Isgur and M. B. Wise, *Weak decays of heavy mesons in the static quark approximation*, Phys. Lett. **B232** (1989) 113.
- [7] N. Isgur and M. B. Wise, *Weak transition form-factors between heavy mesons*, Phys. Lett. **B237** (1990) 527.
- [8] B. Grinstein, *The static quark effective theory*, Nucl. Phys. **B339** (1990) 253.
- [9] A. F. Falk, H. Georgi, B. Grinstein, and M. B. Wise, *Heavy meson form-factors from QCD*, Nucl. Phys. **B343** (1990) 1.
- [10] H. Georgi, *An effective field theory for heavy quarks at low-energies*, Phys. Lett. **B240** (1990) 447.
- [11] Heavy Flavor Averaging Group, Y. Amhis *et al.*, *Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2016*, arXiv:1612.07233, updated results and plots available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [12] K. Hartkorn and H. G. Moser, *A new method of measuring $\Delta(\Gamma)/\Gamma$ in the B_s^0 - \bar{B}_s^0 system*, Eur. Phys. J. **C8** (1999) 381.
- [13] LHCb collaboration, R. Aaij *et al.*, *Measurement of the \bar{B}_s^0 meson lifetime in $D_s^+\pi^-$ decays*, Phys. Rev. Lett. **113** (2014) 172001, arXiv:1407.5873.
- [14] D0 collaboration, V. M. Abazov *et al.*, *Measurement of the B_s^0 lifetime in the flavor-specific decay channel $B_s^0 \rightarrow D_s^- \mu^+ \nu X$* , Phys. Rev. Lett. **114** (2015) 062001, arXiv:1410.1568.
- [15] FOCUS collaboration, J. M. Link *et al.*, *A measurement of the D_s^+ lifetime*, Phys. Rev. Lett. **95** (2005) 052003, arXiv:hep-ex/0504056.
- [16] Fermilab E653 collaboration, K. Kodama *et al.*, *Measurement of the relative branching fraction $\Gamma(D^0 \rightarrow K\mu\nu)/\Gamma(D^0 \rightarrow \mu X)$* , Phys. Rev. Lett. **66** (1991) 1819.
- [17] Particle Data Group, C. Patrignani *et al.*, *Review of particle physics*, Chin. Phys. **C40** (2016) 100001.

- [18] CDF collaboration, A. Abulencia *et al.*, *Observation of B_s^0 - \bar{B}_s^0 oscillations*, Phys. Rev. Lett. **97** (2006) 242003, arXiv:hep-ex/0609040.
- [19] N. T. Leonardo, *Analysis of B_s^0 flavor oscillations at CDF*, PhD thesis, FERMILAB-THESIS-2006-18, (2006).
- [20] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [21] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, arXiv:1412.6352.
- [22] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [23] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [24] I. Caprini, L. Lellouch, and M. Neubert, *Dispersive bounds on the shape of $B^0 \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ form factors*, Nucl. Phys. **B530** (1998) 153, arXiv:hep-ph/9712417.
- [25] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, arXiv:1211.3055.
- [26] LHCb collaboration, R. Aaij *et al.*, *Measurements of the B^+ , B^0 , B_s^0 meson and Λ_b^0 baryon lifetimes*, JHEP **04** (2014) 114, arXiv:1402.2554.
- [27] FOCUS collaboration, J. M. Link *et al.*, *New measurements of the D^0 and D^+ lifetimes*, Phys. Lett. **B537** (2002) 192, arXiv:hep-ex/0203037.

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁹, M. Adinolfi⁴⁸, Z. Ajaltouni⁵, S. Akar⁵⁹, J. Albrecht¹⁰, F. Alessio⁴⁰, M. Alexander⁵³, S. Ali⁴³, G. Alkhazov³¹, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁵⁹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An³, L. Anderlini¹⁸, G. Andreassi⁴¹, M. Andreotti^{17,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, F. Archilli⁴³, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁷, M. Artuso⁶¹, E. Aslanides⁶, G. Auriemma²⁶, M. Baalouch⁵, I. Babuschkin⁵⁶, S. Bachmann¹², J.J. Back⁵⁰, A. Badalov³⁸, C. Baesso⁶², S. Baker⁵⁵, V. Balagura^{7,c}, W. Baldini¹⁷, A. Baranov³⁵, R.J. Barlow⁵⁶, C. Barschel⁴⁰, S. Barsuk⁷, W. Barter⁵⁶, F. Baryshnikov³², M. Baszczyk²⁷, V. Batozskaya²⁹, B. Batsukh⁶¹, V. Battista⁴¹, A. Bay⁴¹, L. Beaucourt⁴, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, A. Beiter⁶¹, L.J. Bel⁴³, V. Bellee⁴¹, N. Belloli^{21,i}, K. Belous³⁷, I. Belyaev³², E. Ben-Haim⁸, G. Bencivenni¹⁹, S. Benson⁴³, S. Beranek⁹, A. Berezhnoy³³, R. Bernet⁴², A. Bertolin²³, C. Betancourt⁴², F. Betti¹⁵, M.-O. Bettler⁴⁰, M. van Beuzekom⁴³, I. Bezshyiko⁴², S. Bifani⁴⁷, P. Billoir⁸, A. Birnkraut¹⁰, A. Bitadze⁵⁶, A. Bizzeti^{18,u}, T. Blake⁵⁰, F. Blanc⁴¹, J. Blouw^{11,†}, S. Blusk⁶¹, V. Bocci²⁶, T. Boettcher⁵⁸, A. Bondar^{36,w}, N. Bondar³¹, W. Bonivento¹⁶, I. Bordyuzhin³², A. Borgheresi^{21,i}, S. Borghi⁵⁶, M. Borisyak³⁵, M. Borsato³⁹, F. Bossu⁷, M. Boubdir⁹, T.J.V. Bowcock⁵⁴, E. Bowen⁴², C. Bozzi^{17,40}, S. Braun¹², T. Britton⁶¹, J. Brodzicka⁵⁶, E. Buchanan⁴⁸, C. Burr⁵⁶, A. Bursche², J. Buytaert⁴⁰, S. Cadeddu¹⁶, R. Calabrese^{17,g}, M. Calvi^{21,i}, M. Calvo Gomez^{38,m}, A. Camboni³⁸, P. Campana¹⁹, D.H. Campora Perez⁴⁰, L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni^{25,j}, R. Cardinale^{20,h}, A. Cardini¹⁶, P. Carniti^{21,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina^{21,i}, L. Castillo Garcia⁴¹, M. Cattaneo⁴⁰, G. Cavallero²⁰, R. Cenci^{24,t}, D. Chamont⁷, M. Charles⁸, Ph. Charpentier⁴⁰, G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, S. Chen⁵⁶, S.-F. Cheung⁵⁷, V. Chobanova³⁹, M. Chruszcz^{42,27}, A. Chubykin³¹, X. Cid Vidal³⁹, G. Ciezarek⁴³, P.E.L. Clarke⁵², M. Clemencic⁴⁰, H.V. Cliff⁴⁹, J. Closier⁴⁰, V. Coco⁵⁹, J. Cogan⁶, E. Cogneras⁵, V. Cogoni^{16,f}, L. Cojocariu³⁰, P. Collins⁴⁰, A. Comerma-Montells¹², A. Contu⁴⁰, A. Cook⁴⁸, G. Coombs⁴⁰, S. Coquereau³⁸, G. Corti⁴⁰, M. Corvo^{17,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴⁰, G.A. Cowan⁵², D.C. Craik⁵², A. Crocombe⁵⁰, M. Cruz Torres⁶², S. Cunliffe⁵⁵, R. Currie⁵², C. D'Ambrosio⁴⁰, F. Da Cunha Marinho², E. Dall'Occo⁴³, J. Dalseno⁴⁸, P.N.Y. David⁴³, A. Davis³, K. De Bruyn⁶, S. De Capua⁵⁶, M. De Cian¹², J.M. De Miranda¹, L. De Paula², M. De Serio^{14,d}, P. De Simone¹⁹, C.T. Dean⁵³, D. Decamp⁴, M. Deckenhoff¹⁰, L. Del Buono⁸, H.-P. Dembinski¹¹, M. Demmer¹⁰, A. Dendek²⁸, D. Derkach³⁵, O. Deschamps⁵, F. Dettori⁵⁴, B. Dey²², A. Di Canto⁴⁰, P. Di Nezza¹⁹, H. Dijkstra⁴⁰, F. Dordei⁴⁰, M. Dorigo⁴¹, A. Dosil Suárez³⁹, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour⁴³, G. Dujany⁵⁶, K. Dungs⁴⁰, P. Durante⁴⁰, R. Dzhelezhyan³⁷, M. Dziewiecki¹², A. Dziurda⁴⁰, A. Dzyuba³¹, N. Déleage⁴, S. Easo⁵¹, M. Ebert⁵², U. Egede⁵⁵, V. Egorychev³², S. Eidelman^{36,w}, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, S. Ely⁶¹, S. Esen¹², H.M. Evans⁴⁹, T. Evans⁵⁷, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, R. Fay⁵⁴, D. Fazzini^{21,i}, D. Ferguson⁵², G. Fernandez³⁸, A. Fernandez Prieto³⁹, F. Ferrari¹⁵, F. Ferreira Rodrigues², M. Ferro-Luzzi⁴⁰, S. Filippov³⁴, R.A. Fini¹⁴, M. Fiore^{17,g}, M. Fiorini^{17,g}, M. Firlej²⁸, C. Fitzpatrick⁴¹, T. Fiutowski²⁸, F. Fleuret^{7,b}, K. Fohl⁴⁰, M. Fontana^{16,40}, F. Fontanelli^{20,h}, D.C. Forshaw⁶¹, R. Forty⁴⁰, V. Franco Lima⁵⁴, M. Frank⁴⁰, C. Frei⁴⁰, J. Fu^{22,g}, W. Funk⁴⁰, E. Furfaro^{25,j}, C. Färber⁴⁰, A. Gallas Torreira³⁹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², M. Gandelman², P. Gandini⁵⁷, Y. Gao³, L.M. Garcia Martin⁶⁹, J. García Pardiñas³⁹, J. Garra Tico⁴⁹, L. Garrido³⁸, P.J. Garsed⁴⁹, D. Gascon³⁸, C. Gaspar⁴⁰, L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck¹², M. Gersabeck⁵⁶, T. Gershon⁵⁰, Ph. Ghez⁴, S. Giani⁴¹, V. Gibson⁴⁹, O.G. Girard⁴¹, L. Giubega³⁰, K. Gizdov⁵², V.V. Gligorov⁸, D. Golubkov³², A. Golutvin^{55,40}, A. Gomes^{1,a}, I.V. Gorelov³³, C. Gotti^{21,i}, E. Govorkova⁴³, R. Graciani Diaz³⁸, L.A. Granado Cardoso⁴⁰, E. Graugés³⁸, E. Graverini⁴², G. Graziani¹⁸, A. Grecu³⁰, R. Greim⁹, P. Griffith¹⁶, L. Grillo^{21,40,i}, B.R. Gruberg Cazon⁵⁷, O. Grünberg⁶⁷, E. Gushchin³⁴, Yu. Guz³⁷,

T. Gys⁴⁰, C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴¹, C. Haen⁴⁰,
 S.C. Haines⁴⁹, B. Hamilton⁶⁰, X. Han¹², S. Hansmann-Menzemer¹², N. Harnew⁵⁷,
 S.T. Harnew⁴⁸, J. Harrison⁵⁶, M. Hatch⁴⁰, J. He⁶³, T. Head⁴¹, A. Heister⁹, K. Hennessy⁵⁴,
 P. Henrard⁵, L. Henry⁶⁹, E. van Herwijnen⁴⁰, M. Heß⁶⁷, A. Hicheur², D. Hill⁵⁷, C. Hombach⁵⁶,
 H. Hopchev⁴¹, Z.-C. Huard⁵⁹, W. Hulsbergen⁴³, T. Humair⁵⁵, M. Hushchyn³⁵, D. Hutchcroft⁵⁴,
 M. Idzik²⁸, P. Ilten⁵⁸, R. Jacobsson⁴⁰, J. Jalocha⁵⁷, E. Jans⁴³, A. Jawahery⁶⁰, F. Jiang³,
 M. John⁵⁷, D. Johnson⁴⁰, C.R. Jones⁴⁹, C. Joram⁴⁰, B. Jost⁴⁰, N. Jurik⁵⁷, S. Kandybei⁴⁵,
 M. Karacson⁴⁰, J.M. Kariuki⁴⁸, S. Karodia⁵³, M. Kecke¹², M. Kelsey⁶¹, M. Kenzie⁴⁹, T. Ketel⁴⁴,
 E. Khairullin³⁵, B. Khanji¹², C. Khurewathanakul⁴¹, T. Kirn⁹, S. Klaver⁵⁶, K. Klimaszewski²⁹,
 T. Klimkovich¹¹, S. Koliiev⁴⁶, M. Kolpin¹², I. Komarov⁴¹, R. Kopecna¹², P. Koppenburg⁴³,
 A. Kosmyntseva³², S. Kotriakhova³¹, A. Kozachuk³³, M. Kozeiha⁵, L. Kravchuk³⁴, M. Kreps⁵⁰,
 P. Krokovny^{36,w}, F. Kruse¹⁰, W. Krzemien²⁹, W. Kucewicz^{27,l}, M. Kucharczyk²⁷,
 V. Kudryavtsev^{36,w}, A.K. Kuonen⁴¹, K. Kurek²⁹, T. Kvaratskheliya^{32,40}, D. Lacarrere⁴⁰,
 G. Lafferty⁵⁶, A. Lai¹⁶, G. Lanfranchi¹⁹, C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷,
 R. Le Gac⁶, J. van Leerdam⁴³, A. Leflat^{33,40}, J. Lefrançois⁷, R. Lefèvre⁵, F. Lemaitre⁴⁰,
 E. Lemos Cid³⁹, O. Leroy⁶, T. Lesiak²⁷, B. Leverington¹², T. Li³, Y. Li⁷, Z. Li⁶¹,
 T. Likhomanenko^{35,68}, R. Lindner⁴⁰, F. Lionetto⁴², X. Liu³, D. Loh⁵⁰, I. Longstaff⁵³,
 J.H. Lopes², D. Lucchesi^{23,o}, M. Lucio Martinez³⁹, H. Luo⁵², A. Lupato²³, E. Luppi^{17,g},
 O. Lupton⁴⁰, A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷, F. Maciuc³⁰, O. Maev³¹, K. Maguire⁵⁶,
 S. Malde⁵⁷, A. Malinin⁶⁸, T. Maltsev³⁶, G. Manca^{16,f}, G. Mancinelli⁶, P. Manning⁶¹,
 J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵, C. Marin Benito³⁸, M. Marinangeli⁴¹,
 P. Marino^{24,t}, J. Marks¹², G. Martellotti²⁶, M. Martin⁶, M. Martinelli⁴¹, D. Martinez Santos³⁹,
 F. Martinez Vidal⁶⁹, D. Martins Tostes², L.M. Massacrier⁷, A. Massafferri¹, R. Matev⁴⁰,
 A. Mathad⁵⁰, Z. Mathe⁴⁰, C. Matteuzzi²¹, A. Mauri⁴², E. Maurice^{7,b}, B. Maurin⁴¹,
 A. Mazurov⁴⁷, M. McCann^{55,40}, A. McNab⁵⁶, R. McNulty¹³, B. Meadows⁵⁹, F. Meier¹⁰,
 D. Melnychuk²⁹, M. Merk⁴³, A. Merli^{22,q}, E. Michielin²³, D.A. Milanes⁶⁶, M.-N. Minard⁴,
 D.S. Mitzel¹², A. Mogini⁸, J. Molina Rodriguez¹, I.A. Monroy⁶⁶, S. Monteil⁵, M. Morandin²³,
 M.J. Morello^{24,t}, O. Morgunova⁶⁸, J. Moron²⁸, A.B. Morris⁵², R. Mountain⁶¹, F. Muheim⁵²,
 M. Mulder⁴³, M. Mussini¹⁵, D. Müller⁵⁶, J. Müller¹⁰, K. Müller⁴², V. Müller¹⁰, P. Naik⁴⁸,
 T. Nakada⁴¹, R. Nandakumar⁵¹, A. Nandi⁵⁷, I. Nasteva², M. Needham⁵², N. Neri^{22,40},
 S. Neubert¹², N. Neufeld⁴⁰, M. Neuner¹², T.D. Nguyen⁴¹, C. Nguyen-Mau^{41,n}, S. Nieswand⁹,
 R. Niet¹⁰, N. Nikitin³³, T. Nikodem¹², A. Nogay⁶⁸, A. Novoselov³⁷, D.P. O'Hanlon⁵⁰,
 A. Oblakowska-Mucha²⁸, V. Obraztsov³⁷, S. Ogilvy¹⁹, R. Oldeman^{16,f}, C.J.G. Onderwater⁷⁰,
 A. Ossowska²⁷, J.M. Otalora Goicochea², P. Owen⁴², A. Oyanguren⁶⁹, P.R. Pais⁴¹,
 A. Palano^{14,d}, M. Palutan^{19,40}, A. Papanestis⁵¹, M. Pappagallo^{14,d}, L.L. Pappalardo^{17,g},
 C. Pappenheimer⁵⁹, W. Parker⁶⁰, C. Parkes⁵⁶, G. Passaleva¹⁸, A. Pastore^{14,d}, M. Patel⁵⁵,
 C. Patrignani^{15,e}, A. Pearce⁴⁰, A. Pellegrino⁴³, G. Penso²⁶, M. Pepe Altarelli⁴⁰, S. Perazzini⁴⁰,
 P. Perret⁵, L. Pescatore⁴¹, K. Petridis⁴⁸, A. Petrolini^{20,h}, A. Petrov⁶⁸, M. Petruzzo^{22,q},
 E. Picatoste Olloqui³⁸, B. Pietrzyk⁴, M. Piekies²⁷, D. Pinci²⁶, A. Pistone²⁰, A. Piucci¹²,
 V. Placinta³⁰, S. Playfer⁵², M. Plo Casasus³⁹, T. Poikela⁴⁰, F. Polci⁸, M. Poli Lener¹⁹,
 A. Poluektov^{50,36}, I. Polyakov⁶¹, E. Polcarpo², G.J. Pomery⁴⁸, S. Ponce⁴⁰, A. Popov³⁷,
 D. Popov^{11,40}, B. Popovici³⁰, S. Poslavskii³⁷, C. Potterat², E. Price⁴⁸, J. Prisciandaro³⁹,
 C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴², G. Punzi^{24,p}, C. Qian⁶³, W. Qian⁵⁰,
 R. Quagliani^{7,48}, B. Rachwal²⁸, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas³⁹,
 M.S. Rangel², I. Raniuk⁴⁵, F. Ratnikov³⁵, G. Raven⁴⁴, F. Redi⁵⁵, S. Reichert¹⁰, A.C. dos Reis¹,
 C. Remon Alepuz⁶⁹, V. Renaudin⁷, S. Ricciardi⁵¹, S. Richards⁴⁸, M. Rihl⁴⁰, K. Rinnert⁵⁴,
 V. Rives Molina³⁸, P. Robbe⁷, A.B. Rodrigues¹, E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁶,
 P. Rodriguez Perez^{56,†}, A. Rogozhnikov³⁵, S. Roiser⁴⁰, A. Rollings⁵⁷, V. Romanovskiy³⁷,
 A. Romero Vidal³⁹, J.W. Ronayne¹³, M. Rotondo¹⁹, M.S. Rudolph⁶¹, T. Ruf⁴⁰, P. Ruiz Valls⁶⁹,
 J.J. Saborido Silva³⁹, E. Sadykhov³², N. Sagidova³¹, B. Saitta^{16,f}, V. Salustino Guimaraes¹,

D. Sanchez Gonzalo³⁸, C. Sanchez Mayordomo⁶⁹, B. Sanmartin Sedes³⁹, R. Santacesaria²⁶,
C. Santamarina Rios³⁹, M. Santimaria¹⁹, E. Santovetti^{25,j}, A. Sarti^{19,k}, C. Satriano^{26,s},
A. Satta²⁵, D.M. Saunders⁴⁸, D. Savrina^{32,33}, S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁵³,
H. Schindler⁴⁰, M. Schlupp¹⁰, M. Schmelling¹¹, T. Schmelzer¹⁰, B. Schmidt⁴⁰, O. Schneider⁴¹,
A. Schopper⁴⁰, H.F. Schreiner⁵⁹, K. Schubert¹⁰, M. Schubiger⁴¹, M.-H. Schune⁷,
R. Schwemmer⁴⁰, B. Sciascia¹⁹, A. Sciubba^{26,k}, A. Semennikov³², A. Sergi⁴⁷, N. Serra⁴²,
J. Serrano⁶, L. Sestini²³, P. Seyfert²¹, M. Shapkin³⁷, I. Shapoval⁴⁵, Y. Shcheglov³¹, T. Shears⁵⁴,
L. Shekhtman^{36,w}, V. Shevchenko⁶⁸, B.G. Siddi^{17,40}, R. Silva Coutinho⁴², L. Silva de Oliveira²,
G. Simi^{23,o}, S. Simone^{14,d}, M. Sirendi⁴⁹, N. Skidmore⁴⁸, T. Skwarnicki⁶¹, E. Smith⁵⁵,
I.T. Smith⁵², J. Smith⁴⁹, M. Smith⁵⁵, I. Soares Lavra¹, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³,
B. Souza De Paula², B. Spaan¹⁰, P. Spradlin⁵³, S. Sridharan⁴⁰, F. Stagni⁴⁰, M. Stahl¹²,
S. Stahl⁴⁰, P. Stefko⁴¹, S. Stefkova⁵⁵, O. Steinkamp⁴², S. Stemmler¹², O. Stenyakin³⁷,
H. Stevens¹⁰, S. Stoica³⁰, S. Stone⁶¹, B. Storaci⁴², S. Stracka^{24,p}, M.E. Stramaglia⁴¹,
M. Straticiu³⁰, U. Straumann⁴², L. Sun⁶⁴, W. Sutcliffe⁵⁵, K. Swientek²⁸, V. Syropoulos⁴⁴,
M. Szczekowski²⁹, T. Szumlak²⁸, S. T'Jampens⁴, A. Tayduganov⁶, T. Tekampe¹⁰,
G. Tellarini^{17,g}, F. Teubert⁴⁰, E. Thomas⁴⁰, J. van Tilburg⁴³, M.J. Tilley⁵⁵, V. Tisserand⁴,
M. Tobin⁴¹, S. Tolk⁴⁹, L. Tomassetti^{17,g}, D. Tonelli²⁴, S. Topp-Joergensen⁵⁷, F. Toriello⁶¹,
R. Tourinho Jadallah Aoude¹, E. Tournefier⁴, S. Tourneur⁴¹, K. Trabelsi⁴¹, M. Traill⁵³,
M.T. Tran⁴¹, M. Tresch⁴², A. Trisovic⁴⁰, A. Tsaregorodtsev⁶, P. Tsopelas⁴³, A. Tully⁴⁹,
N. Tuning⁴³, A. Ukleja²⁹, A. Ustyuzhanin³⁵, U. Uwer¹², C. Vacca^{16,f}, V. Vagnoni^{15,40},
A. Valassi⁴⁰, S. Valat⁴⁰, G. Valenti¹⁵, R. Vazquez Gomez¹⁹, P. Vazquez Regueiro³⁹, S. Vecchi¹⁷,
M. van Veghel⁴³, J.J. Velthuis⁴⁸, M. Veltri^{18,r}, G. Veneziano⁵⁷, A. Venkateswaran⁶¹,
T.A. Verlage⁹, M. Vernet⁵, M. Vesterinen¹², J.V. Viana Barbosa⁴⁰, B. Viaud⁷, D. Vieira⁶³,
M. Vieites Diaz³⁹, H. Viemann⁶⁷, X. Vilasis-Cardona^{38,m}, M. Vitti⁴⁹, V. Volkov³³,
A. Vollhardt⁴², B. Voneki⁴⁰, A. Vorobyev³¹, V. Vorobyev^{36,w}, C. Voß⁹, J.A. de Vries⁴³,
C. Vázquez Sierra³⁹, R. Waldi⁶⁷, C. Wallace⁵⁰, R. Wallace¹³, J. Walsh²⁴, J. Wang⁶¹,
D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷, D. Websdale⁵⁵, A. Weiden⁴², M. Whitehead⁴⁰,
J. Wicht⁵⁰, G. Wilkinson^{57,40}, M. Wilkinson⁶¹, M. Williams⁴⁰, M.P. Williams⁴⁷, M. Williams⁵⁸,
T. Williams⁴⁷, F.F. Wilson⁵¹, J. Wimberley⁶⁰, M.A. Winn⁷, J. Wishahi¹⁰, W. Wislicki²⁹,
M. Witek²⁷, G. Wormser⁷, S.A. Wotton⁴⁹, K. Wraight⁵³, K. Wyllie⁴⁰, Y. Xie⁶⁵, Z. Xing⁶¹,
Z. Xu⁴, Z. Yang³, Z. Yang⁶⁰, Y. Yao⁶¹, H. Yin⁶⁵, J. Yu⁶⁵, X. Yuan^{36,w}, O. Yushchenko³⁷,
K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, L. Zhang³, Y. Zhang⁷, A. Zhelezov¹², Y. Zheng⁶³, X. Zhu³,
V. Zhukov³³, S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁰Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹³School of Physics, University College Dublin, Dublin, Ireland

¹⁴Sezione INFN di Bari, Bari, Italy

¹⁵Sezione INFN di Bologna, Bologna, Italy

¹⁶Sezione INFN di Cagliari, Cagliari, Italy

¹⁷Sezione INFN di Ferrara, Ferrara, Italy

¹⁸Sezione INFN di Firenze, Firenze, Italy

- ¹⁹ *Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*
- ²⁰ *Sezione INFN di Genova, Genova, Italy*
- ²¹ *Sezione INFN di Milano Bicocca, Milano, Italy*
- ²² *Sezione INFN di Milano, Milano, Italy*
- ²³ *Sezione INFN di Padova, Padova, Italy*
- ²⁴ *Sezione INFN di Pisa, Pisa, Italy*
- ²⁵ *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- ²⁶ *Sezione INFN di Roma La Sapienza, Roma, Italy*
- ²⁷ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ²⁸ *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ²⁹ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁰ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³¹ *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- ³² *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- ³³ *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³⁴ *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- ³⁵ *Yandex School of Data Analysis, Moscow, Russia*
- ³⁶ *Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- ³⁷ *Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁸ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ³⁹ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁰ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴¹ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴² *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴³ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴⁴ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴⁵ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵² *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵ *Imperial College London, London, United Kingdom*
- ⁵⁶ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁰ *University of Maryland, College Park, MD, United States*
- ⁶¹ *Syracuse University, Syracuse, NY, United States*
- ⁶² *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶³ *University of Chinese Academy of Sciences, Beijing, China, associated to ³*
- ⁶⁴ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³*
- ⁶⁵ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³*
- ⁶⁶ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁸*
- ⁶⁷ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹²*
- ⁶⁸ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³²*
- ⁶⁹ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ³⁸*
- ⁷⁰ *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to ⁴³*

^a *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

- ^b*Laboratoire Leprince-Ringuet, Palaiseau, France*
- ^c*P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^d*Università di Bari, Bari, Italy*
- ^e*Università di Bologna, Bologna, Italy*
- ^f*Università di Cagliari, Cagliari, Italy*
- ^g*Università di Ferrara, Ferrara, Italy*
- ^h*Università di Genova, Genova, Italy*
- ⁱ*Università di Milano Bicocca, Milano, Italy*
- ^j*Università di Roma Tor Vergata, Roma, Italy*
- ^k*Università di Roma La Sapienza, Roma, Italy*
- ^l*AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- ^m*LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁿ*Hanoi University of Science, Hanoi, Viet Nam*
- ^o*Università di Padova, Padova, Italy*
- ^p*Università di Pisa, Pisa, Italy*
- ^q*Università degli Studi di Milano, Milano, Italy*
- ^r*Università di Urbino, Urbino, Italy*
- ^s*Università della Basilicata, Potenza, Italy*
- ^t*Scuola Normale Superiore, Pisa, Italy*
- ^u*Università di Modena e Reggio Emilia, Modena, Italy*
- ^v*Iligan Institute of Technology (IIT), Iligan, Philippines*
- ^w*Novosibirsk State University, Novosibirsk, Russia*
- [†]*Deceased*