

This is a repository copy of *The impact of out-of-hours admission on patient mortality: longitudinal analysis in a tertiary acute hospital*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/121945/>

Version: Accepted Version

Article:

Han, Lu orcid.org/0000-0001-7198-3380, Sutton, Matt, Clough, Stuart et al. (2 more authors) (2017) The impact of out-of-hours admission on patient mortality: longitudinal analysis in a tertiary acute hospital. *BMJ Quality & Safety*. pp. 1-11. ISSN 2044-5423

<https://doi.org/10.1136/bmjqs-2017-006784>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

BMJ Quality & Safety

The impact of out-of-hours admission on patient mortality: longitudinal analysis in a tertiary acute hospital

Journal:	<i>BMJ Quality & Safety</i>
Manuscript ID	bmjqs-2017-006784.R2
Article Type:	Original Research
Keywords:	Health services research, Healthcare quality improvement, Patient safety, Statistics

SCHOLARONE™
Manuscripts

Abstract

Background

Emergency hospital admission at weekends is associated with an increased risk of mortality. Previous studies have been limited to examining single years and assessing day – not time – of admission. We used an enhanced longitudinal dataset to estimate the ‘weekend effect’ over time and the effect of night-time admission on all-cause mortality rates.

Method

We examined 246,350 emergency spells from a large teaching hospital in England between April 2004 and March 2014. Outcomes included 7-day, 30-day and in-hospital mortality rates. We conducted probit regressions to estimate the impact on the absolute difference in the risk of mortality of two key predictors: i) admission at weekends (7.00pm Friday to 6.59am Monday); ii) night-time admission (7.00pm to 6.59am). Logistic regressions were used to estimate odds ratios for relative mortality risk differences.

Results

Crude 30-day mortality rate decreased from 6.6% in 2004/05 to 5.2% in 2013/14. Adjusted mortality risk was elevated for all out-of-hours periods. The highest risk was associated with admission at weekend night-times: 30-day mortality increased by 0.6 percentage points (adjusted OR: 1.17, 95%CI: 1.10-1.25), 7-day mortality by 0.5 percentage points (adjusted OR: 1.23, 95%CI: 1.12-1.34), and in-hospital mortality by 0.5 percentage points (adjusted OR: 1.14, 95%CI: 1.08-1.21) compared with admission on weekday day-times. Weekend night-time admission was associated with increased mortality risk in 9 out of 10 years, but this was only statistically significant ($p < 0.05$) in 5 out of 10 years.

Conclusions

There is an increased risk of mortality for patients admitted as emergencies both at weekends and during the night-time. These effects are additive, so that the greatest risk of mortality occurs in patients admitted during the night at weekends. This

1
2
3 increased risk appears to be consistent over time, but the effects are small and are not
4 statistically significant in individual hospitals in every year.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

Introduction

Admission to hospital at weekends is consistently associated with poorer patient outcomes, for both emergency and non-emergency conditions [1-5]. Payers and policy makers have responded to these findings with health care system reform on the assumption that reduced service provision underlies the higher risk of adverse outcomes at weekends, and that more consistent hospital care throughout the week will reduce or remove the disparity [6,7]. In England, evidence on the 'weekend effect' has been used to justify controversial attempts to introduce a 'seven-day service' across the National Health Service (NHS), an approach that is supported by evidence suggesting that emergency weekend admission is not associated with poorer outcomes when there is consistent access to early diagnosis and treatments [8-10].

However, much of the evidence on the effect of weekend admission has been limited in key respects. First, it tends to be cross-sectional and there is therefore limited information on the trends in variations over time. Second, it is often based on routinely available data that provides the day but not the hour of admission. As a result, weekends have effectively been defined as 00.00 am Saturday to 11.59 pm Sunday (thereby excluding Friday evenings and Monday mornings) and separate analysis of night-time admissions has not been possible. This is an important omission; if adverse patient outcomes are related to the different working practices and staff availability during weekends, then patients admitted during other out-of-hours periods are likely to be similarly affected. Some studies have utilised specialist data to address this issue, for example finding that at night there is an increased risk of mortality following in-hospital cardiac arrest [11] or coronary artery bypass grafting [12]; an increased risk of in-hospital mortality if admitted to or discharged from ICU units [13]; and a reduced chance of receiving timely intervention following stroke [14,15].

For this study, we analysed emergency admissions to Salford Royal Foundation Trust (SRFT), a large teaching hospital in the Northwest of England providing a complete range of acute services, including specialist tertiary care. SRFT has maintained detailed electronic patient records since 2004, including precise time of admission, and has been a pioneer of extending normal hours of operation and providing

1
2
3 enhanced weekend services. In 2011 it opened an 'emergency village', providing a
4 consultant-led enhanced emergency service seven days a week, and also extended
5 services such as radiology, pathology and pharmacy across the weekend.
6
7

8
9
10 Using the enhanced data available from SRFT, we examined disparities in patient
11 outcomes associated with out-of-hours admission across the full range of clinical
12 specialties. We aimed to answer three key questions: i) What is the size of the
13 weekend effect when weekends are more appropriately defined (i.e. 7.00 pm Friday to
14 6.59 am Monday)? ii) Does the effect vary over time? iii) Is there a 'night-time'
15 effect, with increased risk of all-cause mortality for patients admitted at night-time
16 compared with daytime?
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Data sources and variables

We extracted patient records from SRFT for 10 financial years, from April 2004 to March 2014. This dataset included similar information to nationally available Hospital Statistics of Episode (HES) data: age, sex, time of admission, method and source of admission, primary and secondary diagnoses, specialty, procedures, time of discharge, destination on discharge, consultant codes, and the dates of in- and out-of-hospital deaths. The original data were at consultant episode level (the period when a patient was under the treatment of a consultant) and we created hospital spells (the continuous stay in a single hospital from admission to discharge) by linking episode records from the time of admission to the time of discharge. We excluded maternity admissions throughout to facilitate temporal comparisons, as maternity services ceased to be provided in November 2011. We also excluded patients aged 16 years and under as SRFT does not provide paediatric services (maternal and paediatric cases attending A&E area were assessed and stabilised before being transferred to other treatment centres). For patients with multiple emergency admissions within the last 30 days of life we excluded all but the first admission. The final sample included 246,350 emergency spells, defined as non-elective admissions.

Our primary outcome variable was mortality within 30 days of admission (either in or out of hospital). Sub-analyses of 7-day and all in-hospital (within stay) mortality were conducted for robustness checks. Our main exposure variable was weekend admission. Our dataset included the minute of admission which allowed us to categorise admissions from 7.00 pm on one day to 6.59 am the next day as 'night' and admissions from 7.00 pm on Friday to 6.59 am on Monday as 'weekend', a more precise definition than previous studies, reflecting senior clinicians' working patterns [16]. We expected public holidays to have similar service arrangement as weekends, and we therefore identified emergency admissions on 56 public holidays in the 10-year period and grouped these with weekend admissions. To do this we firstly identified those holidays immediately before or after a weekend and linked the weekend and the holiday as a holiday period. Second, we included in all the public holidays the evening of the day before (from 7.00 pm to 11.59 pm) and the morning of the day after (from 0.00 am to 6.59 am). Patients admitted outside weekends or

1
2
3 holidays were categorised as ‘weekday’ and analysed as a comparison group. Our
4 secondary exposure was all admissions during out-of-hours periods including
5 weekends and the nights of weekdays. We categorised patients by weekday daytime
6 admission (7.00 am - 6.59 pm, Monday to Friday), weekday night-time admission
7 (7.00 pm - 6.59 am, Monday to Thursday), weekend daytime admission (7.00 am -
8 6.59 pm Saturday and Sunday), and weekend night-time admission (7.00 pm – 6.59
9 am, Friday to Sunday). Patients admitted during normal working hours (weekday –
10 day group) were treated as the reference category.
11
12
13
14
15
16

17
18 We adjusted for patient case-mix by including variables available in the
19 administrative hospital data. Collectively these variables provide a good fit for
20 modelling the risk of mortality [17], although they do not include direct measures of
21 severity of illness. We adjusted for patient demographics using age categories on
22 admission, gender, and ethnicity. Socioeconomic characteristics were measured using
23 area deprivation for place of residence in quintiles, based on the 2010 Index of
24 Multiple Deprivation (IMD) at Lower Layer Super Output Area (LSOA) level [18].
25 For each patient we calculated a Charlson Index score to account for the presence and
26 the severity of comorbidities [19]. We updated the weights of included conditions
27 according to their association with the risk of mortality estimated using recent data
28 from the UK [20]. We made further adjustments for patient complexity using
29 indicators for primary diagnosis summary groups [21], the total number of different
30 diagnoses, the total number of different procedures during the admission, the use of
31 palliative care during the spell, and the total number of emergency admissions in the
32 year prior to the index admission date. We also controlled for the method and source
33 of admission, using the most common method or source as the reference category. We
34 identified those patients discharged to other health care providers (both public and
35 private) and adjusted this for 30-day and 7-day mortalities (this variable does not
36 predict in-hospital death). We accounted for seasonal impact by including dummy
37 variables for the month of admission. In the pooled analysis, we included year
38 dummies to capture unobserved factors that varied by financial year.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 *Statistical analysis*

55 We fitted probit models to estimate the extent to which the probability of death was
56 associated with our exposure variables, after controlling for observed patient
57
58
59
60

characteristics. In the pooled analysis on the impact of out-of-hours admission, we assumed a latent propensity of death as a linear function of our exposure and controlling variables as outlined in equation (1)

$$y_i^* = \alpha + \beta_1 \text{out-of-hours}_i + \beta_2 X_i + \gamma_{year} + \delta_{month} + \varepsilon_i \quad (1),$$

where y_i^* was the estimated propensity of death for spell i ; *out-of-hours* _{i} was a categorical variable indicating the type of out-of-hours period for admission i ; X_i was a vector of variables measuring patient demographics and complexity; γ_{year} was a vector of dummies controlling for unobserved year-specific fixed effect; δ_{month} was a vector of admission months adjusting for seasonal impact; the error term ε_i was assumed to follow a standard normal distribution and was independent from the explanatory variables. The observed outcome for admission i , Y_i depended on the value of y_i^* , so that

$$Y_i = \begin{cases} 1 & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases} \quad (2).$$

For the analysis of individual financial year, we removed the year dummies, γ_{year} , from equation (1), allowing the coefficients of explanatory variables to vary with time. We applied variance-covariance matrices clustered around consultant (first episode within each spell) to account for any correlation within management by the same consultant. In this non-linear model, the estimated coefficient of out-of-hours admission should not be interpreted as its effect size, because this also depends on the values of other predictors in the model. We therefore reported the average marginal effects (AME) which took into account the values of other predictors across the sample and computed the average changes in the risk of mortality associated with out-of-hours admission in absolute terms. We also estimated the same risk of mortality models using logistic regressions and reported odds ratios to show the relative differences in the mortality risk for comparability with previous studies. All analyses were conducted using STATA Version 14.2.

Results

There were a total of 246,350 emergency admissions for the study period from April 2004 to March 2014. 5.9% of admissions resulted in death within 30 days, 2.6% in death within 7 days, and 4.6% in death within the hospital stay. Crude 30-day mortality rates decreased over time, from 6.6% in 2004/5 to 5.2% in 2013/14 (**table A1 and figure A1, appendix**).

About one third of spells (81,621) were admitted during weekends and holidays. The demographic composition of weekend and weekday patients groups was similar (**table 1**). For both groups, average age was around 58.2 years, just under 50% of patients were male and 90% were white. The proportion of patients who lived in the most deprived fifth of areas was slightly higher for the weekend admission group (18.3% for weekend compared with 17.9% for weekday). Both groups were similar in terms of complexity: on average, each hospital stay had 6 different diagnoses, 1.5 procedures performed, one emergency admission in the year prior to the index admission date, and a Charlson index score of 5.

Patients admitted at night were on average younger, and more likely to be male and from the most deprived fifth of areas compared with patients admitted during the day. Patients admitted at night also had on average more diagnosis codes and previous emergency admissions per spell than patients admitted during the day, although the absolute differences were marginal.

Crude mortality for out-of-hours admissions

There were substantial differences in mortality rates between patients admitted at weekends compared with weekdays. The crude 30-day mortality rate for weekend admissions was 12.3% higher in relative terms than for weekday admissions (rates of 6.4% and 5.7% respectively); the 7-day mortality rate was 20.8% higher (rates of 2.9% and 2.4% respectively); and the in-hospital mortality rate was 11.1% higher (rates of 5.0% and 4.5% respectively) (**table 1**).

Compared to normal working hours (daytime, weekdays) crude mortality rates were higher for patients admitted during all out-of-hours periods. Rates were highest for

1
2
3 weekend daytimes. Compared with baseline crude 30-day mortality rates of 5.7%,
4 mortality rates for patients admitted at weekend daytime were 17.5% higher (at 6.7%)
5 and rates for weekend nighttime were 7.0% higher (at 6.1%). We observed similar
6 patterns for 7-day and in-hospital mortality.
7
8

9
10
11 *Table 1 about here.*
12

13
14
15 *Adjusted mortality for out-of-hours admissions*

16 **Table 2** summarises the average marginal effects for the key predictors of mortality.
17 Higher mortality was associated with increasing age, comorbidity, number of
18 diagnoses, and socioeconomic deprivation. Mortality was also higher for patients who
19 were male, white (compared to black and Asian), received palliative care, or were
20 admitted via a GP. Mortality was strongly associated with year of admission, with the
21 risk declining over time.
22
23

24
25
26
27
28 After adjusting for other factors, admission on weekends and public holidays had an
29 independent and statistically significant association with 30-day mortality, with an
30 estimated average marginal effect of 0.4 percentage points across the study period
31 (adjusted OR: 1.10, 95%CI: 1.06-1.15, **table A2, appendix**). The risk of both 7-day
32 and in-hospital mortality increased by 0.3 percentage points for admission on
33 weekends and holidays (7-day adjusted OR: 1.12, 95%CI: 1.07-1.18; in-hospital
34 adjusted OR: 1.08, 95%CI: 1.02-1.15), compared with weekday admission.
35 Sensitivity analyses suggested that the effects of case-mix variables were robust
36 across the three outcome measures.
37
38
39
40
41
42
43
44

45 *Table 2 about here.*
46

47
48 In the pooled analysis, all out-of-hours admission periods were associated with higher
49 risk of mortality after adjusting for confounders, compared with admission during
50 normal working hours (**table 3**). Weekend night-time admission had the greatest
51 impact on mortality: patients in this group had a 0.6 percentage points higher risk of
52 death within 30 days (adjusted OR: 1.17, 95%CI: 1.10-1.25, **table A3, appendix**).
53 Weekend daytime and weekday night-time admission was also associated with
54 increased risk of mortality within 30 days (weekend-day AME: 0.4 percentage points,
55
56
57
58
59
60

1
2
3 adjusted OR: 1.11, 95%CI: 1.04-1.18; weekday-night AME: 0.3 percentage points,
4 adjusted OR: 1.09, 95%CI: 1.01-1.17). Results for 7-day and in-hospital mortality
5 were similar: patients admitted at weekend night-times had the highest risk of death,
6 followed by weekend daytime and weekday night-time admissions.
7
8
9

10
11 *Table 3 about here*
12

13
14
15 *Variation in impact over time*

16 The impact of weekend admission by financial year is reported in **table 4**. After case-
17 mix adjustment, weekend admission was positively associated with 30-day mortality
18 in nine out ten years, however this association was only statistically significant
19 (p<0.05) in three out of the ten years. The estimated average marginal effect was 0.6
20 percentage points for 2005/06 (adjusted OR: 1.17, 95%CI: 1.02-1.34, **table A4,**
21 **appendix**), 0.6 percentage points for 2008/09 (adjusted OR: 1.14, 95%CI: 1.02-1.27),
22 and 0.9 percentage points for 2012/13 (adjusted OR: 1.29, 95%CI: 1.15-1.45). 7-day
23 mortality rates were elevated for weekend and holiday admissions in 2005/06,
24 2011/12 and 2012/13 (p<0.05), whereas in-hospital mortality was not significantly
25 associated with weekend admission in any year.
26
27
28
29
30
31
32
33

34 Results for out-of-hours admission by financial year are given in **table 4**. Admission
35 during all three out-of-hours periods was associated with increased adjusted risk of
36 mortality compared to weekday daytime admission in most years, although the pattern
37 and size of impact varied and was not always statistically significant. Consistent with
38 the pooled analysis, weekend night-time admission was associated with the greatest
39 increased risk of mortality: there was a significantly increased risk of 30-day mortality
40 in 2004/05, 2006/07, 2008/09, 2011/12 and 2012/13. There was a significantly
41 increased risk of mortality for weekend daytime admissions in 2012/13 and for
42 weekday night-time admissions in 2004/05, 2006/07 and 2008/09.
43
44
45
46
47
48
49
50
51
52

53 *Table 4 about here.*
54
55
56
57
58
59
60

Discussion

In this study we analysed a quarter of a million emergency admissions to a large teaching hospital over a ten-year period. As with previous studies, we found that crude mortality rates were higher for patients admitted at weekends and holidays, compared with patients admitted on weekdays. We also found that risk-adjusted mortality rates were higher not only for patients admitted at weekends but also for patients admitted at night-time. Compared with weekday daytime admissions, adjusted risk of 30-day mortality was 0.3 percentage points higher for weekday night-time admissions (adjusted OR: 1.09), 0.4 percentage points higher for weekend daytime admissions (adjusted OR: 1.11) and 0.6 percentage points higher for weekend night-time admissions (adjusted OR: 1.17).

Strengths and weaknesses

This study addressed limitations in the existing literature in two main respects. First, by using exact time of admission we could define weekends (and holidays) more precisely by including Friday evenings and Monday mornings, and could examine night-time admissions separately from daytime admissions for all clinical groups. The effect of night-time admission has been analysed mainly in disease-specific studies [11-15], with only few studies based on hospital-level data [22]. Second, we were able to analyse the impact of our exposure variables longitudinally over a 10-year period, and could quantify impacts in each financial year. We were therefore able to relax the assumption that risk factors have constant effects over time.

The study was subject to several limitations. First, it was based on patient records from a single hospital in the Northwest of England, albeit one that has been a pioneer in data collection and quality improvement. Heterogeneities between hospitals in terms of size, performance, financial status, and patient demographics could affect the generalizability of our results. However, the focus on a single provider also confers advantages, including relatively consistent coding practices and population characteristics. The availability of longitudinal data for this provider enabled us to examine the effect of out-of-hours over time, but the restricted sample size limited the power of our study to detect differences in outcomes between groups, particularly for analyses of individual years. Second, we adjusted the risk of mortality for admission

1
2
3 method and source but could not adjust for discharge destination due to the high
4 correlation with the dependent variable (mortality). We therefore only accounted for
5 the patients discharged to health providers (both public and private) and compared
6 them with patients discharged to other destinations. Third, as with previous studies
7 based on analysis of routinely collected hospital data, we had no direct measure of
8 illness severity and our risk adjustment models were likely to be confounded by
9 unmeasured severity. With the available information, we constructed several variables
10 to adjust for patient complexity, including the Charlson index score, the numbers of
11 diagnoses and procedures in a spell. These variables might be under-recorded and less
12 accurately coded during out-of-hours periods, which could create a potential bias.
13
14
15
16
17
18
19
20

21 *Main findings*

22 Mortality rates for emergency admissions throughout the week declined over time,
23 reflecting the wider fall in national mortality rates – attributable to net improvements
24 in living conditions, lifestyles and healthcare – over the study period [23]. In the
25 context of these falling mortality rates over time, we found a persistently increased
26 risk of mortality for patients admitted to hospital outside of normal hours – a pattern
27 that has previously been found for stroke admissions [24]. It might be expected that as
28 overall mortality rates fall, rates for different time periods during the week would
29 converge. We found no such temporal trend, but the smaller sample sizes for
30 individual years means that our study may have been under-powered to detect one.
31 The reasons for both the improvements in survival for overall emergency admissions
32 and the apparent persistence of elevated mortality rates for out-of-hours admissions
33 warrant further investigation. As with the weekend effect itself, improving survival
34 for admitted patients over time may reflect both changes in the admitted population
35 and differences in quality of care, but the explanation must be consistent with the
36 persistence of an increased risk of mortality for patients admitted out-of-hours. In the
37 context of SRFT, the introduction of enhanced emergency services at weekends in
38 2011 had no apparent effect on excess mortality rates for out-of-hours admissions,
39 although we did not formally test for such an effect. This has implications for current
40 initiatives to introduce seven day services across the NHS and the potential impact of
41 these service changes on out-of-hours mortality [25].
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Our finding of increased risk of mortality at night-times across the full range of
4 emergency admissions suggests that more patients are affected by service variations
5 across the week than has been estimated in previous studies based on
6 weekend/weekday comparisons [1-5]. This finding is consistent with the results of
7 studies on outcomes for cardiovascular disease [11-15], which suggest that patient
8 outcomes are worse for acute events and invasive procedures occurring at night, not
9 just at weekends. However, as with previous studies we cannot exclude the possibility
10 that reduced capacity out-of-hours leads to the selection of a sicker patient population,
11 and that this explains increased risk of mortality for both weekend and night-time
12 admissions. Evidence from previous studies suggests there is a higher threshold for
13 admission at weekends [26] and that increased risk of mortality for out-of-hours
14 admissions is reduced after adjusting for proxy [27] measures of severity. If similar
15 patterns apply at night-time, this would suggest that our findings reflect a reduced
16 capacity for the hospital to admit less severely ill patients at nights and weekends,
17 rather than a reduced level of care at these times.
18
19
20
21
22
23
24
25
26
27
28
29

30 *Conclusion*

31 In addition to patients admitted to hospital as an emergency at the weekend, patients
32 admitted at night also experience higher mortality rates. These 'weekend' and 'night-
33 time' effects are additive, with the highest risk of death for patients admitted at night
34 on weekends. Recent UK policy has been moving towards the creation of a seven-day
35 service [28], in part to address the perceived increase risk of adverse outcomes in
36 patients admitted to hospital at weekends. This has generated conflict between a
37 government attempting to increase weekend staffing levels whilst containing costs,
38 and a medical profession resistant to these changes [29]. For government policy to be
39 consistent, it would also need to address adverse outcomes across all out-of-hours
40 periods, including night-times on weekdays. Extending normal hours of operation
41 could be beneficial if it leads to improved access and better patient outcomes, but
42 clear evidence on cost-effectiveness will be needed to justify such major system
43 changes.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Freemantle N, Richardson M, Wood J, et al. Weekend hospitalization and additional risk of death: An analysis of inpatient data. *Journal of the Royal Society of Medicine*. 2012;105(2):74-84.
2. Aylin P, Alexandrescu R, Jen M, et al. Day of week of procedure and 30 day mortality for elective surgery: retrospective analysis of hospital episode statistics. *British Medical Journal*. 2013;346:f2424.
3. McIsaac DI, Bryson GL, van Walraven C. Elective, Major Noncardiac Surgery on the Weekend A Population-based Cohort Study of 30-day Mortality. *Medical Care*. 2014;52(6):557-64.
4. Concha OP, Gallego B, Hillman K, et al. Do variations in hospital mortality patterns after weekend admission reflect reduced quality of care or different patient cohorts? A population-based study. *BMJ Quality & Safety*. 2014;23(3):215-22.
5. Freemantle N, Ray D, McNulty D, et al. Increased mortality associated with weekend hospital admission: a case for expanded seven day services? *British Medical Journal*. 2015;351:h4596.
6. National Advisory Group on the Safety of Patients in England. *A Promise to Learn – A Commitment to Act: Improving the Safety of Patients in England*. London: Crown Copyright, 2013.
7. Royal College of Paediatrics and Child Health. *Consultant Delivered Care: An Evaluation of New Ways of Working in Paediatrics*. London: RCPCH, 2012.
8. Schmulewitz L, Proudfoot A, Bell D. The impact of weekends on outcome for emergency patients. *Clinical Medicine*. 2005;5(6):621-5.
9. Albright K, Raman R, Ernstrom K, et al. Can comprehensive stroke centers erase the 'weekend effect'? *Cerebrovascular Diseases*. 2009;27(2):107-13.
10. McKinney J, Deng Y, Kasner S, et al. Myocardial Infarction Data Acquisition System Study Group. Comprehensive stroke centers overcome the weekend versus weekday gap in stroke treatment and mortality. *Stroke*. 2011;42(9):2403-9.
11. Robinson E, Smith G, Power G, et al. Risk-adjusted survival for adults following in-hospital cardiac arrest by day of week and time of day: observational cohort study. *BMJ Quality & Safety*. 2015; doi:10.1136/bmjqs-2015-004223.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
12. Coumbe A, John R, Kuskowski M, et al. Variation of mortality after coronary artery bypass surgery in relation to hour, day and month of procedure. *BMC Cardiovascular Disorders*. 2011;11:63.
 13. Laupland KB, Shahpori R, Kirkpatrick AW, et al. Hospital mortality among adults admitted to and discharged from intensive care on weekends and evenings. *Journal of Critical Care*. 2008; 23:317–324
 14. Turner M, Barber M, Dodds H, et al. Stroke patients admitted within normal working hours are more likely to achieve process standards and to have better outcomes. *Journal of Neurology, Neurosurgery and Psychiatry*. 2016;87:138-143.
 15. Bray B, Cloud G, James M, et al. Weekly variation in health-care quality by day and time of admission: a nationwide, registry-based, prospective cohort study of acute care. *Lancet* 2016; doi:10.1016/S0140-6736(16)30443-3
 16. Bland D. *BMA Survey of Consultant Working Patterns and on Call Services*. London: BMA, 2014.
 17. Bottle A, Gaudoin R, Goudie R, et al. Can valid and practical risk-prediction or case-mix adjustment models, including adjustment for comorbidity, be generated from English hospital administrative data (Hospital Episode Statistics)? A national observational study. *Health Services and Delivery Research*. 2014; 2(40).
 18. Department of Communities and Local Government. *The English Indices of Deprivation 2010*. Accessed from: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2010>
 19. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Disease*. 1987;40(5):373-83.
 20. Aylin P, Bottle A, Jen M, et al. *HSMR Mortality Indicators (Technical Document)*. 2010. Accessed from: <https://www1.imperial.ac.uk/resources/3321CA24-A5BC-4A91-9CC9-12C74AA72FDC/>
 21. Hospital Episode Statistics Analysis, Health and Social Care Information Centre. *Hospital Episode Statistics – Admitted Patient Care 2012-13*. 2013.

- 1
2
3 22. Coiera E, Wang Y, Magrabi F, et al. Predicting the cumulative risk of death
4 during hospitalization by modeling weekend, weekday and diurnal mortality
5 risks. *BMC Health Services Research*. 2014; 14:226
6
7
8 23. Office for National Statistics. *Mortality in the United Kingdom 1983-2013*.
9 2014. Available from:
10 <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/mortalityintheunitedkingdom/19832013>. Accessed 9 July
11 2017.
12
13 24. McKinney J, Deng Y, Kasner S, et al. Comprehensive stroke centers
14 overcome the weekend versus weekday gap in stroke treatment and mortality.
15 *Stroke*. 2011;42: 2403–2409.
16
17 25. NHS Improvement. *Seven Day Services in the NHS*. Available from:
18 <https://improvement.nhs.uk/resources/seven-day-services/> . Accessed 11 July
19 2017.
20
21 26. Meacock R, Anselmi L, Kristensen SR, et al. Higher mortality rates amongst
22 emergency patients admitted to hospital at weekends reflect a lower
23 probability of admission. *Journal of Health Services Research &*
24 *Policy*.2016;DOI:10.1177/1355819616649630.
25
26 27. Anselmi L, Meacock R, Kristensen SR, Doran T, Sutton M. Arrival by
27 ambulance explains variation in mortality by time of admission: retrospective
28 study of admissions to hospital following emergency department attendance in
29 England. *BMJ Quality & Safety*. 2016: 10.1136/bmjqs-2016-005680.
30
31 28. NHS England, *Seven Days a Week Forum. Summary of Initial Findings*.2013.
32 Available from:
33 <https://www.england.nhs.uk/wp-content/uploads/2013/12/forum-summary-report.pdf> . Accessed 23 June 2016.
34
35 29. Goddard A. Lessons to be learned from the UK Junior Doctors' Strike. *JAMA*.
36 2016: 10.1001/jama.2016.12029.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Patient characteristics by admission time, 2004/05-2013/14.

Admission time of the week	Weekday 7am Mon - 6.59pm Fri	Weekend ¹ 7pm Fri - 6.59am Mon	Weekday-day 7am - 6.59pm Mon - Fri	Weekday-night 7pm - 6.59am Mon - Thu	Weekend-day ¹ 7am - 6.59pm Sat & Sun	Weekend-night ¹ 7pm - 6.59am Fri - Sun
Admissions, n (%)	164,729 (66.87%)	81,621 (33.13%)	101,725 (41.29%)	63,004 (25.57%)	33,409 (13.56%)	48,212 (19.57%)
Age, mean (SD)	58.26 (21.17)	58.18 (21.82)	58.84 (20.93)	57.33 (21.53)	59.91 (21.69)	56.97 (21.84)
Gender, male (%)	79,736 (48.40%)	40,215 (49.27%)	48,742 (47.92%)	30,994 (49.19%)	16,264 (48.68%)	23,951 (49.68%)
Ethnicity, white (%)	148,215 (89.98%)	73,069 (89.52%)	91,784 (90.23%)	56,431 (89.57%)	29,964 (89.69%)	43,105 (89.41%)
Deprivation, most deprived quintile (%)	29,403 (17.85%)	14,973 (18.34%)	17,694 (17.39%)	11,709 (18.58%)	5,884 (17.61%)	9,089 (18.85%)
Diagnosis per spell, mean (SD)	5.96 (4.20)	6.10 (4.19)	5.85 (4.20)	6.13 (4.19)	6.07 (4.22)	6.12 (4.17)
Procedures per spell, mean (SD)	1.47 (2.50)	1.45 (2.54)	1.46 (2.46)	1.47 (2.56)	1.47 (2.51)	1.43 (2.55)
Previous emergency admission², mean (SD)	0.95 (2.03)	1.01 (2.16)	0.89 (1.94)	1.04 (2.17)	0.97 (2.06)	1.03 (2.22)
Previous emergency admission³, mean (SD)	2.46 (2.64)	2.60 (2.80)	2.36 (2.54)	2.60 (2.77)	2.53 (2.67)	2.64 (2.89)
Charlson index, mean (SD)	5.03 (8.14)	5.10 (8.22)	5.01 (8.10)	5.08 (8.21)	5.31 (8.34)	4.96 (8.14)
Death in 30 days, n (%)	9,366 (5.69%)	5,180 (6.35%)	5,765 (5.67%)	3,601 (5.72%)	2,241 (6.71%)	2,939 (6.10%)
Death in 7 days, n (%)	3,965 (2.41%)	2,368 (2.90%)	2,319 (2.28%)	1,646 (2.61%)	1,016 (3.04%)	1,352 (2.80%)
Death in hospital, n (%)	7,373 (4.48%)	4,054 (4.97%)	4,525 (4.45%)	2,848 (4.52%)	1,764 (5.28%)	2,290 (4.75%)

1. Including public holidays.

2. In 365 days prior to the admission date.

3. For those having at least one emergency admission in the last 365 days.

Table 2 Adjusted risk of mortality 2004/05-2013/14, probit regressions with average marginal effects.

Variables	30-day mortality		7-day mortality		In-hospital mortality	
	AME ²	Std.Err. ³	AME	Std.Err.	AME	Std.Err.
Admission time						
Weekday	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Weekend ¹	0.004***	(0.001)	0.003***	(0.001)	0.003***	(0.001)
Case-mix variables						
Age band 17-25	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Age band 26-35	-0.001	(0.002)	-0.001	(0.001)	-0.003	(0.002)
Age band 36-45	0.005**	(0.002)	0.001	(0.001)	0.001	(0.002)
Age band 46-55	0.012***	(0.002)	0.005***	(0.001)	0.005**	(0.002)
Age band 56-65	0.025***	(0.003)	0.012***	(0.002)	0.015***	(0.003)
Age band 66-75	0.033***	(0.003)	0.015***	(0.002)	0.022***	(0.003)
Age band 76-85	0.051***	(0.003)	0.023***	(0.002)	0.038***	(0.003)
Age band 85+	0.079***	(0.004)	0.033***	(0.003)	0.067***	(0.005)
Gender-male	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Gender-female	-0.003***	(0.001)	-0.001	(0.001)	-0.001	(0.001)
Gender-not stated	-0.005	(0.014)	-0.001	(0.011)	0.001	(0.016)
Ethnicity-White	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Ethnicity-mixed	-0.013	(0.009)	-0.001	(0.006)	-0.010	(0.008)
Ethnicity-Asian	-0.015***	(0.004)	-0.007***	(0.002)	-0.013***	(0.003)
Ethnicity-Black	-0.016***	(0.005)	-0.011***	(0.003)	-0.015***	(0.005)
Ethnicity-other	0.016***	(0.005)	0.013***	(0.004)	0.018***	(0.005)
Ethnicity-not stated	0.032***	(0.007)	0.022***	(0.005)	0.030***	(0.006)
IMD quintile-1 (most affluent)	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
IMD quintile-2	0.001	(0.002)	-0.000	(0.001)	-0.000	(0.001)
IMD quintile-3	0.002*	(0.001)	0.002	(0.001)	0.001	(0.001)
IMD quintile-4	0.002	(0.001)	-0.000	(0.001)	-0.000	(0.001)

1							
2							
3							
4							
5	IMD quintile-5 (most deprived)	0.004***	(0.001)	0.002*	(0.001)	0.003**	(0.001)
6	IMD quintile-missing	0.012***	(0.002)	0.005***	(0.001)	0.010***	(0.002)
7							
8	Charlson index	0.002***	(0.000)	0.001***	(0.000)	0.001***	(0.000)
9							
10	No. of diagnosis	0.002***	(0.000)	-0.000	(0.000)	0.003***	(0.000)
11	No. of procedures	-0.001*	(0.000)	-0.002***	(0.000)	0.001***	(0.000)
12	Palliative care	0.150***	(0.013)	0.038***	(0.005)	0.107***	(0.009)
13	No. of emergency admissions in the previous 1 year (365 days)	-0.002***	(0.000)	-0.000**	(0.000)	-0.001***	(0.000)
14							
15	Admission method-emergency A+E	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
16	Admission method-emergency transfer from other provider	-0.015***	(0.003)	-0.013***	(0.002)	-0.014***	(0.003)
17	Admission method-emergency domicile	-0.009	(0.017)	0.000	(0.000)	-0.026***	(0.009)
18	Admission method-emergency GP refer	0.006**	(0.002)	-0.007***	(0.002)	0.002	(0.003)
19	Admission method-emergency outpatient	-0.018***	(0.003)	-0.018***	(0.002)	-0.017***	(0.003)
20	Admission method-emergency ante natal	0.025	(0.026)	0.010	(0.025)	0.000	(0.000)
21	Admission method-emergency post natal	0.000	(0.000)	0.000	(0.000)	0.000	(0.000)
22	Admission method-non-emergency transfer from other provider	-0.015***	(0.004)	-0.015***	(0.003)	-0.016***	(0.004)
23							
24	Transfer to other hospitals	-0.020***	(0.005)	-0.037***	(0.004)		
25							
26							
27	Financial year 2004/05	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
28	Financial year 2005/06	-0.012***	(0.004)	-0.002	(0.002)	-0.015***	(0.004)
29	Financial year 2006/07	-0.011***	(0.003)	-0.003*	(0.002)	-0.020***	(0.003)
30	Financial year 2007/08	-0.010**	(0.004)	0.001	(0.003)	-0.023***	(0.006)
31	Financial year 2008/09	-0.023***	(0.004)	-0.005	(0.003)	-0.042***	(0.005)
32	Financial year 2009/10	-0.029***	(0.005)	-0.006*	(0.003)	-0.048***	(0.005)
33	Financial year 2010/11	-0.035***	(0.004)	-0.009**	(0.004)	-0.059***	(0.005)
34	Financial year 2011/12	-0.039***	(0.004)	-0.010***	(0.004)	-0.064***	(0.005)
35	Financial year 2012/13	-0.037***	(0.004)	-0.009***	(0.003)	-0.062***	(0.005)
36	Financial year 2013/14	-0.042***	(0.004)	-0.011***	(0.004)	-0.067***	(0.005)
37							
38	Diagnosis group dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
39	Admission source dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
40	Admission month dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
41							
42							
43							
44							
45							
46							
47							
48							
49							

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

C statistic	0.88	0.87	0.90
Pseudo R2	0.28	0.23	0.31
Observations	241,338	237,091	242,693

-
- 1. Including public holidays.
 - 2. Average marginal effect. *** p<0.01, ** p<0.05, * p<0.1.
 - 3. Robust standard errors corrected for clustering around consultant in parentheses.

Confidential: For Review Only

Table 3 Adjusted risk of mortality 2004/05-2013/14, probit regressions with average marginal effects.

Variables	30-day mortality		7-day mortality		In-hospital mortality	
	AME ²	Std.Err. ³	AME	Std.Err.	AME	Std.Err.
Admission time						
Weekday day	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Weekday night	0.003***	(0.001)	0.003***	(0.001)	0.003***	(0.001)
Weekend day ¹	0.004***	(0.001)	0.003***	(0.001)	0.003**	(0.002)
Weekend night ¹	0.006***	(0.001)	0.005***	(0.001)	0.005***	(0.001)
Case-mix variables						
Age band 17-25	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Age band 26-35	-0.001	(0.002)	-0.001	(0.001)	-0.003	(0.002)
Age band 36-45	0.005**	(0.002)	0.001	(0.001)	0.001	(0.002)
Age band 46-55	0.012***	(0.002)	0.005***	(0.001)	0.005**	(0.002)
Age band 56-65	0.025***	(0.003)	0.012***	(0.002)	0.015***	(0.003)
Age band 66-75	0.033***	(0.003)	0.016***	(0.002)	0.022***	(0.003)
Age band 76-85	0.051***	(0.003)	0.023***	(0.002)	0.038***	(0.003)
Age band 85+	0.080***	(0.004)	0.033***	(0.003)	0.067***	(0.005)
Gender-male	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Gender-female	-0.003***	(0.001)	-0.001	(0.001)	-0.001	(0.001)
Gender-not stated	-0.005	(0.014)	-0.001	(0.011)	0.001	(0.016)
Ethnicity-White	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Ethnicity-mixed	-0.013	(0.009)	-0.001	(0.006)	-0.010	(0.008)
Ethnicity-Asian	-0.015***	(0.004)	-0.007***	(0.002)	-0.013***	(0.003)
Ethnicity-Black	-0.016***	(0.005)	-0.011***	(0.003)	-0.015***	(0.005)
Ethnicity-other	0.016***	(0.005)	0.013***	(0.004)	0.018***	(0.005)
Ethnicity-not stated	0.032***	(0.007)	0.022***	(0.005)	0.030***	(0.006)
IMD quintile-1 (most affluent)	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
IMD quintile-2	0.001	(0.002)	-0.000	(0.001)	-0.000	(0.001)

IMD quintile-3	0.002*	(0.001)	0.002	(0.001)	0.001	(0.001)
IMD quintile-4	0.002	(0.001)	-0.000	(0.001)	-0.000	(0.001)
IMD quintile-5 (most deprived)	0.004***	(0.001)	0.002*	(0.001)	0.003**	(0.001)
IMD quintile-missing	0.012***	(0.002)	0.005***	(0.001)	0.010***	(0.002)
Charlson index	0.002***	(0.000)	0.001***	(0.000)	0.001***	(0.000)
No. of diagnosis	0.002***	(0.000)	-0.000	(0.000)	0.003***	(0.000)
No. of procedures	-0.001*	(0.000)	-0.002***	(0.000)	0.001***	(0.000)
Palliative care	0.150***	(0.013)	0.038***	(0.005)	0.107***	(0.009)
No. of emergency admissions in the previous 1 year (365 days)	-0.002***	(0.000)	-0.000**	(0.000)	-0.001***	(0.000)
Admission method-emergency A+E	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Admission method-emergency transfer from other provider	-0.015***	(0.003)	-0.013***	(0.002)	-0.013***	(0.003)
Admission method-emergency domicile	-0.008	(0.017)	0.000	(0.000)	-0.025***	(0.009)
Admission method-emergency GP refer	0.007***	(0.002)	-0.006***	(0.002)	0.003	(0.003)
Admission method-emergency outpatient	-0.017***	(0.003)	-0.017***	(0.002)	-0.016***	(0.003)
Admission method-emergency ante natal	0.027	(0.026)	0.011	(0.025)	0.000	(0.000)
Admission method-emergency post natal	0.000	(0.000)	0.000	(0.000)	0.000	(0.000)
Admission method-non-emergency transfer from other provider	-0.014***	(0.004)	-0.014***	(0.003)	-0.015***	(0.004)
Transfer to other hospitals	-0.020***	(0.005)	-0.037***	(0.004)		
Financial year 2004/05	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Financial year 2005/06	-0.012***	(0.004)	-0.002	(0.002)	-0.015***	(0.004)
Financial year 2006/07	-0.011***	(0.003)	-0.003*	(0.002)	-0.020***	(0.003)
Financial year 2007/08	-0.010**	(0.004)	0.001	(0.003)	-0.023***	(0.006)
Financial year 2008/09	-0.023***	(0.004)	-0.005	(0.003)	-0.042***	(0.005)
Financial year 2009/10	-0.029***	(0.005)	-0.006*	(0.003)	-0.049***	(0.005)
Financial year 2010/11	-0.036***	(0.004)	-0.009**	(0.004)	-0.059***	(0.005)
Financial year 2011/12	-0.039***	(0.004)	-0.010***	(0.004)	-0.064***	(0.005)
Financial year 2012/13	-0.038***	(0.004)	-0.009***	(0.003)	-0.063***	(0.005)
Financial year 2013/14	-0.042***	(0.004)	-0.012***	(0.004)	-0.067***	(0.005)
Diagnosis group dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Admission source dummies	<i>Yes</i>	<i>Yes</i>	<i>Yes</i>
Admission month dummies	<i>Yes</i>	<i>Yes</i>	<i>Yes</i>
C statistic	0.88	0.87	0.90
Pseudo R2	0.28	0.23	0.31
Observations	241,338	237,091	242,693

1. Including public holidays.
 2. Average marginal effect. *** p<0.01, ** p<0.05, * p<0.1.
 3. Robust standard errors corrected for clustering around consultant in parentheses.

Confidential: For Review Only

Table 4 Adjusted risk of mortality by financial year, probit regressions with average marginal effects.

Year	N	Weekday 7am Mon - 6.59pm Fri	Weekend ¹ 7pm Fri – 6.59am Mon	Weekday-day 7am - 6.59pm Mon - Fri	Weekday-night 7pm - 6.59am Mon - Thu	Weekend-day ¹ 7am - 6.59pm Sat & Sun	Weekend-night ¹ 7pm - 6.59am Fri - Sun
----- 30-day mortality -----							
2004/05	20,664		0.004 (0.004) ²		0.010*** (0.003)	0.004 (0.004)	0.011** (0.005)
2005/06	20,254		0.006** (0.003)		0.002 (0.004)	0.008* (0.004)	0.005 (0.004)
2006/07	20,991		0.003 (0.003)		0.010*** (0.004)	-0.000 (0.004)	0.012*** (0.004)
2007/08	21,577		0.004 (0.002)		0.002 (0.003)	0.003 (0.004)	0.006* (0.003)
2008/09	21,202	<i>Ref.</i>	0.006** (0.002)	<i>Ref.</i>	0.007** (0.003)	0.005 (0.004)	0.012*** (0.003)
2009/10	22,115		-0.002 (0.003)		0.001 (0.003)	0.001 (0.006)	-0.003 (0.003)
2010/11	23,642		0.006 (0.004)		-0.000 (0.004)	0.007 (0.005)	0.005 (0.006)
2011/12	24,518		0.004* (0.002)		0.000 (0.003)	0.002 (0.004)	0.005** (0.002)
2012/13	24,546		0.009*** (0.003)		0.001 (0.003)	0.010** (0.004)	0.009*** (0.003)
2013/14	27,162		0.002 (0.002)		0.002 (0.002)	0.002 (0.003)	0.004 (0.003)
----- 7-day mortality -----							
2004/05	18,526		0.005* (0.003)		0.009** (0.003)	0.007* (0.003)	0.009** (0.004)
2005/06	17,646		0.005** (0.002)		0.002 (0.003)	0.009** (0.004)	0.003 (0.003)
2006/07	18,826		0.001 (0.003)		0.009*** (0.003)	0.000 (0.004)	0.008** (0.003)
2007/08	19,308		0.001 (0.002)		0.002 (0.003)	0.001 (0.004)	0.003 (0.003)
2008/09	19,797	<i>Ref.</i>	0.003 (0.002)	<i>Ref.</i>	0.002 (0.002)	-0.000 (0.004)	0.008*** (0.003)
2009/10	20,353		-0.000 (0.002)		0.006** (0.003)	0.003 (0.003)	0.002 (0.003)
2010/11	21,906		0.002 (0.003)		0.004* (0.002)	0.002 (0.002)	0.004 (0.005)
2011/12	20,610		0.007*** (0.002)		-0.002 (0.003)	0.005 (0.003)	0.007** (0.003)
2012/13	22,127		0.005*** (0.002)		-0.001 (0.002)	0.005* (0.003)	0.005* (0.003)
2013/14	24,716		0.001 (0.002)		0.004** (0.002)	0.002 (0.003)	0.003 (0.002)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

In-hospital mortality

2004/05	20,454		0.003 (0.003)		0.009** (0.004)	0.003 (0.004)	0.008* (0.005)
2005/06	20,231		0.004 (0.003)		0.004 (0.005)	0.004 (0.004)	0.007 (0.004)
2006/07	20,605		0.002 (0.002)		0.008** (0.004)	0.000 (0.004)	0.010** (0.004)
2007/08	21,763		0.003 (0.002)		-0.002 (0.003)	0.001 (0.004)	0.004 (0.003)
2008/09	21,130	<i>Ref.</i>	0.002 (0.003)	<i>Ref.</i>	0.003 (0.003)	0.002 (0.004)	0.004 (0.003)
2009/10	21,847		0.004 (0.003)		0.003 (0.003)	0.005 (0.006)	0.005* (0.003)
2010/11	23,072		0.002 (0.002)		0.004 (0.003)	0.004 (0.003)	0.005 (0.004)
2011/12	23,921		0.003* (0.002)		-0.002 (0.003)	0.002 (0.004)	0.003 (0.002)
2012/13	23,808		0.004 (0.002)		-0.001 (0.002)	0.004 (0.003)	0.002 (0.003)
2013/14	27,403		0.001 (0.002)		0.005** (0.002)	0.004 (0.003)	0.003 (0.003)

1. Including public holidays.
 2. Average marginal effect (***) p<0.01, ** p<0.05, * p<0.1) with robust standard errors corrected for clustering around consultant in parentheses. Probit regressions adjusted for age, gender, ethnicity, deprivation, Charlson index, number of diagnoses, number of procedures, palliative care, number of emergency admissions in last year, admission method, admission source, transfer-out dummy, admission month and primary diagnosis groups.

Appendix

Table A1 Crude mortality rates, 2004/5 to 2013/14.

Year	Admissions	Overall	Weekday	Weekend ¹	Weekday-day	Weekday-night	Weekend-day ¹	Weekend-night ¹
30-day mortality								
2004/05	22,001	6.61	6.45	6.96	6.32	6.72	7.28	6.69
2005/06	22,356	5.98	5.76	6.46	5.89	5.54	7.01	6.05
2006/07	22,946	6.40	6.19	6.85	5.95	6.62	6.73	6.93
2007/08	23,328	6.35	6.01	7.02	6.02	6.01	7.13	6.95
2008/09	23,194	6.36	6.04	7.05	5.90	6.25	7.27	6.89
2009/10	24,381	5.61	5.55	5.73	5.66	5.38	6.04	5.52
2010/11	25,821	5.91	5.72	6.27	5.59	5.90	6.96	5.80
2011/12	26,291	5.38	5.14	5.86	5.13	5.14	6.17	5.65
2012/13	26,276	5.64	5.34	6.20	5.29	5.42	6.74	5.84
2013/14	29,756	5.17	4.97	5.56	5.08	4.81	5.99	5.28
Total	246,350	5.90	5.69	6.35	5.67	5.72	6.71	6.10
7-day mortality								
2004/05	22,001	2.75	2.54	3.17	2.33	2.98	3.24	3.11
2005/06	22,356	2.74	2.56	3.14	2.56	2.56	3.55	2.83
2006/07	22,946	2.73	2.65	2.91	2.34	3.19	2.83	2.96
2007/08	23,328	2.88	2.72	3.20	2.61	2.89	3.29	3.13
2008/09	23,194	2.75	2.54	3.19	2.45	2.68	3.04	3.30
2009/10	24,381	2.44	2.33	2.66	2.14	2.63	2.86	2.53
2010/11	25,821	2.57	2.50	2.72	2.32	2.75	2.84	2.64
2011/12	26,291	2.37	2.11	2.87	2.10	2.13	3.20	2.66
2012/13	26,276	2.46	2.24	2.88	2.14	2.39	3.10	2.74
2013/14	29,756	2.19	2.04	2.49	1.91	2.22	2.58	2.43
Total	246,350	2.57	2.41	2.90	2.28	2.61	3.04	2.80
In-hospital mortality								

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

2004/05	22,001	5.93	5.81	6.19	5.83	5.77	6.57	5.87
2005/06	22,356	5.35	5.19	5.69	5.17	5.23	5.95	5.49
2006/07	22,946	5.30	5.17	5.58	4.91	5.61	5.49	5.64
2007/08	23,328	5.22	4.98	5.71	5.00	4.93	6.05	5.48
2008/09	23,194	5.07	4.85	5.51	4.73	5.05	5.86	5.26
2009/10	24,381	4.52	4.29	4.99	4.28	4.31	5.20	4.86
2010/11	25,821	4.34	4.22	4.57	4.05	4.45	4.90	4.36
2011/12	26,291	3.69	3.51	4.05	3.53	3.48	4.41	3.82
2012/13	26,276	3.92	3.79	4.16	3.78	3.81	4.45	3.97
2013/14	29,756	3.68	3.50	4.03	3.47	3.55	4.45	3.75
Total	246,350	4.64	4.48	4.97	4.45	4.52	5.28	4.75

1. Including public holidays.

Confidential: For Review Only

Figure A1 Crude 30-day mortality rates, 2004/05 to 2013/14.




Table A2 Adjusted risk of mortality 2004/05-2013/14, logistic regressions with odds ratios.

Variables	30-day mortality		7-day mortality		In-hospital mortality	
	Odds ratio ²	95% CI ³	Odds ratio	95% CI	Odds ratio	95% CI
Admission time						
Weekday	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Weekend ¹	1.104	[1.057, 1.154]	1.122	[1.069, 1.179]	1.083	[1.021, 1.149]
Case-mix variables						
Age band 17-25	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Age band 26-35	0.962	[0.685, 1.349]	0.890	[0.639, 1.241]	0.842	[0.628, 1.128]
Age band 36-45	1.484	[1.086, 2.028]	1.247	[0.931, 1.671]	1.216	[0.913, 1.619]
Age band 46-55	2.161	[1.522, 3.069]	1.719	[1.218, 2.425]	1.585	[1.144, 2.196]
Age band 56-65	3.362	[2.240, 5.048]	2.608	[1.666, 4.082]	2.357	[1.582, 3.511]
Age band 66-75	4.139	[2.773, 6.177]	3.081	[1.998, 4.752]	3.025	[2.062, 4.436]
Age band 76-85	6.074	[4.076, 9.050]	4.100	[2.663, 6.312]	4.518	[3.073, 6.641]
Age band 85+	9.454	[6.312, 14.159]	5.429	[3.416, 8.629]	7.753	[5.137, 11.699]
Gender						
Gender-male	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Gender-female	0.934	[0.880, 0.992]	0.980	[0.920, 1.044]	0.990	[0.942, 1.040]
Gender-not stated	0.858	[0.401, 1.832]	0.962	[0.377, 2.458]	1.037	[0.501, 2.147]
Ethnicity						
Ethnicity-White	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Ethnicity-mixed	0.677	[0.341, 1.344]	0.956	[0.513, 1.784]	0.708	[0.364, 1.376]
Ethnicity-Asian	0.614	[0.462, 0.816]	0.655	[0.464, 0.926]	0.634	[0.481, 0.836]
Ethnicity-Black	0.555	[0.376, 0.818]	0.469	[0.260, 0.848]	0.532	[0.339, 0.835]
Ethnicity-other	1.448	[1.148, 1.825]	1.646	[1.262, 2.146]	1.534	[1.234, 1.906]
Ethnicity-not stated	1.930	[1.512, 2.463]	2.122	[1.636, 2.751]	2.035	[1.613, 2.567]
IMD quintiles						
IMD quintile-1 (most affluent)	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
IMD quintile-2	1.010	[0.930, 1.096]	0.984	[0.909, 1.066]	0.972	[0.900, 1.051]
IMD quintile-3	1.045	[0.980, 1.115]	1.059	[0.970, 1.156]	1.007	[0.947, 1.071]
IMD quintile-4	1.033	[0.963, 1.108]	0.971	[0.886, 1.063]	0.972	[0.906, 1.043]
IMD quintile-5 (most deprived)	1.090	[1.025, 1.159]	1.066	[0.980, 1.159]	1.064	[0.998, 1.135]

IMD quintile-missing	1.300	[1.194, 1.417]	1.219	[1.094, 1.359]	1.291	[1.189, 1.401]
Charlson index	1.040	[1.036, 1.045]	1.037	[1.032, 1.042]	1.034	[1.030, 1.039]
No. of diagnosis	1.037	[1.021, 1.053]	0.987	[0.967, 1.008]	1.094	[1.076, 1.114]
No. of procedures	0.977	[0.960, 0.994]	0.911	[0.876, 0.948]	1.035	[1.016, 1.054]
Palliative care	6.744	[5.558, 8.182]	2.831	[2.268, 3.534]	5.608	[4.610, 6.823]
No. of emergency admissions in the previous 1 year (365 days)	0.953	[0.936, 0.969]	0.976	[0.958, 0.995]	0.967	[0.948, 0.987]
Admission method-emergency A+E	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Admission method-emergency transfer from other provider	0.642	[0.544, 0.756]	0.479	[0.378, 0.606]	0.655	[0.543, 0.791]
Admission method-emergency domicile	0.824	[0.339, 2.005]	1.000	[1.000, 1.000]	0.454	[0.177, 1.165]
Admission method-emergency GP refer	1.124	[1.010, 1.252]	0.702	[0.581, 0.848]	1.038	[0.912, 1.182]
Admission method-emergency outpatient	0.591	[0.494, 0.708]	0.321	[0.225, 0.457]	0.575	[0.452, 0.732]
Admission method-emergency ante natal	1.764	[0.583, 5.331]	1.579	[0.243, 10.251]	1.000	[1.000, 1.000]
Admission method-emergency post natal	1.000	[1.000, 1.000]	1.000	[1.000, 1.000]	1.000	[1.000, 1.000]
Admission method-non-emergency transfer from other provider	0.661	[0.518, 0.843]	0.441	[0.323, 0.603]	0.623	[0.487, 0.796]
Transfer to other hospitals	0.584	[0.471, 0.726]	0.177	[0.137, 0.229]		
Financial year 2004/05	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Financial year 2005/06	0.788	[0.665, 0.933]	0.914	[0.779, 1.072]	0.754	[0.656, 0.866]
Financial year 2006/07	0.805	[0.707, 0.916]	0.913	[0.804, 1.037]	0.681	[0.603, 0.767]
Financial year 2007/08	0.812	[0.690, 0.956]	1.072	[0.847, 1.357]	0.626	[0.507, 0.771]
Financial year 2008/09	0.615	[0.521, 0.726]	0.855	[0.679, 1.078]	0.398	[0.321, 0.493]
Financial year 2009/10	0.533	[0.440, 0.646]	0.829	[0.643, 1.070]	0.335	[0.266, 0.423]
Financial year 2010/11	0.453	[0.369, 0.555]	0.744	[0.533, 1.038]	0.233	[0.184, 0.295]
Financial year 2011/12	0.403	[0.326, 0.498]	0.698	[0.510, 0.955]	0.192	[0.154, 0.240]
Financial year 2012/13	0.420	[0.346, 0.511]	0.720	[0.548, 0.945]	0.202	[0.162, 0.252]
Financial year 2013/14	0.367	[0.293, 0.460]	0.655	[0.478, 0.897]	0.169	[0.130, 0.220]
Diagnosis group dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
Admission source dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
Admission month dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

C statistic	0.88	0.88	0.90
Pseudo R2	0.28	0.23	0.31
Observations	241,338	237,091	242,693

-
- 1. Including public holidays.
 - 2. Estimated by logistic regression.
 - 3. 95% confidence interval corrected for clustering around consultant.

Confidential: For Review Only

Table A3 Adjusted risk of mortality 2004/05-2013/14, logistic regressions with odds ratios.

Variables	30-day mortality		7-day mortality		In-hospital mortality	
	Odds ratio ²	95% CI ³	Odds ratio	95% CI	Odds ratio	95% CI
Admission time						
Weekday day	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Weekday night	1.087	[1.012, 1.167]	1.160	[1.061, 1.268]	1.088	[1.029, 1.150]
Weekend day ¹	1.110	[1.041, 1.183]	1.155	[1.069, 1.249]	1.095	[1.010, 1.187]
Weekend night ¹	1.168	[1.095, 1.245]	1.225	[1.121, 1.338]	1.140	[1.077, 1.208]
Case-mix variables						
Age band 17-25	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Age band 26-35	0.963	[0.687, 1.350]	0.891	[0.640, 1.242]	0.843	[0.629, 1.129]
Age band 36-45	1.485	[1.087, 2.029]	1.247	[0.931, 1.672]	1.216	[0.913, 1.619]
Age band 46-55	2.163	[1.523, 3.073]	1.720	[1.219, 2.428]	1.585	[1.144, 2.197]
Age band 56-65	3.373	[2.247, 5.061]	2.620	[1.675, 4.098]	2.363	[1.586, 3.521]
Age band 66-75	4.155	[2.786, 6.195]	3.100	[2.012, 4.777]	3.035	[2.069, 4.451]
Age band 76-85	6.095	[4.094, 9.073]	4.122	[2.680, 6.338]	4.532	[3.084, 6.660]
Age band 85+	9.502	[6.354, 14.208]	5.473	[3.453, 8.673]	7.789	[5.165, 11.747]
Gender-male	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Gender-female	0.934	[0.880, 0.991]	0.979	[0.919, 1.043]	0.990	[0.942, 1.040]
Gender-not stated	0.858	[0.401, 1.834]	0.962	[0.372, 2.487]	1.036	[0.500, 2.148]
Ethnicity-White	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
Ethnicity-mixed	0.677	[0.339, 1.350]	0.955	[0.509, 1.790]	0.708	[0.363, 1.379]
Ethnicity-Asian	0.613	[0.462, 0.814]	0.654	[0.463, 0.925]	0.633	[0.480, 0.835]
Ethnicity-Black	0.555	[0.376, 0.819]	0.470	[0.260, 0.850]	0.532	[0.338, 0.838]
Ethnicity-other	1.443	[1.147, 1.815]	1.636	[1.259, 2.126]	1.528	[1.230, 1.900]
Ethnicity-not stated	1.932	[1.513, 2.468]	2.125	[1.637, 2.758]	2.037	[1.614, 2.572]
IMD quintile-1 (most affluent)	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
IMD quintile-2	1.009	[0.930, 1.095]	0.984	[0.910, 1.066]	0.972	[0.900, 1.051]
IMD quintile-3	1.045	[0.980, 1.115]	1.059	[0.970, 1.156]	1.007	[0.947, 1.070]

1							
2							
3							
4							
5	IMD quintile-4	1.032	[0.962, 1.107]	0.970	[0.886, 1.062]	0.972	[0.906, 1.043]
6	IMD quintile-5 (most deprived)	1.089	[1.024, 1.159]	1.065	[0.979, 1.158]	1.063	[0.997, 1.134]
7	IMD quintile-missing	1.298	[1.191, 1.414]	1.216	[1.091, 1.356]	1.289	[1.187, 1.399]
8							
9	Charlson index	1.040	[1.036, 1.045]	1.037	[1.032, 1.042]	1.034	[1.030, 1.039]
10							
11	No. of diagnosis	1.037	[1.021, 1.053]	0.987	[0.967, 1.008]	1.094	[1.076, 1.114]
12	No. of procedures	0.976	[0.960, 0.993]	0.911	[0.876, 0.948]	1.034	[1.016, 1.053]
13	Palliative care	6.745	[5.562, 8.180]	2.829	[2.267, 3.531]	5.608	[4.610, 6.822]
14	No. of emergency admissions in the previous 1 year (365 days)	0.953	[0.936, 0.969]	0.976	[0.958, 0.995]	0.967	[0.948, 0.987]
15							
16	Admission method-emergency A+E	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
17	Admission method-emergency transfer from other provider	0.653	[0.554, 0.770]	0.493	[0.389, 0.624]	0.666	[0.551, 0.805]
18	Admission method-emergency domicile	0.832	[0.341, 2.033]	1.000	[1.000, 1.000]	0.459	[0.179, 1.177]
19	Admission method-emergency GP refer	1.145	[1.022, 1.282]	0.725	[0.597, 0.881]	1.057	[0.927, 1.206]
20	Admission method-emergency outpatient	0.605	[0.506, 0.724]	0.335	[0.236, 0.476]	0.589	[0.464, 0.749]
21	Admission method-emergency ante natal	1.814	[0.611, 5.388]	1.636	[0.260, 10.278]	1.000	[1.000, 1.000]
22	Admission method-emergency post natal	1.000	[1.000, 1.000]	1.000	[1.000, 1.000]	1.000	[1.000, 1.000]
23	Admission method-non-emergency transfer from other provider	0.673	[0.526, 0.860]	0.455	[0.333, 0.621]	0.633	[0.495, 0.810]
24							
25	Transfer to other hospitals	0.583	[0.470, 0.725]	0.176	[0.137, 0.228]		
26							
27							
28	Financial year 2004/05	<i>Ref.</i>		<i>Ref.</i>		<i>Ref.</i>	
29	Financial year 2005/06	0.786	[0.663, 0.931]	0.910	[0.776, 1.069]	0.752	[0.654, 0.864]
30	Financial year 2006/07	0.802	[0.704, 0.914]	0.908	[0.801, 1.031]	0.678	[0.601, 0.765]
31	Financial year 2007/08	0.809	[0.687, 0.952]	1.066	[0.842, 1.350]	0.623	[0.505, 0.769]
32	Financial year 2008/09	0.613	[0.519, 0.724]	0.852	[0.675, 1.074]	0.396	[0.320, 0.491]
33	Financial year 2009/10	0.531	[0.439, 0.643]	0.825	[0.639, 1.064]	0.334	[0.265, 0.421]
34	Financial year 2010/11	0.450	[0.368, 0.552]	0.738	[0.530, 1.026]	0.232	[0.183, 0.293]
35	Financial year 2011/12	0.402	[0.326, 0.495]	0.693	[0.508, 0.946]	0.192	[0.154, 0.239]
36	Financial year 2012/13	0.418	[0.345, 0.507]	0.715	[0.545, 0.938]	0.201	[0.162, 0.251]
37	Financial year 2013/14	0.366	[0.293, 0.458]	0.651	[0.476, 0.891]	0.169	[0.130, 0.219]
38							
39	Diagnosis group dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
40	Admission source dummies	<i>Yes</i>		<i>Yes</i>		<i>Yes</i>	
41							
42							
43							
44							
45							
46							
47							
48							
49							

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Admission month dummies	<i>Yes</i>	<i>Yes</i>	<i>Yes</i>
C statistic	0.88	0.88	0.90
Pseudo R2	0.28	0.24	0.31
Observations	241,338	237,091	242,693

-
- 1. Including public holidays.
 - 2. Estimated by logistic regression.
 - 3. 95% confidence interval corrected for clustering around consultant.

Confidential: For Review Only

Table A4 Adjusted risk of mortality by financial year, logistic regressions with odds ratios.

Year	N	Weekday 7am Mon - 6.59pm Fri	Weekend ¹ 7pm Fri - 6.59am Mon	Weekday-day 7am - 6.59pm Mon - Fri	Weekday-night 7pm - 6.59am Mon - Thu	Weekend-day ¹ 7am - 6.59pm Sat & Sun	Weekend-night ¹ 7pm - 6.59am Fri - Sun
30-day mortality							
2004/05	20,664		1.068 [0.917, 1.244] ²		1.250 [1.080, 1.446]	1.078 [0.891, 1.303]	1.223 [0.984, 1.521]
2005/06	20,254		1.166 [1.015, 1.340]		1.064 [0.859, 1.318]	1.237 [1.021, 1.499]	1.157 [0.943, 1.420]
2006/07	20,991		1.064 [0.941, 1.204]		1.224 [1.048, 1.430]	0.997 [0.812, 1.225]	1.285 [1.092, 1.511]
2007/08	21,577		1.099 [0.996, 1.212]		1.049 [0.913, 1.207]	1.079 [0.923, 1.260]	1.152 [1.002, 1.324]
2008/09	21,202	<i>Ref.</i>	1.137 [1.022, 1.265]	<i>Ref.</i>	1.170 [1.012, 1.351]	1.093 [0.923, 1.295]	1.314 [1.134, 1.522]
2009/10	22,115		0.970 [0.829, 1.134]		1.030 [0.870, 1.220]	1.026 [0.743, 1.417]	0.951 [0.817, 1.107]
2010/11	23,642		1.145 [0.947, 1.384]		0.990 [0.817, 1.200]	1.188 [0.945, 1.492]	1.105 [0.821, 1.487]
2011/12	24,518		1.078 [0.970, 1.199]		1.011 [0.872, 1.172]	1.037 [0.836, 1.287]	1.121 [0.984, 1.277]
2012/13	24,546		1.287 [1.146, 1.446]		1.026 [0.888, 1.185]	1.330 [1.114, 1.587]	1.282 [1.088, 1.510]
2013/14	27,162		1.040 [0.934, 1.158]		1.052 [0.947, 1.168]	1.033 [0.870, 1.227]	1.084 [0.936, 1.255]
7-day mortality							
2004/05	18,526		1.195 [0.986, 1.448]		1.350 [1.071, 1.702]	1.304 [1.009, 1.685]	1.361 [1.026, 1.806]
2005/06	17,646		1.181 [1.008, 1.385]		1.079 [0.867, 1.344]	1.314 [1.017, 1.698]	1.137 [0.920, 1.406]
2006/07	18,826		1.038 [0.856, 1.259]		1.357 [1.115, 1.651]	1.020 [0.750, 1.386]	1.310 [1.077, 1.593]
2007/08	19,308		1.041 [0.899, 1.205]		1.085 [0.882, 1.335]	1.055 [0.818, 1.362]	1.092 [0.904, 1.318]
2008/09	19,797	<i>Ref.</i>	1.137 [0.968, 1.336]	<i>Ref.</i>	1.083 [0.890, 1.317]	1.003 [0.760, 1.323]	1.321 [1.067, 1.634]
2009/10	20,353		1.003 [0.845, 1.191]		1.304 [1.057, 1.609]	1.149 [0.870, 1.518]	1.109 [0.875, 1.404]
2010/11	21,906		1.069 [0.850, 1.344]		1.182 [0.973, 1.438]	1.083 [0.870, 1.349]	1.209 [0.805, 1.816]
2011/12	20,610		1.289 [1.093, 1.520]		0.944 [0.740, 1.205]	1.234 [0.964, 1.581]	1.275 [0.959, 1.695]
2012/13	22,127		1.271 [1.100, 1.468]		1.004 [0.838, 1.203]	1.289 [1.018, 1.633]	1.262 [1.035, 1.539]
2013/14	24,716		1.021 [0.855, 1.220]		1.230 [1.011, 1.496]	1.093 [0.815, 1.464]	1.135 [0.901, 1.430]
In-hospital mortality							
2004/05	20,454		1.059 [0.922, 1.216]		1.211 [1.031, 1.423]	1.099 [0.923, 1.309]	1.160 [0.945, 1.425]
2005/06	20,231	<i>Ref.</i>	1.122 [0.969, 1.300]	<i>Ref.</i>	1.107 [0.880, 1.394]	1.119 [0.942, 1.330]	1.024 [0.977, 1.485]
2006/07	20,605		1.055 [0.939, 1.185]		1.207 [0.984, 1.480]	0.990 [0.809, 1.213]	1.261 [1.045, 1.521]
2007/08	21,763		1.081 [0.960, 1.217]		0.969 [0.841, 1.116]	1.038 [0.865, 1.246]	1.091 [0.929, 1.280]

2008/09	21,130	1.038 [0.897, 1.201]	1.069 [0.932, 1.226]	1.029 [0.850, 1.244]	1.098 [0.926, 1.301]
2009/10	21,847	1.127 [0.960, 1.323]	1.088 [0.942, 1.256]	1.153 [0.836, 1.590]	1.178 [1.002, 1.383]
2010/11	23,072	1.075 [0.939, 1.231]	1.120 [0.911, 1.377]	1.112 [0.948, 1.306]	1.146 [0.900, 1.459]
2011/12	23,921	1.090 [0.966, 1.230]	0.931 [0.777, 1.115]	1.057 [0.834, 1.340]	1.056 [0.915, 1.219]
2012/13	23,808	1.131 [0.990, 1.291]	0.958 [0.839, 1.094]	1.145 [0.939, 1.398]	1.085 [0.906, 1.299]
2013/14	27,403	1.032 [0.902, 1.180]	1.187 [1.014, 1.390]	1.175 [0.944, 1.462]	1.064 [0.869, 1.302]

1. Including public holidays.

2. Odds ratio estimated by logistic regression with 95% confidence interval corrected for clustering around consultant in brackets. Logistic regressions adjusted for age, gender, ethnicity, deprivation, Charlson index, number of diagnoses, number of procedures, palliative care, number of emergency admissions in last year, admission method, admission source, transfer-out dummy, admission month and primary diagnosis groups.