



This is a repository copy of *The sustainability of water resources in High Mountain Asia in the context of recent and future glacier change*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/122297/>

Version: Accepted Version

---

**Article:**

Rowan, A.V. [orcid.org/0000-0002-3715-5554](https://orcid.org/0000-0002-3715-5554), Quincey, D.J., Gibson, M.J. et al. (5 more authors) (2017) The sustainability of water resources in High Mountain Asia in the context of recent and future glacier change. *The Himalayan Cryosphere: Past and Present, Special Publications*, 462. pp. 189-204. ISSN 0305-8719

<https://doi.org/10.1144/SP462.12>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# The sustainability of water resources in High Mountain Asia in the context of recent and future glacier change

Ann V. Rowan<sup>1\*</sup>, Duncan J. Quincey<sup>2</sup>, Morgan J. Gibson<sup>3</sup>, Neil F. Glasser<sup>3</sup>, Matthew J. Westoby<sup>4</sup>, Tristram D.L. Irvine-Fynn<sup>3</sup>, Phillip R. Porter<sup>5</sup>, Michael J. Hambrey<sup>3</sup>

<sup>1</sup>*Department of Geography, University of Sheffield, Winter Street, Sheffield, S10 2TN, UK*

\*Corresponding author; email: [a.rowan@sheffield.ac.uk](mailto:a.rowan@sheffield.ac.uk)

<sup>2</sup>*School of Geography, University of Leeds, LS2 9JT, UK*

<sup>3</sup>*Centre for Glaciology, Department of Geography and Earth Sciences, Aberystwyth University, SY23 3DB, UK*

<sup>4</sup>*School of Engineering and Environment, Northumbria University, Newcastle upon Tyne, NE1 8ST*

<sup>5</sup>*University of Hertfordshire, Hatfield, Hertfordshire. AL10 9AB, UK*

## Abstract

High Mountain Asia contains the largest volume of glacier ice outside the Polar regions, and contain the headwaters of some of the largest rivers in central Asia. These glaciers are losing mass at a mean rate of between  $-0.18$  m and  $-0.5$  m water equivalent per year. While glaciers in the Himalaya are generally shrinking, those in the Karakoram have experienced a slight mass gain. Both changes have occurred in response to rising air temperatures due to Northern Hemisphere climatic change. In the Westerly influenced Indus catchment, glacier meltwater makes up a large proportion of the hydrological budget, and loss of glacier mass will ultimately lead to a decrease in water supplies. In the monsoon-influenced Ganges and Brahmaputra catchments, the contribution of glacial meltwater is relatively small compared to the Indus, and the decrease in annual water supplies will be less dramatic. Therefore, enhanced glacier melt will increase river flows until the middle of the 21<sup>st</sup> Century, but in the longer-term into the latter part of this century, river flows will decline as glaciers shrink. Declining meltwater supplies may be compensated by increases in precipitation, but this could exacerbate the risk of flooding.

## 1. Introduction

Millions of people rely on glaciers in the Himalaya, Karakoram and Hindu Kush mountains, collectively referred to as High Mountain Asia, as a water resource. These glaciers form the headwaters of the largest rivers in Asia, including the Indus, the Ganges and the Brahmaputra Rivers (Fig. 1) and as such the mountains are often referred to as the ‘water towers of Asia’ (Immerzeel *et al.* 2010). High Mountain Asia contains the largest glacierised area outside the

39 Polar regions (Bolch *et al.* 2012) and glaciers here are highly sensitive to climate change  
40 (Solomina *et al.* 2016). Glaciers in the Himalaya are predominantly shrinking (Kääb *et al.*  
41 2012), and rates of glacier mass loss, although spatially variable, have accelerated since the  
42 1990s (Bolch *et al.* 2012). If a constant rate of glacier mass loss after 1975 is assumed, then  
43 predictions indicate thinning of 9–28 m water equivalent (w.e.) between 2010 and 2035,  
44 which is sufficient to result in the disappearance of many smaller glaciers across the  
45 mountain range (Cogley 2011). The catchments supplied by rivers draining High Mountain  
46 Asia are located in developing countries that use this water primarily for agriculture and  
47 hydroelectric power generation, and are extremely vulnerable to changes in their water  
48 supply (Kaser *et al.* 2010; Pritchard, 2017). Predictions are needed of how Asian water  
49 supplies are likely to change due to continued glacier mass loss in response to recent and  
50 future climate change. We therefore need to improve understanding of both the contribution  
51 of glaciers to the hydrological budgets of these large catchments, and discover how these  
52 glaciers are responding to recent and future climate change (e.g. Lutz *et al.* 2014; Brun *et al.*  
53 2017).

54

55 Although only 1–3% of the area of the Indus, Ganges and Brahmaputra catchments are  
56 glacierised, these densely-populated catchments rely on glacial meltwater (Immerzeel *et al.*  
57 2010). The contribution of glaciers to runoff varies regionally; from 18.8% in the Dudh  
58 Koshi catchment, which is a major tributary of the Ganges, up to 80.6% in the Hunza  
59 catchment that drains into the Indus (Lutz *et al.* 2014). The Indus and the Ganges provide  
60 important water supplies that are used to irrigate over 140,000 km<sup>2</sup> of agricultural land, and  
61 the largest irrigation network in the world is contained in the Indus catchment (Immerzeel *et al.*  
62 2010). In particular, the importance of glacier meltwater relative to other water sources  
63 (e.g. precipitation, snow melt, groundwater) for regional hydrological budgets has only  
64 recently been documented (Immerzeel *et al.* 2010, Lutz *et al.* 2014). In the monsoon-  
65 influenced Central and Eastern Himalaya the majority of annual precipitation occurs during  
66 the warm summer monsoon months (June to September) (Bookhagen & Burbank 2010). The  
67 high summer rainfall and snowfall roughly coincide with the timing of the majority of glacier  
68 ablation (Benn & Lehmkuhl 2000), so that the relative contribution of glacial melt to river  
69 flows is minimised compared to regions to the west where summers are generally dry (Kaser  
70 *et al.* 2010). In the Western Himalaya, Karakoram and Hindu Kush mountains, where the  
71 majority of precipitation occurs as winter snowfall, glacier melt plays a much more important  
72 role in regulating seasonal river flows, with a relatively larger proportion of the annual flow  
73 in the Indus resulting directly from glacier melt compared to that in the Ganges and  
74 Brahmaputra catchments (Lutz *et al.* 2014).

75

76 The impact of climate change on these vital river flows from the Himalaya is, however, not  
77 straightforward. For example, climatic warming may cause glaciers to lose mass and release a  
78 greater volume of meltwater each year, but may also result in increased orographic  
79 precipitation that could sustain or enhance flows and trigger a gradual or abrupt change in the  
80 seasonality of peak flows (Immerzeel *et al.* 2010). Understanding regional and catchment-  
81 scale hydrological budgets and predicting how they will vary under a changing climate  
82 therefore requires coupling our understanding of glaciers processes with climate model  
83 forecasts. The Intergovernmental Panel on Climate Change (IPCC) climate scenarios for the  
84 21<sup>st</sup> Century are used for this purpose, as they are compiled from comparison of an ensemble  
85 of state-of-the-art climate model outputs (Collins *et al.* 2013). Many regional hydrological  
86 models contain large uncertainties as they do not capture the processes that affect individual  
87 glaciers, and so detailed catchment-scale models validated by field data are also required to  
88 better constrain future hydrological changes (e.g. Ragetli *et al.* 2016).

89

90 Here we review recent and predicted glacier and hydrological change across High Mountain  
91 Asia (Fig. 1). Glaciers on the Tibetan Plateau are excluded from this review, as this region is  
92 strongly influenced by weather systems originating from the Arctic rather than the Asian  
93 monsoons, and show markedly different behaviour compared to these glaciers. The long-term  
94 response to climate change of Tibetan glaciers is described by Owen *et al.* (2008) and Owen  
95 (2009). We first summarise the current knowledge of the state of glaciers in High Mountain  
96 Asia, then discuss these changes in the context of observed longer-term glacier change since  
97 the late Holocene (the last 2,000 years). We then compare predictions of glacier change with  
98 current and future climate change and consider the likely impacts of these changes on  
99 regional hydrological budgets.

100

## 101 **2. The current state of glaciers in High Mountain Asia**

102 Glaciers in High Mountain Asia are discussed in terms of their location in sub-regional areas  
103 that are defined from the major regional climate controls (after Bolch *et al.* 2012; Fig. 1).  
104 From east to west these regions are: the monsoon-influenced Eastern and Central Himalaya,  
105 and the mid-latitude Western Himalaya, Karakoram and Hindu Kush ranges influenced by  
106 westerly weather systems. These areas follow the boundaries of the major river catchments  
107 within the high mountains, with the Western Himalaya, Karakoram and Hindu Kush draining  
108 into the Indus, the Central Himalaya into the Ganges, and the Eastern Himalaya and some of  
109 the Tibetan Plateau forming the headwaters of the Brahmaputra River.

110

### 111 **2.1 Glacier extent and volume**

112 The Himalaya and Karakoram mountains contain 32,353 glaciers with a total glacierised area  
113 of about 41,000 km<sup>2</sup> equivalent to 6% of the global glacierised area (Arendt *et al.* 2015).

114 Until recently, relatively little was known about the total number and size of glaciers in the  
115 Himalaya because perennial snow cover, debris-covered ice and ice-cored moraine impeded  
116 identification of glacier outlines from satellite observations. Improvements in satellite remote  
117 sensing imagery have allowed identification of the majority of glacier outlines, which are  
118 compiled in the 6<sup>th</sup> Randolph Glacier Inventory (released in July 2017) and cover most of  
119 High Mountain Asia (Arendt *et al.* 2015). Global glacier inventories comprising glacier area  
120 boundaries drawn by the glaciological community are now sufficiently complete to estimate  
121 the glacierised extents, but data describing other important characteristics such as ice  
122 thickness are highly spatially variable and are limited by the small number of field  
123 observations. Glacier volume is more difficult to measure than area, as ice thickness is also  
124 unknown for most of the range. Estimated mean ice thickness for all glaciers in the Global  
125 Land Ice Measurements from Space database are low compared to typical values from  
126 individual glaciers derived from field data. Mean ice thickness is estimated to be 86 m for the  
127 Himalaya, and 172 m for the Karakoram (<http://glims.colorado.edu/glacierdata/> accessed on  
128 30/09/16), although the uncertainty associated with these values is undoubtedly large (Cogley  
129 2011; Frey *et al.* 2014) and likely biased by the majority of measurements being obtained  
130 from smaller glaciers.

131  
132 More robust *in situ* ice thickness measurements have only been made for a handful of  
133 glaciers, using ground-penetrating radar or radio-echo sounding surveys. Access to large  
134 high-altitude glaciers can be challenging and so field observations are frequently made at  
135 more accessible glaciers. Accessible glaciers are generally at lower altitudes, smaller than the  
136 majority of the population, and have higher rates of mass loss than larger glaciers at higher  
137 altitude. These glaciers are not necessarily representative of regional-scale behaviour, and  
138 therefore field measurements often contain a bias that may skew understanding of regional  
139 mass balance trends (Fujita & Nuimura 2011). Ice thicknesses for three glaciers in Nepal in  
140 the Central Himalaya ranged from less than 20 near the terminus to 440 m near the icefall for  
141 the largest, Khumbu Glacier (glacier area is 39.5 km<sup>2</sup>; Gades *et al.* 2000), 20–157 m for  
142 Lirung Glacier (13.5 km<sup>2</sup>; Kadota *et al.* 1997) to 51–86 m for the smallest, Glacier AX010  
143 (0.6 km<sup>2</sup>; Kadota *et al.* 1997). Ice thickness was 124–270 m for Chhota Shigri Glacier (15.7  
144 km<sup>2</sup>) in the Western Himalaya (Azam *et al.* 2012). These values represent the centreline ice  
145 thickness, and in each case the glacier cross-section thins towards the valley sides (*cf.* Azam  
146 *et al.* 2012).

147  
148 Across the Himalaya, 14–18% of the glacierised area is debris covered (Kääb *et al.* 2012), the  
149 extent of which increases to the east to reach 36% in the Everest region of Nepal (Thakuri *et*  
150 *al.* 2014). Supraglacial debris thickness typically increases down-glacier, as englacial and  
151 supraglacial debris transport concentrates sediment previously incorporated into the ice (Fig.

152 2) (Rowan *et al.* 2015). The thickness of supraglacial debris layers can exceed several metres  
153 (Nicholson & Benn 2013, Thakuri *et al.* 2014; Soncini *et al.* 2016). Supraglacial debris  
154 modifies glacier mass balance, either through enhancing ablation due to the decreased albedo  
155 of debris compared to ice, or by reducing ablation by insulating the glacier surface; feedbacks  
156 which are largely dependent on the thickness of the debris layer (Østrem 1959, Evatt *et al.*  
157 2015, Nicholson & Benn 2006). The threshold between the enhancement or attenuation of  
158 ablation by supraglacial debris occurs at a critical thickness of around 0.05 m, as  
159 demonstrated both from field (Rounce *et al.* 2015) and laboratory measurements  
160 (Reznichenko *et al.* 2010). The influence of variations in debris thickness on ablation was  
161 demonstrated at Khumbu Glacier, where rates of surface lowering are highest mid-glacier just  
162 below the icefall where the surface is either debris free or only thinly mantled compared to  
163 the heavily debris-mantled terminus where ablation rates are an order of magnitude lower  
164 (Nakawo *et al.* 1999, Adhikari & Huybrechts 2009, Owen *et al.* 2009).

165

## 166 **2.2 State of glacier mass balance and their Equilibrium Line Altitudes**

167 Glacier mass balance is highly variable across High Mountain Asia. This variability in mass  
168 balance has been identified directly using the traditional glaciological method where stakes  
169 are inserted into the glacier surface to measure ice ablation and snow accumulation (e.g.  
170 Soncini *et al.* 2016). Equilibrium Line Altitude (ELA; the point on a glacier at which  
171 accumulation and ablation are balanced) can be estimated from the areal extent and  
172 hypsometry combined with climate data (Benn & Lehmkuhl 2000). The ELA method has the  
173 advantage of allowing reconstructions of past glacier mass balance from geological evidence  
174 of glacier extent (Benn *et al.* 2005, Owen & Benn 2005). Regional glacier mass balance can  
175 be estimated from geodetic methods using multi-temporal satellite imagery that measures  
176 changes in glacier surface elevations (e.g. Bolch *et al.* 2011, Kääb *et al.* 2012). Remote  
177 sensing can also be used to measure snowline altitudes at the end of the ablation season, from  
178 which the ELA can be derived (e.g. Harper & Humphrey 2003). Mass balance is more  
179 difficult to measure for debris-covered glaciers than for clean-ice glaciers due to the rapid  
180 variations in rates of ablation that occur across the glacier surface, influenced by the thermal  
181 conductivity of the debris layer, which is controlled by factors including debris thickness,  
182 debris grain size, porosity and water content (Benn *et al.* 2012).

183

184 Mass balance has only been measured directly for a small number of glaciers (Fig. 3), and the  
185 longest of the continuous records cover only 10 years. Mass balance for Shaune Garang  
186 Glacier in the Himachal Pradesh in northern India was  $-0.36 \text{ m w.e. a}^{-1}$  between 1981 and  
187 1991 (Pratap *et al.* 2015). Measurements of mass balance for small glaciers in the Central  
188 Himalaya indicate the extreme sensitivity of the monsoon-influenced glaciers to air  
189 temperature. Mass balance measurements through the 1978 monsoon for Glacier AX010

190 indicate that a 0.5°C decrease in mean summer air temperature would result in a transition  
191 between positive and negative mass balance (Ageta *et al.* 1980). Three annual ablation stake  
192 surveys indicate a mass balance of  $-1.6 \text{ m w.e. a}^{-1}$  between 2003 and 2014 for Gangju La  
193 Glacier, a small clean-ice glacier in Bhutan in the Eastern Himalaya (Tshering & Fujita  
194 2016). Mass balance modelling for the partially debris-covered Langtang Glacier in the  
195 Central Himalaya simulated a mass balance of  $-0.11 \text{ m w.e. a}^{-1}$  between 1987 and 1997  
196 (Sharma & Owen 1996). Mean present-day ELA calculated from snowline elevations in the  
197 Annapurna region of western Nepal was  $\sim 5050 \text{ m}$  (Harper & Humphrey 2003) and in eastern  
198 Nepal the ELA ranged from 5300 m in the Langtang Valley to 5600 m in the Khumbu Valley  
199 (Kayastha & Harrison 2008).

200

201 The complete mass budget of all glaciers in High Mountain Asia between 2000 and 2016 was  
202 recently calculated from remote topographic measurements as  $-0.18 \pm 0.04 \text{ m w.e. a}^{-1}$  (Brun  
203 *et al.* 2017). This value is slightly lower than that given by glaciological mass balance records  
204 (summarised by Bolch *et al.* 2012) which gave a regional mass budget of about  $-0.3 \text{ m w.e. a}^{-1}$   
205 between the 1960s and 1990s, becoming increasingly more negative during the last two  
206 decades and similar to the global mean (around  $-0.5 \text{ m w.e. a}^{-1}$ ). Remote sensing studies have  
207 previously indicated a slightly more negative regional mass balance between 2003 and 2008  
208 of  $-0.21 \pm 0.05 \text{ m w.e. a}^{-1}$ , lower than the global average due to the slightly positive mass  
209 budget in the Karakoram (Kääb *et al.* 2012)—the so-called ‘Karakoram anomaly’ (Gardelle  
210 *et al.* 2012). Karakoram glaciers have recently gained mass due to rising air temperatures  
211 delivering more winter snowfall from the Arabian Gulf (Kapnick *et al.* 2014). A large  
212 proportion of surge-type glaciers are found in the Karakoram, and this dynamic behavior can  
213 also result in short-term mass gain (Quincey *et al.* 2011; Quincey *et al.* 2015).

214

### 215 **2.3 The “debris-cover anomaly”**

216 The ablation areas of many glaciers in the Himalaya are covered with rock debris, which is  
217 deposited on glacier surfaces as a result of erosion and mass wasting of the surrounding  
218 landscape. Supraglacial debris affects mass balance and complicates understanding of the  
219 response of these debris-covered glaciers to climate change (Scherler *et al.* 2011). There are  
220 four main sources of debris on the surface of Himalayan glaciers: (1) rockfall debris which is  
221 angular in character; (2) mixed rock- and ice-avalanche debris, which is texturally similar,  
222 but which is entrained as prominent debris layers within the glacier (Fig. 2d); (3) material  
223 resulting from collapse from over-steepened moraines which is characterised by sandy  
224 boulder gravel and is typically sub-rounded to angular; (4) debris derived from the base of the  
225 glacier that has been transported to the surface by thrusting or shear from the bed, such as at  
226 the base of an icefall. This debris is a mixture of silt, sand and gravel, with some boulders  
227 bearing striations. These four lithofacies become intimately mixed on the surface of debris-

228 covered glaciers due to local slope movements from uneven ablation (Figure 2c) (Hambrey *et*  
229 *al.* 2008). Many Himalayan glaciers are also bounded by prominent latero-terminal moraine  
230 systems. These moraines are comprised of a mixture of basally worked and rockfall debris,  
231 which texturally are typically sandy boulder-gravels. Downwasting of glaciers since the Little  
232 Ice Age (LIA) (Rowan, 2017) have left ice-cored moraines up to a hundred metres above the  
233 glacier surface, which result in an unstable inner moraine face that is unstable and prone to  
234 collapse (Hambrey *et al.* 2008).

235

236 Debris-covered Himalayan glaciers tend to lose mass by surface lowering rather than  
237 terminus recession (Rowan *et al.* 2015, Quincey *et al.* 2009, Bolch *et al.* 2011). Surface  
238 lowering causes these debris-covered glaciers to develop very low or even reversed long-  
239 profile topographic gradients through their ablation areas, which promotes the formation of  
240 supraglacial water bodies. These ponds and lakes influence the seasonal transport of water  
241 through the glacial system, and can expand and coalesce to form substantial supraglacial or  
242 proglacial, moraine-dammed lakes that may eventually pose a potential flood hazard (Watson  
243 *et al.* 2016, Thompson *et al.* 2012). Such features are commonly bordered by steep, debris-  
244 free ice cliffs, which progressively backwaste, and, if connected to a supraglacial pond or  
245 lake, may undergo thermoerosional notch development at the ice-water interface, promoting  
246 the onset of calving processes (Fig. 2) (Hambrey *et al.* 2008, Thompson *et al.* 2016).

247

248 Satellite observations of glacier mass change suggest that debris-covered glaciers in the  
249 Himalaya and Karakoram may be losing mass at the same rate as those glaciers with clean-  
250 ice (debris-free) surfaces. This ‘debris-cover anomaly’ could be due to enhancement of  
251 ablation at ice cliff faces. Although the exposure of clean ice at these ice cliffs can  
252 dramatically enhance local ablation rates (Miles *et al.* 2016, Brun *et al.* 2016, Reid & Brock  
253 2014) field observations from Changri Nup Glacier in the Everest region suggest that, despite  
254 the presence of these ablation ‘hotspots’, a continuous or near-continuous mantle of  
255 supraglacial debris reduces net ablation, such that glacier-wide mass loss is less than would  
256 be the case for an equivalent clean-ice glacier (Vincent *et al.* 2016). To fully understand the  
257 effect of ice cliffs on ablation from debris-covered glaciers, these features and their evolution  
258 need to be incorporated into glacier-wide surface energy balance modelling (e.g. Buri *et al.*  
259 2016; Brun *et al.* 2016).

260

261

### 262 **3. Changes in glacier volume during the Late Holocene**

263 Changes in the areal extent and volume of glaciers over the last 2,000 years can be inferred  
264 from moraines that indicate the position of glacier margins, historical observations made by  
265 climbing expeditions, and field and satellite measurements of glacier geometries.

266

#### 267 **3.1 Late Holocene (2,000 years ago to present)**

268 Many glaciers in High Mountain Asia have advanced and receded two or three times during  
269 the last 2,000 years in response to climate change (Owen & Dortch 2014, Murari *et al.* 2014,  
270 Rowan 2017) and followed the global trend of glacier recession and shrinkage since about  
271 1850 (Thompson *et al.* 2006). The last period of regional glacier advance was the LIA which  
272 peaked between 1300 and 1600, although glaciers remained close to their LIA limits until the  
273 20<sup>th</sup> Century (Rowan 2017). These observations are based on geochronological data for  
274 moraines compiled from studies using radiocarbon (<sup>14</sup>C) dating (e.g. Muller 1961,  
275 Röthlisberger & Geyh 1986) and terrestrial cosmogenic nuclide dating (e.g. Owen 2009,  
276 Murari *et al.* 2014). More recent applications of these techniques generally provide more  
277 accurate results due to improvements in laboratory measurement protocols. Regional glacier  
278 volume change in the Himalaya over decadal to centennial timescales occurred in response to  
279 hemispheric changes in air temperature (Solomina *et al.* 2016, Rowan 2017). However,  
280 variations in the timing and extent of glacier volume change across this range are primarily  
281 driven by millennial-scale east–west and north–south variations in atmospheric circulation  
282 regimes (National Research Council 2012). The characteristics of local weather systems,  
283 particularly precipitation distribution, are also important and probably governed by  
284 precession-scale insolation cycles (Thompson *et al.* 2006). Consequently, moraine ages  
285 indicate spatial variability in the amount and timing of glacier mass loss due to variations in  
286 the timing and intensity of monsoonal and Westerly snowfall across the region (Rowan 2017,  
287 Owen 2009).

288

#### 289 **3.2 20<sup>th</sup> and 21<sup>st</sup> Centuries (1900 to present)**

290 Changes in glacier length and area during the early part of the 20<sup>th</sup> Century are described by  
291 historical accounts from early climbing expeditions. These records are based on visual  
292 comparison of the state of these glaciers to those in the European Alps, which has led to  
293 misinterpretation of ongoing glacier volume change (Grove 2004). Measurements of changes  
294 in length and area are of limited use for estimating the mass change of debris-covered  
295 glaciers that lose mass by surface lowering rather than terminus recession (e.g. Bolch *et al.*  
296 2011). Geochronological techniques such as <sup>14</sup>C and terrestrial cosmogenic nuclide dating do  
297 not currently operate at sufficient temporal resolution to describe the ages of moraines  
298 formed in the last 100 years. However, changes in glacier volume over small areas can be

299 accurately detected by comparing multi-temporal aerial and satellite topographic data,  
300 including historical imagery from the Corona and HEXAGON satellites that date back to the  
301 1950s (e.g. Bolch *et al.* 2011; Berthier *et al.* 2014). Measurements of the gravitational field of  
302 the Earth's surface (the Gravity Recovery and Climate Experiment; GRACE; Tapley *et al.*  
303 2004) combined with topographic data can be used to estimate changes in glacier mass across  
304 a broad spatial area (e.g. Moiwu *et al.* 2011) but with large uncertainties (Gardner *et al.*  
305 2013).

306

307 Analyses of ice cores demonstrate a sharp decrease in accumulation on low-latitude (25–  
308 35°N) Himalayan glaciers, and an increase in ice volume at higher latitudes (35–70°N) on the  
309 Tibetan Plateau driven by variability in monsoon intensity and timing since 1950 (Hou *et al.*  
310 2002). However, glacier mass loss at high elevations has exceeded that which could be  
311 attributed to change in monsoon intensity alone (Mölg *et al.* 2012). Mass balance is most  
312 negative in the Eastern Himalaya, and becomes less negative to the north and in the northern  
313 and eastern parts of the Karakoram where some glaciers showed slightly positive mass  
314 balances between 1999 and 2008 (Gardelle *et al.* 2012). The opposing trends in glacier mass  
315 balance between the Karakoram and the Eastern Himalaya over the last 50 years are  
316 attributed to spatial variations in the rates of change in temperature and precipitation  
317 (Gardelle *et al.* 2012, Nakawo *et al.* 1999), as rising Northern Hemisphere air temperatures  
318 deliver winter snowfall from the Arabian Gulf further into the range (Kapnick *et al.* 2014).  
319 Climate warming appears to have accelerated the mass loss from glaciers in the Himalaya  
320 after 1995, reflecting the high sensitivity of the regional energy balance to small changes in  
321 climate (Cogley 2011). Over the same period, a slight gain in mass has been observed for  
322 glaciers in the Karakoram, attributed to a greater influence of Westerly winter snowfall  
323 (Gardelle *et al.* 2012, Yao *et al.* 2012), although not all glaciers in the Karakoram gained  
324 mass in the last 40 years (Sarikaya *et al.* 2012).

325

#### 326 **4. Predictions of future glacier change**

327 Predictions of how glaciers will continue to change from the present day requires  
328 quantifying, specifically: current glacier mass balances, the response time over which glaciers  
329 will reach equilibrium with climate, and how the climate will change over the period of  
330 interest. For the 21<sup>st</sup> and 22<sup>nd</sup> Centuries, climate model ensembles such as those produced by  
331 the IPCC (Collins *et al.* 2013) give a range of possible warming values for future emissions  
332 scenarios, which are useful for forcing meteorological and glacier modelling. Glacier models  
333 are often somewhat less sophisticated than these climate models and operate at different  
334 spatial scales, particularly in representing the dynamics of mountain glaciers, in which the  
335 processes controlling the flow of ice through steep rugged terrain and where feedbacks with  
336 often extreme topography and orographic meteorology are poorly documented. The rate of

337 regional glacier change in High Mountain Asia may also be enhanced when compared to  
338 lower-altitude glacierised regions, as Northern Hemisphere warming is enhanced at altitudes  
339 above 5000 m (Xu *et al.* 2016) where the ELAs of the many large glaciers are located (Benn  
340 & Lehmkuhl, 2000). To better understand the impact of climate change on glacier mass  
341 balance, meteorological variables on a smaller spatial scale than the entire region are needed.

342

343 Predictions of glacier response to future climate change can be made either by extrapolating  
344 from observations of recent glacier change and present-day glacier characteristics, or by  
345 applying numerical glacier–climate models. These glacier models vary widely in their level  
346 of sophistication and complexity depending on the required application, but generally can  
347 either extrapolate from observed trends in the relationship between glacier mass balance and  
348 climate, or replicate the physical processes by which glacier change occurs and be forced by  
349 changing climate conditions. Numerical modelling of glacier mass balance forced by detailed  
350 simulations of mesoscale meteorology has been undertaken to better understand the  
351 atmospheric controls on Zhadang Glacier in central Tibet (Mölg *et al.* 2012), but is still in  
352 development for regional applications. Glacier-climate models can be used to make  
353 catchment-scale and regional-scale predictions of the contribution of glacial meltwater to  
354 hydrological budgets, and their contribution in the context of water supplied by precipitation  
355 or groundwater flow (e.g. Lutz *et al.* 2014). However, predictions based on numerical  
356 modelling must also consider the range of uncertainties associated with the data used to drive  
357 models. Many of these climatic and glaciological variables, such as the relationship between  
358 air temperature and ablation beneath supraglacial debris, or the subglacial conditions  
359 controlling ice flow, are poorly constrained both at present and in terms of future change in  
360 the Himalaya (Rowan *et al.* 2015).

361

362 Precipitation in the monsoon-influenced Eastern and Central Himalaya is predicted to  
363 increase by up to 10% during the 21<sup>st</sup> Century (IPCC scenario A2; Collins *et al.* 2013).  
364 Although this increase in precipitation would mean that widespread droughts are unlikely,  
365 with warmer air temperatures a greater proportion of precipitation will fall as rain rather than  
366 snow and will melt glacier ice (Meehl *et al.* 2007). The mass balances of the majority of  
367 glaciers in High Mountain Asia are out of equilibrium with present-day climate, as is the case  
368 for glaciers worldwide. A degree day model of the Eastern Himalaya based on a 20-year  
369 climate record demonstrated that loss of 25% of the glacierised area could occur with only  
370 1°C warming from present (Rupper *et al.* 2012). Under an extreme scenario of 2.5°C  
371 warming by the end of the 21<sup>st</sup> Century, the glacierised area of Bhutan would be reduced by  
372 50%, and the contribution of meltwater flux to annual hydrological budgets would become  
373 negligible (Rupper *et al.* 2012). Catchment-scale hydrological modelling of the Langtang  
374 catchment in Nepal, a typical high-altitude valley in the Central Himalaya, suggest a loss of

375 35–55% of the total glacierised area by 2100, with the contribution of areal loss from debris-  
376 covered glaciers only 25–33% over the same period (Ragettli *et al.* 2016).

377

### 378 **5. Impacts on water resources with future glacier change**

379 Until 2050, if only the contribution of glaciers to the hydrological budget is considered, river  
380 flows are likely to rise during the monsoon an effect called the ‘deglaciation discharge  
381 dividend’ (Fig. 4) (Kaser *et al.* 2010). River flows will reach ‘peak water’ then decline as  
382 glacier mass is dramatically reduced and rivers have a greater dependence on precipitation  
383 and snow melt (Soncini *et al.* 2016). ‘Peak water’ in monsoon-influenced regions is predicted  
384 to occur by the mid-21<sup>st</sup> Century, as identified by hydrological modelling of glacier  
385 meltwater production (Lutz *et al.* 2014). In Nepal, glacier mass loss is predicted to increase  
386 downstream water supplies during the first half of the 21<sup>st</sup> Century compared to 2001–2010,  
387 as the additional meltwater released each year will boost river flows. Water supplies are then  
388 either predicted to decline or remain stable depending on how the monsoon changes during  
389 this period, as the predicted 10% decrease in meltwater runoff could be compensated by a  
390 similar increase in precipitation (Ragettli *et al.* 2016). The contribution of glacier meltwater  
391 to future river flows may increase slightly during the monsoon due to enhanced ice melt, but  
392 decrease overall by 4% by 2050 as glacier mass rapidly declines (Soncini *et al.* 2016).  
393 Hydrological modelling predicts a decline in the glacial meltwater contribution to catchment  
394 hydrological budgets over the next century; by 2065 the change in mean catchment water  
395 supply is likely to be –8% in the Indus, –18% in the Ganges, and –20% in the Brahmaputra  
396 (Immerzeel *et al.* 2010). These decreases in meltwater supply are likely to be compensated, at  
397 least partially, by increasing rainfall of +25% in the Indus and Brahmaputra and +8% in the  
398 Ganges. However, these projections should be treated with caution, since changes in the  
399 monsoon are currently difficult to represent in predictive climate models (Immerzeel *et al.*  
400 2010).

401

402 Short-term increases in river flow as rainfall becomes a more important constituent of the  
403 hydrological budget are likely to increase the risk of regional flooding (Ragettli *et al.* 2016).  
404 However, the magnitude and timing of peak flows relative to the present day are generally  
405 unknown. The expansion and coalescence of supraglacial melt ponds to form larger  
406 supraglacial or proglacial lakes bounded by terminal and lateral moraines presents an  
407 additional risk in the form of the hazard posed by potentially catastrophic glacial lake  
408 outburst floods (Benn *et al.* 2012). These sudden-onset floods generally arise from the  
409 breaching of an impounding moraine, and are capable of generating peak flood discharges  
410 that can exceed seasonal high flow floods by over an order of magnitude (Cenderelli and  
411 Wohl, 2001). Large glacial lakes may also be considered as an intermediate storage  
412 component in the hydrological cascade of glacierised (and generally deglaciating) catchments

413 and effectively regulate the downstream transmission of glacial meltwater (Carrivick and  
414 Tweed, 2013). An anticipated increase in the number and extent of glacial lakes as a result of  
415 climate change is of concern, especially when considered in the context of the rapidly  
416 expanding Asian hydropower sector, which is likely to become increasingly exposed to  
417 climatically controlled glacial flood hazards (Schwanghart *et al.* 2016), and modified  
418 hydrological regimes.

419

420 Beyond 2050, sustained glacier mass loss will result in declining water supplies and possible  
421 shifting of seasonal river flows, as spring meltwater will no longer sufficiently compensate  
422 for the pre-monsoon dry season (Immerzeel *et al.* 2010). River flow will decline most  
423 dramatically in the Indus where a significant proportion of the annual hydrograph is derived  
424 from glacier melt (Fig. 5) (Lutz *et al.* 2014). Future river flows will depend on changes in the  
425 amount and timing of precipitation, and highly seasonal river flows are likely to change their  
426 timing compared to the present day, possibly resulting in enhanced spring flows (Immerzeel  
427 *et al.* 2010). Total hydrological budgets are likely to decline dramatically by 2100, with  
428 extreme scenarios predicting a 26% decrease in flow predicted by 2100 for the Everest  
429 region, due to glaciers losing over 50% of their volume compared to 2012–2014 (Soncini *et*  
430 *al.* 2016).

431

## 432 **6. Improving understanding of glacier response to climate change**

433 Whilst there are large uncertainties about the current state of and the ongoing changes  
434 experienced by glaciers, conclusions can nevertheless be drawn about important controls on  
435 their response to climate change to make predictions of their future state (Bolch *et al.* 2012).  
436 These predictions often contain large uncertainties, due to a lack of available data with which  
437 to evaluate models, and the suitability of existing glacier models which have often been  
438 developed for application to Polar ice sheets rather than glaciers flowing through steep,  
439 mountainous terrain. Many factors control the response of mountain glaciers to climate  
440 change (Fig. 6), and spatial and temporal variability in these controls can be challenging to  
441 represent in numerical models. Some key areas for possible future research to reduce these  
442 uncertainties are described here.

443

### 444 ***6.1 Modification of glacier response to climate change by catchment geomorphology***

445 The dynamics and hydrology of individual glaciers are governed by characteristics such as  
446 glacier aspect, size, altitude, and hypsometry, collectively known as morphometry. These  
447 morphometric factors exert a significant control on the dynamics and mass balance of  
448 mountain glaciers (Quincey *et al.* 2009). Moreover, the pronounced interaction between high  
449 topography and atmospheric circulation systems such as the Indian summer monsoon results  
450 in distinctive local mesoscale meteorological patterns (Bookhagen & Burbank 2010). This

451 interaction between the atmosphere, landscape and cryosphere can produce catchment-scale  
452 variations in energy and mass balance that cause adjacent glaciers to exhibit different  
453 responses to the same change in climate (e.g. Glasser *et al.* 2009). For this reason, a coupled  
454 mesoscale–energy balance modeling approach represents an important advance in the  
455 understanding of glacier–climate interactions in High Mountain Asia (Mölg *et al.* 2012).  
456 Robust dynamic or statistical methods are needed to downscale climate model outputs to a  
457 scale relevant to glacier mass balance that accounts for mountainous topography (e.g.  
458 Reichert *et al.* 2002). Furthermore, these relationships may need to be reconsidered for  
459 application using future climatologies (Meehl *et al.* 2007). The degree-day model  
460 applications, such as that used by Rupper *et al.* (2012) are useful to predict regional  
461 glaciological and hydrological changes with climate variations. However, the modification of  
462 glacier–climate relationships by factors such as surface debris cover and glacier morphometry  
463 requires further exploration and the acquisition of field data for model evaluation and testing.

464

## 465 ***6.2 Sensitivity of debris-covered glaciers to climate change***

466 As glaciers lose mass, debris accumulates on their surfaces, and as a result, the debris-  
467 covered glacierised area worldwide is increasing. Ablation under a supraglacial debris layer  
468 is primarily controlled by its thickness, but to a lesser extent by debris properties including  
469 lithology, moisture content and porosity (Benn *et al.* 2012). These parameters are spatially  
470 and temporally variable, due to variations in input, transport and exhumation of debris to the  
471 glacier surface in space and time (Anderson & Anderson, 2016; Gibson *et al.* 2017). Much  
472 recent work has focused on determining spatial variability in debris thickness, either remotely  
473 using thermal satellite imagery (e.g. Mihalcea *et al.* 2008; Foster *et al.* 2012; Rounce &  
474 McKinney, 2014) or directly using ground-penetrating radar (e.g. McCarthy *et al.* 2017) and,  
475 in some cases, the impact of this spatial variation on glacial hydrology (Minora *et al.* 2015;  
476 Soncini *et al.* 2016). Many of these inputs are validated with minimal field measurements of  
477 debris thickness, but such validation would greatly extent the scope of predictions that could  
478 be made considering debris-covered glacier change. Few studies currently consider the  
479 influence of spatial variation in moisture content (e.g. Collier *et al.* 2014), porosity (e.g. Evatt  
480 *et al.* 2015), albedo (e.g. Nicholson & Benn, 2013) or aerodynamic roughness length (e.g.  
481 Rounce *et al.* 2015; Miles *et al.* 2017; Quincey *et al.* 2017).

482

483 Predictions of glacier mass balance modified by debris cover requires distributed surface  
484 energy balance models that consider variations in debris cover across the glacier surface and  
485 through time, and ideally simulate the interaction of free air and moisture with the porosity of  
486 debris layers (e.g. Evatt *et al.* 2015, Collier *et al.* 2014). However, suitable models are few,  
487 often only consider one important variable (e.g. debris thickness or porosity), and are mainly  
488 driven by empirical relationships derived from limited field data (e.g. Mihalcea *et al.* 2008).

489 Therefore, to comprehensively understand the influence of a debris layer on ablation, further  
490 field data is needed to quantify the extent of variations in debris parameters and to develop  
491 understanding of heat flux through supraglacial debris. It is potentially possible to use field  
492 measurements of debris thickness to calibrate and validate a method of classification of this  
493 variable from remote observations, which would greatly extend the scope of predictions that  
494 could be made considering debris-covered glacier change. Furthermore, as field-based  
495 research tends to focus on relatively accessible, often smaller lower altitude glaciers, a remote  
496 calibration method for debris thickness has the potential to dramatically advance knowledge  
497 of how these glaciers behave in response to climatic forcing.

498

### 499 ***6.3 Modification of glacier response to climate by glacier dynamics***

500 Feedbacks between glacier mass balance and dynamics control the magnitude and timing of  
501 the response of individual glaciers to climate change. Ice flow also drives processes that  
502 affect glacier mass. For example, the transport of debris to the glacier surface from englacial  
503 or subglacial storage can cause the supraglacial debris layer to thicken and thereby reduce  
504 ablation. In contrast, ice flow stagnation may promote the development and expansion of  
505 supraglacial or moraine-dammed lakes, which promotes widespread calving of the lake-  
506 terminating glacier tongue and thereby accelerating mass loss (Gardelle *et al.* 2011). Debris  
507 cover frequently causes glacier tongues to lose mass by surface lowering rather than terminus  
508 recession, which in turn affects dynamics, since ice flow tends to stagnate with the loss of  
509 driving stress (Quincey *et al.* 2009). Commonly, supraglacial or proglacial lake formation  
510 coincides with this process, and once a lake crosses a threshold of depth of about 80 m deep  
511 the lake-marginal glacier ice starts to calve (Quincey *et al.* 2007, Robertson *et al.* 2012) with  
512 potentially dramatic consequences for glacier dynamics and mass balance. Under IPCC 21<sup>st</sup>  
513 Century climate change scenarios, proglacial lakes are increasingly likely to pose a potential  
514 hazard to human life through an increased risk of sudden-onset glacial lake outburst floods.  
515 However, the timing and magnitude of lake formation and growth are difficult to predict.

516

517 Rates of recent proglacial lake expansion have been quantified from satellite imagery and are  
518 particularly well-studied in the Khumbu Himal in the Central Himalaya (Watson *et al.* 2016)  
519 and the Lunana region of Bhutan in the Eastern Himalaya (Fujita *et al.* 2008). The primary  
520 process by which lakes expand appears to be via the subaerial loss of mass at the active  
521 calving front rather than the ablation of subaqueous ice, including ice at the lake bottom  
522 (Fujita *et al.* 2009). Controls on calving rates for glaciers terminating in freshwater rather  
523 than marine settings differ, and are dominated by wave fetch and lake temperature (Sakai *et*  
524 *al.* 2009). However, prediction of future change in these variables and the impact on lake  
525 development has not been investigated. Proglacial lake growth has typically accelerated with  
526 rapid glacier recession since the 1960s (Bajracharya & Mool 2010), but the impact of lake

527 formation on future glacier change is poorly understood. Although it seems intuitive that  
528 lake-terminating glaciers should recede more rapidly than equivalent land-terminating  
529 glaciers, few conclusive data are yet available to confirm this (e.g. King *et al.* 2017).

530

## 531 **7. Conclusions**

532 Glaciers in High Mountain Asia are changing rapidly in response to recent climate change,  
533 with many glaciers losing mass at accelerated rates since the 1990s. Sustained glacier mass  
534 loss is predicted to continue through the 21<sup>st</sup> and 22<sup>nd</sup> Centuries even in the absence of further  
535 climate warming from the present day. In the monsoon-influenced Eastern and Central  
536 Himalaya, the majority of glacier mass loss occurs during the warm summer months (June to  
537 September), which coincides with the timing of maximum precipitation at high altitudes. As a  
538 result, the glacier meltwater contribution to the Ganges and Brahmaputra catchments is  
539 relatively less important than for the Indus catchment, which drains the Westerly influenced  
540 Western Himalaya, Karakoram and Hindu Kush and has a smaller rainfall component of the  
541 annual hydrological budget. Seasonal river flows in the Eastern and Central Himalaya are  
542 therefore unlikely to decrease during the next 50 years and may even increase slightly, the  
543 ‘deglaciation discharge dividend’, as climate models predict increased monsoon precipitation  
544 with warming during this century, and because accelerated glacier melt will provide  
545 additional water over the same period. In the Western Himalaya, Karakoram and Hindu  
546 Kush, river flows are much more dependent on glacier mass change than in the catchments  
547 further to the east. The future of these glaciers is a less clear, as those in the Karakoram  
548 appear to have recently experienced a slightly increase in ice mass which may be due to the  
549 increased extent of winter snowfall resulting from warming air temperatures. However, in the  
550 Western Himalaya, glacier mass loss appears similar to that elsewhere in the mountain range,  
551 suggesting that the hydrological budget of the Indus is likely to be severely affected by  
552 climate change.

553

554 Beyond the mid-21<sup>st</sup> Century, when large volumes of glacier ice have been lost, the  
555 hydrological budget of catchments in the monsoon-influenced regions will be more  
556 dependent on the timing and availability of monsoon precipitation than glacier melt.  
557 Therefore, water availability is unlikely to decrease in the short term, as a warming climate  
558 will result in decreasing glacier runoff compensated by increased monsoon precipitation, with  
559 little change in the seasonality of river flows. In the Westerly influenced regions, glacier  
560 mass loss will likely lead to decreased river flows, and although glaciers in the Karakoram  
561 are at present showing slightly positive mass balances, this trend is unlikely to continue with  
562 sustained climate warming in the longer-term. As rainfall becomes a more important  
563 component of the total hydrological budget, the risk of flooding is predicted to increase as  
564 rainfall is transferred much more rapidly into rivers than snow that accumulates as glacier ice.

565 In the longer term, by the start of the 22<sup>nd</sup> Century, the predicted loss of over 50% of glacier  
566 volume and the complete removal of smaller, lower-altitude glaciers across High Mountain  
567 Asia is likely to result in a widespread decline in water supplies, which will have a dramatic  
568 impact on the large populations relying on these glacier-fed rivers.

569

### 570 **Acknowledgements**

571 We thank Tobias Bolch and Arthur Lutz for sharing regional and catchment boundaries and  
572 model outputs used to draw Figures 1 and 5.

573

574

### 575 **Figure captions**

576 Figure 1. High Mountain Asia, showing the Himalaya, Karakoram and Hindu Kush regions  
577 defined by Bolch *et al.* (2012) and major rivers. The location of glaciers for which  
578 glaciological mass balance records exist and the length of the record in years (Bolch *et al.*  
579 2012; Pratap *et al.* 2016; Soncini *et al.* 2016; Tshering & Fujita 2016), and Vincent *et al.*  
580 (2016), and the location of automatic weather stations (AWS) collecting data at high altitudes  
581 are also indicated. The names of individual glaciers or glacierised regions referred to in the  
582 text are highlighted in bold.

583

584 Figure 2. The surface features of Khumbu Glacier in the Everest region of Nepal, showing (a)  
585 the debris-covered ablation area, looking to the south, (b) the Khumbu Icefall marking the  
586 transition between the clean-ice accumulation area and the debris-covered ablation area, (c  
587 and d) typical ice cliffs and supraglacial ponds in the ablation area showing englacial debris  
588 layers within the ice, likely resulting from ice-rock avalanching (note figures circled in red  
589 for scale).

590

591 Figure 3. Direct measurements of mass balance for six glaciers in the Himalaya between  
592 1992 and 2012, showing; (a) annual mass balance, (b) cumulative annual mass balance, and  
593 (c) cumulative mass balance normalised by glacier terminus altitude. Note that the data for  
594 Kangwure Glacier between 1994 and 2008 (dashed line) are reconstructed mass balance  
595 values derived from meteorological data using the relationship between *in situ* mass balance  
596 measurements made in other years. Redrawn from Yao *et al.* (2012). See Figure 1 for the  
597 locations of these glaciers.

598

599 Figure 4. Schematic diagram showing hypothetical changes in glacier volume and meltwater  
600 release from mountain glaciers in response to regional climate warming over a period  
601 equivalent to their last advance and recession, from the Little Ice Age maximum through the  
602 present day and 21<sup>st</sup> Century.

603

604 Figure 5. Annual hydrographs for headwaters of the major Himalayan catchments produced  
605 using a hydrological model for a present-day reference period of 1998–2007, showing the  
606 contribution to the total hydrological budget from glacier melt, snow melt, rainfall–runoff  
607 and base (groundwater) flow [redrawn from Lutz *et al.* (2014)], for (a) the Indus in the  
608 westerly influenced Western Himalaya, (b) the Ganges in the transition between the westerly  
609 and monsoon-influenced Central Himalaya, and (c) the Brahmaputra in the monsoon-  
610 influenced Central and Eastern Himalaya. (d) shows the catchment boundaries used to make  
611 these calculations, with topographic imagery from Google OpenLayers.

612

613 Figure 6. Climatic, glaciological and landscape space–time controls on glacier and climate  
614 change in the Himalaya and Karakoram.

615

616

617

618

## 619 **References**

- 620 Adhikari, S. & Huybrechts, P. 2009. Numerical modelling of historical front variations and  
621 the 21st-century evolution of glacier AX010, Nepal Himalaya. *Annals of Glaciology*,  
622 **50**, 27–34.
- 623 Ageta, Y., Ohata, T., Tanaka, Y., Ikegami, K., & Higuchi, K. 1980. Mass Balance of  
624 Glacier AX010 in Shorong Himal, East Nepal during the Summer Monsoon Season.  
625 *Seppyo*, **41**, 34–41.
- 626 Anderson, L.S. & Anderson, R.S., 2016. Modeling debris-covered glaciers: response to  
627 steady debris deposition. *The Cryosphere*, **10**, 1105–1124.
- 628 Arendt, A., A. Bliss, T. Bolch, J.G. Cogley, A.S. Gardner, J.-O. Hagen, R. Hock, M. Huss,  
629 G. Kaser, C. Kienholz, W.T. Pfeffer, G. Moholdt, F. Paul, V. & Radić, *et al.*, 2015,  
630 Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 5.0.  
631 Global Land Ice Measurements from Space, Boulder Colorado, USA. Digital Media.
- 632 Azam, M.F., Wagnon, P., Ramanathan, A., Vincent, C., Sharma, P., Arnaud, Y., Linda, A.,  
633 Pottakkal, J.G., Chevallier, P., Singh, V.B. & Berthier, E., 2012. From balance to  
634 imbalance: a shift in the dynamic behaviour of Chhota Shigri glacier, western  
635 Himalaya, India. *Journal of Glaciology*, **58**, 315–324.
- 636 Bajracharya, S.R. & Mool, P. 2010. Glaciers, glacial lakes and glacial lake outburst floods in  
637 the Mount Everest region, Nepal. *Annals of Glaciology*, **50**, 81–86.
- 638 Benn, D.I. & Lehmkuhl, F. 2000. Mass balance and equilibrium-line altitudes of glaciers in  
639 high-mountain environments. *Quaternary International*, **65**, 15–29.

- 640 Benn, D.I., Owen, L.A., Osmaston, H.A., Seltzer, G.O., Porter, S.C. & Mark, B. 2005.  
641 Reconstruction of equilibrium-line altitudes for tropical and sub-tropical glaciers.  
642 *Quaternary International*, **138**, 8–21.
- 643 Benn, D.I., Bolch, T., Hands, K., Gulley, J., Luckman, A., Nicholson, L.I., Quincey, D.,  
644 Thompson, S., Toumi, R. & Wiseman, S., 2012. Response of debris-covered glaciers  
645 in the Mount Everest region to recent warming, and implications for outburst flood  
646 hazards. *Earth Science Reviews*, **114**, 156–174.
- 647 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á.P., Pitte, P., Le Meur, E.,  
648 Masiokas, M., Ruiz, L., Pálsson, F., Belart, J.M.C. & Wagnon, P., 2014. Glacier  
649 topography and elevation changes derived from Pléiades sub-meter stereo images.  
650 *The Cryosphere*, **8**, 2275–2291.
- 651 Bolch, T., Pieczonka, T. & Benn, D.I. 2011. Multi-decadal mass loss of glaciers in the  
652 Everest area (Nepal Himalaya) derived from stereo imagery. *The Cryosphere*, **5**,  
653 349–358.
- 654 Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S.,  
655 Fujita, K., Scheel, M. & Bajracharya, S., 2012. The state and fate of Himalayan  
656 glaciers. *Science*, **336**, 310-314.
- 657 Bookhagen, B. & Burbank, D. 2010. Towards a complete Himalayan hydrologic budget: The  
658 spatiotemporal distribution of snow melt and rainfall and their impact on river  
659 discharge. *Journal of Geophysical Research*, **115**, F03019.
- 660 Brun, F., Buri, P., Miles, E.S., Wagnon, P., Steiner, J., Berthier, E., Ragetti, S.,  
661 Kraaijenbrink, P., Immerzeel, W.W. & Pellicciotti, F. 2016. Quantifying volume loss  
662 from ice cliffs on debris-covered glaciers using high-resolution terrestrial and aerial  
663 photogrammetry. *Journal of Glaciology*, 1–12.
- 664 Brun, F., Berthier, E., Wagnon, P., Kääb, A. & Treichler, D. 2017. A spatially resolved  
665 estimate of High Mountain Asia glacier mass balances from 2000 to 2016, *Nature*  
666 *Geoscience*, **482**, 514–7.
- 667 Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P. & Pellicciotti, F. 2016. A  
668 physically based 3 - D model of ice cliff evolution over debris-covered glaciers.  
669 *Journal of Geophysical Research: Earth Surface*, **121**, 2471-2493.
- 670 Carrivick, J.L. & Tweed, F.S. 2013. Proglacial lakes: character, behaviour and geological  
671 importance. *Quaternary Science Reviews*, **78**, 34-52.
- 672 Cenderelli, D.A. & Wohl, E.E. 2001. Peak discharge estimates of glacial-lake outburst  
673 floods and "normal" climatic floods in the Mount Everest region, Nepal.  
674 *Geomorphology*, **1**, 57-90.
- 675 Cogley, J.G. 2011. Himalayan Glaciers in 2010 and 2035. In: Singh, V. P., Singh, P. &  
676 Haritashya, U. (Eds.) *Encyclopedia of Snow, Ice and Glaciers*. Springer, Dordrecht.

- 677 Collier, E., Nicholson, L.I., Brock, B.W., Maussion, F., Essery, R. & Bush, A.B.G. 2014.  
678 Representing moisture fluxes and phase changes in glacier debris cover using a  
679 reservoir approach. *The Cryosphere*, **8**, 1429–1444.
- 680 Collins, M., Knutti, R. & Arblaster, J. 2013. Long-term Climate Change: Projections,  
681 Commitments and Irreversibility. In: Climate Change 2013: The Physical Science  
682 Basis. Contribution of Working Group I to the Fifth Assessment Report of the  
683 Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner,  
684 M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley  
685 (Eds.)]. Cambridge University Press, Cambridge, 1–108.
- 686 Evatt, G.W., Abrahams, I.D., Heil, M. & Mayer, C. 2015. Glacial melt under a porous debris  
687 layer. *Journal of Glaciology*, **61**, 825–836.
- 688 Foster, L.A. Brock, B.W., Cutler, M.E.J. & Diotri, 2012. A physically based method for  
689 estimating supraglacial debris thickness from thermal band remote-sensing data.  
690 *Journal of Glaciology*, **58**, 677–691.
- 691 Frey, H., Machguth, H., Huss, M., Huggel, C., Bajracharya, S., Bolch, T., Kulkarni, A.,  
692 Linsbauer, A., Salzmann, N. & Stoffel, M. 2014. Estimating the volume of glaciers in  
693 the Himalaya–Karakoram region using different methods, *The Cryosphere*, **8**, 2313–  
694 2333.
- 695 Fujita, K. & Nuimura, T. 2011. Spatially heterogeneous wastage of Himalayan glaciers.  
696 *Proceedings of the National Academy of Sciences*, **108**, 14011–14014.
- 697 Fujita, K., Suzuki, R., Nuimura, T. & Sakai, A. 2008. Performance of ASTER and SRTM  
698 DEMs, and their potential for assessing glacial lakes in the Lunana region, Bhutan  
699 Himalaya. *Journal of Glaciology*, **54**, 220–228.
- 700 Fujita, K., Sakai, A., Nuimura, T., Yamaguchi, S. & Sharma, R.R. 2009. Recent changes in  
701 Imja Glacial Lake and its damming moraine in the Nepal Himalaya revealed by *in*  
702 *situ* surveys and multi-temporal ASTER imagery. *Environmental Research Letters*, **4**,  
703 045205.
- 704 Gades, A., Conway, H., Nereson, N., Naito, N. & Kadota, T. 2000. Radio echo-sounding  
705 through supraglacial debris on Lirung and Khumbu Glaciers, Nepal Himalayas.  
706 *Publications of the International Association of Hydrological Sciences*, **264**, 13–24.
- 707 Gardner, A.S., Moholdt, G., Cogley, J.G., Wouters, B., Arendt, A.A., Wahr, J., Berthier, E.,  
708 Hock, R., Pfeffer, W.T., Kaser, G. & Ligtenberg, S.R., 2013. A reconciled estimate  
709 of glacier contributions to sea level rise: 2003 to 2009. *Science*, **340**, 852–857.
- 710 Gardelle, J., Arnaud, Y. & Berthier, E., 2011. Contrasted evolution of glacial lakes along the  
711 Hindu Kush Himalaya mountain range between 1990 and 2009. *Global and*  
712 *Planetary Change*, **75**, 47–55.
- 713 Gardelle, J., Berthier, E. & Arnaud, Y. 2012. Slight mass gain of Karakoram glaciers in the  
714 early twenty-first century. *Nature Geoscience*, **5**, 322–325.

- 715 Gibson, M. J., Glasser, N. F., Quincey, D. J., Mayer, C., Rowan, A. V. & Irvine-Fynn, T. D.  
716 L. 2017. Temporal variations in supraglacial debris distribution on Baltoro Glacier,  
717 Karakoram between 2001 and 2012. *Geomorphology*, **295**, 572–585.
- 718 Glasser, N.F., Harrison, S. & Jansson, K.N. 2009. Topographic controls on glacier sediment-  
719 landform associations around the temperate North Patagonian Icefield. *Quaternary*  
720 *Science Reviews*, **28**, 2817–2832.
- 721 Grove, J.M. 2004. *The Little Ice Age: Ancient and Modern*, 2nd Ed. Routledge.
- 722 Hambrey, M.J., Quincey, D.J., Glasser, N.F., Reynolds, J.M., Richardson, S.J. & Clemmens,  
723 S. 2008. Sedimentological, geomorphological and dynamic context of debris-mantled  
724 glaciers, Mount Everest (Sagarmatha) region, Nepal. *Quaternary Science Reviews*,  
725 **27**, 2361–2389.
- 726 Harper, J. & Humphrey, N. 2003. High altitude Himalayan climate inferred from glacial ice  
727 flux. *Geophysical Research Letters*, **30**, 1764.
- 728 Hou, S., Qin, D., Yao, T., Zhang, D. & Chen, T. 2002. Recent change of the ice core  
729 accumulation rates on the Qinghai-Tibetan Plateau. *Chinese Science Bulletin*, **47**,  
730 1746–1749.
- 731 Immerzeel, W.W., van Beek, L.P.H. & Bierkens, M.F.P. 2010. Climate Change Will Affect  
732 the Asian Water Towers. *Science*, **328**, 1382–1385.
- 733 Kadota, T., Fujita, K., Seko, K., Kayastha, R.B. & Ageta, Y. 1997. Monitoring and  
734 prediction of shrinkage of a small glacier in the Nepal Himalaya. *Annals of*  
735 *Glaciology*, **24**, 90–94.
- 736 Kapnick, S.B., Delworth, T.L., Ashfaq, M., Malyshev, S. & Milly, P.C.D., 2014. Snowfall  
737 less sensitive to warming in Karakoram than in Himalayas due to a unique seasonal  
738 cycle. *Nature Geoscience*, **7**, 834–840.
- 739 Kaser, G., Großhauser, M. & Marzeion, B. 2010. Contribution potential of glaciers to water  
740 availability in different climate regimes. *Proceedings of the National Academy of*  
741 *Sciences*, **107**, 20223–20227.
- 742 Kayastha, R.B. & Harrison, S.P. 2008. Changes of the equilibrium-line altitude since the  
743 Little Ice Age in the Nepalese Himalaya. *Annals of Glaciology*, **48**, 93–99.
- 744 Käab, A., Berthier, E., Nuth, C., Gardelle, J. & Arnaud, Y. 2012. Contrasting patterns of  
745 early twenty-first-century glacier mass change in the Himalayas. *Nature*, **488**, 495–  
746 498.
- 747 King, O., Quincey, D.J. Carrivick, J.L. & Rowan, A.V. 2017. Spatial variability in mass loss  
748 of glaciers in the Everest region, central Himalayas, between 2000 and 2015. *The*  
749 *Cryosphere*, **11**, 407–426.
- 750 Lutz, A.F., Immerzeel, W.W., Shrestha, A.B. & Bierkens, M.F.P. 2014. Consistent increase  
751 in High Asia's runoff due to increasing glacier melt and precipitation. *Nature Climate*  
752 *Change*, **4**, 587–592.

- 753 McCarthy, M., Pritchard, H., Willis, I. & King, E. 2017. Ground-penetrating radar  
754 measurements of debris thickness on Lirung Glacier, Nepal. *Journal of Glaciology*,  
755 **63**, 543–555.
- 756 Meehl, G., Stocker, T. & Collins, W.D. 2007. Global Climate Projections. In: Climate  
757 Change 2007: The Physical Science Basis. Contribution of Working Group I to the  
758 Fourth Assessment Report of the Intergovernmental Panel on Climate Change  
759 [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor  
760 and H.L. Miller (Eds.)]. Cambridge University Press, Cambridge, United Kingdom  
761 and New York, NY, USA.
- 762 Mihalcea, C., Mayer, C., Diolaiuti, G., D’Agata, C., Smiraglia, C., Lambrecht, A.,  
763 Vuillermoz, E. & Tartari, G., 2008. Spatial distribution of debris thickness and  
764 melting from remote-sensing and meteorological data, at debris-covered Baltoro  
765 glacier, Karakoram, *Annals of Glaciology*, **48**, 49-57.
- 766 Miles, E.S., Pellicciotti, F., Willis, I.C., Steiner, J.F., Buri, P. & Arnold, N.S. 2016. Refined  
767 energy-balance modelling of a supraglacial pond, Langtang Khola, Nepal. *Annals of*  
768 *Glaciology*, **57**, 29–40.
- 769 Miles, E. S., Steiner, J. F., & Brun, F. 2017. Highly variable aerodynamic roughness length (  
770  $z_0$ ) for a hummocky debris-covered glacier. *Journal of Geophysical Research:*  
771 *Atmospheres*, 1–58.
- 772 Minora, U., Senese, A., Bocchiola, D., Soncini, A., D’agata, C., Ambrosini, R., Mayer, C.,  
773 Lambrecht, A., Vuillermoz, E., Smiraglia, C. & Diolaiuti, G., 2015. A simple model  
774 to evaluate ice melt over the ablation area of glaciers in the Central Karakoram  
775 National Park, Pakistan. *Annals of Glaciology*, **48**, 49–57.
- 776 Mölg, T., Maussion, F., Yang, W. & Scherer, D. 2012. The footprint of Asian monsoon  
777 dynamics in the mass and energy balance of a Tibetan glacier. *The Cryosphere*, **6**,  
778 1445–1461.
- 779 Moiwo, J. P., Yang, Y., Tao, F., Lu, W., & Han, S. (2011). Water storage change in the  
780 Himalayas from the Gravity Recovery and Climate Experiment (GRACE) and an  
781 empirical climate model. *Water Resources Research*, **47**, W07521.
- 782 Muller, F. 1961. Khumbu moraine series, Nepal. *Radiocarbon*, **3**, 16.
- 783 Murari, M.K., Owen, L.A., Dortch, J.M., Caffee, M.W., Dietsch, C., Fuchs, M., Haneberg,  
784 W.C., Sharma, M.C. & Townsend-Small, A., 2014. Timing and climatic drivers for  
785 glaciation across monsoon-influenced regions of the Himalayan–Tibetan orogen.  
786 *Quaternary Science Reviews*, **88**, 159-182.
- 787 Nakawo, M., Ybuki, H. & Sakai, A. 1999. Characteristics of Khumbu Glaciers, Nepal  
788 Himalaya: recent change in the debris-covered area. *Annals of Glaciology*, **28**, 118–  
789 122.

- 790 Nicholson, L. & Benn, D. 2006. Calculating ice melt beneath a debris layer using  
791 meteorological data. *Journal of Glaciology*, **52**, 463–470.
- 792 Nicholson, L. & Benn, D.I. 2013. Properties of natural supraglacial debris in relation to  
793 modelling sub-debris ice ablation. *Earth Surface Processes and Landforms*, **38**, 490-  
794 501.
- 795 Østrem, G. 1959. Ice melting under a thin layer of moraine, and the existence of ice cores in  
796 moraine ridges. *Geografiska Annaler*, **41**, 228–230.
- 797 Owen, L.A. 2009. Latest Pleistocene and Holocene glacier fluctuations in the Himalaya and  
798 Tibet. *Quaternary Science Reviews*, **28**, 2150–2164.
- 799 Owen, L.A. & Benn, D.I. 2005. Equilibrium-line altitudes of the Last Glacial Maximum for  
800 the Himalaya and Tibet: an assessment and evaluation of results. *Quaternary*  
801 *International*, **138**, 55–78.
- 802 Owen, L., Caffee, M. & Finkel, R. 2008. Quaternary glaciation of the Himalayan–Tibetan  
803 orogen. *Journal of Quaternary Science*, **23**, 513–531.
- 804 Owen, L.A., Robinson, R., Benn, D.I., Finkel, R.C., Davis, N.K., Yi, C., Putkonen, J., Li, D.  
805 & Murray, A.S., 2009. Quaternary glaciation of Mount Everest. *Quaternary Science*  
806 *Reviews*, **28**, 1412–1433.
- 807 Owen, L.A. & Dortch, J.M. 2014. Quaternary Science Reviews. *Quaternary Science*  
808 *Reviews*, **88**, 14–54.
- 809 Pratap, B., Dobhal, DP, Mehta, M & Bhambri, R 2015. Influence of debris cover and altitude  
810 on glacier surface melting: a case study on Dokriani Glacier, central Himalaya, India.  
811 *Annals of Glaciology*, **56**, 9–16.
- 812 Pratap, B., Dobhal, D. P., Bhambri, R., Mehta, M., & Tewari, V. C. 2016. Four decades of  
813 glacier mass balance observations in the Indian Himalaya. *Regional Environmental*  
814 *Change*, **16**, 643–658.
- 815 Pritchard, H.D. 2017. Asia’s glaciers are a regionally important buffer against drought.  
816 *Nature*, **545**, 169–174.
- 817 Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J.  
818 & Glasser, N.F., 2007. Early recognition of glacial lake hazards in the Himalaya  
819 using remote sensing datasets. *Global and Planetary Change*, **56**, 137-152.
- 820 Quincey, D.J., Luckman, A. & Benn, D. 2009. Quantification of Everest region glacier  
821 velocities between 1992 and 2002, using satellite radar interferometry and feature  
822 tracking. *Journal of Glaciology*, **55**, 596–606.
- 823 Quincey, D.J., Braun, M., Glasser, N.F., Bishop, M.P., Hewitt, K. & Luckman, A. 2011.  
824 Karakoram glacier surge dynamics. *Geophysical Research Letters*, **38**.
- 825 Quincey, D.J., Glasser, N.F., Cook, S.J. & Luckman, A. 2015. Heterogeneity in Karakoram  
826 glacier surges. *Journal of Geophysical Research: Earth Surface*, **120**, 1288-1300.

- 827 Quincey, D., Smith, M., Rounce, D., Ross, A., King, O., & Watson, C.S. 2017. Evaluating  
828 morphological estimates of the aerodynamic roughness of debris covered glacier ice.  
829 *Earth Surface Processes and Landforms*, **49**, 547–44.
- 830 Ragettli, S., Immerzeel, W.W. & Pellicciotti, F. 2016. Contrasting climate change impact on  
831 river flows from high-altitude catchments in the Himalayan and Andes Mountains.  
832 *Proceedings of the National Academy of Sciences*, **113**, 9222–9227.
- 833 Reichert, B.K., Bengtsson, L. & Oerlemans, J. 2002. Recent glacier retreat exceeds internal  
834 variability. *Journal of Climate*, **15**, 3069–3081.
- 835 Reid, T.D. & Brock, B.W. 2014. Assessing ice-cliff backwasting and its contribution to total  
836 ablation of debris-covered Miage glacier, Mont Blanc massif, Italy. *Journal of  
837 Glaciology*, **60**, 3–13.
- 838 Reznichenko, N., Davies, T. & Shulmeister, J. 2010. Effects of debris on ice-surface melting  
839 rates: an experimental study. *Journal of Glaciology*, **56**, 384–394.
- 840 Robertson, C.M., Benn, D.I., Brook, M.S., Fuller, I.C. & Holt, K.A. 2012. Subaqueous  
841 calving margin morphology at Mueller, Hooker and Tasman glaciers in  
842 Aoraki/Mount Cook National Park, New Zealand. *Journal of Glaciology*, **58**, 1037–  
843 1046.
- 844 Rounce, D.R., Quincey, D.J. & McKinney, D.C. 2015. Debris-covered glacier energy  
845 balance model for Imja–Lhotse Shar Glacier in the Everest region of Nepal. *The  
846 Cryosphere*, **9**, 2295–2310.
- 847 Rounce, D.R., & McKinney, D.C. 2014. Debris thickness of glaciers in the Everest area  
848 (Nepal Himalaya) derived from satellite imagery using a nonlinear energy balance  
849 model. *The Cryosphere*, **8**, 1317–1329.
- 850 Rowan, A.V. 2017. The 'Little Ice Age' in the Himalaya: A review of glacier advance driven  
851 by Northern Hemisphere temperature change. *The Holocene*, **27**, 292–308.
- 852 Rowan, A.V., Egholm, D.L., Quincey, D.J. & Glasser, N.F. 2015. Modelling the feedbacks  
853 between mass balance, ice flow and debris transport to predict the response to climate  
854 change of debris-covered glaciers in the Himalaya. *Earth and Planetary Science  
855 Letters*, **430**, 427–438.
- 856 Röthlisberger, R, F. & Geyh, M. 1986. Glacier variations in Himalayas and Karakorum.  
857 *Journal of Glaciology and Geocryology*, **4**, 237–249.
- 858 Rupper, S., Schaefer, J.M., Burgener, L.K., Koenig, L.S., Tsering, K. & Cook, E.R. 2012.  
859 Sensitivity and response of Bhutanese glaciers to atmospheric warming. *Geophysical  
860 Research Letters*, **39**, L19503.
- 861 Sakai, A., Nishimura, K., Kadota, T. & Takeuchi, N. 2009. Onset of calving at supraglacial  
862 lakes on debris-covered glaciers of the Nepal Himalaya. *Journal of Glaciology*, **55**,  
863 909–917.

- 864 Sarikaya, M.A., Bishop, M.P., Shroder, J.F. & Olsenholler, J.A. 2012. Space-based  
865 observations of Eastern Hindu Kush glaciers between 1976 and 2007, Afghanistan  
866 and Pakistan. *Remote Sensing Letters*, **3**, 77–84.
- 867 Scherler, D., Bookhagen, B. & Strecker, M.R. 2011. Spatially variable response of  
868 Himalayan glaciers to climate change affected by debris cover. *Nature Geoscience*, **4**,  
869 156–159.
- 870 Schwanghart, W., Worni, R., Huggel, C., Stoffel, M. & Korup, O. 2016. Uncertainty in the  
871 Himalayan energy-water nexus: estimating regional exposure to glacial lake outburst  
872 floods. *Environmental Research Letters*, **11**, 074005.
- 873 Sharma, M.C. & Owen, L.A. 1996. Quaternary glacial history of NW Garhwal, central  
874 Himalayas. *Quaternary Science Reviews*, **15**, 335–365.
- 875 Solomina, O.N., Bradley, R.S., Jomelli, V., Geirsdottir, A., Kaufman, D.S., Koch, J.,  
876 McKay, N.P., Masiokas, M., Miller, G., Nesje, A. & Nicolussi, K., 2016. Glacier  
877 fluctuations during the past 2000 years. *Quaternary Science Reviews*, **149**, 61–90.
- 878 Soncini, A., Bocchiola, D., Confortola, G., Minora, U., Vuillermoz, E., Salerno, F., Viviano,  
879 G., Shrestha, D., Senese, A., Smiraglia, C. & Diolaiuti, G., 2016. Future hydrological  
880 regimes and glacier cover in the Everest region: The case study of the upper Dudh  
881 Koshi basin. *Science of The Total Environment*, **565**, 1084–1101.
- 882 Tapley, B.D., Bettadpur, S., Ries, J. C., Thompson, P. F., & Watkins, M. M. 2004. The  
883 gravity recovery and climate experiment: Mission overview and early results.  
884 *Geophysical Research Letters*, **31**, L09607.
- 885 Thakuri, S., Salerno, F., Smiraglia, C., Bolch, T., D'Agata, C., Viviano, G. & Tartari, G.  
886 2014. Tracing glacier changes since the 1960s on the south slope of Mt. Everest  
887 (central Southern Himalaya) using optical satellite imagery. *The Cryosphere*, **8**,  
888 1297–1315.
- 889 Thompson, L.G., Mosley-Thompson, E., Davis, M.E., Mashiotto, T.A., Henderson, K.A.,  
890 Lin, P.-N. & Tandong, Y. 2006. Ice core evidence for asynchronous glaciation on the  
891 Tibetan Plateau. *Quaternary International*, **154**, 3–10.
- 892 Thompson, S.S., Benn, D.I., Dennis, K. & Luckman, A. 2012. A rapidly growing moraine-  
893 dammed glacial lake on Ngozumpa Glacier, Nepal. *Geomorphology*, **145**, 1–11.
- 894 Thompson, S., Benn, D.I., Mertes, J. & Luckman, A. 2016. Stagnation and mass loss on a  
895 Himalayan debris-covered glacier: processes, patterns and rates. *Journal of*  
896 *Glaciology*, **62**, 467–485.
- 897 Tshering, P. & Fujita, K. 2016. First in situ record of decadal glacier mass balance (2003–  
898 2014) from the Bhutan Himalaya. *Annals of Glaciology*, **57**, 289–294.
- 899 National Research Council. 2012. Himalayan Glaciers: Climate Change, Water Resources,  
900 and Water Security. Washington, DC: *The National Academies Press*.

- 901 Vincent, C., Wagnon, P., Shea, J.M. & Immerzeel, W.W. 2016. Reduced melt on debris-  
902 covered glaciers: investigations from Changri Nup Glacier, Nepal. *The Cryosphere*,  
903 **10**, 1845–1858.
- 904 Watson, C.S., Quincey, D.J., Carrivick, J.L. & Smith, M.W. 2016. The dynamics of  
905 supraglacial water storage in the Everest region, central Himalaya. *Global and*  
906 *Planetary Change*, **142**, 14–27.
- 907 Xu, Y., Ramanathan, V. & Washington, W.M., 2016. Observed high-altitude warming and  
908 snow cover retreat over Tibet and the Himalayas enhanced by black carbon aerosols.  
909 *Atmospheric Chemistry and Physics*, **16**, 1303–1315.
- 910 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H.,  
911 Xu, B. & Pu, J., 2012. Different glacier status with atmospheric circulations in  
912 Tibetan Plateau and surroundings. *Nature Climate Change*, **2**, 663-667.
- 913











