

This is a repository copy of *Part by part:synthetic biology parts used in solventogenic Clostridia*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/125408/>

Version: Accepted Version

Article:

Gyulev, Ivan, Willson, Benjamin James, Hennessy, Rosanna Catherine et al. (3 more authors) (2018) Part by part:synthetic biology parts used in solventogenic Clostridia. ACS Synthetic Biology. pp. 311-327. ISSN 2161-5063

<https://doi.org/10.1021/acssynbio.7b00327>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Part by part: synthetic biology parts used in**
2 **solventogenic Clostridia**
3

4
5
6
7 **Authors:**

8 **Ivan S. Gyulev^{1*}, Benjamin J. Willson^{1*}, Rosanna C. Hennessy², Preben Krabben³,**
9 **Elizabeth R. Jenkinson⁴, Gavin H. Thomas^{1§}**

10
11
12
13 **Author affiliation:**

14 **1. Department of Biology, University of York, Wentworth Way, York, YO10 5DD, United Kingdom**

15 **2. Microbial Ecology and Biotechnology, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark**


16 **3. 9 Marsh Lane, Didcot, Oxfordshire, OX11 8FD, United Kingdom**

17 **4. Green Biologics Limited, Milton Park, Abingdon, Oxfordshire, OX14 4RU, United Kingdom**

18 *** These authors contributed equally to this work.**

19 **§ - Phone: 44-1904-328678. E-mail: gavin.thomas@york.ac.uk.**

20



21

22 **Graphical Table of Contents.**

23 **ABSTRACT:** The solventogenic *Clostridia* are of interest to the chemical industry
 24 because of their natural ability to produce chemicals such as butanol, acetone and
 25 ethanol from diverse feedstocks. Their use as whole cell factories presents multiple
 26 metabolic engineering targets that could lead to improved sustainability and
 27 profitability of *Clostridium* industrial processes. However, engineering efforts have been
 28 held back by the scarcity of genetic and synthetic biology tools. Over the last decade,
 29 genetic tools to enable transformation and chromosomal modifications have been
 30 developed, but the lack of a broad palette of synthetic biology parts remains one of the
 31 last obstacles to the rapid engineered improvement of these species for bioproduction.
 32 We have systematically reviewed existing parts that have been used in the modification
 33 of solventogenic *Clostridia*, revealing a narrow range of empirically chosen and non-
 34 engineered parts that are in current use. The analysis uncovers elements, such as
 35 promoters, transcriptional terminators and ribosome binding sites where increased
 36 fundamental knowledge is needed for their reliable use in different applications.
 37 Together, the review provides the most comprehensive list of parts used and also
 38 presents areas where an improved toolbox is needed for full exploitation of these
 39 industrially important bacteria.

40

41 **Keywords**

42

43 *Clostridium*, ABE fermentation, solventogenesis, transcription termination, reporters,
 44 promoters

45

46 **Abbreviations**

47 ABE: acetone-butanol-ethanol

48 ACE: allele-coupled exchange

49 aSD: anti Shine-Dalgarno sequence, found at the 3' end of 16S rRNA

50 CDS: coding sequence

51 FbFP: flavin-binding fluorescent protein

52 RBS: ribosome binding site

53 RNAP: RNA-polymerase

54 SD: Shine-Dalgarno sequence

55 UTR: untranslated region

56

57 **Acknowledgements:**

58 We acknowledge the BBSRC for support for Ivan Gyulev through an iCASE PhD
59 studentship with Green Biologics Ltd (BB/M016331/1), Benjamin Willson from an IB
60 Catalyst grant for Project DETOX (BB/N01040X/1) and Rosanna Hennessy for a BBSRC
61 IUK award (BB/L011522/1). We thank Stephen Thorpe for writing a Python script that
62 we used to compare the overlaps of the outputs of the terminator prediction algorithms.


63

64

65

66 One of the earliest industrial biotechnology processes used by man was the production
67 of acetone by bacterial fermentation. The bacterium used was from the genus
68 *Clostridium*, which are Gram-positive, spore forming, obligately anaerobic organisms¹. In
69 addition to acetone, these processes produced butanol and ethanol, leading to the
70 fermentation being given the name 'ABE' for 'acetone-butanol-ethanol'². ABE-producing
71 clostridia typically show a biphasic growth pattern, producing acetic and butyric acids
72 during the early stages of growth, and then undergoing a metabolic 'switch',
73 reassimilating the produced acids and producing solvents³. Driven by the demands of
74 industry, many different ABE-producing clostridial isolates were identified; those
75 strains that were used in industry are primarily represented by four species^{4,5}, namely
76 *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *Clostridium saccharobutylicum*, and
77 *Clostridium saccharoperbutylacetonicum*, although ABE fermentation has been observed
78 in other clostridia⁶⁻⁸. Furthermore, not all butanol-producing clostridia produce the
79 canonical trio of solvent products; for example, *Clostridium pasteurianum* produces
80 butanol and 1,3-propanediol (instead of acetone) from glycerol⁹. While the ABE process
81 had until recently fallen out of favour due to competition from the petrochemical
82 industry, the necessity of identifying alternative fuels has renewed interest in the
83 production of butanol as a potential biofuel candidate and sustainable commodity
84 chemical¹⁰.

85



86

87 **Figure 1.** Phylogenetic tree of 16S sequences from selected organisms from the genus *Clostridium*. The tree was built
 88 using Maximum Likelihood method based on the Tamura-Nei model¹¹. The bootstrap consensus tree inferred from 1000
 89 replicates¹² is taken to represent the evolutionary history of the taxa analyzed¹². Evolutionary analyses were conducted
 90 in MEGA5¹³. Green circles denote butanol-producing species used in industrial biotechnology (IB), red squares mark risk
 91 group 2 species¹⁴ (risk group 3 being highest risk).

92


93 A recent comparative genomics study of many industrial saccharolytic strains (those
94 preferring sugar as a carbon source) revealed that the known solventogens fall within
95 two sister clades: one exemplified by *C. acetobutylicum* and one by *C. beijerinckii*¹⁵
96 (Figure 1). Interestingly, another comparative genomics paper which included more
97 genus-wide species but fewer industrial solventogens supported the split, with the
98 genus' type species *Clostridium butyricum* being more closely related to *C. beijerinckii*
99 and the pathogen *Clostridium tetani* clustering closer to *C. acetobutylicum*¹⁶. Altogether,
100 these findings serve to re-iterate that complex traits within the Clostridia such as
101 pathogenicity are paraphyletic (also see Fig. 1). On the other hand, solventogenesis (of
102 ethanol, butanol or acetone) may be very widespread in the genus, but there has not yet
103 been a definitive comparative study reporting the extent of its conservation to our
104 knowledge, and species and strains certainly vary in their productivity¹⁷. Topics of
105 engineering interest have included improved characteristics such as solvent
106 production^{18,19}, sugar utilisation^{20,21}, growth on alternative feedstocks such as
107 lignocellulose²²⁻²⁵ and the production of alternative products²⁶⁻²⁸.

108

109 **A shortage of biological parts for the solventogenic Clostridia**

110 To manipulate the solventogenic Clostridia using rational engineering, the development
111 of reliable standard genetic parts for use in synthetic biology is essential. The assembly
112 of these parts into synthetic gene cassettes and larger gene clusters means that they
113 must work together in a consistent and predictable manner to become a useful tool for
114 strain engineering. The basic parts of synthetic biology are the minimal sequence
115 elements with biological function in gene expression (Figure 2), including promoters,
116 ribosome binding sites, transcriptional terminators and other factors in the mRNA that
117 affect stability. Also, synthetic biology tools rely on gene reporters to measure levels of
118 gene expression, and these elements are often combined into plasmids which need their
119 own origins of replication. Work in *Escherichia coli*, *Saccharomyces cerevisiae* and other
120 model organisms has been transformed in the past decade by the rapid development,
121 characterization and standardization of parts. However, work in these organisms
122 benefits from a legacy of biological knowledge that is not necessarily available when
123 working in non-model organisms, making the engineering of the latter even more

124 challenging.



125

126 **Figure 2.** Features of genetic constructs depicted using symbols from the SBOLv²⁹. Examples of promoters, ribosomal
127 binding sites, reporters, terminators and replicons are described in the main text.

128

129

130 It is also important to note that, of the existing body of molecular biology research in this
131 genus, the majority has been carried out in *C. acetobutylicum* ATCC 824. Work in this
132 species has benefited from a relatively early development of a reliable electroporation
133 protocol³⁰ and from the publication of the whole genome sequence in 2001, making it
134 the first published clostridial genome³¹. However, *C. acetobutylicum* is only one of many
135 industrially promising solventogenic clostridia. While the engineering of some of these
136 species has historically been hindered by a lack of transformation protocols, these are
137 now available for the transformation of the aforementioned five industrial species<sup>30,32-
138 35</sup>. In several cases, development of transformation protocols has required the
139 circumvention of restriction systems which degrade incorrectly methylated DNA.
140 Indeed, electroporation of *C. acetobutylicum* only became viable with the expression of
141 the *B. subtilis* phage ϕ 3T I methylase in the *E. coli* cloning host; this methylates the
142 sequence GCNGC, which would otherwise be cleaved by the *C. acetobutylicum* Cac824I
143 type II restriction enzyme³⁰. Likewise, transformation of *C. pasteurianum* ATCC 6013
144 (DSM 525) requires the methylation of CGCG sequences, which has been accomplished
145 by the use of the M.FnuDII³³ or M.BepI³⁶ methylases. Other organisms have more
146 complex restriction systems; in *C. saccharobutylicum* NCP 262, which has two type I
147 restriction systems, expression of the methylation and specificity domains of these

148 systems on a plasmid in *E. coli* was sufficient to allow transformation by conjugation³⁴.
149 Some developments have also been made in transforming non-type strains, which may
150 have restriction patterns which differ from those of the type strains. For example, *C.*
151 *pasteurianum* NRRL B-598 is part of the *C. beijerinckii* cluster¹⁵ but requires the use of a
152 *dam*⁻/*dcm*⁻ strain of *E. coli* for successful transformation, suggesting that the type IV
153 system of this strain is particularly important³⁷; conversely, *C. beijerinckii* NCIMB 8052
154 can be transformed with much greater efficiency even with DNA from a *dam*⁺/*dcm*⁺
155 host³⁸. While an analysis of such developments in the entire genus *Clostridium* would be
156 beyond the scope of this review, recent publications by Pyne et al.³⁹ and Minton et al.⁴⁰
157 provide a comprehensive review of the development of *Clostridium* strains for genetic
158 engineering. However, it is certain that the range of genetically tractable *Clostridium*
159 species and strains will expand with future research.

160

161 The resurgent interest in solventogenic *Clostridium* species suggests that synthetic
162 biology tools are needed and there are clear shortages of particular biological parts with
163 characterized activity available for the engineering of these organisms. We believe this
164 warrants a concerted effort to address the weaknesses in the toolbox. Furthermore, with
165 the establishment of transformation protocols throughout the genus *Clostridium*, it is
166 also important to consider the potential of adapting existing parts and tools for use in
167 other, less well-developed solventogens. Here, we assess the currently available parts
168 used in synthetic biology projects, starting first with reporters available to measure gene
169 expression, then the promoters, terminators working at the level of transcription and
170 then post-transcriptional factors including mRNA stability, ribosome binding sites,
171 riboswitches and codon usage. Finally we review the replicons currently being used to
172 assemble synthetic gene clusters and briefly provide an overview of methods to
173 introduce these genetic elements onto the chromosome, which has been reviewed
174 elsewhere⁴⁰. Overall, we aim to summarize and systematize the existing biological parts
175 used in the industrial solventogenic *Clostridium* species in order to highlight areas
176 where more research is required and accelerate progress in creating a full range of
177 synthetic biology tools for manipulating these industrially important organisms.

178

179 **Reporters of gene expression**

180

181 Genetically-encoded reporter systems are the major *in vivo* gene expression
182 measurement techniques available and are required to measure the activity of different
183 parts being tested. The oxygen sensitivity of obligate anaerobes such as the clostridia
184 limits the ease of use of many reporter systems, including some popular fluorescence-
185 based and enzymatically-based reporters, and has led to the development of some novel
186 reporters which are slowly being adopted.

187 *Fluorescent Reporter Proteins*

188
189 The use of fluorescent reporter proteins is now widespread in biology. Successful use
190 requires the correct folding and maturation of the fluorescent protein to enable
191 detection. The level of signal for the fluorescent protein must be sufficiently high to
192 enable accurate detection as there is no signal amplification as seen in enzymatic
193 reporters.

194 GFP-family fluorescent proteins

195
196 The highly engineered family of Green Fluorescent Proteins (GFPs) now includes
197 variants with improved brightness and photostability and with a range of different
198 colours (different excitation and emission maxima)⁴¹. The major limitation of the GFP-
199 like proteins for anaerobes is the requirement of molecular oxygen for chromophore
200 maturation and fluorescence⁴². This excludes the possibility of real-time gene expression
201 monitoring; however, *in vivo* observations are still possible in some oxygen-tolerant
202 clostridia such as *Clostridium perfringens*⁴³, where yellow fluorescent protein (YFP) was
203 synthesized and fluorescence was developed after exposure of live cells to atmospheric
204 oxygen⁴⁴.

205
206 Using a similar principle, it is possible to obtain a snapshot of the protein levels in whole
207 cells by exposing fixed anaerobically grown bacteria to atmospheric O₂ in a process
208 termed 'aerobic fluorescence recovery', enabled by the discovery that GFP-family
209 proteins can undergo maturation in fixed cells. Two studies have demonstrated the
210 technique in *C. difficile* using paraformaldehyde and glutaraldehyde to fix cells
211 expressing codon-optimized variants of CFP (cyan)⁴⁵ and mCherry (red)⁴⁶. This
212 approach works as the fixatives cross-link primary amines (especially lysine residues)
213 which are absent from the GFP-family chromophores. Fixed cells offer an advantage to

214 the imaging of dying live cells as they more accurately represent normal protein
215 localization (gene expression changes are also likely to occur in dying or metabolically
216 stressed cells)⁴⁶. So far, the technique has not been used to systematically quantify gene
217 expression but rather to label proteins and track their intracellular localization^{46,47}.

218 Flavin-binding Fluorescent proteins

219
220 The flavin-binding fluorescent proteins (FbFPs) are a class of alternative fluorescent
221 reporters capable of maturation in anaerobic conditions. The FbFPs are small proteins
222 (~11-15kDa) that have oxygen-independent fluorescent properties, using a flavin-
223 mononucleotide (FMN) cofactor as the chromophore.

224 A pioneering study in 2007 reported the development of three anaerobic fluorescent
225 reporters derived from the LOV (Light-, Oxygen- or Voltage-sensing) domains of
226 bacterial proteins: BsFbFP derived from YtvA of *Bacillus subtilis*, PpFbFP from SB2 of
227 *Pseudomonas putida* plus an *E. coli* codon optimized variant of BsFbFP called EcFbFP⁴².
228 This domain was engineered to emit fluorescence by mutagenesis of a reactive cysteine
229 in the FMN-binding pocket⁴².

230 These proteins are now commercially available from Evocatal GmbH under the
231 trademark name 'evoglow'. Currently listed variants marketed for use in *Clostridium*
232 species include Bs1 (monomeric BsFbFP), Bs2 (dimeric BsFbFP) and Pp1 (PpFbFP).

233 The reporters' functionality in *C. acetobutylicum* was demonstrated by Schulz in 2013;
234 the highest fluorescence levels were observed for a *C. acetobutylicum* codon-optimized
235 Pp1 (referred to in text as "Pp2"), followed by codon-optimized Bs2 (referred to as
236 "Bs3"). Interestingly the same constructs gave inverse results in *E. coli*⁴⁸. While Evocatal
237 GmbH offers C-Pp1 and C-Bs2 *Clostridium* codon-optimized reporters with publicly
238 available nucleotide sequences, it is not clear whether the nucleotide sequences
239 (reflective of the codon optimization approach) are the same as Pp2 and Bs3. A study in
240 2014 reported the placement of the evoglow Pp1 reporter downstream of the *cipP*
241 promoter to monitor growth of *Clostridium cellulolyticum* on cellulose⁴⁹. The
242 functionality of the evoglow Bs2 variant has also been demonstrated in *C.*
243 *pasteurianum*⁵⁰ and in *Clostridium ljungdahlii*⁵¹.

244 Another FbFP example, developed from the *Arabidopsis thaliana* LOV2 domain of the
245 blue-light receptor protein Phot2, is the improved LOV (iLOV) FbFP⁵². The iLOV FbFP
246 has been further modified for enhanced properties, generating variants such as

247 photostable iLOV 2.1 (phiLOV 2.1) and others^{53,54}. While phiLOV 2.1 FbFP has improved
248 photostability, it is still substantially lower than that of GFP. A recent study
249 demonstrated the utility of a *C. difficile* codon optimized phiLOV 2.1 in three *Clostridium*
250 species (expression was driven from an engineered strong *xyl* promoter)⁵⁵. Under the
251 test conditions, the three species exhibited varying levels of fold-increase of
252 fluorescence over their background autofluorescence level: 3.2-fold increase in *C.*
253 *difficile* R20291 (not a *Clostridium sensu stricto* species), 5.6-fold in the pathogenic toxin-
254 producer *Clostridium sordellii* ATCC 9714 and 4.5-fold in the solventogen *C.*
255 *acetobutylicum* ATCC 824⁵⁵. While the FbFP reporters will be detectable when
256 expression is driven from very strong promoters, further improvements in brightness
257 and photostability will make them more generally useful for multiple applications.

258

259 One of the drawbacks of the FbFPs is that all of the available variants are of the same
260 colour; additionally, unfolding caused by translocation has been reported to result in
261 loss of fluorescence due to the loss of the bound flavin cofactor⁵⁴. These limitations
262 prevent the use of FbFPs as the sole reporters for multi-output circuits or secretion.
263 Engineering of FbFPs for different emission spectra has proven to be somewhat
264 complicated; while a simulation study of a theoretical iLOV mutant predicted a red shift
265 increase⁵⁶, experimental results showed that the mutation resulted in an overall
266 brightness decrease and a blue shift of emission⁵⁷. Research is underway to develop
267 more fluorescent derivatives of the LOV domains and to explore the potential for
268 diversifying their colour spectrum^{56,57}. The other limitation of FbFPs is their relative
269 dimness, which can decrease the sensitivity of a reporter system; depending on the
270 measurement instrument used, a promoter's expression level might be below the
271 detection limit. Photostability improvements (as in phiLOV) have resulted in a dimmer
272 mutant⁵³; for applications where a longer exposure time is crucial, the dimmer, stable
273 protein is preferable, but for an automated single measurement application such as flow
274 cytometry, the brighter variant may be more suitable. Overall, all fluorescent protein
275 reporters provide the ability to quantify cell-to-cell heterogeneity of gene expression
276 (when used in flow cytometry or microscopy), which can be very useful, and while there
277 are still some limitations with FbFPs, they have the potential to be a route to reliable *in*
278 *vivo* real-time expression monitoring in *Clostridium*.

279

280 *Enzymatic Reporter Proteins*

281 Enzymatic reporters catalyse a (preferably) unique reaction either *in vivo* or *in vitro* and
282 the specific activity is calculated to estimate protein levels and thus gene expression.
283 Enzymatic reporters require the addition of substrates and cofactors to the assay
284 reaction mixture as well as the production of cell lysate. This often means that enzymatic
285 assays involve more preparation steps than fluorescent proteins but can have adjustable
286 sensitivity by varying substrate levels. Also, *in vitro* lysate assays are, by definition, bulk
287 population measurements.

288 *Chloramphenicol acetyltransferase*

289
290 The first reporter which has been extensively used in the *Clostridium* genus, both *in vivo*
291 and, more quantitatively, *in vitro*, is the chloramphenicol acetyltransferase (CAT)
292 reporter (encoded by *catP*). The system was first developed for use in *C. perfringens* (a
293 medically relevant non-solventogenic bacterium)⁵⁸ and has since been used in *C.*
294 *acetobutylicum*⁵⁹. Chloramphenicol acetyltransferase is an enzyme that catalyzes the
295 covalent attachment of an acetyl group from acetyl-CoA to chloramphenicol⁶⁰, and is the
296 basis of the chloramphenicol resistance marker found in many bacterial vectors.
297 Transfer of the acetyl group exposes the thiol group of CoA, allowing the progress of the
298 reaction to be observed by addition of 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB, also
299 known as Ellman's reagent); this compound reacts with the free thiol, releasing 5-thio-2-
300 nitrobenzoate, which can be detected by measurement of absorbance at 412 nm^{60, 61}.
301 The assay relies on a continuous spectrophotometric rate determination to calculate
302 specific activity. Potential drawbacks include endogenous activity in chloramphenicol
303 resistant strains (which may be resolved by disruption of the resistance gene) as well as
304 high levels of endogenous non-specific coenzyme A transferases⁶² (knockouts of which
305 would be more laborious and would likely have growth and phenotypic effects) and the
306 cost of the substrate acetyl-CoA.

307 *Carbohydrate hydrolases: β -glucuronidase, β -galactosidase, amylase, endoglucanase*

308
309 The CAT assay's drawbacks led to the adaptation of a classical *E. coli* reporter in
310 *Clostridium*: the β -galactosidase enzyme, encoded by *lacZ* (the gene was derived from
311 *Thermoanaerobacter thermosulfurigenes*)⁶². To quantify enzyme activity
312 spectrophotometrically, ortho-nitrophenyl- β -galactoside (ONPG) is used in an *in vitro*
313 assay and an increase in absorbance at 420 nm due to the release of ortho-nitrophenol is

314 measured. Similarly, the GUS reporter system, which utilizes β -glucuronidase (*E. coli*
315 *gusA*), has been used in *C. acetobutylicum* in a fluorimetric assay with a cell lysate⁶³. The
316 fluorimetric assay measures the release of 4-methylumbelliferone (4-MU) after cleavage
317 of 4-MU- β -galactoside or 4-MU- β -glucuronide (by a β -galactosidase or β -glucuronidase,
318 respectively), 4-MU emits light at 460 nm when excited by 365 nm light⁶⁴. Both the β -
319 galactosidase and β -glucuronidase reporter systems benefit from the commercially
320 available range of fluorometric, spectrophotometric and histochemical substrates and
321 kits, making them an improvement over the CAT assay, although they are still not
322 inexpensive.

323 The endogenous *amyP* gene (encoding an amylase expressed during solventogenesis)
324 has been used in *C. acetobutylicum*⁶⁵ as a reporter to study the phenomenon of strain
325 degeneration (loss of solventogenesis) which is often caused by loss of the pSOL1
326 megaplasmid on which *amyP* is located. A codon-optimized amylase (*AmyE^{opt}*) has been
327 used successfully as a secreted reporter in *C. difficile* by addition of a zinc
328 metalloprotease PPEP-1 signal sequence⁶⁶. It is noteworthy that the strain used in the
329 above study was not capable of degrading starch under laboratory conditions; use of
330 amylase as a reporter in amylolytic strains (such as *C. acetobutylicum* ATCC 824) may
331 require knockout of endogenous amylases to increase signal to background ratio.

332 In *C. beijerinckii*, a secreted endoglucanase (*eglA*) was cloned from *C. saccharobutylicum*
333 NCP 262 and used as a reporter for gene expression⁶⁷; assays used were agar plate and
334 cell lysate assays which measured substrate (carboxymethylcellulose, CMC) clearance
335 on plates and product (*p*-nitrophenol from cleavage of *p*-nitrophenyl cellobioside)
336 accumulation in lysate, respectively.

337 *Lipase and Alkaline phosphatase*

338 The lipase encoded by *tliA*, from *Pseudomonas fluorescens* SIK W1, has also been used as
339 a reporter in *C. beijerinckii* NCIMB 8052⁶⁸. Lipases are enzymes which hydrolyse the
340 ester bonds found in long-chain acylglycerols, releasing fatty acids⁶⁹. Activity can thus be
341 assessed by an enzymatic assay measuring the cleavage of *p*-nitrophenyl-esters⁶⁹ such
342 as *p*-nitrophenyl decanoate⁶⁸. However, this reporter was only useable in *C. beijerinckii*
343 as this species has a very low endogenous lipase activity. As such, it may not be suitable
344 in other species, such as *C. acetobutylicum*, which has been observed to show inducible
345 lipase activity⁷⁰.

346 A colorimetric alkaline phosphatase assay was developed and used in *C. difficile* in 2015
347 based on the *phoZ* gene product from *Enterococcus faecalis*⁷¹. Activity of this gene can be
348 determined by a colorimetric assay with *p*-nitrophenyl phosphate as the substrate. Use
349 in solventogenic clostridia could be limited by native phosphatase activity. In order to
350 examine the suitability of *C. difficile* as a host for this reporter, BLAST analysis was used
351 to screen for *phoZ* homologues, and activity towards 5-bromo-4-chloro-3-indolyl
352 phosphate was tested⁷¹. While the four main industrial strains do not have a *phoZ*
353 homologue, homologues can be found in other species such as *C. pasteurianum*
354 (CLPA_RS02340, with 29% identity to *phoZ*).

355 *Luciferase*

356

357 The luciferase (*lucB*) reporter was also used in *C. acetobutylicum* successfully, allowing
358 luminescence detection⁷². However, it also requires oxygen, ATP and luciferin. Notably,
359 the cells used in the luciferase assay were neither lysed nor fixed, but were live cells that
360 were exposed to atmospheric conditions, washed, and kept on ice. This treatment could
361 conceivably introduce changes in gene expression levels prior to measurement. This
362 assay has the lowest background signal level but the requirement for live cell exposure
363 to oxygen may introduce variability. A codon-optimized luciferase (*sLuc*^{opt}) was also
364 successfully secreted in *C. difficile* using the aforementioned zinc metalloprotease signal
365 peptide⁶⁶.

366

367 While there are many reporter choices available to clostridial researchers, we would
368 argue that the multiplicity of reporters used has not helped ease the comparison of data
369 obtained by different laboratories. Altogether, a single reporter has not been established
370 as a community standard; given the drawbacks of each particular system, it is difficult to
371 identify one standout reporter, although our hope would be that improved fluorescence
372 reporters with increased brightness and photostability would be the most useful and
373 enable single cell studies in live cells. Currently we would advocate a choice of more
374 than one reporter, enzymatic and fluorescent, and their systematic and comparative use
375 within the genus. Such an approach could provide insight into the sources of gene
376 expression variability between strains and provide a starting point for future
377 engineering. To achieve this seamlessly, it is advisable to explore translational fusions
378 between reporters; examples from *E. coli* include a FRET pair YFP-FbFP fusion⁷³ and

379 Gemini (lacZ α -GFP) fusion⁷⁴. If successfully applied to the *Clostridium* species, such bi-
380 functional reporters have the potential to become a single standard reporter.
381 Additionally, α -peptide complementation strategies using the *E. coli lacZ* gene in *B.*
382 *subtilis*⁷⁵ and *Mycobacterium smegmatis*⁷⁶ have been reported but a similar approach in
383 Clostridia has not been attempted. Interestingly, the *T. thermosulfurigenes*-derived LacZ
384 protein appears to lack the α -complementation region⁷⁷ and is a dimer⁷⁸ (rather than a
385 tetramer like *E. coli LacZ*).

386 **Transcription**

387 Transcription is the first stage of gene expression and the main stage for regulation of
388 gene expression. Most existing biological parts in *Clostridium* fall within the
389 transcription category and are mostly promoters (summarized in Table 1). In contrast,
390 the termination of transcription has been studied relatively little and we present some
391 analysis on the potential to study and improve the parts available for reliable
392 termination.

393 **Transcription initiation**

394 Promoters are DNA elements that are capable of driving transcription by RNA
395 polymerase (RNAP) of downstream regions. Core promoter architecture in bacteria
396 includes the -10 region or TATAAT box (Pribnow sequence), the -35 region and a spacer
397 (with an optimal length of 17bp in *E. coli*) between the two conserved regions⁷⁹. This
398 motif is recognized by the housekeeping sigma factor that provides sequence specificity
399 to RNAP (called RpoD in *E. coli* and SigA in *B. subtilis*, *C. acetobutylicum* and other
400 solventogens)^{80,81}. Both the consensus sequence and sigma factors bear very significant
401 similarities between Firmicutes and *Escherichia*.

402 In addition, some *E. coli* promoters contain AT-rich UP elements (upstream of -35
403 region) that are responsible for recognition by the carboxy-terminal domain of the
404 RNAP α -subunit⁸², an additional *E. coli* promoter feature is the 'extended -10' region⁸³
405 (upstream of the -10 region and within the spacer). Interestingly, near-consensus
406 promoters (including ones with UP elements) have been found to be significantly more
407 common in Firmicutes (including Clostridia) than in other bacteria⁸⁴, a feature that was
408 not explicable through higher AT-content alone. In Clostridia additional sequences with
409 resemblances to UP elements (termed phased A-tracts) have been described⁸⁵ and a
410 conserved extended -10 region that differs from its *E. coli analogue* has also been
411 proposed⁸⁶. The majority of promoters used in the genetic engineering of clostridia have

412 been identified from the transcriptional units of important metabolic genes and most
413 have been shown to be able to drive strong constitutive expression of a gene of
414 interest⁸⁷. The most commonly used constitutive promoter in *C. acetobutylicum* (Table
415 1) is that of the native thiolase gene (*thlA*)^{88,89}, which has been used for the expression of
416 genes involved in production of solvents such as butanol⁹⁰ and isopropanol²⁶, the
417 transcriptional regulator gene *tetR*⁹¹ (used a minimal promoter variant- *miniPthl*), and,
418 in a modified form, for the expression of cellulosomal scaffoldins⁹² and glycoside
419 hydrolases^{25,93}. The *C. acetobutylicum* thiolase enzyme catalyzes the condensation of
420 acetoacetyl-CoA from acetyl-CoA which are central metabolic intermediates⁸⁸. The *C.*
421 *acetobutylicum thlA* promoter (hereinafter referred to as *thlA*) is a sigA-dependent
422 promoter (as evidenced by its near-consensus -35 and -10 regions⁸⁹); however, in its
423 normally chromosomal context it is also the subject of regulation by the redox-sensing
424 transcriptional repressor Rex⁹⁴. Rex inactivation was found to increase native *thlA*
425 activity in *C. acetobutylicum* about 12-fold⁹⁵. The Rex-binding site has been omitted from
426 the core promoter in commonly used synthetic constructs⁹⁶; yet this binding site is
427 conserved in the promoters of orthologous genes from other solventogens (according to
428 the RegPrecise database⁹⁷ and our promoter region alignments), suggesting that there
429 are additional levels of control for this promoter that could be easily added by addition
430 of the Rex sites if desirable. Indeed, some studies using similar promoters such as *thlA*
431 and *ptb* have reported conflicting strength/activity findings, which may well be due to
432 the cloning of regions of varying length (that likely include regulator binding sites such
433 as Rex) and choosing to include the native RBSs or not (such as the ones reported
434 between promoters in the Schulz 2013⁴⁸ and Girbal 2003⁶³ studies).

435

436 Thiolase promoters have also been used for expression in other clostridia such as *C.*
437 *beijerinckii*⁹⁸ and *C. pasteurianum*⁹⁹. However, gene expression data (RNAseq from *C.*
438 *beijerinckii*) indicates that there are other genes that have higher expression levels than
439 the ones currently used as promoter sources; many of those genes encoded are
440 hypothetical proteins¹⁰⁰. Recently, Yang *et al.*, 2017 constructed a sequence logo of 18
441 promoters previously identified in *C. acetobutylicum* which revealed a strongly
442 conserved -10 region. Using degenerate oligos to mutagenize the core *thlA* promoter
443 elements, the authors were able to generate a promoter library of variable strengths,

444 including mutations that increased the strength of the promoter, suggesting that *thlA*
445 can be improved further using synthetic biology approaches.

446

447 The use of a constitutive promoter may not always be desirable; it may be preferable to
448 use an inducible promoter, allowing controlled expression of a gene of interest. So far,
449 the only naturally inducible promoters exemplified in a solventogenic clostridium are
450 the *Staphylococcus xylosus xylA* promoter, which is repressed by XylR in the absence of
451 D-xylose⁶³, and the *C. perfringens bgaL* promoter, which is repressed by BglR in the
452 absence of lactose¹⁰¹. However, constitutive promoters can be made inducible by
453 addition of operator sites (for transcriptional repressors) or by the addition of binding
454 sites for activators, and several such promoters have been developed for use in
455 solventogenic clostridia. LacI-repressible versions of thiolase and ferredoxin promoters
456 have allowed the construction of clostridial expression constructs for genes where
457 expression in *E. coli* would be toxic^{24,25,93}. In *C. acetobutylicum*, the *fac* promoter
458 (*Clostridium pasteurianum* ferredoxin promoter with *lac* operator) is able to function as
459 an IPTG-inducible system in the presence of LacI, allowing approximately 10-fold
460 induction⁵⁹. A TetR-repressed, anhydrotetracycline-inducible promoter has also been
461 used in *C. acetobutylicum*, generated by the fusion of the chloramphenicol
462 acetyltransferase promoter *pcm* with the tetracycline operator *tetO*⁹¹. This promoter
463 could achieve up to 313-fold induction, although high levels of anhydrotetracycline were
464 inhibitory to growth. However, operators are known to influence the basal activity of the
465 promoter¹⁰². In fact, a recent study in *E. coli* found the core RpoD promoter too sensitive
466 to sequence context and operator insertions to be a suitable target for forward
467 engineering efforts and turned to extracellular sigma factors (ECFs) with T7 RNAP
468 whose promoter core sequences they found to be more insensitive to operator
469 addition¹⁰³. An alternative strategy - CRISPR-mediated repression of transcription - has
470 been demonstrated in several solventogenic species, namely *C. acetobutylicum*¹⁰⁴, *C.*
471 *beijerinckii*⁹⁸ and *C. pasteurianum*⁵⁰,

472

473 Orthogonal expression systems, i.e. promoters that require other non-native elements
474 for activity (commonly an alternative sigma factor or phage polymerase), first found
475 widespread use in the cloning of genes with toxic products. This approach allows the
476 total repression of genes until they are introduced into the organism of interest. A

477 commonly used example in *E. coli* is the T7 promoter, which requires the phage T7
478 polymerase for activity¹⁰⁵. An example from Clostridia is expression driven from the *C.*
479 *difficile tcdB* promoter that is dependent on the native sigma factor TcdR. The *tcdB*
480 promoter is highly active in *C. acetobutylicum*, but only when *tcdR* has been integrated
481 into the genome⁵⁹. This enabled the high-level expression of a *mariner* transposon in *C.*
482 *acetobutylicum* without negative effects on the *E. coli* cloning host. There is a strong case
483 for designing these orthogonal-type systems as simply using a clostridial
484 housekeeping promoter (such as the *thlA* promoter) with a clostridial ribosome binding
485 site would not provide orthogonality in commonly used cloning hosts such as *E. coli* or *B.*
486 *subtilis* due to the degree of conservation between the transcription and translation
487 initiation systems amongst these organisms; on the contrary, as a rule strong clostridial
488 promoters and RBSs (see translation initiation section) often retain their strength in *E.*
489 *coli* whereas the opposite is observed more rarely, this is likely due to the on average
490 higher similarity to the Bacteria-wide consensus translation and transcription initiation
491 signals in *Clostridium*^{86,106}. Interestingly, AT-rich DNA can be toxic to *E. coli* (clostridial
492 genomes are very AT-rich) due to transcriptional activity from spurious intragenic
493 promoters and RNAP titration¹⁰⁷. This is exacerbated by the fact that the *E. coli* extended
494 -10 region, which is sufficient alone for transcription initiation in *E. coli*, could also be
495 present by chance in *Clostridium*-derived sequences, increasing the likelihood of
496 spurious and unpredictable transcription during cloning in *E. coli*.

Table 1. Promoters used in engineering of *C. acetobutylicum*

Promoter	Comments
Constitutive	
<i>thlA</i> (<i>C. acetobutylicum</i>)	Widely used for constitutive gene expression in <i>C. acetobutylicum</i> for example expression on <i>C. beijerinckii ald</i> , <i>C. ljungdahlii bdh</i> ⁹⁰ , <i>C. acetobutylicum adc</i> , <i>ctfA</i> , <i>ctfB</i> , ²⁶ & <i>tetR</i> (<i>miniPthl</i> variant) ⁹¹ . Activity analyzed using Gusa ⁶³ and several FbFP ⁴⁸ reporters
<i>thl</i> (<i>C. perfringens</i>)	Used for expression of <i>srtA</i> genes from <i>C. acetobutylicum</i> , <i>L. monocytogenes</i> , and <i>B. cereus</i> ²⁵ .
<i>fdx</i> (<i>C. sporogenes</i>)	Activity analyzed using CatP reporter ⁵⁹ . Used in the Clostron system for expression of the Ll.LtrB intron ¹⁰⁸ . Used for expression of <i>spoA</i> integrated into the chromosome for complementation of a <i>spoA</i> mutant ¹⁰⁹ .
Controlled	
<i>thl</i> (<i>thl</i> with double <i>lac</i> operator)	<i>lac</i> -repressed version of <i>thlA</i> . Used to express a miniscaffoldin <i>cipc1</i> ⁹² & weakened version for expression of mannanase <i>man5K</i> ⁹³ .
<i>thlOid</i> (<i>thl</i> with single <i>lac</i> operator)	A <i>lac</i> -repressible version, used to drive chromosomal expression of the <i>C. cellulolyticum</i> glycoside hydrolase <i>xyn10A</i> ²⁵ .
<i>fac</i> (single <i>lac</i> operator, derived from <i>C. pasteurianum</i>)	Activity analyzed using CatP reporter ^{59,110} and Pp2 FbFP ⁴⁸ . Formerly used in the Clostron system for expression of the Ll.LtrB intron ¹¹⁰ . Used for expression of <i>codA</i> in the <i>C. acetobutylicum</i> knockout vector pMTL-SC7515 ¹⁰⁹ .
<i>fdxOid</i> (<i>fdx</i> with single <i>lac</i> operator)	Used for chromosomal expression of <i>C. cellulolyticum</i> glycoside hydrolase <i>cel19G</i> ²⁵ . Has RBS from <i>C. acetobutylicum thlA</i> promoter.
<i>facOid</i> (<i>fac</i> with single <i>lac</i> operator)	Used for chromosomal expression of <i>C. cellulolyticum</i> glycoside hydrolase <i>cel48F</i> ²⁵ . Has RBS from <i>C. acetobutylicum thlA</i> promoter.
<i>tcdB</i> (<i>C. difficile</i>)	Developed for an orthogonal expression system, requires the exogenous sigma factor TcdR for function ⁵⁹ .
<i>xylA</i> (<i>S. xylosus</i>)	Repressed by XylR and inducible by D-xylose; assessed with β -glucuronidase reporter giving 17-fold induction with D-xylose ⁶³ .
<i>pcm</i> (<i>tetO1</i> -containing variants)	Repressed by TetR; inducible in the presence of anhydrotetracycline ⁹¹ .
<i>bgaL</i> (<i>C. perfringens</i>)	Repressed by BgaR and inducible by lactose ¹⁰¹ .
<i>xyl/tetO</i> (<i>B. subtilis</i>)	Repressed by TetR ¹¹¹ ; activity in <i>C. acetobutylicum</i> shown by phiLOV 2.1 Opt FbFP expression ⁵⁵
Fermentation phase-specific	
<i>adc</i> (<i>C. acetobutylicum</i>)	Activity assessed using β -glucuronidase ⁶³ , β -galactosidase ⁷² and Pp2 FbFP ⁴⁸ . Primarily active after onset of solventogenesis.
<i>ptb</i> (<i>C. acetobutylicum</i>)	Activity assessed using β -glucuronidase ⁶³ , luciferase ⁷² and Pp2 FbFP ⁴⁸ . Active during acidogenesis (not solventogenesis)
<i>ptb</i> (<i>C. beijerinckii</i>)	Activity assessed through expression of <i>lacI</i> ⁵⁹ .
<i>sol</i> (<i>C. acetobutylicum</i>)	Activity assessed using β -galactosidase & luciferase reporters ⁷² . Weak expression, primarily active during late exponential phase.
<i>bdhA</i> (<i>C. acetobutylicum</i>)	Analyzed using β -galactosidase reporter ⁷² . Comparatively weak expression, primarily active in early exponential phase.
<i>bdhB</i> (<i>C. acetobutylicum</i>)	Analyzed using β -galactosidase reporter ⁷² . Primarily active until onset of solventogenesis.
<i>hydA</i> (<i>C. acetobutylicum</i>)	Activity analyzed using β -glucuronidase ⁶³ and Pp2 FbFP ⁴⁸ . High activity during acidogenesis, decreases to low after phase shift
<i>adhe2</i> (<i>C. acetobutylicum</i>)	Activity analyzed using Pp2 FbFP ⁴⁸ . Strong expression during solventogenesis ^{48,112}

498 **Transcription termination**

499 Bacteria have two distinct mechanisms that function in transcription termination. Both
500 types of terminators are usually located in the 3' end of transcriptional units. Rho-
501 dependent terminators rely on the Rho protein that recognizes a target sequence and
502 causes RNA polymerase to fall off of the template DNA¹¹³. The specific DNA recognition
503 sites, *rut* sites, have been used before in genetic circuits¹¹⁴, but not extensively, perhaps
504 due to the relatively poor mechanistic understanding of the process¹¹⁵. The second
505 mechanism, which also has been reported to be more widespread¹¹⁶, is referred to as
506 Rho-independent or intrinsic termination and relies on the folding of a short GC rich
507 hairpin followed by a poly-U transcribed sequence. The hairpin folding and subsequent
508 transcription of the U-tract causes the polymerase to pause and leads to transcript and
509 RNAP dissociation¹¹⁷. This is also facilitated by the fact that ribo-uracil-deoxyribo-
510 adenine is the weakest nucleotide base pair¹¹⁸. Intrinsic terminators are often found
511 downstream of operons; however, they are also involved in transcription attenuation
512 when present within coding regions or downstream of promoters¹¹⁹.

513

514 While a number of terminators have been used in the construction of clostridial
515 expression constructs, often derived from clostridial genes such as *adc*^{63,120}, *fdx*¹⁰⁹,
516 CD0164¹⁰⁹ and *thlA*^{90,121}, there have been few published analyses of terminator
517 strengths in clostridia. The *C. pasteurianum* *fdx* terminator has been shown to be highly
518 effective at preventing read-through inhibition of the replicon from the *fac* promoter in a
519 clostridial vector⁵⁹, and screening of a selection of terminators in *C. acetobutylicum*²⁵
520 showed that the *E. coli* *rrnB* terminator T1 loop was able to function as an efficient
521 terminator, reducing expression of a downstream gene. However, these analyses only
522 took into consideration the effect of the terminator on a downstream target under the
523 influence of a single promoter. As previously mentioned, the introduction of a
524 terminator may have an effect on expression of an upstream gene by influencing mRNA
525 stability, which is not a desirable feature of a standard part¹²². However, this effect can
526 be quantified in an appropriately designed assay¹²³. Furthermore, the efficiency of
527 termination may increase or decrease depending on promoter activity; this has been
528 recently exploited for the development of a genetic band-pass filter in *E. coli*¹²⁴.
529 Terminator strength has also been shown to be influenced by the hairpin's proximity to
530 a stop codon or when present within a coding region¹²⁵. A recent study demonstrated

531 the gradual increase in termination efficiency as distance between the stop codon and
532 hairpin increased¹²⁶. The ribosome was found to repress transcriptional termination
533 when the stop codon and hairpin were in close proximity – termination repression was
534 relieved when insulator sequences of approximately 30 bp were used¹²⁶. Expanding the
535 range of characterized clostridial terminators is clearly necessary for the development
536 of a clostridial synthetic biology toolbox.

537

538 One approach that can be undertaken is the use of algorithms to extract putative
539 terminator sequences from genomes, also known as ‘part mining’. Several such
540 bioinformatics tools exist; they rely on seed sequences, secondary RNA structure
541 features or both^{116,127-129}. These bioinformatics tools were developed to aid genome
542 annotations by identifying the 3’ ends of operons and may not then detect all the
543 features required by a functional terminator. In order to evaluate the relative difference
544 in sequences retrieved by different programs, we compared the output of three searches
545 performed on the *C. acetobutylicum* ATCC 824 genome (Figure 3A). The three search
546 algorithms that were chosen are as follows: TransTermHP (TTHP)¹²⁸, RNIE¹¹⁶ and
547 WebGeSTer (WG)¹³⁰. TTHP is a widely used tool, while RNIE’s authors state that their
548 tool eliminates false positives in comparison to TTHP¹¹⁶. Finally, WebGeSTer was
549 selected because the authors classify the results into different types of intrinsic
550 terminators based on overall secondary structure. They discovered that canonical U-
551 tract containing intrinsic terminators (termed ‘L-shaped’) form the majority of
552 structures found within Firmicutes, whereas they are a minority in the *E. coli* model
553 where they were first identified¹²⁹.




Figure 3. Application of terminator prediction software to *C. acetobutylicum*. **A.** Outputs from predicted terminator sets derived from the *C. acetobutylicum* genome by application of TransTermHP (TTHP), RNIE and WedGeSTer were compared. **B.** An alignment of predictions of a putative bi-directional terminator (marked by T sign) by all three algorithms (note all three produce separate predictions for plus (+) and minus (-)). **C.** RNA structures predicted by RNAfold^{215,216} for the TransTermHP sequence, drawn with VARNA²¹⁷; it is worth noting that the base-pairing between the A-tract and U-tract may not form until after termination²¹⁸. Terminator lengths: WG: maximum length (max)-48bp, minimum (min) - 28bp, average (avg) - 35.8bp; RNIE: max-45bp, min- 34bp, avg-41.7bp, TTHP: max-89bp, min-41bp, avg-57.2bp.

555 In our analysis, each algorithm was used to produce sets of putative terminator
556 predictions, which were examined to identify overlapping predictions and are presented
557 in Fig. 3A. Using the default settings of the different algorithms, the total number of
558 predictions per program varied considerably, with TTHP and WG each predicting over
559 1000 terminators while RNIE predicted fewer than 300 (Fig. 3A). Despite this much
560 smaller set, only about half of the predictions were shared with either TTHP and WG
561 sets or both, producing a small set of 104 terminators shared by all three methods, an
562 example of which is illustrated in Fig. 3B, an intergenic region separating two
563 converging coding sequences, CA_C0323 (a sensory transduction histidine kinase) and
564 CA_C0324 (a tetratricopeptide-repeats-containing protein). This also illustrates a class
565 of terminators with particular utility in synthetic biology, namely bi-directional
566 terminators, which are capable of terminating transcription from both the plus and
567 minus strands. Our analysis finds that 37 bi-directional terminators are supported by all
568 three algorithms, supporting the concepts that there are plentiful targets in clostridial
569 genomes for 'part mining' and future experimental characterisation of these likely
570 strong terminators. An interesting feature of the bi-directional terminators is their
571 hypersymmetry¹³¹ (can be seen in Fig. 3B). This leads to the possibility to form an
572 extended hairpin between the so-called A-tract (can act as U-tract in the reverse
573 transcriptional orientation) and the U-tract (as shown in Fig. 3C). However, as pausing
574 and termination occur at the U-tract¹¹⁷, the formation of an extended hairpin may not be
575 relevant and does not appear to contribute to strength in all terminators¹³². Also, note
576 that in this example the terminator on the minus strand (also referred to as Crick
577 strand) is predicted to have a more stable hairpin (rG and rU form a basepair while rA
578 and rC do not), whereas terminators from both strands are capable of forming an
579 extended hairpin if fully transcribed as a result of base pairing between the A- and U-
580 tracts. Combined with having a longer U-tract (Fig. 3C), the Crick strand terminator is
581 possibly the stronger one, interestingly WebGeSTer predicts a smaller terminator
582 downstream of the one depicted in Fig. 3A (not shown). A previous analysis of the *C.*
583 *acetobutylicum* genome for terminators using TransTerm¹²⁷, a predecessor of TTHP,
584 revealed the presence of 66 putative bi-directional terminators¹³³, supporting the
585 rationale of combining outputs from multiple algorithms to create a small part list for
586 experimental characterisation.

587

588 Translation

589 Translation initiation

590 Bacterial ribosome-binding sites (RBS) are short sequences located in the 5'
591 untranslated region of messenger RNA (mRNA) transcripts, consisting of a Shine-
592 Dalgarno sequence (SD), polynucleotide spacer, and a translation initiation codon¹³⁴.
593 Commonly, native promoter-RBS combinations have been used (such as in the
594 pMTL80000 vectors). An alternative strategy is to use a native RBS (such as that of the *C.*
595 *acetobutylicum thlA* gene) fused to a new promoter, as in a study from 2016²⁵ that
596 generated several new hybrid promoters. Others have experimented with the length of
597 the spacer^{135,136}. Yet, in contrast to model organisms (*E. coli* and even *B. subtilis*), there
598 are few published comparisons of modified RBSs for use in solventogenic clostridia.
599 Hence the generation and screening of synthetic RBS libraries could be a promising
600 route for optimising the expression of synthetic gene constructs in clostridia.

601 SD sequences provide sequence complementarity for the 3' terminus of the 16S rRNA
602 (known as the anti-Shine-Dalgarno or aSD¹³⁷) which acts as a guide for the ribosome
603 complex enabling mRNA recognition by the translation machinery and translation
604 initiation. While most of the knowledge on bacterial translation initiation comes from *E.*
605 *coli* work, early studies indicated that the Firmicute *B. subtilis* requires a longer
606 complementary region between the SD and the 16S rRNA to achieve comparable
607 expression levels^{138,139}. An early study on translation initiation revealed that Firmicutes
608 have, on average, a higher complementarity of the predicted SD region to the 16S rRNA
609 3' terminus than *E. coli* does¹⁰⁶. Recently, a systematic analysis of SD-aSD pairings in *B.*
610 *subtilis* and in *E. coli* confirmed this trend¹⁴⁰. Replicating these studies in solventogens
611 would provide useful information.

612

613 The spacer is the mRNA region between the SD and the start codon. The aligned spacer
614 (which is the distance between the start codon and the 5' end of the aSD, base-paired to
615 a SD^{137,141,142} as revealed by sequence alignment) is of particular importance for
616 translation initiation¹³⁷. Defining the length of the aligned spacer precisely is difficult, as
617 interpretations of the extent of the aSD region (starting from the 3' 16S rRNA terminus)
618 vary. The 3' end of the 16S rRNA gene (aSDs) of *B. subtilis* is identical to that of *C.*
619 *perfringens*¹⁴³ (as well as identical to those of solventogenic clostridia, based on our
620 sequence analysis of published genomes) but to our knowledge there is no reported

621 experimental validation of the clostridial mature 16S rRNA 3' ends *in vivo*. Spacers in
622 different species may have different optimal lengths; for example, spacers of *Pyrococcus*
623 *abyssi* are, on average, roughly 3 nucleotides longer than those of *E. coli*¹⁴⁴, whereas the
624 spacers in *Bifidobacterium longum*¹⁴⁵ would be considered shorter. According to our
625 definition of the putative clostridial aSD (5' GAUCACCUCCUUUCU 3'), in *C.*
626 *acetobutylicum*, the native RBS of the *thlA* promoter has an 'aligned spacer' of 4 bases.
627 The effect of altering the length of the *thlA* spacer was recently investigated in two
628 studies conducted in *C. acetobutylicum*^{135,136}. Interestingly Yang *et al.*, 2016 showed that
629 a lengthened *thlA* RBS spacer with an extra 6 nucleotides (to a total of 14 bases- aligned
630 spacer of 10bp), encoding a *Sall* restriction site, did not significantly alter reporter
631 expression in comparison to the WT *thlA* RBS. Shortening the spacer below the WT
632 length resulted in a decrease in expression, while further increases over an aligned
633 spacer length of 10 bases (by the addition of a *XbaI* site) removed almost all of the
634 expression¹³⁵.

635 Yang *et al.* (2016) demonstrated the potential benefits of utilising modified spacers in *C.*
636 *acetobutylicum*¹³⁵. Overexpression of the biotin synthesis genes *bioY*, *bioD*, *bioA*, and
637 *bioB* was observed to provide an improvement in growth phenotype and solvent
638 production. The *thlA* promoter was used to drive expression of *bioY*, *bioD* and *bioA*;
639 replacement of the *thlA* RBS spacer with a shortened, less 'effective' variant resulted in a
640 further improvement in growth characteristics. These results demonstrate not only the
641 ability of modified RBS to optimize expression of synthetic pathways, but also the
642 necessity of considering the effects of any alterations to the spacer, e.g. the introduction
643 of restriction sites, when generating synthetic constructs. In 2017, Yang *et al.* generated
644 a library of spacers (a *BamHI* site preceded the start codon in all cases) by starting with
645 an aligned spacer of 2 bases and progressively increasing its length by 2 bases. After
646 testing 11 variants the authors found that an aligned spacer of 4 bases gave the
647 strongest levels of expression followed by 8, 6 and 10 bases in that order. It is
648 conceivable that not only the length but also the sequence of the spacer might influence
649 translation initiation, for example the introduction of a secondary SD within the primary
650 SD's spacer region.

651 Computational tools to design RBSs exist, such as the RBS calculator, which use
652 biophysical models of RBS recognition and translation initiation, including RNA base-
653 pairing between the aSD and SD, spacer length and messenger secondary

654 structure^{146,147}. There are reports that *de novo* design produces more accurate results
655 than translation initiation rate prediction of natural sequences for Gram positives¹⁴⁸.

656

657 **Codon usage during translation**

658 While codon usage is not formally a 'biological part', it is an important feature of coding
659 sequences, the differential frequency of synonymous codons amongst genomes, referred
660 to as codon usage bias^{149,150}, has been shown to strongly influence heterologous protein
661 expression levels^{151,152}.

662 High genomic AT-content is characteristic of the Firmicutes and is reflected in the
663 nucleotide composition of coding sequences. The *Clostridium* species' codon usage
664 differs from that of other Firmicutes as well as the Proteobacterium *E. coli*¹⁵³ and there
665 are also bioinformatically observable variations within the *Clostridium* genus itself¹⁵⁴
666 but the significance of the latter in influencing gene expression has not been
667 experimentally verified to our knowledge. Genetically encoded reporters have been used
668 heterologously within the Firmicutes phylum without codon-optimization, for example
669 *Staphylococci* have been sources of reporters and antibiotic resistance genes for
670 Clostridia. On several occasions researchers have successfully used native reporter
671 genes from *E. coli* (*gusA*)⁶³, *T. thermosulfurigenes* (*lacZ*)⁷² and the firefly *Photinus pyralis*
672 (*lucB*)⁷² (we describe these reporters in more detail in the Enzymatic Reporter Protein
673 section). Codon-optimized genes for *C. difficile* have been used in *C. acetobutylicum*⁵⁵, as
674 well as bespoke *C. acetobutylicum* codon-optimized ones^{24,48}. Researchers have also
675 codon-optimized several GFP-like proteins for use in Firmicutes¹⁵⁵ (with *B. subtilis* in
676 mind) but these have not been used in solventogens to our knowledge.

677 Codon optimization is not a trivial problem and codon optimization strategies vary
678 considerably. The codon-adaptation index (CAI)¹⁵⁰ has been the historical measure of
679 codon usage bias in an organism while there are others such as the codon bias index and
680 the effective number of codons¹⁵⁶. Interestingly, simply improving a heterologous gene's
681 CAI (making it more like a native gene or a highly expressed native gene) has not been
682 found to correlate with expression levels^{157,158}. Efforts have been made to improve the
683 indices describing codon usage bias and translational efficiency (which codon usage bias
684 is thought to reflect) by studying endogenous gene expression^{159,160}. In addition,
685 condition-specific usage tables have also been reported¹⁶¹. In an alternative strategy,
686 Welch and colleagues developed genetic algorithms to select partial least squares

687 regression models which revealed that codons predominantly read by tRNAs that are
688 most highly charged during amino acid starvation were good predictors of expression
689 levels. Based on these results the researchers developed proprietary codon optimization
690 algorithms to maximize protein expression^{158,162} which allowed them to predict
691 expression levels in *E. coli*.

692 While it is clear that there is room for improvement in the heterologous protein
693 expression strategies used in *Clostridium* solventogens, codon optimization strategies
694 themselves are still being developed and the underlying principles are not yet fully
695 understood; progress and existing approaches in the field have been reviewed
696 elsewhere¹⁶³. A good starting point is for researchers to report the details of the codon
697 optimization strategy undertaken when publishing work containing codon-optimized
698 genes. This way data from heterologous protein expression in solventogens can be
699 compared more reliably.

700 **Post-transcriptional control of gene expression**

701

702 Tuning gene expression levels in *Clostridium* species has been achieved using control at
703 the RNA level – by either influencing translation or changing RNA degradation rates.

704 **Controlling mRNA stability**

705 Another potential avenue for optimisation of expression levels is the adjustment of
706 mRNA stability. Altering the stability of an mRNA transcript influences the number of
707 transcripts in the cell, thereby affecting the overall rate of translation. In bacteria, a
708 number of factors are associated with mRNA stability, such as secondary structures,
709 RNase recognition sites and polyadenylation, amongst others¹²². The presence of
710 secondary structures at the 5' end of the mRNA has been observed to provide an
711 increase in mRNA stability¹²². In *E. coli*, these structures prevent the binding of RNase E,
712 an endonuclease which binds single-stranded RNA at the 5' end and then scans for
713 cleavage sites. This property has been exploited in *E. coli* by the generation of libraries of
714 synthetic hairpins for introduction into the 5' untranslated region (UTR)¹⁶⁴. While *C.*
715 *acetobutylicum* has an RNase E homolog, RNase E/G, it is not certain if this behaves in
716 the same way as *E. coli* RNase E, due to having a different domain organisation¹⁶⁵.
717 Nevertheless, *C. acetobutylicum* also has a homolog of RNase Y¹⁶⁶, which fulfils the role of
718 RNase E in *B. subtilis*¹⁶⁷. Correspondingly, the utility of 5' hairpins for protection of
719 mRNA has been demonstrated in clostridia; the introduction of 5' stem-loop sequences
720 was confirmed to increase mRNA stability, reporter expression, and expression of the

721 genes *adhE1* and *adhE2* in both *C. acetobutylicum* and *C. beijerinckii*¹⁶⁸. This effect was
722 much more pronounced during solventogenesis than during acidogenesis. Similarly, the
723 introduction of a terminator hairpin in the 3' UTR can result in improved mRNA stability
724 via inhibition of nuclease activity¹²². Although not fully explored, this principle has been
725 demonstrated in solventogenic clostridia; the expression of a *cat* reporter gene in *C.*
726 *acetobutylicum* was observed to increase by approximately 36% when the downstream
727 *adc* terminator was replaced by a synthetic terminator, BBa_B1010, from the iGEM
728 registry¹⁶⁸. Additionally, a terminator with activity in the reverse orientation prevents
729 the formation of antisense transcripts which are known to reduce protein expression
730 levels¹⁶⁹.

731

732 A completely opposite approach is to reduce mRNA stability by introducing RNase sites
733 into the 3' UTR. This may be useful in the case of proteins which form inclusion bodies
734 when overexpressed. In *E. coli*, sequences derived from the *cat* gene, which contains 28
735 RNase E sites, have been shown to reduce mRNA stability¹⁷⁰. When combined with two
736 poorly soluble heterologous enzymes, the *cat*-derived 3' UTRs were shown to result in
737 an increase in soluble protein, with concomitant increases in enzyme activity. The
738 authors noted that this improvement could not be observed simply by using a weaker
739 promoter; it was proposed that by limiting the stability of the mRNA, the number of
740 proteins that could be produced from a single transcript was decreased, thus limiting
741 the local concentrations of protein during translation.

742

743 **Antisense RNAs and Riboswitches**

744

745 Bacterial antisense RNAs (asRNAs) are transcripts with complementarity to another
746 RNA. Several studies have used asRNAs were used to reduce native gene expression in *C.*
747 *acetobutylicum*^{169,171} and in *C. pasteurianum*¹⁷², demonstrating that multiple routes to
748 regulate gene expression are available in the solventogenic clostridia.

749 Riboregulators are another class of naturally-occurring and generally trans-activating
750 asRNA elements that respond to a signal nucleic acid by Watson-Crick base pairing¹⁷³.
751 They have defined sensor and effector domains and have been rationally designed to
752 repress¹⁷⁴ and activate gene expression¹⁷⁵ in *E. coli*. Toehold switches (a synthetic cis-
753 regulatory subgroup of riboregulators) that activate gene expression in the presence of

754 cognate RNAs rely on sequestering the RBS and start codon¹⁷⁶. Like riboswitches,
755 riboregulators are known to be present in *Clostridium* genomes.

756 Riboswitches are RNA sequences that are able to bind to a soluble ligand, influencing the
757 properties of the RNA. In nature, riboswitches typically contain a binding domain, or
758 'aptamer', and an 'expression platform' which mediates the effect¹⁷⁷. Binding leads to a
759 change in conformation of the RNA, leading to formation of a secondary structure which
760 can act as a terminator or an anti-terminator. While riboswitches are typically found in
761 the 5' UTRs of mRNAs, some have been recently determined to control the expression of
762 antisense RNAs or protein-sequestering small RNAs, while yet others have been shown
763 to control access to recognition sequences such as RNase sites¹⁷⁸. The range of
764 applications has been further developed by the creation of synthetic riboswitches. In
765 bacteria, riboswitches have been developed that can influence translation initiation by
766 inhibiting access to the RBS; ligand binding leads to a conformational change or even to
767 self-cleavage, revealing the RBS and allowing translation¹⁷⁹. Riboswitches are found in
768 all taxa and a number have been characterized in the solventogenic clostridia¹⁸⁰⁻¹⁸².
769 However, riboswitches have not yet been used in the engineering of these organisms. As
770 discussed earlier, only a limited range of inducible systems is available for the
771 engineering of clostridia; the use of synthetic riboswitches could be a promising
772 alternative for the creation of controlled promoters.

773 Replication

774

775 Plasmid origins of replication

776 Four replicons are in routine usage in solventogens: pCB102 (from *C. butyricum*), pBP1
777 (from *C. botulinum*), pCD6 (from *C. difficile*) and pIM13 (from *B. subtilis*)⁹⁶. The pIM13
778 replication origin is thought to replicate via rolling-circle replication¹⁸³ while there is
779 evidence that pCD6 replicates in similar fashion to pIP404^{184,185} (which is either theta or
780 strand-displacement). The replication mechanisms of pCB102 and pBP1 are unknown.
781 In *C. acetobutylicum*, two other replicons that have been used are the pAM β 1 replicon
782 and the pUB110 replicon¹⁸⁶; the pUB110 replicon was found to be somewhat more
783 stable than pIM13, whereas the pAM β 1 replicon was highly unstable. The *C. beijerinckii*
784 filamentous phage CAK1's origin of replication has been used in *C. beijerinckii* strains¹⁸⁷.
785 Additionally, the development of a replicon specific for *C. saccharoperbutylacetonicum*
786 N1-4 was reported in 2007¹⁸⁸, this replicon is identical to the origin of the endogenous
787 plasmid from *C. saccharoperbutylacetonicum* N1-504¹⁵. A thermosensitive origin
788 pWV01ts derived from *Lactococcus lactis cremoris*¹⁸⁹ has been shown to work in both *C.*
789 *ljungdahlii* and *C. acetobutylicum*⁵¹. Segregation and transformation frequencies are
790 available; however, more work is needed to determine copy number and compatibility
791 groups.

792

793 Chromosomal integration

794 The integration of DNA into the genome, while not a 'part' in itself, is an important
795 consideration for synthetic biology projects. Genomic integration has several advantages
796 over plasmid-based expression strategies, including increased stability, removal of the
797 requirement for antibiotic selection, and standardisation of copy number¹⁹⁰⁻¹⁹².
798 However, there are other factors that must be considered when using chromosomal
799 integration. One implication of the integration position is the copy number effect – genes
800 closer to the origin have a higher copy number than ones near the terminus in
801 exponentially dividing cells due to the mechanism of DNA replication. There is
802 contradictory evidence as to the effects of chromosome location and levels of
803 expression^{193,194,195}. The cause of some of the apparent transcriptional insulation of
804 chromosomal genes (and thus promoter independence) was found to be due to the gene
805 silencing activity of the HU-protein in *E. coli*¹⁹⁴, a protein that is also present in

806 Firmicutes¹⁹⁶. Genomic engineering in *E. coli* has been carried out extensively; lambda-
807 Red recombineering¹⁹⁷ is well-established, and the utility of the newly developed
808 CRISPR technique has been demonstrated in this species^{198,199}. However, in the
809 solventogenic clostridia, until recently published examples of genomic integration had
810 been still somewhat limited. An early enabling technology was ClosTron which adapted
811 the L1.LtrB intron for use in *Clostridium*¹¹⁰. Another method for genomic integration is
812 Allele-Coupled Exchange (ACE), as demonstrated in *C. acetobutylicum*^{24,25,190}. This is a
813 homologous recombination-based method, where homology arms with different lengths
814 are used to control the sequence of recombination events, and the second recombination
815 leads to the generation of a selectable phenotype; currently, this involves either the
816 truncation or repair of the *pyrE* gene or the activation of a promoterless antibiotic
817 resistance gene by integration downstream of a strong chromosomal promoter such as
818 *thlA*¹⁹⁰. Thus, one potential drawback to this method is that it only allows integration
819 into a limited selection of loci. This drawback can be mitigated by the ability to carry out
820 multiple rounds of iterative ACE, thereby making further genomic integrations into the
821 same locus.

822

823 Nevertheless, many new developments have been made regarding the genetic
824 manipulation of solventogenic clostridia. A variety of different allelic exchange-based
825 strategies have been exemplified in *C. acetobutylicum*^{101,109,200,201} and *C. beijerinckii*²⁰²;
826 while most of these studies have focused on the generation of in-frame deletions and
827 subsequent complementation, Al-Hinai et al.¹⁰¹ demonstrate the integration of a
828 heterologous gene through gene replacement. Furthermore, the generation of point
829 mutations through recombineering has been demonstrated in *C. acetobutylicum*²⁰³,
830 suggesting that the integration of DNA through this method may be feasible. Finally,
831 mutant selection via CRISPR has been established in almost all of the main solventogenic
832 species, with published examples in *C. beijerinckii*^{121,204}, *C. acetobutylicum*¹⁰⁴, *C.*
833 *pasteurianum*²⁰⁵, and *C. saccharoperbutylacetonicum*²⁰⁶. Future developments are certain
834 to expand the range of genomic modifications that can be made in these organisms.

835

836 **Insulators**

837 A biological part's adjacent sequences can have a profound effect on its behaviour
838 compared to the sequence context in which it was characterized²⁰⁷. This poses a

839 fundamental challenge to the synthetic biology principle of part creation and
840 characterization²⁰⁸. To counteract this issue in reproducibility genetic engineers have
841 started utilizing a new class of parts called insulators²⁰⁹. Several strategies can be
842 undertaken to insulate a part from its genetic context, and these can be split into two
843 main categories: DNA-level insulators (such as simply using flanking buffer zones of
844 sequence without a biological function or secondary structure) and RNA-level
845 insulators. The latter includes post-transcriptional modification of RNA as well as the
846 commonly used flanking double terminators¹²³ that prevent read-through transcription
847 into synthetic gene constructs. Post-transcriptional insulators consist of inclusion of
848 ribozyme-based insulators or using CRISPR-RNA-processing to decouple the 5' UTR
849 from the coding sequence (CDS)^{208,210}.

850 Different parts require different types of insulation in order to achieve maximum
851 reproducibility without compromising features such as strength.

852 Promoters used in synthetic biology projects are often minimal (-35 to -10 region) and
853 without characterized transcriptional start sites; the inclusion of important functional
854 elements such as the UP element has been previously recommended.²¹¹ Accordingly, the
855 addition of an upstream and downstream insulating sequence has increased
856 reproducibility in different genetic contexts²¹². It is worth noting that the strong
857 *Clostridium* promoters that are in widespread use in the *Clostridium* community, *fdx* and
858 *thlA*, are 200 and 150bp respectively. Even though this recombinant *thlA* is longer than a
859 usual minimal promoter, a long 5'UTR contributes to this part's length (recombinant
860 *thlA* is 59bp from its 5' end to the transcriptional start site), while the recombinant 5'
861 end is slightly truncated to exclude a Rex NADH-dependent regulator binding site (as
862 mentioned before). A study that reported the use of a minimal *thlA* promoter in *C.*
863 *acetobutylicum*, *miniPthl* which has a truncated 5' UTR, did not test activity variation in
864 different genetic contexts⁹¹.

865 Work in *E. coli* has demonstrated the utility of 'bicistronic RBS', where a leader peptide
866 is translationally coupled with the CDS of interest, in improving reliability and context
867 independence (downstream gene sequence) when a particular 5' UTR is combined with
868 a new coding sequence²¹². An upstream RBS and start codon initiate translation of the
869 leader peptide, the stop codon of which overlaps with the start of the downstream CDS,
870 while the latter's RBS is positioned with the leader peptide. The ribosomes translating
871 the leader peptide unfold the 5' UTR (of the downstream CDS), preventing it from

872 forming secondary structures with the mRNA of the downstream CDS and thus
873 influencing translation. The same study employed standard transcriptional start site
874 (“+1 promoter”): 5’ UTR junctions to minimize (or insulate against) unforeseen effects of
875 combining promoters with new 5’ UTRs.

876 As mentioned above, the observation of ribosomal repression of transcription
877 termination also necessitates the more widespread use of ‘distance’ insulators of a
878 sequence without emergent function and secondary structure to separate the stop
879 codon and the stem-loop hairpin¹²⁶. Such strategies are yet to be implemented in
880 *Clostridium* engineering projects.

881

882 Discussion

883 The development of biorenewables as an alternative to petroleum-derived commodity
884 chemicals and fuels has resulted in the emergence of new markets²¹³. Increasing
885 productivity, broadening the range of feedstocks, improving tolerance to solvents and
886 by-products are all existing challenges to achieving higher sustainability and ensuring
887 the economic viability of *Clostridium*-derived biorenewables²¹⁴. *Clostridium* species
888 remain important hosts for the biological production of solvents and their further
889 development relies on the adaptation of novel methodologies such as synthetic biology
890 and metabolic engineering. Much progress has been made in the latter with several
891 projects improving industrially relevant strains; however, the availability of biological
892 parts with known behaviour is one of the limiting factors for the rate and scale of
893 research. The reasons for our anticipation are twofold: first, the physical availability of
894 biological parts streamlines assembly of genetic pathways, and second, knowledge about
895 parts’ behaviour is crucial in predicting and analyzing the behaviour of pathways and
896 genetic circuits. We have shown in this review that there are multiple areas where
897 improved knowledge of parts would provide a better toolkit for synthetic biologists
898 using these organisms and consequently improve the rate at which industrial
899 biotechnology and bioenergy process development can improve.

900 References

901

- 902 (1) Breed, R. S., Murray, E. G. D., and Smith, N. R. (1957) Bergey's manual of
903 determinative bacteriology 7th ed. Williams & Wilkins Co., Baltimore.
- 904 (2) Jones, D. T., and Woods, D. R. (1986) Acetone-butanol fermentation revisited.
905 *Microbiol. Rev.* 50, 484–524.
- 906 (3) Gheshlaghi, R., Scharer, J. M., Moo-Young, M., and Chou, C. P. (2009) Metabolic
907 pathways of clostridia for producing butanol. *Biotechnol. Adv.* 27, 764–781.
- 908 (4) Keis, S., Bennett, C. F., Ward, V. K., and Jones, D. T. (1995) Taxonomy and phylogeny
909 of industrial solvent-producing clostridia. *Int. J. Syst. Bacteriol.* 45, 693–705.
- 910 (5) Keis, S., Shaheen, R., and Jones, D. T. (2001) Emended descriptions of *Clostridium*
911 *acetobutylicum* and *Clostridium beijerinckii*, and descriptions of *Clostridium*
912 *saccharoperbutylacetonicum* sp. nov. and *Clostridium saccharobutylicum* sp. nov. *Int. J.*
913 *Syst. Evol. Microbiol.* 51, 2095–2103.
- 914 (6) Hellinger, E. (1947) *Clostridium aurantibutyricum* (n.sp.): A Pink Butyric Acid
915 *Clostridium*. *J. Gen. Microbiol.* 1, 203–210.
- 916 (7) Holt, R. A., Cairns, A. J., and Morris, J. G. (1988) Production of butanol by *Clostridium*
917 *puniceum* in batch and continuous culture. *Appl. Microbiol. Biotechnol.* 27, 319–324.
- 918 (8) Abd-Alla, M. H., Zohri, A. N. A., El-Enany, A. W. E., and Ali, S. M. (2015) Acetone-
919 butanol-ethanol production from substandard and surplus dates by Egyptian native
920 *Clostridium* strains. *Anaerobe* 32, 77–86.
- 921 (9) Sarchami, T., Munch, G., Johnson, E., Kießlich, S., and Rehmann, L. (2016) A Review of
922 Process-Design Challenges for Industrial Fermentation of Butanol from Crude Glycerol
923 by Non-Biphasic *Clostridium pasteurianum*. *Fermentation* 2, 13.
- 924 (10) Tashiro, Y., Yoshida, T., Noguchi, T., and Sonomoto, K. (2013) Recent advances and
925 future prospects for increased butanol production by acetone-butanol-ethanol
926 fermentation. *Eng. Life Sci.* 13, 432–445.
- 927 (11) Tamura, K., and Nei, M. (1993) Estimation of the number of nucleotide substitutions
928 in the control region of mitochondrial DNA in humans and chimpanzees. *Mol. Biol. Evol.*
929 10, 512–26.
- 930 (12) Felsenstein, J. (1985) Confidence Limits on Phylogenies: An Approach Using the
931 Bootstrap. *Evolution (N. Y.)*. 39, 783–791.
- 932 (13) Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011)
933 MEGA5: molecular evolutionary genetics analysis using maximum likelihood,
934 evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–9.

935 (14) ABAS. (2015) Classification of prokaryotes (bacteria and archaea) into risk groups
936 (TRBA 466). *Bundesarbeitsblatt*. <http://www.baua.de> 19.

937 (15) Poehlein, A., Solano, J. D. M., Flitsch, S. K., Krabben, P., Winzer, K., Reid, S. J., Jones, D.
938 T., Green, E., Minton, N. P., Daniel, R., and Dürre, P. (2017) Microbial solvent formation
939 revisited by comparative genome analysis. *Biotechnol. Biofuels* 10, 58.

940 (16) Udaondo, Z., Duque, E., and Ramos, J. L. (2017) The pangenome of the genus
941 *Clostridium*. *Environ. Microbiol.* 19, 2588–2603.

942 (17) Shaheen, R., Shirley, M., and Jones, D. T. (2000) Comparative fermentation studies of
943 industrial strains belonging to four species of solvent-producing clostridia. *J. Mol.*
944 *Microbiol. Biotechnol.* 2, 115–124.

945 (18) Jang, Y.-S., Lee, J. Y., Lee, J., Park, J. H., Im, J. A., Eom, M.-H., Lee, J., Lee, S.-H., Song, H.,
946 Cho, J.-H., Seung, D. Y., and Lee, S. Y. (2012) Enhanced butanol production obtained by
947 reinforcing the direct butanol-forming route in *Clostridium acetobutylicum*. *MBio* 3,
948 e00314-12.

949 (19) Pyne, M. E., Sokolenko, S., Liu, X., Srirangan, K., Bruder, M. R., Aucoin, M. G., Moo-
950 Young, M., Chung, D. A., and Chou, C. P. (2016) Disruption of the reductive 1,3-
951 propanediol pathway triggers production of 1,2-propanediol for sustained glycerol
952 fermentation by *Clostridium pasteurianum*. *Appl. Environ. Microbiol.* 82, 5375–5388.

953 (20) Xiao, H., Li, Z., Jiang, Y., Yang, Y., Jiang, W., Gu, Y., and Yang, S. (2012) Metabolic
954 engineering of D-xylose pathway in *Clostridium beijerinckii* to optimize solvent
955 production from xylose mother liquid. *Metab. Eng.* 14, 569–578.

956 (21) Xiao, H., Gu, Y., Ning, Y., Yang, Y., Mitchell, W. J., Jiang, W., and Yang, S. (2011)
957 Confirmation and elimination of xylose metabolism bottlenecks in glucose
958 phosphoenolpyruvate-dependent phosphotransferase system-deficient *Clostridium*
959 *acetobutylicum* for simultaneous utilization of glucose, xylose, and arabinose. *Appl.*
960 *Environ. Microbiol.* 77, 7886–7895.

961 (22) Mingardon, F., Chanal, A., Tardif, C., and Fierobe, H. P. (2011) The issue of secretion
962 in heterologous expression of *Clostridium cellulolyticum* cellulase-encoding genes in
963 *Clostridium acetobutylicum* ATCC 824. *Appl. Environ. Microbiol.* 77, 2831–2838.

964 (23) Chanal, A., Mingardon, F., Bauzan, M., Tardif, C., and Fierobe, H. P. (2011) Scaffoldin
965 modules serving as “cargo” domains to promote the secretion of heterologous
966 cellulosomal cellulases by *Clostridium acetobutylicum*. *Appl. Environ. Microbiol.* 77, 6277–
967 6280.

968 (24) Kovács, K., Willson, B. J., Schwarz, K., Heap, J. T., Jackson, A., Bolam, D. N. N., Winzer,
969 K., and Minton, N. P. (2013) Secretion and assembly of functional mini-cellulosomes
970 from synthetic chromosomal operons in *Clostridium acetobutylicum* ATCC 824.
971 *Biotechnol. Biofuels* 6, 117.

972 (25) Willson, B. J., Kovács, K., Wilding-Steele, T., Markus, R., Winzer, K., and Minton, N. P.
973 (2016) Production of a functional cell wall-anchored minicellulosome by recombinant
974 *Clostridium acetobutylicum* ATCC 824. *Biotechnol. Biofuels* 9, 109.

975 (26) Collas, F., Kuit, W., Clément, B., Marchal, R., López-Contreras, A. M., and Monot, F.
976 (2012) Simultaneous production of isopropanol, butanol, ethanol and 2,3-butanediol by
977 *Clostridium acetobutylicum* ATCC 824 engineered strains. *AMB Express* 2, 45.

978 (27) González-Pajuelo, M., Meynial-Salles, I., Mendes, F., Andrade, J. C., Vasconcelos, I.,
979 and Soucaille, P. (2005) Metabolic engineering of *Clostridium acetobutylicum* for the
980 industrial production of 1,3-propanediol from glycerol. *Metab. Eng.* 7, 329–336.

981 (28) Cai, X., and Bennett, G. N. (2011) Improving the *Clostridium acetobutylicum* butanol
982 fermentation by engineering the strain for co-production of riboflavin. *J. Ind. Microbiol.*
983 *Biotechnol.* 38, 1013–1025.

984 (29) Quinn, J. Y., Cox, R. S., Adler, A., Beal, J., Bhatia, S., Cai, Y., Chen, J., Clancy, K.,
985 Galdzicki, M., Hillson, N. J., Le Novère, N., Maheshwari, A. J., McLaughlin, J. A., Myers, C. J.,
986 Umesh, P., Pocock, M., Rodriguez, C., Soldatova, L., Stan, G. B. V., Swainston, N., Wipat, A.,
987 and Sauro, H. M. (2015) SBOL Visual: A Graphical Language for Genetic Designs. *PLoS*
988 *Biol.* 13, e1002310.

989 (30) Mermelstein, L. D., and Papoutsakis, E. T. (1993) *In vivo* methylation in *Escherichia*
990 *coli* by the *Bacillus subtilis* phage ϕ 3T I methyltransferase to protect plasmids from
991 restriction upon transformation of *Clostridium acetobutylicum* ATCC 824. *Appl. Environ.*
992 *Microbiol.* 59, 1077–1081.

993 (31) Nölling, J., Breton, G., Omelchenko, M. V, Makarova, K. S., Zeng, Q., Gibson, G., Hong
994 Mei Lee, Dubois, J., Qiu, D., Hitti, J., Aldredge, T., Ayers, M., Bashirzadeh, R., Bochner, H.,
995 Boivin, M., Bross, S., Bush, D., Butler, C., Caron, A., Caruso, A., Cook, R., Daggett, P.,
996 Deloughery, C., Egan, J., Ellston, D., Engelstein, M., Ezedi, J., Gilbert, K., Goyal, A., Guerin, J.,
997 Ho, T., Holtham, K., Joseph, P., Keagle, P., Kozlovsky, J., LaPlante, M., LeBlanc, G., Lumm,
998 W., Majeski, A., McDougall, S., Mank, P., Mao, J. I., Nocco, D., Patwell, D., Phillips, J., Pothier,
999 B., Prabhakar, S., Richterich, P., Rice, P., Rosetti, D., Rossetti, M., Rubenfield, M., Sachdeva,
1000 M., Snell, P., Spadafora, R., Spitzer, L., Shimer, G., Thomann, H. U., Vicaire, R., Wall, K.,

1001 Wang, Y., Weinstock, K., Lai Peng Wong, Wonsey, A., Xu, Q., Zhang, L., Wolf, Y. I., Tatusov,
1002 R. L., Sabathe, F., Doucette-Stamm, L., Soucaille, P., Daly, M. J., Bennett, G. N., Koonin, E. V,
1003 and Smith, D. R. (2001) Genome sequence and comparative analysis of the solvent-
1004 producing bacterium *Clostridium acetobutylicum*. *J. Bacteriol.* *183*, 4823–4838.

1005 (32) Oultram, J. D., Loughlin, M., Swinfield, T.-J. J., Brehm, J. K., Thompson, D. E., and
1006 Minton, N. P. (1988) Introduction of plasmids into whole cells of *Clostridium*
1007 *acetobutylicum* by electroporation. *FEMS Microbiol. Lett.* *56*, 83–88.

1008 (33) Pyne, M. E., Moo-Young, M., Chung, D. A., and Chou, C. P. (2013) Development of an
1009 electrotransformation protocol for genetic manipulation of *Clostridium pasteurianum*.
1010 *Biotechnol. Biofuels* *6*, 50.

1011 (34) Lesiak, J. M., Liebl, W., and Ehrenreich, A. (2014) Development of an *in vivo*
1012 methylation system for the solventogen *Clostridium saccharobutylicum* NCP 262 and
1013 analysis of two endonuclease mutants. *J. Biotechnol.* *188*, 97–99.

1014 (35) Herman, N. A., Li, J., Bedi, R., Turchi, B., Liu, X., Miller, M. J., and Zhang, W. (2017)
1015 Development of a High-Efficiency Transformation Method and Implementation of
1016 Rational Metabolic Engineering for the Industrial Butanol Hyperproducer *Clostridium*
1017 *saccharoperbutylacetonicum* Strain N1-4. *Appl. Environ. Microbiol.* *83*, AEM.02942-16.

1018 (36) Grosse-Honebrink, A., Schwarz, K. M., Wang, H., Minton, N. P., and Zhang, Y. (2017)
1019 Improving gene transfer in *Clostridium pasteurianum* through the isolation of rare
1020 hypertransformable variants. *Anaerobe* *48*, 203–205.

1021 (37) Kolek, J., Sedlar, K., Provaznik, I., and Patakova, P. (2016) Dam and Dcm
1022 methylations prevent gene transfer into *Clostridium pasteurianum* NRRL B-598:
1023 development of methods for electrotransformation, conjugation, and sonoporation.
1024 *Biotechnol. Biofuels* *9*, 14.

1025 (38) Oh, Y. H., Eom, G. T., Kang, K. H., Choi, J. W., Song, B. K., Lee, S. H., and Park, S. J.
1026 (2015) Optimized Transformation of Newly Constructed *Escherichia coli*-Clostridia
1027 Shuttle Vectors into *Clostridium beijerinckii*. *Appl. Biochem. Biotechnol.* *177*, 226–236.

1028 (39) Pyne, M. E., Bruder, M., Moo-Young, M., Chung, D. A., and Chou, C. P. (2014)
1029 Technical guide for genetic advancement of underdeveloped and intractable *Clostridium*.
1030 *Biotechnol. Adv.* *32*, 623–641.

1031 (40) Minton, N. P., Ehsaan, M., Humphreys, C. M., Little, G. T., Baker, J., Henstra, A. M.,
1032 Liew, F., Kelly, M. L., Sheng, L., Schwarz, K., and Zhang, Y. (2016) A roadmap for gene
1033 system development in *Clostridium*. *Anaerobe* *41*, 104–112.

1034 (41) Shaner, N. C., Steinbach, P. A., and Tsien, R. Y. (2005) A guide to choosing
1035 fluorescent proteins. *Nat. Methods* 2, 905–909.

1036 (42) Drepper, T., Eggert, T., Circolone, F., Heck, A., Krauß, U., Guterl, J.-K., Wendorff, M.,
1037 Losi, A., Gärtner, W., and Jaeger, K.-E. (2007) Reporter proteins for *in vivo* fluorescence
1038 without oxygen. *Nat. Biotechnol.* 25, 443–445.

1039 (43) Jean, D., Briolat, V., and Reysset, G. (2004) Oxidative stress response in *Clostridium*
1040 *perfringens*. *Microbiology* 150, 1649–1659.

1041 (44) Hartman, A. H., Liu, H., and Melville, S. B. (2011) Construction and Characterization
1042 of a Lactose-Inducible Promoter System for Controlled Gene Expression in *Clostridium*
1043 *perfringens*. *Appl. Environ. Microbiol.* 77, 471–478.

1044 (45) Ransom, E. M., Williams, K. B., Weiss, D. S., and Ellermeier, C. D. (2014)
1045 Identification and characterization of a gene cluster required for proper rod shape, cell
1046 division, and pathogenesis in *Clostridium difficile*. *J. Bacteriol.* 196, 2290–2300.

1047 (46) Ransom, E. M., Ellermeier, C. D., and Weiss, D. S. (2015) Use of mCherry red
1048 fluorescent protein for studies of protein localization and gene expression in *Clostridium*
1049 *difficile*. *Appl. Environ. Microbiol.* 81, 1652–1660.

1050 (47) Ransom, E. M., Weiss, D. S., and Ellermeier, C. D. (2016) Use of mCherryOpt
1051 fluorescent protein in *Clostridium difficile*. *Methods Mol. Biol.* 1476, 53–67.

1052 (48) Schulz, F. (2013) Fluoreszenzproteine in *Clostridium acetobutylicum* - Ein neues *in*
1053 *vivo* Reportersystem. Universität Rostock.

1054 (49) Li, Y., Xu, T., Tschaplinski, T. J., Engle, N. L., Yang, Y., Graham, D. E., He, Z., and Zhou, J.
1055 (2014) Improvement of cellulose catabolism in *Clostridium cellulolyticum* by sporulation
1056 abolishment and carbon alleviation. *Biotechnol. Biofuels* 7, 25.

1057 (50) Bruder, M. R., Pyne, M. E., Moo-Young, M., Chung, D. A., and Chou, C. P. (2016)
1058 Extending CRISPR-Cas9 technology from genome editing to transcriptional engineering
1059 in the genus *Clostridium*. *Appl. Environ. Microbiol.* 82, 6109–6119.

1060 (51) Molitor, B., Kirchner, K., Henrich, A. W., Schmitz, S., and Rosenbaum, M. A. (2016)
1061 Expanding the molecular toolkit for the homoacetogen *Clostridium ljungdahlii*. *Sci. Rep.*
1062 6, 31518.

1063 (52) Chapman, S., Faulkner, C., Kaiserli, E., Garcia-Mata, C., Savenkov, E. I., Roberts, A. G.,
1064 Oparka, K. J., and Christie, J. M. (2008) The photoreversible fluorescent protein iLOV
1065 outperforms GFP as a reporter of plant virus infection. *Proc. Natl. Acad. Sci.* 105, 20038–
1066 20043.

1067 (53) Christie, J. M., Hitomi, K., Arvai, A. S., Hartfield, K. a., Mettlen, M., Pratt, A. J., Tainer, J.
1068 a., and Getzoff, E. D. (2012) Structural tuning of the fluorescent protein iLOV for
1069 improved photostability. *J. Biol. Chem.* 287, 22295–22304.

1070 (54) Gawthorne, J. A., Reddick, L. E., Akpunarlieva, S. N., Beckham, K. S. H., Christie, J. M.,
1071 Alto, N. M., Gabrielsen, M., and Roe, A. J. (2012) Express Your LOV: An Engineered
1072 Flavoprotein as a Reporter for Protein Expression and Purification. *PLoS One* 7, e52962.

1073 (55) Buckley, A. M., Jukes, C., Candlish, D., Irvine, J. J., Spencer, J., Fagan, R. P., Roe, A. J.,
1074 Christie, J. M., Fairweather, N. F., and Douce, G. R. (2016) Lighting Up *Clostridium difficile*:
1075 Reporting Gene Expression Using Fluorescent Lov Domains. *Sci. Rep.* 6, 23463.

1076 (56) Khrenova, M. G., Nemukhin, A. V., and Domratcheva, T. (2015) Theoretical
1077 characterization of the flavin-based fluorescent protein iLOV and its Q489K mutant. *J.*
1078 *Phys. Chem. B* 119, 5176–5183.

1079 (57) Davari, M. D., Kopka, B., Wingen, M., Bocola, M., Drepper, T., Jaeger, K. E.,
1080 Schwaneberg, U., and Krauss, U. (2016) Photophysics of the LOV-Based Fluorescent
1081 Protein Variant iLOV-Q489K Determined by Simulation and Experiment. *J. Phys. Chem. B*
1082 120, 3344–3352.

1083 (58) Bullifent, H. L., Moir, A., and Titball, R. W. (1995) The construction of a reporter
1084 system and use for the investigation of *Clostridium perfringens* gene expression. *FEMS*
1085 *Microbiol. Lett.* 131, 99–105.

1086 (59) Zhang, Y., Grosse-Honebrink, A., and Minton, N. P. (2015) A universal mariner
1087 transposon system for forward genetic studies in the genus *Clostridium*. *PLoS One* 10,
1088 e0122411.

1089 (60) Shaw, W. V. (1975) Chloramphenicol acetyltransferase from chloramphenicol-
1090 resistant bacteria. *Methods Enzymol.* 43, 737–755.

1091 (61) Roberts, M. C., Swenson, C. D., Owens, L. M., and Smith, A. L. (1980) Characterization
1092 of chloramphenicol-resistant *Haemophilus influenzae*. *Antimicrob. Agents Chemother.* 18,
1093 610–615.

1094 (62) Tummala, S. B., Welker, N. E., and Papoutsakis, E. T. (1999) Development and
1095 characterization of a gene expression reporter system for *Clostridium acetobutylicum*
1096 ATCC 824. *Appl. Environ. Microbiol.* 65, 3793–3799.

1097 (63) Girbal, L., Mortier-Barrière, I., Raynaud, F., Rouanet, C., Croux, C., and Soucaille, P.
1098 (2003) Development of a sensitive gene expression reporter system and an inducible
1099 promoter-repressor system for *Clostridium acetobutylicum*. *Appl. Environ. Microbiol.* 69,

1100 4985–4988.

1101 (64) Moberg, L. J. (1985) Fluorogenic assay for rapid detection of *Escherichia coli* in food.
1102 *Appl. Environ. Microbiol.* 50, 1383–1387.

1103 (65) Sabathé, F., Croux, C., Cornillot, E., and Soucaille, P. (2002) *amyP*, a reporter gene to
1104 study strain degeneration in *Clostridium acetobutylicum* ATCC 824. *FEMS Microbiol. Lett.*
1105 210, 93–98.

1106 (66) Oliveira Paiva, A. M., Friggen, A. H., Hossein-Javaheri, S., and Smits, W. K. (2016) The
1107 Signal Sequence of the Abundant Extracellular Metalloprotease PPEP-1 Can Be Used to
1108 Secrete Synthetic Reporter Proteins in *Clostridium difficile*. *ACS Synth. Biol.* 5, 1376–
1109 1382.

1110 (67) Quixley, K. W., and Reid, S. J. (2000) Construction of a reporter gene vector for
1111 *Clostridium beijerinckii* using a *Clostridium* endoglucanase gene. *J. Mol. Microbiol.*
1112 *Biotechnol.* 2, 53–7.

1113 (68) Oh, Y. H., Eom, G. T., Kang, K. H., Joo, J. C., Jang, Y. A., Choi, J. W., Song, B. K., Lee, S. H.,
1114 and Park, S. J. (2016) Construction of heterologous gene expression cassettes for the
1115 development of recombinant *Clostridium beijerinckii*. *Bioprocess Biosyst. Eng.* 39, 555–
1116 563.

1117 (69) Gilham, D., and Lehner, R. (2005) Techniques to measure lipase and esterase
1118 activity *in vitro*. *Methods* 36, 139–147.

1119 (70) Hassan, E. A., Abd-Alla, M. H., Bagy, M. M. K., and Morsy, F. M. (2015) In situ
1120 hydrogen, acetone, butanol, ethanol and microdiesel production by *Clostridium*
1121 *acetobutylicum* ATCC 824 from oleaginous fungal biomass. *Anaerobe* 34, 125–131.

1122 (71) Edwards, A. N., Pascual, R. A., Childress, K. O., Nawrocki, K. L., Woods, E. C., and
1123 McBride, S. M. (2015) An alkaline phosphatase reporter for use in *Clostridium difficile*.
1124 *Anaerobe* 32, 98–104.

1125 (72) Feustel, L., Nakotte, S., and Dürre, P. (2004) Characterization and development of
1126 two reporter gene systems for *Clostridium acetobutylicum*. *Appl. Environ. Microbiol.* 70,
1127 798–803.

1128 (73) Potzkei, J., Kunze, M., Drepper, T., Gensch, T., Jaeger, K.-E., and Buechs, J. (2012)
1129 Real-time determination of intracellular oxygen in bacteria using a genetically encoded
1130 FRET-based biosensor. *BMC Biol.* 10, 28.

1131 (74) Martin, L., Che, A., and Endy, D. (2009) Gemini, a bifunctional enzymatic and
1132 fluorescent reporter of gene expression. *PLoS One* 4.

1133 (75) Haima, P., van Sinderen, D., Bron, S., and Venema, G. (1990) An improved β -
1134 galactosidase α -complementation system for molecular cloning in *Bacillus subtilis*. *Gene*
1135 93, 41–47.

1136 (76) Knipfer, N., Nooruddin, L., and Shrader, T. E. (1998) Development of an alpha-
1137 complementation system for mycobacterial promoter analysis. *Gene* 217, 69–75.

1138 (77) Volkov, I. Y., Lunina, N. A., Berezina, O. V., Velikodvorskaya, G. A., and Zverlov, V. V.
1139 (2005) *Thermoanaerobacter ethanolicus* gene cluster containing the α - and β -
1140 galactosidase genes *melA* and *lacA* and properties of recombinant LacA. *Mol. Biol.* 39,
1141 799–805.

1142 (78) Huber, R. E., Roth, N. J., and Bahl, H. (1996) Quaternary structure, Mg²⁺
1143 interactions, and some kinetic properties of the β -galactosidase from
1144 *Thermoanaerobacterium thermosulfurigenes* EM1. *J. Protein Chem.* 15, 621–629.

1145 (79) Browning, D. F., and Busby, S. J. W. (2004) The regulation of bacterial transcription
1146 initiation. *Nat. Rev. Microbiol.* 2, 57–65.

1147 (80) Helmann, J. D., and Chamberlin, M. J. (1988) Structure and function of bacterial
1148 sigma factors. *Annu. Rev. Biochem.* 57, 839–872.

1149 (81) Sauer, U., Santangelo, J. D., Treuner, A., Buchholz, M., and Dürre, P. (1995) Sigma
1150 factor and sporulation genes in *Clostridium*. *FEMS Microbiol. Rev.* 17, 331–340.

1151 (82) Estrem, S. T., Gaal, T., Ross, W., and Gourse, R. L. (1998) Identification of an UP
1152 element consensus sequence for bacterial promoters. *Biochemistry* 95, 9761–9766.

1153 (83) Kumar, A., Malloch, R. A., Fujita, N., Smillie, D. A., Ishihama, A., and Hayward, R. S.
1154 (1993) The minus 35-recognition region of *Escherichia coli* sigma 70 is inessential for
1155 initiation of transcription at an “extended minus 10” promoter. *J. Mol. Biol.* 232, 406–18.

1156 (84) Sinoquet, C., Demey, S., and Braun, F. (2008) Large-scale computational and
1157 statistical analyses of high transcription potentialities in 32 prokaryotic genomes.
1158 *Nucleic Acids Res.* 36, 3332–3340.

1159 (85) Katayama, S., Matsushita, O., Jung, C. M., Minami, J., and Okabe, A. (1999) Promoter
1160 upstream bent DNA activates the transcription of the *Clostridium perfringens*
1161 phospholipase C gene in a low temperature-dependent manner. *EMBO J.* 18, 3442–3450.

1162 (86) Graves, M. C., and Rabinowitz, J. C. (1986) *In vivo* and *in vitro* transcription of the
1163 *Clostridium pasteurianum* ferredoxin gene. Evidence for “extended” promoter elements
1164 in gram-positive organisms. *J. Biol. Chem.* 261, 11409–15.

1165 (87) Pyne, M. E. (2014, April) Development of genetic tools for metabolic engineering of

1166 *Clostridium pasteurianum*. University of Waterloo.

1167 (88) Wiesenborn, D. P., Rudolph, F. B., and Papoutsakis, E. T. (1988) Thiolase from
1168 *Clostridium acetobutylicum* ATCC 824 and Its Role in the Synthesis of Acids and Solvents.
1169 *Appl Env. Microbiol* 54, 2717–2722.

1170 (89) Stim-Herndon, K. P., Petersen, D. J., and Bennett, G. N. (1995) Characterization of an
1171 acetyl-CoA C-acetyltransferase (thiolase) gene from *Clostridium acetobutylicum* ATCC
1172 824. *Gene*, pp 81–85.

1173 (90) Bormann, S., Baer, Z. C., Sreekumar, S., Kuchenreuther, J. M., Dean Toste, F., Blanch,
1174 H. W., and Clark, D. S. (2014) Engineering *Clostridium acetobutylicum* for production of
1175 kerosene and diesel blendstock precursors. *Metab. Eng.* 25, 124–130.

1176 (91) Dong, H., Tao, W., Zhang, Y., and Li, Y. (2012) Development of an
1177 anhydrotetracycline-inducible gene expression system for solvent-producing
1178 *Clostridium acetobutylicum*: A useful tool for strain engineering. *Metab. Eng.* 14, 59–67.

1179 (92) Perret, S., Casalot, L., Fierobe, H. P., Tardif, C., Sabathe, F., Belaich, J. P., and Belaich,
1180 A. (2004) Production of Heterologous and Chimeric Scaffoldins by *Clostridium*
1181 *acetobutylicum* ATCC 824. *J. Bacteriol.* 186, 253–257.

1182 (93) Mingardon, F., Perret, S., Bélaïch, A., Tardif, C., Bélaïch, J. P., and Fierobe, H. P. (2005)
1183 Heterologous production, assembly, and secretion of a minicellulosome by *Clostridium*
1184 *acetobutylicum* ATCC 824. *Appl. Environ. Microbiol.* 71, 1215–1222.

1185 (94) Wietzke, M., and Bahl, H. (2012) The redox-sensing protein Rex, a transcriptional
1186 regulator of solventogenesis in *Clostridium acetobutylicum*. *Appl. Microbiol. Biotechnol.*
1187 96, 749–761.

1188 (95) Zhang, L., Nie, X., Ravcheev, D. A., Rodionov, D. A., Sheng, J., Gu, Y., Yang, S., Jiang, W.,
1189 and Yang, C. (2014) Redox-responsive repressor rex modulates alcohol production and
1190 oxidative stress tolerance in *Clostridium acetobutylicum*. *J. Bacteriol.* 196, 3949–3963.

1191 (96) Heap, J. T., Pennington, O. J., Cartman, S. T., and Minton, N. P. (2009) A modular
1192 system for *Clostridium* shuttle plasmids. *J. Microbiol. Methods* 78, 79–85.

1193 (97) Novichkov, P. S., Kazakov, A. E., Ravcheev, D. A., Leyn, S. A., Kovaleva, G. Y., Sutormin,
1194 R. A., Kazanov, M. D., Riehl, W., Arkin, A. P., Dubchak, I., and Rodionov, D. A. (2013)
1195 RegPrecise 3.0--a resource for genome-scale exploration of transcriptional regulation in
1196 bacteria. *BMC Genomics* 14, 745.

1197 (98) Wang, Y., Zhang, Z.-T., Seo, S.-O., Lynn, P., Lu, T., Jin, Y.-S., and Blaschek, H. P. (2016)
1198 Gene transcription repression in *Clostridium beijerinckii* using CRISPR-dCas9. *Biotechnol.*

1199 *Bioeng. 113*, 2739–2743.

1200 (99) Pyne, M. E., Moo-Young, M., Chung, D. A., and Chou, C. P. (2014) Expansion of the
1201 genetic toolkit for metabolic engineering of *Clostridium pasteurianum*: chromosomal
1202 gene disruption of the endogenous CpaAI restriction enzyme. *Biotechnol. Biofuels* 7, 163.

1203 (100) Wang, Y., Li, X., Mao, Y., and Blaschek, H. P. (2012) Genome-wide dynamic
1204 transcriptional profiling in *Clostridium beijerinckii* NCIMB 8052 using single-nucleotide
1205 resolution RNA-Seq. *BMC Genomics* 13, 102.

1206 (101) Al-Hinai, M. A., Fast, A. G., and Papoutsakis, E. T. (2012) Novel system for efficient
1207 isolation of *Clostridium* double-crossover allelic exchange mutants enabling markerless
1208 chromosomal gene deletions and DNA integration. *Appl. Environ. Microbiol.* 78, 8112–
1209 8121.

1210 (102) Garcia, H. G., Sanchez, A., Boedicker, J. Q., Osborne, M., Gelles, J., Kondev, J., and
1211 Phillips, R. (2012) Operator sequence alters gene expression independently of
1212 transcription factor occupancy in bacteria. *Cell Rep.* 2, 150–161.

1213 (103) Zong, Y., Zhang, H. M., Lyu, C., Ji, X., Hou, J., Guo, X., Ouyang, Q., and Lou, C. (2017)
1214 Insulated transcriptional elements enable precise design of genetic circuits. *Nat.*
1215 *Commun.* 8, 52.

1216 (104) Li, Q., Chen, J., Minton, N. P., Zhang, Y., Wen, Z., Liu, J., Yang, H., Zeng, Z., Ren, X.,
1217 Yang, J., Gu, Y., Jiang, W., Jiang, Y., and Yang, S. (2016) CRISPR-based genome editing and
1218 expression control systems in *Clostridium acetobutylicum* and *Clostridium beijerinckii*.
1219 *Biotechnol. J.* 11, 961–972.

1220 (105) Tabor, S., and Richardson, C. C. (1985) A bacteriophage T7 RNA
1221 polymerase/promoter system for controlled exclusive expression of specific genes (T7
1222 DNA polymerase/T7 gene 5 protein/proteolysis/ β -lactamase/rifampicin). *Biochemistry*
1223 24, 1074–1078.

1224 (106) McLaughlin, J. R., Murray, C. L., and Rabinowitz, J. C. (1981) Unique features in the
1225 ribosome binding site sequence of the gram-positive *Staphylococcus aureus* beta-
1226 lactamase gene. *J. Biol. Chem.* 256, 11283–91.

1227 (107) Lamberte, L. E., Baniulyte, G., Singh, S. S., Stringer, A. M., Bonocora, R. P., Stracy, M.,
1228 Kapanidis, A. N., Wade, J. T., and Grainger, D. C. (2017) Horizontally acquired AT-rich
1229 genes in *Escherichia coli* cause toxicity by sequestering RNA polymerase. *Nat. Microbiol.*
1230 2, 16249.

1231 (108) Heap, J. T., Kuehne, S. A., Ehsaan, M., Cartman, S. T., Cooksley, C. M., Scott, J. C., and

1232 Minton, N. P. (2010) The ClosTron: Mutagenesis in *Clostridium* refined and streamlined.
1233 *J. Microbiol. Methods* 80, 49–55.

1234 (109) Ehsaan, M., Kuit, W., Zhang, Y., Cartman, S. T., Heap, J. T., Winzer, K., and Minton, N.
1235 P. (2016) Mutant generation by allelic exchange and genome resequencing of the
1236 biobutanol organism *Clostridium acetobutylicum* ATCC 824. *Biotechnol. Biofuels* 9, 4.

1237 (110) Heap, J. T., Pennington, O. J., Cartman, S. T., Carter, G. P., and Minton, N. P. (2007)
1238 The ClosTron: A universal gene knock-out system for the genus *Clostridium*. *J. Microbiol.*
1239 *Methods* 70, 452–464.

1240 (111) Geissendörfer, M., and Hillen, W. (1990) Regulated expression of heterologous
1241 genes in *Bacillus subtilis* using the Tn10 encoded *tet* regulatory elements. *Appl. Microbiol.*
1242 *Biotechnol.* 33, 657–63.

1243 (112) Fontaine, L., Meynial-Salles, I., Girbal, L., Yang, X., Croux, C., and Soucaille, P. (2002)
1244 Molecular characterization and transcriptional analysis of *adhE2*, the gene encoding the
1245 NADH-dependent aldehyde/alcohol dehydrogenase responsible for butanol production
1246 in alcohologenic cultures of *Clostridium acetobutylicum* ATCC 824. *J. Bacteriol.* 184, 821–
1247 830.

1248 (113) Ciampi, M. S. (2006) Rho-dependent terminators and transcription termination.
1249 *Microbiology* 152, 2515–2528.

1250 (114) Liu, C. C., Qi, L., Lucks, J. B., Segall-Shapiro, T. H., Wang, D., Mutalik, V. K., and Arkin,
1251 A. P. (2012) An adaptor from translational to transcriptional control enables predictable
1252 assembly of complex regulation. *Nat. Methods* 9, 1088–1094.

1253 (115) Banerjee, S., Chalissery, J., Bandey, I., and Sen, R. (2006) Rho-dependent
1254 transcription termination: more questions than answers. *J. Microbiol.* 44, 11–22.

1255 (116) Gardner, P. P., Barquist, L., Bateman, A., Nawrocki, E. P., and Weinberg, Z. (2011)
1256 RNIE: Genome-wide prediction of bacterial intrinsic terminators. *Nucleic Acids Res.* 39,
1257 5845–5852.

1258 (117) Gusarov, I., and Nudler, E. (1999) The mechanism of intrinsic transcription
1259 termination. *Mol. Cell* 3, 495–504.

1260 (118) Sugimoto, N., Nakano, S. ichi, Katoh, M., Matsumura, A., Nakamuta, H., Ohmichi, T.,
1261 Yoneyama, M., and Sasaki, M. (1995) Thermodynamic Parameters To Predict Stability of
1262 RNA/DNA Hybrid Duplexes. *Biochemistry* 34, 11211–11216.

1263 (119) Naville, M., and Gautheret, D. (2010) Premature terminator analysis sheds light on
1264 a hidden world of bacterial transcriptional attenuation. *Genome Biol.* 11, R97.

1265 (120) Ravagnani, A., Jennert, K. C. B., Steiner, E., Grünberg, R., Jefferies, J. R., Wilkinson, S.
1266 R., Young, D. I., Tidswell, E. C., Brown, D. P., Youngman, P., Gareth Morris, J., and Young,
1267 M. (2000) Spo0A directly controls the switch from acid to solvent production in solvent-
1268 forming clostridia. *Mol. Microbiol.* *37*, 1172–1185.

1269 (121) Wang, Y., Zhang, Z. T., Seo, S. O., Lynn, P., Lu, T., Jin, Y. S., and Blaschek, H. P. (2016)
1270 Bacterial Genome Editing with CRISPR-Cas9: Deletion, Integration, Single Nucleotide
1271 Modification, and Desirable “Clean” Mutant Selection in *Clostridium beijerinckii* as an
1272 Example. *ACS Synth. Biol.* *5*, 721–732.

1273 (122) Carrier, T. A., and Keasling, J. D. (1997) Controlling messenger RNA stability in
1274 bacteria: Strategies for engineering gene expression. *Biotechnol. Prog.* *13*, 699–708.

1275 (123) Chen, Y.-J., Liu, P., Nielsen, A. A. K., Brophy, J. A. N., Clancy, K., Peterson, T., and
1276 Voigt, C. A. (2013) Characterization of 582 natural and synthetic terminators and
1277 quantification of their design constraints. *Nat. Methods* *10*, 659–664.

1278 (124) Lin, M. T., Wang, C. Y., Xie, H. J., Cheung, C. H. Y., Hsieh, C. H., Juan, H. F., Chen, B.
1279 Sen, and Lin, C. (2016) Novel Utilization of Terminators in the Design of Biologically
1280 Adjustable Synthetic Filters. *ACS Synth. Biol.* *5*, 365–374.

1281 (125) Abe, H., Abo, T., and Aiba, H. (1999) Regulation of intrinsic terminator by
1282 translation in *Escherichia coli*: transcription termination at a distance downstream.
1283 *Genes Cells* *4*, 87–97.

1284 (126) Li, R., Zhang, Q., Li, J., and Shi, H. (2015) Effects of cooperation between translating
1285 ribosome and RNA polymerase on termination efficiency of the Rho-independent
1286 terminator. *Nucleic Acids Res.* *44*, 2554–2563.

1287 (127) Lesnik, E. A., Sampath, R., Levene, H. B., Henderson, T. J., McNeil, J. A., and Ecker, D.
1288 J. (2001) Prediction of rho-independent transcriptional terminators in *Escherichia coli*.
1289 *Nucleic Acids Res.* *29*, 3583–94.

1290 (128) Kingsford, C. L., Ayanbule, K., and Salzberg, S. L. (2007) Rapid, accurate,
1291 computational discovery of Rho-independent transcription terminators illuminates their
1292 relationship to DNA uptake. *Genome Biol.* *8*, R22.

1293 (129) Mitra, A., Angamuthu, K., Jayashree, H. V., and Nagaraja, V. (2009) Occurrence,
1294 divergence and evolution of intrinsic terminators across Eubacteria. *Genomics* *94*, 110–
1295 116.

1296 (130) Mitra, A., Kesarwani, A. K., Pal, D., and Nagaraja, V. (2011) WebGeSTer DB-A
1297 transcription terminator database. *Nucleic Acids Res.* *39*, D129-35.

1298 (131) Wright, J. J., Kumar, A., and Hayward, R. S. (1992) Hypersymmetry in a
1299 transcriptional terminator of *Escherichia coli* confers increased efficiency as well as
1300 bidirectionality. *EMBO J.* 11, 1957–64.

1301 (132) Yang, M. Te, Scott, H. B., and Gardner, J. F. (1995) Transcription termination at the
1302 *thr* attenuator: Evidence that the adenine residues upstream of the stem and loop
1303 structure are not required for termination. *J. Biol. Chem.* 270, 23330–23336.

1304 (133) Paredes, C. J., Rigoutsos, I., and Papoutsakis, T. (2004) Transcriptional
1305 organization of the *Clostridium acetobutylicum* genome. *Nucleic Acids Res.* 32, 1973–
1306 1981.

1307 (134) Shultzaberger, R. K., Bucheimer, R. E., Rudd, K. E., and Schneider, T. D. (2001)
1308 Anatomy of *Escherichia coli* ribosome binding sites. *J. Mol. Biol.* 313, 215–228.

1309 (135) Yang, Y., Lang, N., Yang, G., Yang, S., Jiang, W., and Gu, Y. (2016) Improving the
1310 performance of solventogenic clostridia by reinforcing the biotin synthetic pathway.
1311 *Metab. Eng.* 35, 121–128.

1312 (136) Yang, G., Jia, D., Jin, L., Jiang, Y., Wang, Y., Jiang, W., and Gu, Y. (2017) Rapid
1313 Generation of Universal Synthetic Promoters for Controlled Gene Expression in Both
1314 Gas-Fermenting and Saccharolytic *Clostridium* Species. *ACS Synth. Biol.* 6, 1672–1678.

1315 (137) Chen, H., Bjercknes, M., Kumar, R., and Jay, E. (1994) Determination of the optimal
1316 aligned spacing between the Shine - Dalgarno sequence and the translation initiation
1317 codon of *Escherichia coli* mRNAs. *Nucleic Acids Res.* 22, 4953–4957.

1318 (138) Hager, P. W., and Rabinowitz, J. C. (1985) Translational specificity in *Bacillus*
1319 *subtilis*. *Mol. Biol. bacilli* 2, 1–32.

1320 (139) Band, L., and Henner, D. J. (1984) *Bacillus subtilis* requires a “stringent” Shine-
1321 Dalgarno region for gene expression. *DNA* 3, 17–21.

1322 (140) Abolbaghaei, A., Silke, J. R., and Xia, X. (2017) How Changes in Anti-SD Sequences
1323 Would Affect SD Sequences in *Escherichia coli* and *Bacillus subtilis*. *G3 (Bethesda)*. 7,
1324 1607–1615.

1325 (141) Dunn, J. J., and Studier, F. W. (1981) Nucleotide sequence from the genetic left end
1326 of bacteriophage T7 DNA to the beginning of gene 4. *J. Mol. Biol.* 148, 303–330.

1327 (142) Ringquist, S., Shinedling, S., Barrick, D., Green, L., Binkley, J., Stormo, G. D., and
1328 Gold, L. (1992) Translation initiation in *Escherichia coli*: sequences within the ribosome-
1329 binding site. *Mol. Microbiol.* 6, 1219–1229.

1330 (143) Garnier, T., Canard, B., and Cole, S. T. (1991) Cloning, mapping, and molecular

1331 characterization of the rRNA operons of *Clostridium perfringens*. *J. Bacteriol.* *173*, 5431–
1332 5438.

1333 (144) Ma, J., Campbell, A., and Karlin, S. (2002) Correlations between Shine-Dalgarno
1334 sequences and gene features such as predicted expression levels and operon structures.
1335 *J. Bacteriol.* *184*, 5733–5745.

1336 (145) He, J., Sakaguchi, K., and Suzuki, T. (2012) Determination of the ribosome-binding
1337 sequence and spacer length between binding site and initiation codon for efficient
1338 protein expression in *Bifidobacterium longum* 105-A. *J. Biosci. Bioeng.* *113*, 442–444.

1339 (146) Salis, H. M. (2011) The ribosome binding site calculator. *Methods Enzymol.* *498*,
1340 19–42.

1341 (147) Salis, H. M., Mirsky, E. A., and Voigt, C. A. (2009) Automated design of synthetic
1342 ribosome binding sites to control protein expression. *Nat. Biotechnol.* *27*, 946–950.

1343 (148) Reeve, B., Martinez-Klimova, E., De Jonghe, J., Leak, D. J., and Ellis, T. (2016) The
1344 *Geobacillus* Plasmid Set: A Modular Toolkit for Thermophile Engineering. *ACS Synth. Biol.*
1345 *5*, 1342–1347.

1346 (149) Ikemura, T. (1985) Codon usage and tRNA content in unicellular and multicellular
1347 organisms. *Mol. Biol. Evol.* *2*, 13–34.

1348 (150) Sharp, P. M., and Li, W. H. (1987) The codon adaptation index—a measure of
1349 directional synonymous codon usage bias, and its potential applications. *Nucleic Acids*
1350 *Res.* *15*, 1281–1295.

1351 (151) Baca, A. M., and Hol, W. G. J. (2000) Overcoming codon bias: A method for high-
1352 level overexpression of *Plasmodium* and other AT-rich parasite genes in *Escherichia coli*.
1353 *Int. J. Parasitol.* *30*, 113–118.

1354 (152) Novy, R., Drott, D., Yaeger, K., and Mierendorf, R. (2001) Overcoming the codon
1355 bias of *E. coli* for enhanced protein expression. *Innovations* *12*, 4–6.

1356 (153) Mondal, S. K., Kundu, S., Das, R., and Roy, S. (2016) Analysis of phylogeny and
1357 codon usage bias and relationship of GC content, amino acid composition with
1358 expression of the structural *nif* genes. *J. Biomol. Struct. Dyn.* *34*, 1649–1666.

1359 (154) Musto, H., Romero, H., and Zavala, A. (2003) Translational selection is operative
1360 for synonymous codon usage in *Clostridium perfringens* and *Clostridium acetobutylicum*.
1361 *Microbiology* *149*, 855–863.

1362 (155) Sastalla, I., Chim, K., Cheung, G. Y. C., Pomerantsev, A. P., and Leppla, S. H. (2009)
1363 Codon-optimized fluorescent proteins designed for expression in low-GC gram-positive

1364 bacteria. *Appl. Environ. Microbiol.* 75, 2099–2110.

1365 (156) Wright, F. (1990) The “effective number of codons” used in a gene. *Gene* 87, 23–29.

1366 (157) Kudla, G., Murray, A. W., Tollervey, D., and Plotkin, J. B. (2009) Coding-Sequence
1367 Determinants of Gene Expression in *Escherichia coli*. *Science* (80-.). 324, 255–258.

1368 (158) Welch, M., Govindarajan, S., Ness, J. E., Villalobos, A., Gurney, A., Minshull, J., and
1369 Gustafsson, C. (2009) Design parameters to control synthetic gene expression in
1370 *Escherichia coli*. *PLoS One* 4, e7002.

1371 (159) Xia, X. (2007) An improved implementation of codon adaptation index. *Evol.*
1372 *Bioinforma.* 3, 53–58.

1373 (160) Xia, X. (2015) A major controversy in codon-Anticodon adaptation resolved by a
1374 new codon usage index. *Genetics* 199, 573–579.

1375 (161) Lanza, A. M., Curran, K. A., Rey, L. G., and Alper, H. S. (2014) A condition-specific
1376 codon optimization approach for improved heterologous gene expression in
1377 *Saccharomyces cerevisiae*. *BMC Syst. Biol.* 8, 33.

1378 (162) Welch, M., Villalobos, A., Gustafsson, C., and Minshull, J. (2009) You’re one in a
1379 googol: optimizing genes for protein expression. *J. R. Soc. Interface* 6, S467–S476.

1380 (163) Plotkin, J. B., and Kudla, G. (2011) Synonymous but not the same: the causes and
1381 consequences of codon bias. *Nat. Rev. Genet.* 12, 32–42.

1382 (164) Carrier, T. A., and Keasling, J. D. (1999) Library of synthetic 5’ secondary
1383 structures to manipulate mRNA stability in *Escherichia coli*. *Biotechnol. Prog.* 15, 58–64.

1384 (165) Condon, C., and Putzer, H. (2002) The phylogenetic distribution of bacterial
1385 ribonucleases. *Nucleic Acids Res.* 30, 5339–5346.

1386 (166) Durand, S., Tomasini, A., Braun, F., Condon, C., and Romby, P. (2015) sRNA and
1387 mRNA turnover in Gram-positive bacteria. *FEMS Microbiol. Rev.* 39, 316–330.

1388 (167) Lehnik-Habrink, M., Newman, J., Rothe, F. M., Solovyova, A. S., Rodrigues, C.,
1389 Herzberg, C., Commichau, F. M., Lewis, R. J., and Stülke, J. (2011) RNase Y in *Bacillus*
1390 *subtilis*: A natively disordered protein that is the functional equivalent of RNase E from
1391 *Escherichia coli*. *J. Bacteriol.* 193, 5431–5441.

1392 (168) Lee, J., Jang, Y. S., Papoutsakis, E. T., and Lee, S. Y. (2016) Stable and enhanced gene
1393 expression in *Clostridium acetobutylicum* using synthetic untranslated regions with a
1394 stem-loop. *J. Biotechnol.* 230, 40–43.

1395 (169) Desai, R. P., and Papoutsakis, E. T. (1999) Antisense RNA strategies for metabolic
1396 engineering of *Clostridium acetobutylicum*. *Appl. Environ. Microbiol.* 65, 936–945.

1397 (170) Song, J.-W., Woo, J.-M., Jung, G. Y., Bornscheuer, U. T., and Park, J.-B. (2016) 3'-UTR
1398 engineering to improve soluble expression and fine-tuning of activity of cascade
1399 enzymes in *Escherichia coli*. *Sci. Rep.* 6, 29406.

1400 (171) Tummala, S. B., Junne, S. G., and Papoutsakis, E. T. (2003) Antisense RNA
1401 downregulation of coenzyme A transferase combined with alcohol-aldehyde
1402 dehydrogenase overexpression leads to predominantly alcohologenic *Clostridium*
1403 *acetobutylicum* fermentations. *J. Bacteriol.* 185, 3644–3653.

1404 (172) Pyne, M., Moo-Young, M., Chung, D., and Chou, C. (2015) Antisense-RNA-Mediated
1405 Gene Downregulation in *Clostridium pasteurianum*. *Fermentation* 1, 113–126.

1406 (173) Storz, G., Altuvia, S., and Wassarman, K. M. (2005) AN ABUNDANCE OF RNA
1407 REGULATORS. *Annu. Rev. Biochem.* 74, 199–217.

1408 (174) Mutalik, V. K., Qi, L., Guimaraes, J. C., Lucks, J. B., and Arkin, A. P. (2012) Rationally
1409 designed families of orthogonal RNA regulators of translation. *Nat. Chem. Biol.* 8, 447–
1410 454.

1411 (175) Isaacs, F. J., Dwyer, D. J., Ding, C., Pervouchine, D. D., Cantor, C. R., and Collins, J. J.
1412 (2004) Engineered riboregulators enable post-transcriptional control of gene
1413 expression. *Nat. Biotechnol.* 22, 841–847.

1414 (176) Green, A. A., Silver, P. A., Collins, J. J., and Yin, P. (2014) Toehold switches: De-novo-
1415 designed regulators of gene expression. *Cell* 159, 925–939.

1416 (177) Garst, A. D., Edwards, A. L., and Batey, R. T. (2011) Riboswitches: Structures and
1417 mechanisms. *Cold Spring Harb. Perspect. Biol.* 3, 1–13.

1418 (178) Mellin, J. R., and Cossart, P. (2015) Unexpected versatility in bacterial
1419 riboswitches. *Trends Genet.* 31, 150–156.

1420 (179) Wittmann, A., and Suess, B. (2012) Engineered riboswitches: Expanding
1421 researchers' toolbox with synthetic RNA regulators. *FEBS Lett.* 586, 2076–2083.

1422 (180) Saad, N. Y., Schiel, B., Brayé, M., Heap, J. T., Minton, N. P., Dürre, P., and Becker, H. D.
1423 (2012) Riboswitch (T-box)-mediated control of tRNA-dependent amidation in
1424 *Clostridium acetobutylicum* rationalizes gene and pathway redundancy for asparagine
1425 and asparaginyl-tRNA^{Asn} synthesis. *J. Biol. Chem.* 287, 20382–20394.

1426 (181) André, G., Even, S., Putzer, H., Burguière, P., Croux, C., Danchin, A., Martin-
1427 Verstraete, I., and Soutourina, O. (2008) S-box and T-box riboswitches and antisense
1428 RNA control a sulfur metabolic operon of *Clostridium acetobutylicum*. *Nucleic Acids Res.*
1429 36, 5955–5969.

1430 (182) Kim, P. B., Nelson, J. W., and Breaker, R. R. (2015) An ancient riboswitch class in
1431 bacteria regulates purine biosynthesis and one-carbon metabolism. *Mol. Cell* 57, 317–
1432 328.

1433 (183) Projan, S. J., Monod, M., Narayanan, C. S., and Dubnau, D. (1987) Replication
1434 properties of pIM13, a naturally occurring plasmid found in *Bacillus subtilis*, and of its
1435 close relative pE5, a plasmid native to *Staphylococcus aureus*. *J. Bacteriol.* 169, 5131–
1436 5139.

1437 (184) Garnier, T., and Cole, S. T. (1988) Identification and Molecular Genetic Analysis of
1438 Replication Functions of the Bacteriocinogenic Plasmid pIP404 from *Clostridium*
1439 *perfringens*. *Plasmid* 19, 151–160.

1440 (185) Purdy, D., O’Keeffe, T. A. T., Elmore, M., Herbert, M., McLeod, A., Bokori-Brown, M.,
1441 Ostrowski, A., and Minton, N. P. (2002) Conjugative transfer of clostridial shuttle vectors
1442 from *Escherichia coli* to *Clostridium difficile* through circumvention of the restriction
1443 barrier. *Mol. Microbiol.* 46, 439–452.

1444 (186) Lee, S. H., Kwon, M. A., Choi, S., Kim, S., Kim, J., Shin, Y. A., and Kim, K. H. (2015) A
1445 new shuttle plasmid that stably replicates in *Clostridium acetobutylicum*. *J. Microbiol.*
1446 *Biotechnol.* 25, 1702–1708.

1447 (187) Li, Y., Blaschek, H. P., and Tan, D. (2002) Molecular characterization and utilization
1448 of the CAK1 filamentous viruslike particle derived from *Clostridium beijerinckii*. *J. Ind.*
1449 *Microbiol. Biotechnol.* 28, 118–26.

1450 (188) Nakayama, S., Irie, R., Kosaka, T., Matsuura, K., Yoshino, S., and Furukawa, K.
1451 (2007) New host-vector system in solvent-producing *Clostridium*
1452 *saccharoperbutylacetonicum* strain N1-4. *J. Gen. Appl. Microbiol.* 53, 53–6.

1453 (189) Maguin, E., Duwat, P., Hege, T., Ehrlich, D., and Gruss, A. (1992) New
1454 thermosensitive plasmid for gram-positive bacteria. *J. Bacteriol.* 174, 5633–5638.

1455 (190) Heap, J. T., Ehsaan, M., Cooksley, C. M., Ng, Y. K., Cartman, S. T., Winzer, K., and
1456 Minton, N. P. (2012) Integration of DNA into bacterial chromosomes from plasmids
1457 without a counter-selection marker. *Nucleic Acids Res.* 40, e59.

1458 (191) Gu, P., Yang, F., Su, T., Wang, Q., Liang, Q., and Qi, Q. (2015) A rapid and reliable
1459 strategy for chromosomal integration of gene(s) with multiple copies. *Sci. Rep.* 5, 9684.

1460 (192) Kuhlman, T. E., and Cox, E. C. (2010) Site-specific chromosomal integration of
1461 large synthetic constructs. *Nucleic Acids Res.* 38, e92.

1462 (193) Bryant, J. A., Sellars, L. E., Busby, S. J. W., and Lee, D. J. (2014) Chromosome

1463 position effects on gene expression in *Escherichia coli* K-12. *Nucleic Acids Res.* 42, 11383–
1464 11392.

1465 (194) Berger, M., Gerganova, V., Berger, P., Rapiteanu, R., and Lisicovas, V. (2016) Genes
1466 on a Wire : The Nucleoid- Associated Protein HU Insulates Transcription Units in
1467 *Escherichia coli*. *Nat. Publ. Gr.* 6, 1–12.

1468 (195) Sauer, C., Syvertsson, S., Bohorquez, L. C., Cruz, R., Harwood, C. R., Van Rij, T., and
1469 Hamoen, L. W. (2016) Effect of Genome Position on Heterologous Gene Expression in
1470 *Bacillus subtilis*: An Unbiased Analysis. *ACS Synth. Biol.* 5, 942–947.

1471 (196) Anuchin, A. M., Goncharenko, A. V, Demidenok, O. I., and Kaprelyants, A. S. (2011)
1472 Histone-like proteins of bacteria (review). *Appl. Biochem. Microbiol.* 47, 580–585.

1473 (197) Sharan, S. K., Thomason, L. C., Kuznetsov, S. G., and Court, D. L. (2009)
1474 Recombineering: a homologous recombination-based method of genetic engineering.
1475 *Nat. Protoc.* 4, 206–223.

1476 (198) Li, Y., Lin, Z., Huang, C., Zhang, Y., Wang, Z., Tang, Y. jie, Chen, T., and Zhao, X.
1477 (2015) Metabolic engineering of *Escherichia coli* using CRISPR-Cas9 mediated genome
1478 editing. *Metab. Eng.* 31, 13–21.

1479 (199) Jiang, Y., Chen, B., Duan, C., Sun, B., Yang, J., and Yang, S. (2015) Multigene editing
1480 in the *Escherichia coli* genome via the CRISPR-Cas9 system. *Appl. Environ. Microbiol.* 81,
1481 2506–2514.

1482 (200) Croux, C., Nguyen, N.-P.-T., Lee, J., Raynaud, C., Saint-Prix, F., Gonzalez-Pajuelo, M.,
1483 Meynial-Salles, I., and Soucaille, P. (2016) Construction of a restriction-less, marker-less
1484 mutant useful for functional genomic and metabolic engineering of the biofuel producer
1485 *Clostridium acetobutylicum*. *Biotechnol. Biofuels* 9, 23.

1486 (201) Yoo, M., Croux, C., Meynial-Salles, I., and Soucaille, P. (2016) Elucidation of the
1487 roles of *adhE1* and *adhE2* in the primary metabolism of *Clostridium acetobutylicum* by
1488 combining in-frame gene deletion and a quantitative system-scale approach. *Biotechnol.*
1489 *Biofuels* 9, 92.

1490 (202) Zhang, N., Shao, L., Jiang, Y., Gu, Y., Li, Q., Liu, J., Jiang, W., and Yang, S. (2015) I-SceI-
1491 mediated scarless gene modification via allelic exchange in *Clostridium*. *J. Microbiol.*
1492 *Methods* 108, 49–60.

1493 (203) Dong, H., Tao, W., Gong, F., Li, Y., and Zhang, Y. (2014) A functional *recT* gene for
1494 recombineering of *Clostridium*. *J. Biotechnol.* 173, 65–67.

1495 (204) Wang, Y., Zhang, Z. T., Seo, S. O., Choi, K., Lu, T., Jin, Y. S., and Blaschek, H. P. (2015)

1496 Markerless chromosomal gene deletion in *Clostridium beijerinckii* using CRISPR/Cas9
1497 system. *J. Biotechnol.* 200, 1–5.

1498 (205) Pyne, M. E., Bruder, M. R., Moo-Young, M., Chung, D. A., and Chou, C. P. (2016)
1499 Harnessing heterologous and endogenous CRISPR-Cas machineries for efficient
1500 markerless genome editing in *Clostridium*. *Sci. Rep.* 6, 25666.

1501 (206) Wang, S., Dong, S., Wang, P., Tao, Y., and Wang, Y. (2017) Genome editing in
1502 *Clostridium saccharoperbutylacetonicum* N1-4 with the CRISPR-Cas9 system. *Appl.*
1503 *Environ. Microbiol.* 83.

1504 (207) Kosuri, S., Goodman, D. B., Cambray, G., Mutalik, V. K., Gao, Y., Arkin, A. P., Endy, D.,
1505 and Church, G. M. (2013) Composability of regulatory sequences controlling
1506 transcription and translation in *Escherichia coli*. *Proc. Natl. Acad. Sci.* 110, 14024–14029.

1507 (208) Lou, C., Stanton, B., Chen, Y.-J., Munsky, B., and Voigt, C. A. (2012) Ribozyme-based
1508 insulator parts buffer synthetic circuits from genetic context. *Nat. Biotechnol.* 30, 1137–
1509 1142.

1510 (209) Davis, J. H., Rubin, A. J., and Sauer, R. T. (2011) Design, construction and
1511 characterization of a set of insulated bacterial promoters. *Nucleic Acids Res.* 39, 1131–
1512 1141.

1513 (210) Qi, L., Haurwitz, R. E., Shao, W., Doudna, J. A., and Arkin, A. P. (2012) RNA
1514 processing enables predictable programming of gene expression. *Nat. Biotechnol.* 30,
1515 1002–1006.

1516 (211) Nielsen, A. A. K., Segall-Shapiro, T. H., and Voigt, C. A. (2013) Advances in genetic
1517 circuit design: Novel biochemistries, deep part mining, and precision gene expression.
1518 *Curr. Opin. Chem. Biol.* 17, 878–892.

1519 (212) Mutalik, V. K., Guimaraes, J. C., Cambray, G., Lam, C., Christoffersen, M. J., Mai, Q.-A.,
1520 Tran, A. B., Paull, M., Keasling, J. D., Arkin, A. P., and Endy, D. (2013) Precise and reliable
1521 gene expression via standard transcription and translation initiation elements. *Nat.*
1522 *Methods* 10, 354–360.

1523 (213) Bardhan, S. K., Gupta, S., Gorman, M. E., and Haider, M. A. (2015) Biorenewable
1524 chemicals: Feedstocks, technologies and the conflict with food production. *Renew.*
1525 *Sustain. Energy Rev.* 51, 506–520.

1526 (214) Papoutsakis, E. T. (2008) Engineering solventogenic clostridia. *Curr. Opin.*
1527 *Biotechnol.* 19, 420–9.

1528 (215) Lorenz, R., Bernhart, S. S. H., Höner zu Siederdisen, C., Tafer, H., Flamm, C.,

1529 Stadler, P. F. P., and Hofacker, I. I. L. (2011) ViennaRNA Package 2.0. *Algorithms Mol. Biol.*
1530 6, 26.

1531 (216) Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R., and Hofacker, I. L. (2008) The
1532 Vienna RNA websuite. *Nucleic Acids Res.* 36, W70-4.

1533 (217) Darty, K., Denise, A., and Ponty, Y. (2009) VARNA: Interactive drawing and editing
1534 of the RNA secondary structure. *Bioinformatics* 25, 1974–1975.

1535 (218) Wilson, K. S., and Hippel, P. H. Von. (1995) Transcription termination at intrinsic
1536 terminators: The role of the RNA hairpin (*Escherichia coli*/RNA polymerase/rho-
1537 independent termination). *Biochemistry* 92, 8793–8797.

1538