

This is a repository copy of *Variation in the species richness of parasitoid wasps (Ichneumonidae: Pimplinae and Rhyssinae) across sites on different continents.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/125769/>

Version: Accepted Version

Article:

Gomez, Israel, Saaksjarvi, Ilari, Mayhew, Peter John orcid.org/0000-0002-7346-6560 et al. (6 more authors) (2018) Variation in the species richness of parasitoid wasps (Ichneumonidae: Pimplinae and Rhyssinae) across sites on different continents. *Insect conservation and diversity*. ISSN 1752-458X

<https://doi.org/10.1111/icad.12281>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



**Variation in the species richness of parasitoid wasps
(Ichneumonidae: Pimplinae and Rhyssinae) across sites on
different continents**

Journal:	<i>Insect Conservation and Diversity</i>
Manuscript ID	ICDIV-16-0190.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	23-Oct-2017
Complete List of Authors:	Gomez, Israel; Biodiversity and Environmental Sciences Saaksjarvi, Ilari; University of Turku, Zoological Museum, Department of Biology Mayhew, Peter; University of York, Department of Biology Pollet, Marc; INBO, Research Group Species Diversity Rey Del Castillo, Carmen Nieves-Aldrey, José; Museo Nacional de Ciencias Naturales, Departamento de Biodiversidad y Biología Evolutiva Broad, Gavin; The Natural History Museum, Department of Entomology Roininen, Heikki Tuomisto, Hanna; Turun Yliopisto
Keywords:	Amazonia, Biodiversity, Biogeography, Peru, Rhyssinae, Rain forest, Pimplinae, Latitudinal gradient

Variation in the species richness of parasitoid wasps (Ichneumonidae: Pimplinae and Rhyssinae) across sites on different continents

Gómez, Israel C.^{1,8}, Sääksjärvi, Ilari E.¹, Mayhew, Peter J.², Pollet, Marc³, Rey del Castillo, Carmen⁴, Nieves-Aldrey, Jose-Luis⁴, Broad, Gavin R.⁵, Roininen Heikki⁶ & Tuomisto, Hanna⁷

¹Zoological Museum, Biodiversity Unit, FI-20014 University of Turku, Finland.

²Department of Biology, University of York, Heslington, York YO10 5DD, UK

³Department of Entomology, Royal Belgian Institute of Natural Sciences (RBINS), B-1000 Brussels; Research Group Terrestrial Ecology (TEREC), University of Ghent (UGent), B-9000 Ghent; and Research Group Species Diversity (SPECDIV), Research Institute for Nature and Forest (INBO), B-1070 Brussels, all Belgium.

⁴Departamento de Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales (CSIC) Madrid, Spain.

⁵Department of Life Sciences, the Natural History Museum, Cromwell Road, London SW7 5BD, UK.

⁶Department of Biology, University of Eastern Finland. FI-80101 Joensuu, Finland.

⁷Department of Biology, FI-20014 University of Turku, Finland.

⁸Corresponding author: Gómez, Israel C. (israel.gomez@yahoo.com)

Phone+358 2333 5447

Running title: Species-richness patterns in parasitoid wasps

ABSTRACT

1 The old idea that parasitoid wasps (Ichneumonidae) show an inverse latitudinal diversity gradient has recently been challenged, but how ichneumonid species richness varies across the globe is still not well understood. We carried out field inventories in 21 sites on three continents to clarify this question, focusing on the subfamilies Pimplinae and Rhyssinae. Our total sampling effort was 628 Malaise trap months and the total catch exceeded 65,000 individuals. Our main focus was in two intensively inventoried areas in Amazonia, together yielding 257 Malaise trap months and 26,390 ichneumonid individuals.

2 To expand the scope and assess global species diversity patterns of the Pimplinae and Rhyssinae, we compiled published species lists from a total of 97 study localities around the world. The highest observed species richness in any locality, 105 species, was found in one of our field sites in Peruvian Amazonia. None of the other localities reported more than 70 species, even the ones with a sampling effort comparable to ours.

3 Despite the local thoroughness of our field inventories in Amazonia, data analyses indicated that a substantial proportion of the parasitoid wasp species occurring in each site remained unobserved.

4 The highest local species richness values were reported from the tropics. However, parasitoid wasps are still too sparsely sampled to draw solid conclusions about whether or not their species richness follows a particular latitudinal trend, and if so, where their richness peaks.

Key words. Amazonia, biodiversity, biogeography, latitudinal gradient, Peru, Pimplinae, rain forest, Rhyssinae.

INTRODUCTION

The parasitoid wasp family Ichneumonidae is the most species-rich hymenopteran family, and it is among the most diverse animal groups on Earth. Interestingly, it has been suggested that the Ichneumonidae is less species-rich in tropical regions than at higher latitudes, i.e. that the family presents an inverse latitudinal diversity gradient (e.g. Owen & Owen, 1974; Timms *et al.* 2015). Over the years, various hypotheses have been proposed to explain why ichneumonids are relatively species-poor in the tropics (Santos & Quicke, 2011). However, the high number of newly described taxa in recent taxonomic publications shows that tropical ichneumonids are still incompletely known (e.g. Aguiar & Ramos, 2011; Broad *et al.*, 2011; Santos & Aguiar, 2013). Therefore, further large-scale and long-term studies are needed, especially in the tropics, to improve our knowledge about global ichneumonid diversity patterns.

Amazonia is the largest rainforest area in the world, but also one of the least known in terms of its insect fauna. Although western Amazonia is famous for its high diversity in several animal and plant groups (Gentry, 1988, 1992; Robins, 1996; Robbins & Opler, 1997; Patterson *et al.*, 2006), some early authors proposed that ichneumonid diversity is low in Amazonia (Porter, 1978). These studies were based on small data sets, and the situation has changed in the 2000s, as new studies on Amazonian ichneumonid diversity have challenged the old paradigm (Sääksjärvi *et al.*, 2004; Gómez *et al.*, 2015; Veijalainen *et al.*, 2013).

In this study, we compare observations of local species richness of pimelines and rhyssines across latitudes on the basis of field sampling we carried out in 21 sites on three continents.

The two most intensively studied areas were in Peruvian Amazonia, where the sampled localities were selected so as to span different forest types (Sääksjärvi *et al.*, 2004; Gómez *et*

1
2
3 *al.*, 2014, 2015). Most sampling was done with Malaise traps, but in the two Peruvian areas,
4
5 we also used other sampling methods for comparison.
6
7

8 The main aims of this study are 1) to document how species accumulation of pimpline and
9
10 rhyssine wasps relates to sampling effort in our two well-sampled Amazonian sites, and 2) to
11
12 assess global patterns in pimpline and rhyssine species richness using data from both our own
13
14 inventories and from publications of other researchers.
15
16
17
18
19
20
21
22

23 MATERIAL AND METHODS

24
25
26 STUDY ORGANISMS.—The parasitoid wasp family Ichneumonidae currently includes
27
28 about 40 subfamilies, 1,600 genera and 24,000 described species (Aguiar *et al.*, 2013),
29
30 making it the most species-rich family within Hymenoptera. Here we focus on two of the
31
32 most extensively studied subfamilies in the Neotropical region, Pimplinae and Rhyssinae
33
34 (e.g. Porter, 1978; Gauld, 1991; Gauld *et al.*, 1998, 2002; Sääksjärvi *et al.*, 2004; Gómez *et*
35
36 *al.*, 2009, 2014, 2015; Gómez & Yabar, 2015).
37
38

39 The subfamily Pimplinae is a relatively large, cosmopolitan group with about 78 genera
40
41 (Pham, 2013) and more than 1,500 described species worldwide (Yu *et al.*, 2012). This is the
42
43 most biologically diverse of all ichneumonid subfamilies and it is associated with a wide
44
45 range of hosts (Gauld, 1991). The species are mainly idiobiont ectoparasitoids of immature
46
47 stages of holometabolan insects or idiobiont endoparasitoids of lepidopteran or hymenopteran
48
49 pupae. However, the species of the *Polysphincta* genus-group are koinobiont ectoparasitoids
50
51 of immature and mature spiders (Gauld & Dubois, 2006).
52
53
54
55
56
57
58
59
60

1
2
3 The subfamily Rhyssinae is a relatively small, cosmopolitan group with eight genera
4
5 worldwide and more than 250 described species (Gómez *et al.*, 2015). According to Kamath
6
7 & Gupta (1972), the species-richness of Rhyssinae is highest in the lowland rainforests of
8
9 South-East Asia. In the Neotropical region the species are believed, on the basis of the known
10
11 biology of a few Old World species, to be idiobiont ectoparasitoids of immature stages of
12
13 endopterygote insects boring in wood (Gauld, 1991).
14
15

16
17
18
19
20 STUDY SITES.— We carried out intensive long-term sampling in two sites in Peruvian
21
22 Amazonia, Allpahuayo-Mishana in the north and Los Amigos in the south. In addition, we
23
24 sampled 19 sites outside Amazonia. The distribution of these sites is shown in Fig. 1.
25
26 Elevation, geographical coordinates and sampling effort of each locality are given in
27
28 Supplementary materials (Table S1).
29
30

31
32 Allpahuayo-Mishana National Reserve (3° 57-58' S, 73° 25-26' W) is located in northern
33
34 Peruvian Amazonia in the department of Loreto. The climate is tropical, with mean annual
35
36 precipitation of about 3000 mm and mean annual temperature of 26 °C (Marengo, 1998).
37
38 Elevation is between 100 and 180 m above sea level and the prevailing vegetation is moist
39
40 tropical forest according to the Holdridge (1967) classification. The environmental conditions
41
42 have been described in Kalliola & Flores Paitán (1998). Our ichneumonid sampling was
43
44 carried out in non-inundated areas (*terra firme*), where the terrain is divided among three
45
46 main soil types. Flat low-lying areas often have soils derived from the marine Pebas
47
48 formation of Miocene age. These soils are clayey in texture, relatively rich in nutrients, and
49
50 mostly brownish-grey in colour (Kalliola & Flores Paitan, 1998). The tops of the plateau-like
51
52 hills often have white sand soils, which are nutrient-poor and white or grey in colour. Loamy
53
54 soils are probably derived from old river terraces (Kalliola & Flores Paitan, 1998). The
55
56
57
58
59
60

1
2
3 forests growing on the clayey to loamy soils have a classical rain forest physiognomy with
4
5 large trees and different canopy layers, whereas the white sand forests have slenderer trees
6
7 and a less complex canopy structure (Anderson, 1981; Tuomisto & Ruokolainen, 1994).
8
9

10 Los Amigos Conservation Concession (12° 33-34' S, 70° 04-05' W) is located in the
11
12 southern Peruvian Amazonia in the department of Madre de Dios close to the Andean-
13
14 Amazonian interface. Mean annual precipitation is about 2 770 mm and annual mean
15
16 temperature 23 °C with monthly mean temperatures varying 20-26 °C (Gómez, 2009). This is
17
18 somewhat drier, cooler and more seasonal than in Allpahuayo-Mishana. Elevation varies
19
20 between 230 and 270 m above sea level and the vegetation is moist tropical forest according
21
22 to the Holdridge (1967) classification. For description of the environmental conditions, see
23
24 Wilson & Sandoval (1996). The Los Amigos watershed comprises sediments dated from the
25
26 Neogene and Quaternary that have been deposited during the last 20 million years (Pino,
27
28 2007). Due to the proximity of the Andes, the soils are a mosaic of clayey, sandy, silty and
29
30 gravelly materials, but extremely nutrient-poor soils such as white sands have not been
31
32 reported for the area. Our ichneumonid sampling covered both non-inundated and seasonally
33
34 inundated forests. In general, *terra firme* soils are sandier, more acidic and poorer in nutrients
35
36 than those located in the floodplains (Hovikoski *et al.*, 2005).
37
38
39
40
41

42 Seven of our other 19 sampling localities are located in temperate South America between the
43
44 regions of Maule (35°S) and Los Lagos (42°S) in Central Chile. One sampling locality is in
45
46 Central America on Coiba island off the Pacific coast of Panama (7°N). This is a national
47
48 park and the vegetation is moist tropical forest according to Holdridge's classification.
49
50

51 Eight sampling localities were established in the temperate region of Spain, between 38°N
52
53 and 42°N; for details see Nieves-Aldrey (1995).
54
55
56
57
58
59
60

1
2
3 Three localities are situated in the Canary Islands in Isla de la Palma (28°N) and were mostly
4 dominated by forests of the endemic Canarian pine (*Pinus canariensis* Chr. Sm. ex DC; for
5 survey details see Domingo Quero *et al.*, 2003). One sampling locality is in central Africa,
6 the Kibale National Park in Uganda.
7
8
9
10

11
12
13
14
15 SAMPLING METHODS.—Our primary sampling method consisted of long-term Malaise
16 trapping. A Malaise trap is an open tent-like structure widely used in entomological studies,
17 and ichneumonid surveys in particular, because it is efficient in collecting insects that are
18 strong flyers (Nieves-Aldrey & Rey del Castillo, 1991, Sääksjärvi *et al.*, 2004; Mayhew *et*
19 *al.*, 2009). Thanks to the popularity of Malaise traps, it was possible to find comparable data
20 from earlier parasitoid studies, as the sampling effort can be conveniently expressed in terms
21 of Malaise trap months (MTM). One MTM corresponds to one trap collecting in the field for
22 a period of one month (or a combination of more traps that cumulatively operate for one
23 month).
24
25
26
27
28
29
30
31
32
33
34
35

36 In the Peruvian Allpahuayo-Mishana, we combined data from three separate field campaigns.
37 The first one ran from August 1998 to January 1999 with 12 traps and had a total sampling
38 effort of 41 MTM. The second ran from January 2000 to January 2001 with 15 traps and had
39 a total sampling effort of 144 MTM. Both of these campaigns have been described by
40 Sääksjärvi *et al.* (2004). The last campaign was conducted by 14 Malaise traps, four of which
41 were active in April-December 2011 and ten in October-December 2011. The total sampling
42 effort of these traps was 45 MTM, and details have been described by Gómez *et al.* (2015).
43
44
45
46
47
48
49
50
51 In Los Amigos, nine Malaise traps were in operation in May-July 2008 with a total sampling
52 effort of 27 MTM. The total sampling effort in Allpahuayo-Mishana and Los Amigos
53
54
55
56
57
58
59
60

1
2
3 combined was 257 MTM. To our knowledge, this is the most extensive sampling of
4
5 ichneumonids in Amazonia.
6
7

8
9
10 To find out if Malaise trapping might be systematically biased and miss some species
11
12 entirely, we applied four additional sampling methods in Allpahuayo-Mishana and one in Los
13
14 Amigos (Table 1) 1) Yellow pan traps (Moericke traps). During 23 weeks, a total of 100 traps
15
16 were set out in Allpahuayo-Mishana for a period of 24 hours every week. In each weekly
17
18 sampling, the traps were spread across the different forest types. 2) Hand netting with a
19
20 standard entomological net. This was done weekly for 13 weeks in Los Amigos and for 23
21
22 weeks in Allpahuayo-Mishana. Each netting session consisted of four hours of random
23
24 sweeping in the vegetation of different forest types. 3) Larval rearing. A total of 3,298
25
26 lepidopteran larvae were collected in the forests of Allpahuayo-Mishana and reared
27
28 individually in one-litre plastic jars. 4) Pupal rearing. A total of 113 lepidopteran pupae were
29
30 collected in the forests of Allpahuayo-Mishana and reared individually in one-litre plastic
31
32 jars.
33
34
35
36

37
38 The number of localities and sampling effort outside Amazonia is as follows: seven in
39
40 Central Chile (45.4 MTM), one in Panama (1.4 MTM), seven in Spain (57.4 MTM), three on
41
42 the Canary Islands (4.8 MTM) and one in Uganda (252 MTM). Samples from all localities
43
44 were prepared and identified in the Zoological Museum of the University of Turku (ZMUT)
45
46 and the National Museum of Natural Sciences, Madrid (CSIC).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

GLOBAL COMPARISON OF PIMPLINAE AND RHYSSINAE SPECIES-RICHNESS.—

To evaluate global patterns in Pimplinae and Rhyssinae species richness we searched the literature for ichneumonid inventories that could be compared with our field data. Because the number of observed species is dependent on sampling effort, we only considered quantitative inventories that had used Malaise traps and in addition recorded the duration of sampling and/or the number of captured Pimplinae and Rhyssinae individuals. We found 76 studies fulfilling these criteria. In most cases, relevant information for our analysis was not available in the original publication but was kindly provided by the authors of the studies (see Table S1).

All the inventories were carried out in delimited sites with little altitudinal variation. In most cases, the variation within a locality was less than 200 meters. It was important to ensure that all data points represented single localities; if some had included longer altitudinal gradients than others, drawing conclusions about latitudinal patterns in local species richness would not be appropriate. Some published inventories only included data on the subfamily Pimplinae. Because this subfamily is much more abundant and species-rich than Rhyssinae (Gauld, 1991) we used these data as well but identify them in the results.

DATA ANALYSIS.—Our first aim was to document species accumulation patterns as a function of sampling effort in the Peruvian sites, for which time series data were available. Each sampling interval (originally 1 or 2 weeks) from each trap was used as a separate sampling unit. Species accumulation curves were obtained in two different ways. Firstly, the actual observed species accumulation was documented by aggregating consecutive sampling intervals from the beginning of a trap's operation to its end. Secondly, sampling units were combined in a random order, within habitat types only, to observe patterns of species accumulation at the site level.

1
2
3 To take into account species abundance patterns, we calculated the effective number of
4 species, also known as true diversity. This equals the reciprocal of the weighted mean of the
5 proportional species abundances, with the species abundances themselves used as the weights
6 and the parameter q defining which mean is used (Tuomisto, 2010). We used two different
7 means, the geometric mean (corresponding to $q=1$) and the arithmetic mean ($q=2$). The
8 former gives a diversity value that equals the exponential of the well-known Shannon index,
9 and the latter gives a value that equals the inverse Simpson index.
10
11

12 We used the Chao-1 estimator to assess how many additional species could be expected at a
13 site, given the number of species observed only once or twice with a given sampling effort.
14 The estimated number of additional species was then compared with the corresponding
15 sampling effort to check for any indication that species sampling was approaching
16 completeness.
17
18

19 Division of the Peruvian data to subsets of different sampling efforts made it possible to
20 compare richness values from elsewhere in the world with Peruvian data of similar sampling
21 effort. Here we focus on the effect of sampling effort only, and do not take into account
22 possible effects of seasonality. The effect of habitat heterogeneity was minimized by only
23 pooling samples taken in the same forest type. This was done to ensure that the Peruvian
24 species richness values are not inflated in relation to the other sites, for which information on
25 possible habitat heterogeneity was not available. Species accumulation was plotted against
26 both sampling effort in the strict sense (the length of the sampling interval in MTM) and
27 sampling efficiency (the number of Pimplinae and Rhyssinae individuals captured). In
28 addition, we analysed latitudinal and elevational trends in local species richness after taking
29 into account the effect of sampling efficiency. This was done by first fitting a second-order
30 polynomial regression of number of species against number of individuals (both
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 logarithmically transformed), and then using the residuals from this regression as the
4
5 response variable in linear regression against latitude or elevation. All analyses were done in
6
7 the R environment using either the 'vegan' package (Oksanen *et al.*, 2013) or code written by
8
9 HT.
10

11 12 13 14 15 16 17 RESULTS

18
19
20
21
22 The 21 localities we sampled in different parts of the world produced a total catch of
23
24 approximately 65,000 ichneumonid individuals, of which 2,698 individuals representing 293
25
26 species were Pimplinae or Rhyssinae. In Allpahuayo-Mishana, all sampling methods together
27
28 yielded a total of 24,378 ichneumonid individuals representing 19 subfamilies. The subfamily
29
30 Pimplinae was represented by 1,516 individuals (92 species) and the subfamily Rhyssinae by
31
32 81 individuals (13 species; Table S2). In Los Amigos, a total of 2,012 ichneumonid
33
34 individuals representing 18 subfamilies were captured. The subfamily Pimplinae was
35
36 represented by 106 individuals (42 species) and the subfamily Rhyssinae by 15 individuals
37
38 (nine species; Table S2).
39
40
41

42
43 The two Peruvian sites had 16 genera of Pimplinae, 9 (56%) of which were shared, and a
44
45 single shared genus of Rhyssinae (Table S2). The species compositions of the two sites were
46
47 rather different: of the 128 Pimplinae and Rhyssinae species observed, 81 species (63%) were
48
49 only collected in Allpahuayo-Mishana, 23 species (18%) only in Los Amigos, and only 24
50
51 species (19%) were shared between both localities. In other words, the overall similarity of
52
53 the species compositions in the two sites equaled 0.19 if measured with the Jaccard index and
54
55 0.32 if measured with the Sørensen index. It is noteworthy that half of the species that were
56
57
58
59
60

1
2
3 observed in Los Amigos were not observed in Allpahuayo-Mishana in spite of the massive
4
5 sampling effort (230 MTM) in the latter.
6
7

8 In both Peruvian sites, the Malaise trap proved to be the most efficient collecting technique,
9
10 as all the Pimplinae and Rhyssinae species encountered in the inventory were found in
11
12 Malaise traps. The other methods only yielded additional individuals of some of the species.
13
14

15 In the 2011 inventory in Allpahuayo-Mishana, Malaise trapping captured 1,961 ichneumonid
16
17 individuals, which was 74% of the total captured with all methods together in that year (2,640
18
19 individuals). Yellow pan trapping yielded 511 ichneumonid individuals (19% of the total)
20
21 and hand netting produced a catch of 142 individuals (5% of the total). Larval rearing
22
23 produced 22 individuals and pupal rearing only four individuals. Given that 3,411
24
25 lepidopteran larvae and pupae were reared, this indicates that only a small proportion of them
26
27 were parasitized by ichneumonids.
28
29
30
31

32 Overall, Cryptinae was the most abundant subfamily with 1,168 individuals (44% of the
33
34 total), whereas Pimplinae with 205 individuals and Rhyssinae with 19 individuals represented
35
36 smaller proportions of the total catch (5% and 0.7%, respectively). The individual Malaise
37
38 traps differed greatly in how many individuals and species they captured per unit time, but
39
40 there did not seem to be any systematic trend in this either among habitat types locally or
41
42 between the two Peruvian sites (Fig. 2A, B). When species accumulation was related to the
43
44 number of individuals captured, much more uniform trends were observed across the traps.
45
46 There was only a modest tendency for species accumulation to slow down with increasing
47
48 sampling efficiency (Fig. 2C).
49
50
51

52 The Chao-1 estimates indicated that even at the largest sample sizes within habitat types,
53
54 many species still remained unobserved (Fig. 3). The estimated number of unseen species
55
56
57
58
59
60

1
2
3 was only weakly related to the number of individuals already observed, although a decreasing
4
5 tendency emerged when the sample size became larger than about 200 individuals. A linearly
6
7 decreasing trend was found if the estimated number of unseen species was expressed as a
8
9 percentage of the observed number of species (3B). Nevertheless, even at the largest sample
10
11 sizes available to us, it was still estimated that the number of species would increase by 10-
12
13 30% if more individuals were sampled.

14
15
16
17 The global sampling effort for Ichneumonidae that is both known to us and reports adequate
18
19 data for comparison (Table S1) totals 3,267 MTM. Costa Rica is the best-studied country
20
21 with data available from 17 localities with a combined sampling effort of 1,266 MTM (39 %
22
23 of the world total). Our Peruvian study contributed 19 % of the world total (628 MTM). In
24
25 general, most of the study sites were at relatively low altitudes with only 10 of the 97
26
27 localities (about 10 % of total MTM) occurring above 2000 m elevation.

28
29
30
31 Our Malaise traps in Peruvian Amazonia were about as efficient in capturing Pimplinae and
32
33 Rhyssinae individuals as the traps for which results have been reported in the literature: the
34
35 number of individuals caught with any given number of Malaise traps months in our data was
36
37 firmly within the limits reported from other sites with a similar number of Malaise trap
38
39 months (Fig. 4A). In contrast, the number of species caught in the Peruvian traps was
40
41 generally higher than the numbers of species reported for the same number of Malaise trap
42
43 months elsewhere (Fig. 4B). This difference became even clearer when the number of species
44
45 was shown against number of individuals: for any given number of individuals captured, the
46
47 number of species observed was always highest in the Peruvian samples (Fig. 5A).

48
49
50
51 All the sites with more than 50 observed species were in the tropics, and the upper limit of
52
53 local species richness clearly decreased towards the poles. However, sites with very few
54
55 observed species (even taking into account sampling effort) were found at all latitudes (Fig.
56
57
58
59
60

1
2
3 6). Therefore, no consistent overall gradient in species richness could be seen in relation to
4
5 latitude. With elevation, the relationship was even weaker.
6
7

8
9 When some species are more abundant than others, the effective number of species (=species
10 diversity) becomes smaller than the actually observed number of species. This effect was
11 clearly visible in our Peruvian data (Fig. 5). Differences between the habitat types in species
12 accumulation were small, but traps in white sand forests tended to yield fewer species for a
13 given number of individuals than traps in the other habitats. Interestingly, the traps in
14 secondary forests tended to have relatively high species richness, but these samples were
15 dominated by a few species, which resulted in clearly decreasing diversity with increasing
16 value of q . The species diversity accumulation curves also showed a distinct tendency to level
17 off at a much earlier stage than the species-richness accumulation curve did.
18
19
20
21
22
23
24
25
26
27

28
29 The general impression from plotting observed species richness against latitude was that the
30 highest numbers of species are found in the tropics (Fig. 6A). This impression was confirmed
31 for the lowland sites when the effect of collecting efficiency (number of individuals) was
32 taken into account, but no such trend was seen among the montane sites (Fig. 6B). Within the
33 tropical sites, species richness seemed to decrease with elevation, but no trend was observed
34 in the non-tropical sites (Fig. 6C).
35
36
37
38
39
40
41
42
43
44

45 **DISCUSSION**

46
47
48 PARASITOID WASP SPECIES-RICHNESS.—Allpahuayo-Mishana is currently the site
49 with the highest known species-richness for pimpline and rhyssine wasps in the world, with
50 105 species. Our current estimate is that 27 of these (25.8%) represent already described
51 species, whereas 78 species (74.2%) are new to science.
52
53
54
55
56
57
58
59
60

1
2
3 The total sampling effort in Allpahuayo-Mishana was 240 MTM, which is high in global
4 comparison (Table S1). However, the most intensively sampled locality in the world known
5 to us is Santa Rosa in Costa Rica, where researchers have observed 40 species with a
6 sampling effort of 576 MTM (Gaston & Gauld, 1993). Only four other sites can boast a
7 sampling effort exceeding 100 MTM for Pimplinae and Rhyssinae, and none of these has
8 reported more than 26 species (Table S1): Cerro Hacha in Costa Rica, Ria Lagartos in
9 Mexico, Cabañeros in Spain and Kibale in Uganda. The two Peruvian localities show
10 consistently higher numbers of Pimplinae and Rhyssinae species than most other sites when
11 comparisons are made among subsamples based on an equivalent sampling effort (MTM;
12 Fig. 4B). The difference becomes even more evident when comparisons are based on
13 sampling efficiency (number of individuals; Fig. 5A).

14
15
16 Peruvian Amazonia is known to have high habitat heterogeneity (Tuomisto *et al.*, 1995), and
17 it has very high species-richness of trees (Gentry 1988, 1992), butterflies (Robins 1996,
18 Robbins & Opler, 1997) and birds (Patterson *et al.*, 2006).

19
20
21 Pimplinae and Rhyssinae, and idiobiont ichneumonids in general, have already been
22 documented to be species-rich groups in the tropics (Gauld, 1991; Sime & Brower 1998;
23 Quicke, 2012). These subfamilies are among the best-known groups of parasitoids in the
24 Neotropics, but our results indicate that their species richness in Amazonia is even higher
25 than previously reported by Sääksjärvi *et al.* (2004).

26
27
28 Recent findings have demonstrated that the subfamily composition of idiobiont and
29 koinobiont ichneumonids is rather similar in temperate and tropical regions (Veijalainen *et*
30 *al.*, 2013). Furthermore, also high numbers of koinobiont species have been discovered
31 recently from the tropics (Broad *et al.*, 2011; Khalaim & Broad, 2013). These findings
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 together with our results suggest that species richness of both idiobiont and koinobiont
4
5 ichneumonids is still mainly undiscovered and most likely very high in the tropics.
6
7

8 The functional composition of the pimpline and rhyssine faunas in our Peruvian Amazonian
9
10 study localities (Table S2) is rather similar to that in other tropical study localities (e.g. those
11
12 mentioned by Gauld, 1991). Idiobiont parasitoids are clearly dominating, with the genus
13
14 *Neotheronia* being especially abundant in the Neotropics.
15
16

17
18
19
20
21 LATITUDINAL SPECIES-RICHNESS GRADIENT.—Our results lend no support to the
22
23 anomalous latitudinal species-richness gradient that was originally proposed by Owen &
24
25 Owen (1974), at least for the subfamilies Pimplinae and Rhyssinae. We found a lot of
26
27 variation in species richness within latitudes but variable sampling effort makes the results
28
29 difficult to interpret. If anything, the observed maxima of local species richness follow the
30
31 usual species-richness gradient, i.e. the observed numbers of species peak in the tropics and
32
33 decrease towards the poles. Owen & Owen (1974) examined samples collected in Northern
34
35 Europe and in the Afrotropical region. They observed that “a quite extraordinary number of
36
37 species was found once” in their samples, which suggests that the parasitoid wasp
38
39 communities were only superficially sampled. Janzen & Pond (1975) and Janzen *et al.* (1976)
40
41 agreed with the phenomenon of reverse latitudinal gradient but did not demonstrate that their
42
43 sampling efforts were representative enough to support such broad generalizations. As a
44
45 result, the phenomenon was popularized although it was based on data obtained from only a
46
47 handful of study localities that were not even distributed along a proper latitudinal gradient.
48
49
50
51

52
53 In our northern Peruvian site, two years of sampling with several Malaise traps was
54
55 insufficient to obtain a complete representation of the species in any one locality. Such an
56
57
58
59
60

1
2
3 outcome is expected when sampling hyperdiverse organisms (Colwell & Coddington, 1994).
4
5 Together with the results from all the parasitoid studies analysed here, this leads us to agree
6
7 with the proposal that the main reason why low latitudes have been thought to be relatively
8
9 species-poor for ichneumonids is insufficient sampling effort (e. g. Hespeneide, 1978;
10
11 Morrison *et al.*, 1979; Sime & Brower 1998; Sääksjärvi *et al.*, 2004; Santos & Quicke, 2011).
12
13

14 LOCAL SPECIES COMPOSITION IN PERUVIAN AMAZONIA.—The Malaise traps
15
16 differed considerably in how many species they captured (Fig. 2). This was not related to
17
18 habitat type in any obvious way, although in Allpahuayo-Mishana there was a weak tendency
19
20 for traps in white sand forests to capture fewer species per unit time than traps in the other
21
22 kinds of forest did (Fig. 4B). This would agree with earlier reports that white sand forests
23
24 have fewer plant and animal species than other non-inundated forest types do (Anderson,
25
26 1981; Tuomisto & Ruokolainen, 1994). This, together with the simpler forest structure, may
27
28 lead to this habitat type providing fewer niches for parasitoids. The relatively open canopy in
29
30 white sand forests also makes them rather sunny and possibly dry, which is thought to be
31
32 unfavourable for ichneumonids (Gauld, 1991). However, this cannot be the only explanation
33
34 since secondary forest had higher species-richness than the other habitats. This parallels
35
36 results from Brazilian Amazonia, where moth species richness in secondary forest was
37
38 consistently higher than in primary forest (Barlow *et al.*, 2007). The degree of dominance
39
40 was also high in secondary forests: diversity (which takes species abundances into account)
41
42 was clearly lower in secondary forests than in old-growth forests on loamy soils (Fig 5B),
43
44 although there appeared to be no obvious difference in species richness (Fig 5A).
45
46
47
48
49

50 It is noteworthy that half of the Pimplinae and Rhyssinae species that were observed in Los
51
52 Amigos were not found in Allpahuayo-Mishana, and less than 20% of the observed species
53
54 were shared between the two sites. This may partly be due to undersampling, as the species
55
56
57
58
59
60

1
2
3 accumulation curves indicate that part of the species pools remained unsampled in both areas.
4
5 However, it is likely that there is also a real difference between their ichneumonid faunas,
6
7 which could be due to at least two causes. Firstly, the two sites are more than 1000 km apart,
8
9 and there might already be biogeographical differences between their regional species pools.
10
11 Secondly, the habitat types sampled in the two sites were different: sampling in Los Amigos
12
13 was mostly in floodplain forests, whereas sampling in Allpahuayo-Mishana was in non-
14
15 inundated forests. Therefore, any specialists of floodplain forests would have been missed in
16
17 Allpahuayo-Mishana, even though they might exist in the general area. Conversely,
18
19 specialists of white sand forests could not have been observed in Los Amigos, as such forests
20
21 are not known to occur in the area. Such possibilities open up interesting questions about the
22
23 ecology and biogeography of ichneumonid wasps.
24
25
26
27
28
29
30

31 SAMPLING EFFORT.—There was considerable variation among individual Malaise traps in
32
33 how many specimens and species they captured per unit time. Indeed, it is obvious that the
34
35 precise placement of a Malaise trap may have a considerable impact on how many insects it
36
37 captures. On the other hand, the number of individuals captured was a good predictor of the
38
39 number of species observed (Fig. 5A). This suggests that when the interest is in comparing
40
41 species-richness or diversity values among sites, it is more important to ensure comparable
42
43 sampling efficiency (number of individuals captured) than comparable sampling effort (time
44
45 spent sampling).
46
47
48

49 The total sampling effort in all our field localities (the red dots in Fig. 1) was 628 MTM, of
50
51 which 257 MTM was focused on Peruvian Amazonia. The species accumulation curves for
52
53 the different habitats showed that our sampling was nowhere near complete (Fig. 3–4;
54
55 Colwell & Coddington, 1994). This was true even for the most intensively studied habitat
56
57
58
59
60

1
2
3 type, namely white-sand forest, which was sampled by 93 MTM in Allpahuayo-Mishana.
4
5 Indeed, species-richness was still expected to increase by 10-30% if sampling were continued
6
7 (Fig. 4). This parallels earlier results from Costa Rica, where only 32 Pimplinae and
8
9 Rhyssinae species were known before the study of Gauld (1991). After extensive sampling
10
11 programs this number increased to a total of 195 species (Gauld *et al.*, 1998). Likewise,
12
13 sampling conducted over a decade increased the number of Pimplinae species known from
14
15 Vietnam from 39 to 122 (Pham 2013). In both of these tropical countries, it took several years
16
17 of sampling to produce a significant increase in the species richness. In Peru, a total of 128
18
19 Pimplinae and Rhyssinae species were recorded here from the two Amazonian localities, but
20
21 more can be expected.
22
23
24

25
26 Most studies on parasitoid wasps have been carried out in the northern hemisphere. This has
27
28 left the southern hemisphere poorly known, including the recognised diversity hotspots there
29
30 (Orme *et al.*, 2005). This situation may also be one reason for the fact that the known
31
32 ichneumonid genera are heavily concentrated in the northern hemisphere (Quicke, 2012). In
33
34 addition, most studies have focused on lowlands (Table S1), so information from mid and
35
36 high altitude areas is sparse. Both of these biases in sampling need to be considered when
37
38 studying the distributional patterns of ichneumonids.
39
40
41
42

43
44 Morrison *et al.*, (1979) lamented that the sampling effort was insufficient in tropical and
45
46 temperate regions to provide a clear understanding of species-richness patterns of these
47
48 parasitoids, and we suggest that the situation has remained similar to the present day. Species
49
50 accumulation curves from our intensively sampled Peruvian sites suggest that most of the
51
52 study localities for which we have found published data (Table S1) are still too superficially
53
54 sampled to allow a reliable estimation of local ichneumonid species richness to be made.
55
56
57
58
59
60

1
2
3 Although the two Peruvian Amazonian localities seem to be the most species-rich sites in the
4 world for the subfamilies Pimplinae and Rhyssinae, this may just be a consequence of the
5 extensive sampling effort in them. The situation may well change when other sites become
6 more intensively sampled. However, in the light of the currently available data, it is plausible
7 that the species richness of Ichneumonidae (or at least Pimplinae and Rhyssinae) follows the
8 usual latitudinal trend of highest local diversity in the lowland tropics.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

ACKNOWLEDGEMENTS

The Ministry of Agriculture and the Ministry of Environment of Peru provided the collecting and export permits for the Peruvian samples. We appreciate the support of the Instituto de Investigaciones de la Amazonía Peruana (IIAP) for the field studies made in Allpahuayo-Mishana. The colleagues and the friends of the “Centro de Investigaciones Allpahuayo and Los Amigos (CICRA)” station helped considerably during the field work. Olavi Kurina, Carol Castillo and Tapani Hopkins helped with the preparation of the Afrotropical material. Michael Sharkey, Eric Chapman, (projects NSF DEB 9972024 and 0205982, cited as Sharkey, 2006), Rafael Menjivar, Gergely Várkonyi, Enrique Ruíz-Cancino, Claire Villemant and Santiago Bordera kindly provided data on the field inventories conducted by them. We thank A. Bennett for helpful comments on the manuscript. The study was partly funded by the Kone Foundation, Finland (a grant awarded to the project: Biodiversity and multiple trophic interactions lead by Ilari E. Sääksjärvi). Isrrael C. Gómez thanks the support of the following foundations: the Amazon Conservation Association (Peru), Conservation International Foundation-Peru and the Turku University Foundation. Marc Pollet is indebted to the Leopold III Fund (Belgium) for providing support for the Chilean survey. Field work in Los Queules and Huinay (Chile) was funded by research projects FONDECYT 1050745 and "Ayudas para la realización de estancias de investigación en el centro científico de la Fundación Huinay" (CSIC-Endesa), grants to J. L. Nieves-Aldrey.

LITERATURE CITED

- 1
2
3
4
5
6
7
8 Aguiar, A. & Ramos, A. (2011) Revision of *Digonocryptus* Viereck (Hymenoptera:
9 Ichneumonidae: Cryptinae), with twenty six new taxa and cladistic interpretation of two
10 species complexes. *Zootaxa*, 2846, 1-98.
- 11
12
13
14 Aguiar A., Deans, A.R., Engel, M.S., Forshage, M., Huber, J.T., Jennings J.T., Johnson, N.
15 F., Lelej A.S., Longino, J.T., Lohrmann V., Mikó, I., Ohl, M., Rasmussen, C., Taeger A. &
16
17 Yu, D. S. K. (2013) Hymenoptera In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of
18
19 Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). *Zootaxa*,
20
21 3703, 51-62.
- 22
23
24
25
26
27 Anderson, A.B. (1981) White-sand vegetation of Brazilian Amazonia. *Biotropica*, 13, 199–
28
29 210.
- 30
31
32 Barlow, J., Gardner, T.A., Araujo, I.S., Ávila, T.C., Bonaldo, A.B., Costa, J.E., Esposito,
33
34 M.C., Ferreira, L.V., Hawes, J., Hernandez, M.I., Hoogmoed, M.S., Leite, R.N., LO-Man-
35
36 Hung, N.F., Malcolm J.R., Martins, M.B., Mestre, L.A.M., Miranda-Santos, R., Nunes-
37
38 Gutjahr, A.L., Overal, W.L., Parry, L., Peters, S.L., Ribeiro, M.A., Da Silva, M.N.F., Da
39
40 Silva Motta, C. & Peres, C.A. (2007) Quantifying the biodiversity value of tropical primary,
41
42 secondary, and plantation forests. *Proceedings of the National Academy of Sciences*, 104,
43
44 18555-18560.
- 45
46
47
48 Bartlett, R., Pickering, J., Gauld, I.D. & Windsor, D. (1999) Estimating global biodiversity:
49
50 tropical beetles and wasps send different signals. *Ecological Entomology*, 24, 118-121.
51
52
53
54
55
56
57
58
59
60

1
2
3 Broad, G.R., Sääksjärvi, I.E., Veijalainen, A. & Notton, D.G. (2011) Three new genera of
4
5 Banchinae (Hymenoptera: Ichneumonidae) from Central and South America. *Journal of*
6
7 *Natural History*, 45, 1311-1329.
8

9
10 Castillo, P.M., Ruiz, E., Coronado, J., Khalaim, A. & Myartseva S. (2014) Ichneumonidae
11
12 (Hymenoptera: Ichneumonidae) en un bosque de Quercus de Tamaulipas, Mexico.
13
14 *Dugesiana*, 21, 161-174.
15
16

17
18 Colwell, R.K. & Coddington, J.A. (1994) Estimating terrestrial biodiversity through
19
20 extrapolation. *Philosophical Transactions of the Royal Society, Series B*, 345, 101-118.
21
22

23 Di Giovanni, F., Cerretti, P., Mason, F., Minari, E. & Marini, L. (2015), Vertical stratification
24
25 of ichneumonid wasp communities: the effects of forest structure and life-history traits. *Insect*
26
27 *Science*, 22, 688–699.
28
29

30 Domingo Quero, T., Alonso-Zarazaga, M.A., Sánchez Ruiz, A., Araujo Armero, R., Navas
31
32 Sánchez, A., García Becerra, R., Nebreda, M., Sánchez Ruiz, M., Fontal-Cazalla, F. &
33
34 Nieves-Aldrey, J.L. (2003) Inventariando la Biodiversidad en el Parque Nacional de la
35
36 Caldera de Taburiente (La Palma, Islas Canarias, España): Novedades científicas. *Graellsia*,
37
38 59 (2-3), 45-68.
39
40

41 Gaston, K.J. & Gauld, I.D. (1993) How many species of Pimplines (Hymenoptera:
42
43 Ichneumonidae) are there in Costa Rica? *Journal of tropical Ecology*, 9(4), 491-499.
44
45
46

47 Gauld, I.D. (1991) The Ichneumonidae of Costa Rica, 1. *Memoirs of the American*
48
49 *Entomological Institute*, 47, 1-589.
50
51

52 Gauld, I.D., Hanson, P. & Ugalde, J.A. (1998) Guía de los Pimplinae de Costa Rica
53
54 (Hymenoptera: Ichneumonidae). *Revista de Biología Tropical*, 46, 1-189.
55
56
57
58
59
60

1
2
3 Gauld, I.D., Menjívar, R., González, M.O. & Monro, A. (2002) Guía para la Identificación de
4 los Pimplinae de cafetales bajo sombra de El Salvador (Hymenoptera: Ichneumonidae). *The*
5 *Natural History Museum*, London, 1-76.
6
7

8
9
10 Gauld, I.D. & Dubois, J. (2006) Phylogeny of the *Polysphincta* group of genera
11 (Hymenoptera: Ichneumonidae; Pimplinae): a taxonomic revision of spider ectoparasitoids.
12 *Systematic Entomology*, 31, 529-564.
13
14

15
16
17
18 Gentry, A. (1988) Tree species: richness of Upper Amazonian forest. *Proceedings of the*
19 *National Academy of Sciences of USA*, 85, 156-159.
20
21

22
23 Gentry, A. (1992) Tropical Forest Biodiversity: Distributional Patterns and Their
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conservational Significance. *Oikos*, 63, 19-28.

Gomez, I. C. (2009) Diversity and distribution of pimpline parasitoids wasp (Hymenoptera;
Ichneumonidae; Pimplinae) in the Peruvian Amazonia. MSc Dissertation. University of
Turku. 1-50.

Gómez, I.C., Sääksjärvi, I.E., Veijalainen, A. & Broad, G.R. (2009) Two new species of
Xanthopimpla (Hymenoptera, Ichneumonidae) from Western Amazonia, with a revised key
to the Neotropical species of the genus. *ZooKeys*, 14, 55-65.

Gómez, I.C., Sääksjärvi, I.E., Broad, G.R., Puhakka, L., Peña, C., Castillo, C. & Padua, D.
(2014) The Neotropical species of *Xanthopimpla* Saussure (Hymenoptera, Ichneumonidae).
Zootaxa, 3774 (1), 057-073.

Gómez, I.C., Sääksjärvi, I.E., Broad, G.R., Puhakka, L., Peña, C., Castillo, C. & Bordera, S.
(2015) The Peruvian Amazonian species of *Epirhyssa* (Hymenoptera: Ichneumonidae:
Rhyssinae), with notes on tropical species richness. *Zootaxa*, 3937 (2), 311-336.

- 1
2
3 Gómez, I.C., & Yabar, E. (2015). Description of the first species of *Scambus* Hartig
4
5 (Hymenoptera, Ichneumonidae) from Peru, with a key to the Neotropical species. *Zootaxa*,
6
7 3956 (3), 437-443.
8
9
10 Gonzalez-Moreno, A. & Bordera, S. (2012) The Ichneumonidae (Hymenoptera:
11
12 Ichneumonoidea) of Ria Largatos Biosphere Reserve, Yucatan, Mexico. *Zootaxa*, 3230, 1-51.
13
14
15 Hespeneide, H.A. (1978) Are there fewer parasitoids in the tropics? *American Naturalist*,
16
17 113, 766-769.
18
19
20 Holdridge, L. R. (1967) Life Zone Ecology. *Tropical Science Centre*, San Jose, Costa Rica.
21
22 1-206.
23
24
25 Hovikoski, J., Räsänen, M., Gingras, M., Roddaz, M., Brusset, S. Hermoza, W., Romero, L.
26
27 & Lertora, K. (2005) Miocene semidiurnal tidal rhythmites in Madre de Dios, Peru. *Geology*,
28
29 33(3), 177-180.
30
31
32
33 Idris, A.B. & Kee, S.S. (2002) Horizontal and Vertical Diversity of Ichneumonid Wasps
34
35 (Hymenoptera: Ichneumonidae) in the Sungkai Wildlife Forest Reserve in Perak, Malaysia.
36
37 *Journal Asia-Pacific Entomology*, 5(1), 85-89.
38
39
40
41 Idris, A. B., & Hainidah, J. (2003) Diversity of Ichneumonid Wasps in the logged over forest
42
43 of Langat Basin in Selangor, Malaysia. *Journal of Biological Sciences*, 3, 259-270.
44
45
46 Janzen, D.H. & Pond, C.M. (1975) A comparison, by sweep sampling, of the arthropod fauna
47
48 of secondary vegetation in Michigan, England and Costa Rica. *Royal Entomological Society*
49
50 *of London*, 127, 33-50.
51
52
53
54
55
56
57
58
59
60

1
2
3 Janzen, D. H., Ataroff, M. Farinas M., Reyes, S. Rincon, N., Soler A., Soriano, P. & Vera,
4
5 M. (1976) Changes in the arthropod community along an elevational transect in the
6
7 Venezuelan Andes. *Biotropica*, 8, 193-203.
8
9

10 Kalliola, R., & Flores-Paitan S. (1998) Geologia y Desarrollo Amazonico. *Annales*
11
12 *Universitatis Turkuensis Ser A II*, University of Turku, Finland, 114, 35-57.
13
14

15 Kamath, M.K. & Gupta, V.K. (1972) *Ichneumonologia Orientalis*, 2. The tribe Rhyssini
16
17 (Hymenoptera, Ichneumonidae). *Oriental insects Monograph*, 2, 1-300.
18
19

20 Khalaim, A.I. & Broad G.R. (2013) Tersilochinae (Hymenoptera: Ichneumonidae) of Costa
21
22 Rica, part 2. Genera *Megalochus* gen. nov. and *Stethantyx* Townes. *Zootaxa*, 3693, 221-266.
23
24

25 Khalaim, A.I. & Ruiz-Cancino, E. (2013) A new species of *Megarhyssa* Ashmead
26
27 (Hymenoptera: Ichneumonidae: Rhyssinae) from tropical Mexico with a key to Mexican
28
29 species. *Zootaxa*, 36(2), 195-200.
30
31

32 Kumagai, A.F. (2002) Os Ichneumonidae (Hymenoptera) da Estacao Ecologica da
33
34 Universidad Federal de Minas Gerais, Belo Horizonte, com enfase nas especies de Pimplinae.
35
36 *Revista Brasileira de Entomologia*, 46(2), 189-194.
37
38

39 Kumagai, A.F. & Graf, V. (2002) Biodiversidade de Ichneumonidae (Hymenoptera) e
40
41 monitoramento das especies de Pimplinae e Poemeniinae do Capao da Imbuia, Curitiba,
42
43 Parana. *Revista brasileira de Zoologia*, 19(2), 445-452.
44
45
46
47

48 Marengo, J. (1998): Climatologia en la zona de Iquitos, Peru. In Kalliola, R & Flores Paitan,
49
50 S. Geologia y Desarrollo Amazonico. *Annales Universitatis Turkuensis Ser A II*, University of
51
52 Turku, Finland, 114, 35-57.
53
54
55
56
57
58
59
60

1
2
3 Mayhew, P.J., Dytham, C., Shaw, M.R. & Fraser, S.E.M. (2009) Collections of ichneumonid
4 wasps (Subfamilies Diacritinae, Diplazoninae, Pimplinae and Poemeniinae) from woodlands
5 near York and their implications for conservation planning. *The Naturalist*, 134, 3-24.
6
7
8

9
10 Mazon, M., Rodriguez, A. & Bordera S. (2009) Estudio de la fauna de Ichneumonidae
11 idiobiontes (Hymenoptera) en un ecosistema de montaña mediterránea, I. Subfamilias
12 Pimplinae y Xoridinae. *Boletín de la Asociación Española de Entomología*, 33, 123-137.
13
14
15

16
17
18 Mazon, M., & Bordera, S. (2014) Diversity of Ichneumonidae (Insecta: Hymenoptera) in a
19 protected area of Central Spain: What are we protecting? *Insect Conservation and Diversity*,
20 7: 393-484.
21
22
23

24
25 Mohammadi-Khoramabadi, A., Talebi A. & Zwakhals, K. (2013) A study of the subfamily
26 Pimplinae (Hymenoptera: Ichneumonidae) in the north of Iran, with eleven new species
27 records. *Entomofauna*, 34, 29-56.
28
29
30

31
32 Morrison, G., Auerbach, M. & McCoy, E.D. (1979) Anomalous diversity of tropical
33 parasitoids: a general phenomenon? *American Naturalist*, 114, 303-307.
34
35
36

37 Nieves-Aldrey, J. L. (1995) Abundancia, diversidad y dinámica temporal de cinípidos en
38 dos hábitats del centro de España (Hymenoptera, Cynipidae). En: *Avances en Entomología*
39 *Ibérica*. Com. organiz. VI Congreso Iber. Entom (eds). Madrid. pp 113-136.
40
41
42

43
44 Nieves-Aldrey, J.L. & Rey del Castillo, C. (1991) Ensayo preliminar sobre la captura de
45 insectos por medio de una trampa Malaise en la sierra de Guadarrama (España) con especial
46 referencia a los himenópteros (Insecta, Hymenoptera). *Ecología*, 5, 383-403.
47
48
49

50
51 Nieves-Aldrey, J. L., Fontal F., Garrido-Torres A & Rey Del Castillo C. (2003) Inventario
52 de Hymenoptera (Hexapoda) en El Ventorrillo: un rico enclave de Biodiversidad en la Sierra
53 de Guadarrama. *Graellsia*, 59, 25-43.
54
55
56
57
58
59
60

- 1
2
3 Oksanen, J., Guillaume Blanchet F., Kindt, R. Legendre, P. Minchin, P. R. O'hara, R. B.
4
5 Simpson, G. L. Solymos, P. Stevens, M. H. H. & Wagner H. (2013) vegan: Community
6
7 Ecology Package. R package version 2.0-7. <http://CRAN.R-project.org/package=vegan>
8
9
10 Orme, C.D., Davies L.R.G., Burgess M., Eigenbrod F., Pickup N., Olson, V.A. Webster, A.J
11
12 Ding T., Rasmussen P.C., Ridgel R.S., Stattersfield A.J., Bennett P.M., Blackburn T.M.,
13
14 Gaston, K.J. & Owens I.P.F. (2005) Global hotspots of species richness are not congruent
15
16 with endemism or threat. *Nature*, 436, 1016–1019.
17
18
19
20 Owen, D.F. & Owen, J.M. (1974) Species diversity in temperate and tropical Ichneumonidae.
21
22 *Nature*, 249, 583-584.
23
24
25 Owen, J., Townes, H. & Townes, M. (1981) Species diversity of Ichneumonidae and
26
27 Serphidae (Hymenoptera) in an English suburban garden. *Biological Journal of the Linnean*
28
29 *Society*, 16, 315-336.
30
31
32
33 Patterson, B.D., Stotz, D.F. & Solari, S. (2006) Mammals and Birds of the Manu Biosphere
34
35 Reserve, Peru. *Fieldiana. Zoology*, 10, 1-49.
36
37
38 Peck, R. W., Banko P.C., Schwarzfeld M., Euaparadorn M. & Brinck K.W. (2008) Alien
39
40 dominance of the parasitoid wasp community along an elevation gradient on Hawai'i Island.
41
42 *Biological Invasions*, 10, 1441-1455.
43
44
45 Pérez-Urbina, B., Correa-Sandoval, A. Ruíz-Cancino, E. Kasparyan, D.R. Coronado-
46
47 Blanco, J. & Horta-Vega. V. (2010) Diversidad de Ichneumonidae (Hymenoptera) en el
48
49 Cañón del Novillo, Victoria, Tamaulipas, México. *Entomotropica*, 25: 83-97.
50
51
52
53 Pham, N.T. (2013) Taxonomy and distributional patterns of Pimplinae (Hymenoptera:
54
55 Ichneumonidae) from Vietnam. PhD Dissertation. 1-281. University of Bonn, Germany.
56
57
58
59
60

1
2
3 Pino, D. (2007) Distribution and Habitat Preferences of Lecythidaceae in the Los Amigos
4 Uplands, South Western Amazonia Peru. MSc Dissertation. 1-84. University of Göttingen,
5 Germany.
6
7
8

9
10 Porter, C.C. (1970) A revision of the South American species of *Coccygomimus*
11 (Hymenoptera, Ichneumonidae). *Studia Entomologica*, 13, 1-92.
12

13
14
15 Porter, C.C. (1978) A revision of the genus *Epirhyssa* (Hymenoptera, Ichneumonidae). *Studia*
16 *Entomologica*, 20, 297-318.
17

18
19
20 Porter, C. C. 1997. Guía de los géneros de Ichneumonidae en la región Neantartica del sur de
21 Sudamérica. *Acta Zoologica Lilloana*, 42, 1-216.
22

23
24
25
26 Quicke, D.L.J. (2012) We Know Too Little about Parasitoid Wasp Distributions to Draw Any
27 Conclusions about Latitudinal Trends in Species Richness, Body Size and Biology. *PLoS*
28 *ONE*, 7(2): e32101.
29

30
31
32 Rathcke, B.J. & Price, P.W. (1976) Anomalous diversity of tropical ichneumonid parasitoids:
33 a predation hypothesis. *American Naturalis*, 110, 889-893.
34

35
36
37
38 Robbins, R.K., Lamas, G., Mielke, O.H., Harvey, D.J. & Casagrande, M.M. (1996)
39 Taxonomic composition and ecological structure of the species-rich butterfly community at
40 Pakitza, Parque Nacional del Manu, Peru. 217-252 pp. In Wilson and Sandoval, eds., La
41 biodiversidad sureste del Peru: Manu. Editorial Horizonte, Lima, Peru.
42
43
44

45
46
47 Robbins, R.K. & Opler, P.A. (1997) Butterfly diversity and a preliminary comparison with
48 bird and mammal Diversity. In Reaka-Kudla, Wilson D.E. & Wilson E.O. eds. Biodiversity
49 II. Joseph Henry Press, Washington, D.C 69-82.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Robert, T. (2011) Méthodologie d'échantillonnage des Hyménoptères Ichneumonidae en
4 milieu forestier. Cas de la chênaie charmaie sur Plateau lorrain. Ministère de l'enseignement
5 supérieur et de la recherche. école pratique des hautes études. *Sciences de la Vie et de la*
6 *Terre*, France.

7
8
9
10
11
12 Rome, Q. (2006) Impact de la gestion forestière sur la biodiversité des Ichneumonides de la
13 hêtraie pyrénéenne. Master 2 Systématique, Évolution, Paléontologie parcours Expertise
14 Faune Flore et Gestion du Patrimoine Naturel. Université Pierre et Marie CURIE. France.

15
16
17
18
19
20 Ruiz-Guerra, B., Hanson, P., Guevara, R. & Dirz, R. (2013) Richness and Abundance of
21 Ichneumonidae in a Fragmented Tropical Rain Forest. *Neotropical Entomology*. DOI
22 10.1007/s13744-013-0143-3.

23
24
25
26
27 Santos, A. & Quicke, D. (2011) Large-scale diversity patterns of parasitoid insects.
28 *Entomological science*, 14, 371-382.

29
30
31
32 Santos, B. & Aguiar, A. (2013) Phylogeny and revision of *Messatoporus* Cushman
33 (Hymenoptera, Ichneumonidae, Cryptinae), with descriptions of sixty five new species.
34 *Zootaxa*, 3634 (1), 1-284.

35
36
37
38
39
40 Sharkey, M.J. (2006) Two new genera of Agathidinae (Hymenoptera: Braconidae) with a key
41 to the genera of the New World. *Zootaxa*, 1185, 37-51.

42
43
44
45
46 Sime, K. & Brower, A. (1998) Explaining the latitudinal gradient anomaly in ichneumonid
47 species richness: evidence from butterflies. *Journal of Animal Ecology*, 67, 387-399.

48
49
50
51
52
53
54
55
56
57
58
59
60 Steinbauer, M.J., Short M.W. & Schmidt, S. (2006) The influence of architectural and
vegetational complexity in eucalypt plantations on communities of native wasp parasitoids:
Towards silviculture for sustainable pest management. *Forest Ecology and Management*,
233, 153-164.

- 1
2
3 Sääksjärvi, I.E., Haataja, S., Neuvonen, S., Gauld, I.D., Jussila, R., Salo, J. & Marmol-
4
5 Burgos, A. (2004) High local species richness of parasitic wasps (Hymenoptera:
6
7 Ichneumonidae; Pimplinae and Rhyssinae) from the lowland rain forests of Peruvian
8
9 Amazonia. *Ecological Entomology*, 29, 735-743.
10
11
12 Tanque, R.L. (2009) Pimplinae, Poemeniinae e Rhyssinae (Hymenoptera: Ichneumonidae) na
13
14 Unidade Ambiental de Peti (Cemig)-MG. Dissertação de mestrado, Universidade Federal de
15
16 Lavras, Lavras, Brasil.
17
18
19
20 Tanque, R.L., & Frieiro-Costa F.A. (2011) Pimplinae (Hymenoptera, Ichneumonidae) em um
21
22 fragmento de Cerrado na Reserva Biológica Unilavras/Boqueirão, Ingaí, Minas Gerais,
23
24 Brasil. *Biota Neotropica*, 11, 169-171.
25
26
27
28 Timms, L.L., Schwarzfeld, M. & Sääksjärvi, I.E. (2015) Extending understanding of
29
30 latitudinal patterns in parasitoid wasp diversity. *Insect Conservation and Diversity*, 9, 74-86.
31
32
33 Townes, H. & M. Townes. (1966) A catalogue and reclassification of Neotropical
34
35 Ichneumonidae. *Memoirs of the American Entomological Institute*, 8, 1-367.
36
37
38 Tuomisto, H. & Ruokolainen K. (1994) Distribution of Pteridophyta and Melastomataceae
39
40 along an edaphic gradient in an Amazonian rain forest. *Journal of Vegetation Science*, 5, 25-
41
42 34.
43
44
45 Tuomisto, H., Ruokolainen, K., Kalliola, R., Linna, A. Danjoy, W. & Rodriguez, Z. (1995)
46
47 Dissecting Amazon Biodiversity. *Science*, 269, 63-66.
48
49
50
51 Tuomisto, H. (2010) A consistent terminology for quantifying species diversity? Yes, it does
52
53 exist. *Oecologia*, 164, 853-860.
54
55
56
57
58
59
60

1
2
3 Van Noort, S. (2004) Ichneumonid (Hymenoptera: Ichneumonoidea) diversity across an
4 elevational gradient on Monts Doudou in Southwestern Gabon. *Memoirs of the California*
5 *Academy of Sciences*, 28, 187-216.
6
7

8
9
10 Várkonyi, G. & Roslin, V. (2013) Freezing cold yet diverse: dissecting a high-Arctic
11 parasitoid community associated with Lepidoptera hosts. *Canadian Entomology*, 145, 193-
12 218.
13
14

15
16
17 Veijalainen, A., Sääksjärvi, I.E., Erwin, T.L., Gómez, I.C. & Longino, J.T. (2013) Subfamily
18 composition of Ichneumonidae (Hymenoptera) from western Amazonia: insights into
19 diversity of tropical parasitoid wasps. *Insect Conservation Diversity*, 6, 28-37.
20
21

22
23
24 Veijalainen, A., Sääksjärvi, I.E. Tuomisto, H., Broad G.R., Bordera, S. & Jussila R. 2014.
25 Altitudinal trends in species richness and diversity of Mesoamerican parasitoid wasps
26 (Hymenoptera: Ichneumonidae) *Insect Conservation Diversity*, 7, 496-507.
27
28

29
30
31 Wilson, D.E. & Sandoval, A. (1996) *Manu - The Biodiversity of Southeastern Peru*.
32 Smithsonian Institution and Editorial Horizonte, Lima, Peru.
33
34

35
36
37 Yu, D.S.K., Achterberg, C. & Horstmann, K. (2012) Taxapad 2012, Ichneumonoidea 2011.
38 Database on flash-drive. www.taxapad.com , Ottawa, Ontario, Canada.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURES AND TABLES

Fig. 1 Sites with information on Pimplinae and Rhyssinae species richness. Red circles represent localities inventoried for this study, blue circles localities for which data were compiled from the literature.




Figure 2. Accumulation of individuals (A) and species (B, C) as a function of sampling duration (A, B) and the number of individuals captured (C) in Malaise traps in Peruvian Amazonia. Each line corresponds to one trap and displays the observed accumulation of individuals or species during the entire operational period. A-M refers to Allpahuayo-Mishana (northern Peru), LA to Los Amigos (southern Peru).




Figure 3. Estimated number (A) and percentage (B) of additional Pimplinae and Rhyssinae species expected if more individuals could have been added to samples currently available from Peruvian Amazonia. Each circle corresponds to the total catch in a number of original 1-2-week samples that were combined randomly but always within the same habitat type. A-M refers to Allpahuayo-Mishana (northern Peru), LA to Los Amigos (southern Peru).




Figure 4. Accumulation of Pimplinae and Rhyssinae individuals (A) and species (B) as a function of sampling effort (Malaise trap months) in Peruvian Amazonia (A-M and LA) and in other parts of the world (Literature). Symbols in grey correspond to samples that only reported Pimplinae. Open circles correspond to lowland sites (<1000 m elevation) and closed circles to montane sites (>1000 m elevation). Circle size is proportional to absolute value of latitude. For Allpahuayo-Mishana and Los Amigos (A-M and LA, respectively), each symbol corresponds to a combination of randomly chosen 1-2-week samples from the same habitat type. Some literature sites present in B are missing from A because they did not report number of individuals.




Figure 5. Accumulation of species richness (A) and diversity (B, C) of Pimplinae and Rhyssinae wasps as a function of the number of individuals captured in Malaise traps in Peruvian Amazonia (A-M and LA) and in other parts of the world (Literature). Diversity at $q=1$ equals the exponential of the Shannon index and diversity at $q=2$ the reciprocal of the Simpson index. Second-order polynomial regression lines are shown in (A) separately for our Peruvian data (black) and for data from other parts of the world (grey). For detailed explanation of the symbols, see Fig. 4.





Figure 6. (A) Number of Pimplinae and Rhyssinae species in samples obtained at different latitudes with circle sizes reflecting sampling effort (Malaise trap months). Open symbols correspond to lowland sites (<1000 m) and closed symbols to montane sites (>1000 m). (B) The residuals from a second-order polynomial regression of number of species against number of individuals, both logarithmically transformed with log base 10, plotted against absolute latitude (a residual of -1.0 indicates that the observed number of species is one-tenth of what would be expected, given the number of individuals). Two regression lines are shown to test separately for latitudinal trends in lowland and montane sites (cut-off elevation 1000 m). Symbol size reflects elevation, with montane sites highlighted with a slightly thicker line. (C) Like (B), but residuals plotted against elevation, and regressions are given separately for tropical and non-tropical sites (cut-off latitude 25 degrees). Symbol size reflects absolute latitude. In all three panels, coloured symbols correspond to a random subsample of the data from our Peruvian sites; colour scheme is the same as in Fig. 4. Each of the other study localities shown in Table S1 is indicated with a black or grey symbol (for studies reporting both Pimplinae and Rhyssinae vs only Pimplinae, respectively).



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1. Summary of the sampling methods and sampling effort in ichneumonid inventories in two Peruvian Amazonian sites.

Site	Allpahuayo-Mishana			Los Amigos
Collection year	1998	2000	2011	2008
Malaise sampling	41MTM	144MTM	45MTM	27MTM
Yellow pan trap	–	–	76.7 traps month	–
Hand netting	–	–	92 hours	52 hours
Larval rearing	–	–	3298	–
Pupal rearing	–	–	113	–

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17 **SUPPORTING INFORMATION**
18
19

20
21
22
23 **Additional Supporting Information may be found in the online version of this article:**
24

25
26 **Table S1** Species-richness of Pimplinae and Rhyssinae in 97 sites sampled with Malaise traps around the world. MTM: Malaise trap months;
27
28 C.T: Caldera de Taburiente; S.N.: Sierra Nevada; P.N.: Parque Nacional; R.N. Reserva Nacional; *: Studies that reported only Pimplinae and
29
30 may not include Rhyssinae.
31
32
33
34
35
36

Country	Locality	Species	Individuals	MTM	Indiv./MTM	Elevation	Latitude	Longitude	Reference
---------	----------	---------	-------------	-----	------------	-----------	----------	-----------	-----------

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

						(m)			
Australia	Ginninderra	5	22	9.1	0.55	588-615	-35.15	149.03	Steinbauer <i>et al.</i> , (2006)
Belize	Las Cuevas	25	400	18	22.2	600	16.73	-88.98	*Bartlett <i>et al.</i> , (1999)
Brazil	Boqueirão	26	100	8	12.5	1100	-21.33	-44.98	*Tanque & Frieiro 2011
Brazil	Capão da Imbuia	47	620	70	8.9	900	-25.42	-49.23	Kumagai & Graf 2002
Brazil	Belo Horizonte	30	306	12	25.5	850	-19.87	-43.97	Kumagai 2002
Brazil	Santa Barbara	38	465	24	19.3	770-800	-19.88	-43.37	Tanque <i>et al.</i> , 2009
Chile	Alerce Andino	2	2	0.3	6.7	129-141	-41.58	-72.58	In this study
Chile	Conguillio	2	46	1.2	38.3	1195-1264	-38.63	-71.65	In this study
Chile	Huinay, Comau Fjord	2	104	4.25	24.47	45	-42.38	-72.42	In this study
Chile	Ñuble	2	5	1.3	3.8	754-765	-36.93	-71.55	In this study
Chile	Puyehue	1	2	0.5	4	367-558	-40.73	-72.32	In this study
Chile	R. N. Les Queules	4	44	37.17	1.18	420	-35.99	-72.71	In this study
Chile	Valle las Trancas	1	1	0.64	1.6	1220	-36.9	-71.48	In this study
Chile	Rio Clarillo	3	36	8	4.5	1600	-33.75	-70.42	Porter, 1997

Colombia	Amacayacu	8	–	50	–	150-210	-3.68	-70.25	* Sharkey 2006
Colombia	Iguaque high	27	–	15.83	–	3300-3420	6.25	-72.71	* Sharkey 2006
Colombia	Iguaque low	19	–	8.1	–	2600	5.71	-73.46	* Sharkey 2006
Colombia	Iguaque mid	52	–	20.7	–	2820-2920	5.41	-73.45	* Sharkey 2006
Colombia	Planada	10	–	59	–	1800-1930	1.5	-77.53	* Sharkey 2006
Colombia	S. N. de Santa Marta high	11	–	7.4	–	2300-2500	10.8	-73.65	* Sharkey 2006
Colombia	S. N. de Santa Marta low	3	–	2.67	–	1500-1700	10.76	-73.91	* Sharkey 2006
Colombia	S. N. de Santa Marta mid	15	–	8.93	–	2200-2220	10.82	-73.88	* Sharkey 2006
Costa Rica	16 km W Guapiles	26	87	18	4.8	400	10.15	-83.92	Gaston & Gauld 1993
Costa Rica	Casa Maritza	54	1289	75	17.2	600-650	10.95	-85.5	Gaston & Gauld 1993
Costa Rica	Cerro de Hacha	26	602	144	4.2	300-400	11	-85.55	Gaston & Gauld 1993
Costa Rica	Cerro de la Muerte	14	420	42	10	2700-2800	9.63	-83.8	Gaston & Gauld 1993
Costa Rica	Estacion Cacao	67	1239	84	14.8	1000-1100	10.93	-85.47	Gaston & Gauld 1993
Costa Rica	Estacion Pitilla	57	517	72	7.2	600-700	11	-85.43	Gaston & Gauld 1993
Costa Rica	Finca San Gabriel	21	76	6	12.7	650-700	10.88	-85.4	Gaston & Gauld 1993

Costa Rica	Golfo Dulce Forest	32	383	45	8.5	100-200	8.75	-83.33	Gaston & Gauld 1993
Costa Rica	La Selva	25	152	12	12.7	50-75	10.43	-84.02	Gaston & Gauld 1993
Costa Rica	Monteverde	23	64	3	21.3	1300-1350	10.3	-84.8	Gaston & Gauld 1993
Costa Rica	Rio Agres	18	93	3	31	1500	9.88	-84.13	Gaston & Gauld 1993
Costa Rica	San Antonio de Escazu	47	590	84	7	1300	9.9	-84.13	Gaston & Gauld 1993
Costa Rica	Santa Rosa	40	1579	576	2.7	250-300	10.85	-85.62	Gaston & Gauld 1993
Costa Rica	San Vito	28	88	18	4.9	1200	8.8	-82.97	Gaston & Gauld 1993
Costa Rica	Vara Blanca	14	281	15	18.7	2100	10.15	-84.13	Gaston & Gauld 1993
Costa Rica	Villa Mills	8	144	54	2.7	3000-3100	9.57	-83.72	Gaston & Gauld 1993
Costa Rica	Zurqui de Moravia	29	102	15	6.8	1600	10.07	-84	Gaston & Gauld 1993
El Salvador	Ahuachapan	29	-	1	-	660-800	13.8	-89.85	* Gauld <i>et al.</i> , 2002
El Salvador	La Libertad	37	-	1	-	1000	13.65	-89.37	* Gauld <i>et al.</i> , 2002
El Salvador	San Vicente	22	-	1	-	1100	13.47	-88.52	* Gauld <i>et al.</i> , 2002
El Salvador	Santa Ana	24	-	1	-	1450	14.37	-89.27	* Gauld <i>et al.</i> , 2002
El Salvador	Usulután	20	-	1	-	1480	13.47	-88.52	* Gauld <i>et al.</i> , 2002

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

France	Bezange à bois mort	32	2412	36	67	260	48.75	6.45	Robert 2011
France	Parroy à bois mort	29	1909	36	53	270	48.63	6.63	Robert 2011
France	Gar-Cagire	21	169	13.5	12.5	1328-1500	42.95	0.7	Rome 2006
Gabon	Monts Doudou	3	4	3	1.3	110-630	-2.22	10.38	Van Noort 2004
Greenland	Northeast National Park	1	–	21	–	598-791	75.95	-29.93	Várkonyi & Roslin, 2013
Guatemala	Atlitan	6	17	0.6	28.3	1575-1685	14.52	-91.19	Veijalainen <i>et al.</i> , 2014
Guatemala	Montaña Chiclera	4	4	0.67	6	210-245	14.95	-88.87	Veijalainen <i>et al.</i> , 2014
Honduras	Cusuco	13	39	0.67	58.2	1210-1260	15.55	-88.24	Veijalainen <i>et al.</i> , 2014
Honduras	Guisayote	4	9	0.67	13.4	2090-2190	14.45	-89.07	Veijalainen <i>et al.</i> , 2014
Honduras	La Muralla	7	21	0.67	31.3	1410-1530	15.07	-86.74	Veijalainen <i>et al.</i> , 2014
Iran	Arangeh	4	5	16.6	0.3	1891	35.92	51.08	Mohammadi <i>et al.</i> , 2013
Iran	Astaneh-e-Ashrafieh	5	55	16.6	3.3	-1	37.36	49.95	Mohammadi <i>et al.</i> , 2013
Iran	Ghazichak	10	53	16.6	3.2	1803	36.75	50.32	Mohammadi <i>et al.</i> , 2013
Iran	Karaj	4	18	16.6	1.1	1278	35.77	50.93	Mohammadi <i>et al.</i> , 2013

Iran	Orkom	12	64	16.6	3.9	1201	36.75	50.3	Mohammadi <i>et al.</i> , 2013
Iran	Sarziarat	3	8	16.6	0.5	1980	35.92	51.1	Mohammadi <i>et al.</i> , 2013
Iran	Shahrestanak	5	7	16.6	0.4	2305	35.95	51.37	Mohammadi <i>et al.</i> , 2013
Iran	Shahryar	1	3	16.6	0.2	1168	35.67	50.93	Mohammadi <i>et al.</i> , 2013
Iran	Ziaz	9	28	16.6	1.7	537	36.87	50.22	Mohammadi <i>et al.</i> , 2013
Italy	Bosco della Fontana	25	3945	56	70.4	25	45.2	10.73	Di Giovanni <i>et al.</i> , 2014
Malaysia	Langat basin	28	84	27	3.1	50-200	2.73	101.44	Idris & Hainidah, 2003
Malaysia	Sungkai	8	22	21	1	60	3.97	101.25	Idris & Kee, 2002
Mexico	Ria Lagartos	11	181	168	1.1	0-4	21.6	-88.17	Gonzalez & Bordera, 2012
Mexico	Cañon del Novillo	12	99	12	8.3	420	23.68	-99.2	Perez-Urbina <i>et al.</i> , 2010
Mexico	Rancho Santa Elena	13	263	12	21.9	750	23.62	-99.2	Castillo <i>et al.</i> , 2014
Mexico	Los Tuxtlas	6	14	13	1.07	76-159	18.58	-95.05	Ruiz-Guerra <i>et al.</i> , 2013
Nicaragua	Cerro Jesus	11	31	0.76	40.8	1440-1780	13.97	-86.18	Veijalainen <i>et al.</i> , 2014
Nicaragua	Datanlí-El Diablo	18	45	0.7	64.3	1330-1440	13.12	-85.87	Veijalainen <i>et al.</i> , 2014

Nicaragua	Saslaya low	5	5	0.8	6.3	290-370	13.73	-84.98	Veijalainen <i>et al.</i> , 2014
Panama	Barro Colorado	28	952	54	17.6	100	9.15	-79.85	*Bartlett <i>et al.</i> , 1999
Panama	P. N. Isla de Coiba	2	3	1.43	2.1	40	7.63	-81.73	In this study
Peru	Allpahuayo-Mishana	105	1597	240	6.7	110-180	-3.95	-73.43	In this study
Peru	Los Amigos	51	121	27	4.5	260	-12.57	-70.08	In this study
Spain	El Cabril	5	18	12.5	1.44	300	38.05	-5.42	In this study
Spain	Roblehondo (Cazorla)	14	51	5.5	9.27	1280	37.95	-2.88	In this study
Spain	Hoyo de Manzanares	13	50	5	10	970	40.62	-3.91	In this study
Spain	Llueva	26	219	8	27.38	100	43.33	-3.55	In this study
Spain	Mirambell	30	326	17	19.18	900	41.8	2.19	In this study
Spain	Mojácar	2	15	4	3.75	137	37.12	-1.85	In this study
Spain	San Roque de Riomiera	9	28	5.4	5.19	570	43.23	-2.71	In this study
Spain	El Pardo	11	63	12	5.25	750	40.54	-3.73	In this study
Spain	Carrasqueta	6	48	46	1.04	940-1300	38.6	-0.48	Mazon <i>et al.</i> , 2009
Spain	Mariola	5	106	46	2.3	900-1200	38.75	-0.5	Mazon <i>et al.</i> , 2009

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Spain	El Ventorrillo	33	428	12.6	34	1450	40.75	-4.02	Nieves-Aldrey <i>et al.</i> , 2003
Spain	Cabañeros	22	236	130	1.8	800	39.37	-4.48	Mazon <i>et al.</i> , 2014
Spain (C. Islands)	Isla de la Palma C.T. Barranco de Las Traves	1	1	1.76	0.57	1050	28.72	-17.89	In this study
Spain (C. Islands)	Isla de la Palma C.T. Roque de la Cumbrecita	2	5	0.74	6.75	1297-1377	28.7	-17.86	In this study
Spain (C. Islands)	Isla de la Palma C.T. Roque de los Muchachos	2	10	2.27	4.4	2250	28.75	-17.88	In this study
Uganda	Kibale	17	50	252	0.2	1240-1375	0.48	30.38	In this study
UK	York	41	1836	60	30.6	15	53.95	-1.07	Mayhew <i>et al.</i> , 2009
UK	Leicester	19	158	20	7.9	120	52.67	-1.18	Owen <i>et al.</i> , 1981

USA (Hawaii)	Bryson's	2	57	39.8	1.4	285	19.45	-154.92	Peck <i>et al.</i> , 2008
USA (Hawaii)	Solomon's Waterhole	5	143	52.1	2.7	1792	19.53	-155.32	Peck <i>et al.</i> , 2008

Table S2 Abundance, species richness and functional type of the Pimplinae and Rhyssinae genera found in the Peruvian localities Allpahuayo-Mishana and Los Amigos. The functional types follow Gauld (1991) and Sääksjärvi *et al.* (2004) and are: IDC = idiobiont parasitoids of deeply concealed hosts (hosts located e.g. within wood); IWC = idiobiont parasitoids of weakly concealed hosts (exposed hosts or hosts within e.g. leaf rolls); IES / Pseudo = idiobiont ectoparasitoids of spiders, or pseudoparasitoids of spider egg sacs; and 4) KES = koinobiont ectoparasitoids of spiders.

Subfamily	Genus	Allpahuayo-Mishana		Los Amigos		Functional type
		Individuals	Species	Individuals	Species	
Rhyssinae	<i>Epirhyssa</i>	81	13	15	9	IDC
Pimplinae	<i>Anastelgis</i>	11	1	5	1	IDC
Pimplinae	<i>Dolichomitus</i>	44	4	12	5	IDC
Pimplinae	<i>Amazopimpla</i>	9	2	1	1	IWC
Pimplinae	<i>Clydonium</i>	1	1	–	–	IWC

Pimplinae	<i>Zonopimpla</i>	3	1	3	3	IWC
Pimplinae	<i>Neotheronia</i>	881	41	51	20	IWC
Pimplinae	<i>Nomosphaeria</i>	10	3	–	–	IWC
Pimplinae	<i>Pimpla</i>	95	1	14	1	IWC
Pimplinae	<i>Xanthopimpla</i>	137	6	3	2	IWC
Pimplinae	<i>Zaglyptus</i>	39	3	2	1	IES / Pseudo
Pimplinae	<i>Tromatobia</i>	–	–	2	1	IES / Pseudo
Pimplinae	<i>Acrotaphus</i>	103	8	2	1	KES
Pimplinae	<i>Hymenoepimecis</i>	146	13	8	5	KES
Pimplinae	<i>Polysphincta</i>	15	3	–	–	KES
Pimplinae	<i>Ticapimpla</i>	12	3	–	–	KES
Pimplinae	<i>Zatypota</i>	10	2	–	–	KES
Pimplinae	genus unknown	–	–	3	1	KES
Total		1597	105	121	51	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49