

This is a repository copy of *Fate, uptake, and distribution of nanoencapsulated pesticides in soil–earthworm systems and implications for environmental risk assessment*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/126473/>

Version: Accepted Version

Article:

Firdaus, Mohd Anuar Mohd, Agatz, Annika orcid.org/0000-0003-3228-8822, Hodson, Mark E orcid.org/0000-0002-8166-1526 et al. (2 more authors) (2018) Fate, uptake, and distribution of nanoencapsulated pesticides in soil–earthworm systems and implications for environmental risk assessment. *Environmental Toxicology and Chemistry*. pp. 1420-1429. ISSN 1552-8618

<https://doi.org/10.1002/etc.4094>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

2
3 **Fate, Uptake and Distribution of Nanoencapsulated Pesticides in Soil-Earthworm**
4 **Systems and Implications for Environmental Risk Assessment**

5 Mohd Firdaus M. A.^{1,2}, Agatz. A.¹, Hodson M. E.¹, Omar S. A. Al-Khazrajy^{1,3}, Boxall A. B. A.^{1*}

6 ¹Environment Department, Wentworth Way, University of York, Heslington, York, YO10
7 5NG, United Kingdom

8 ²Department of Land Management, Faculty of Agriculture, 43400 UPM Serdang, Selangor
9 Darul Ehsan, Malaysia

10 ³Department of Chemistry, College of Ibn al-Haitham, University of Baghdad, Baghdad, Iraq

11 * - corresponding author, email: Alistair.boxall@york.ac.uk

12 **Abstract**


13 Nanopesticides are novel plant protection products offering numerous benefits. As
14 nanoparticles behave differently from dissolved chemicals, environmental risks of these
15 materials could differ from conventional pesticides. Here we used soil-earthworm systems to
16 compare the fate and uptake of analytical grade bifenthrin to that of bifenthrin in traditional and
17 nano-encapsulated formulations. Apparent sorption coefficients for bifenthrin in the nano-
18 treatments were up to 3.8 times lower than in the non-nano treatments whereas dissipation
19 half-lives of the nano-treatments were up to two time longer. Earthworms in the nano-
20 treatments accumulated around 50% more bifenthrin than those in the non-nano treatments.
21 In the non-nano treatments, most of the accumulated material was found in the earthworm
22 tissue while in the nano-treatments, the majority resided in the gut. Evaluation of toxicokinetic
23 modelling approaches showed that models incorporating the release rate of bifenthrin from
24 the nanocapsule and distribution within the earthworm provided the best estimations of uptake
25 from the nanoformulations. Overall, our findings indicate that the risks of nanopesticides may

26 be different from conventional formulations. The modelling presented here provides a starting
27 point for assessing risks of these materials but needs to be further developed to better
28 consider the behaviour of the nanoencapsulated pesticide within the gut system.

29 Keywords: Nanopesticides; Synthetic pyrethroids; Nanoencapsulation; Earthworms;
30 Toxicokinetic modelling, *Eisenia fetida*, *Lumbricus terrestris*

31

32 Graphical abstract



33

34

35

36 **Introduction**

37 Recently, novel pesticide products have been developed that employ nanotechnology (Kah et
38 al., 2013; Kah, 2015). These so called 'nanopesticides' comprise either nanoparticulate forms
39 of a pesticide active ingredient or nanocapsules containing an active ingredient (a.i.).
40 Nanopesticides offer a range of advantages over conventional pesticides in that they may
41 increase efficacy of the a.i. and/or enhance the environmental and human health safety
42 profiles of the products (Kah et al., 2013; Kookana et al., 2014). However, there is recognition
43 that the application of nanotechnology could also have negative and unanticipated impacts on
44 the environment so it is also possible that nanopesticides could pose a greater risk than
45 equivalent conventional pesticide products. The applicability of existing environmental risk
46 assessment approaches for pesticides to nanoformulations has also been questioned (Kah,
47 2015).

48 One group of organisms that will be exposed to nanopesticides are terrestrial invertebrates
49 such as earthworms. Earthworms are known to bio-magnify inorganic and organic soil
50 contaminants, including pesticides, polycyclic aromatic hydrocarbons, brominated flame
51 retardants, and metals (Heikens et al., 2001; Matscheko et al., 2002; Langdon et al., 2005).
52 Earthworms being at the base of a food chain hold an integral position. Uptake and
53 accumulation of contaminants into earthworms not only poses a risk to the earthworm directly,
54 but bioaccumulation and contaminant transfer through the food chain to top predators such as
55 birds has the potential to result in secondary poisoning (Spurgeon and Hopkin, 1996).

56 Data for other non-pesticide nanoparticles shows that these materials can be taken up by
57 earthworms (Kwak and Youn-Joo, 2005). Investigations determining distribution of
58 nanoparticles show that highest concentrations of accumulated materials are associated with
59 the earthworm gut (Unrine et al., 2010; Waissi-Leinonen et al. 2012). Adverse effects have
60 also been reported in earthworms following exposure to carbon-based and metal and metal
61 oxide nanoparticles (Kwak and Youn-Joo, 2005; Scott-Fordsmand et al. 2008).

62 To date, the focus of research into bioconcentration and impacts of nanoparticles on
63 earthworms has been on metals and metal oxides (Kwak and Youn-Joo, 2005), carbon
64 nanotubes (Petersen et al., 2008, Petersen et al., 2011, Scott-Fordsmand et al. 2008) and
65 fullerenes (Li et al. 2010, Kelsey and White 2013). To the best of our knowledge, no-one has
66 explored the uptake of nanopesticides, even though it is inevitable that earthworms will be
67 exposed to these products during use. Therefore, here we investigate the effects of
68 nanoencapsulation on the fate, uptake, depuration and distribution of a pesticide a.i. in soil-
69 earthworm systems. The nanoencapsulated materials used in the study were developed by
70 Vive Crop Protection Inc and comprise bifenthrin encapsulated in a polymer nanoparticle with
71 the aim to better target the active ingredient to the pest species. We compare the fate, uptake
72 and distribution of the analytical grade a.i. with that of conventional and nanoformulated
73 products for the two earthworm species *Eisenia fetida* and *Lumbricus terrestris*. The findings
74 are used to explore the suitability of existing and novel toxicokinetic models to better
75 characterise the environmental risks of nanoencapsulated substances in the future.

76 **Materials and methods**

77 *Chemicals, soils and organisms*

78 Analytical PESTANAL[®] grade bifenthrin was purchased from Sigma-Aldrich (Dorset, UK),
79 formulated bifenthrin (Capture LFR) was obtained from FMC Corporation (Philadelphia, USA).
80 Two nanoencapsulated formulations of bifenthrin (Nano A and B) were obtained from Vive
81 Crop Protection Inc. (Toronto, Canada). The precise make-up of these materials is proprietary
82 but both formulations employ an acrylate copolymer to encapsulate the bifenthrin but contain
83 different co-formulants. Acetonitrile (99.9%) was purchased from Fisher Scientific
84 (Loughborough, UK). Details of the bifenthrin treatments are provided in the Supporting
85 Information.

86 A sandy loam soil was obtained from Landlook (Midlands, UK). Prior to use, the soil was air
87 dried, sieved to ≤ 2 mm to ensure homogeneity within the soil matrix and stored at room
88 temperature. Characteristics of the study soil are provided in the Supporting Information.

89 *Eisenia fetida* and *Lumbricus terrestris* were obtained from Blades Biological Ltd. (Kent, UK).
90 The earthworms were cultured in a medium of horse manure and peat (50:50) for *E. fetida*,
91 and in moist soil for *L. terrestris*. They were kept moist with deionized water under laboratory
92 conditions (20 ± 3 °C). The horse manure used in this culture was collected from horses that
93 were not under medication to avoid any toxic effects on the earthworms. *E. fetida* were fed
94 twice weekly with homogenized mashed potato powder which was added to the surface of the
95 culture and *L. terrestris* were fed with dead birch leaves distributed on the surface of the moist
96 soil.

97 *Uptake and depuration studies*

98 Uptake and depuration experiments followed OECD Guideline 317 'Bioaccumulation in
99 Terrestrial Oligochaetes' and used only *E. fetida* (OECD, 2010). Experiments were performed
100 in glass jars at a concentration of 10 µg/g of active ingredient where each jar contained 50 ±
101 1 g of test soil and kept in an incubator at 20 ± 2 °C, using a 16:8 light/dark cycle. Assuming
102 a mixing depth of 20 cm, concentrations expected in the environment from the use of Capture
103 would be expected to range from 35 - 100 µg g⁻¹. The test concentration was one order of
104 magnitude lower than the concentration we used previously to assess the toxicity of the
105 different bifenthrin treatments to *E. fetida*. At 100 µg g⁻¹, no mortality, a slight increase in
106 growth and a small decrease in cocoon production were observed (Anuar, unpublished data).
107 Before the earthworms were exposed to the different treatments, they were acclimated to the
108 experimental conditions in the incubator for 48 h using non-treated soil. The different bifenthrin
109 treatments were then mixed with the soil using deionized water as solvent carrier to achieve
110 a moisture content between 60-70% of the maximum water holding capacity (MWHC). Treated
111 soil was left for 24 h before adding the earthworms.

112 For each bifenthrin treatment (analytical grade, conventional and two nanoformulations), 45
113 glass jars of treated soil were prepared. At the start of the uptake phase, one mature adult *E.*
114 *fetida* with a visible clitellum was added to each glass jar. Glass jars were then covered with
115 garden fleece (to prevent earthworms from escaping while allowing sufficient air supply to be

116 maintained) attached with an elastic band. The uptake phase of the experiment lasted for up
117 to 21 d with triplicate samples being taken at 0 and 6 h and 1, 3, 7, 10, 14, 21 d. *E. fetida* in
118 the remaining glass jars were then transferred to clean soil for up to another 21 d of depuration
119 with samples being taken at 6 h and 1, 3, 7, 10, 14, 21 d after transfer. At each time point in
120 both phases, the earthworm weight and mortality were recorded. Soil moisture content in each
121 glass jars was monitored throughout both phases, and adjusted, where necessary, by adding
122 deionized water so that it remained between 60-70% of the MWHC. The pH of the soils was
123 measured at the beginning and end of the uptake phase and at the end of the depuration
124 phase. Earthworms were fed weekly with mashed potato powder.

125 Once samples were collected, earthworms were removed, rinsed with deionized water, blotted
126 dry, weighed and then placed for 48 h on moist filter papers to allow the earthworms to purge
127 their gut contents (Dalby et al., 1996). The moist filter papers were changed twice a day (in
128 the morning and evening). The earthworms were then frozen prior to analysis. Soil samples
129 were taken for chemical analysis and to extract soil pore water.

130 *Distribution of bifenthrin in earthworms*

131 The distribution of bifenthrin following exposure to the different treatments was assessed using
132 both *E. fetida* and *L. terrestris*. Experiments were performed at the same concentration and
133 conditions as used in the uptake and depuration studies. *E. fetida*, were exposed to 50 ± 1 g
134 of soil treated with each treatment or soil only while *L. terrestris* were exposed to 350 ± 5 g of
135 treated soil or soil only. The duration of the uptake phase was 10 d while the depuration phase
136 lasted for 7 d. There were six replicates per treatment and sampling point. Soil, faeces and
137 earthworm samples were taken at the end of each phase for analysis.

138 The removed earthworms were placed on a dissecting tray with their dorsal side facing
139 upwards. Using a pair of dissecting scissors, an opening cut was made below the clitellum. A
140 straight line cut was made from the opening cut down to the posterior. The cut was made
141 carefully and not too deeply to avoid damage to the internal organs. The skin was pulled apart
142 using forceps and pinned back using dissecting pins. The earthworms were then separated

143 into skin, gut and other tissue (hereafter referred to as 'tissue') for *L. terrestris*. Separation of
144 *E. fetida* tissues proved challenging so it was only possible to separate these samples into gut
145 + tissue and skin. Prior to analysis, samples were washed with distilled water and centrifuged
146 at 3000 rpm for 15 min. Samples and washing water were analysed separately for bifenthrin
147 residues.

148 *Sample extraction and HPLC analysis*

149 Soil (5 ± 0.5 g) was extracted by adding 15 mL acetonitrile and then shaking the mixture on
150 an orbital shaker ($250 \text{ oscillations min}^{-1}$) at room temperature (20 ± 2 °C) for 2 h. Samples
151 were then allowed to settle and 2 mL aliquots of supernatant were taken for analysis. Soil pore
152 water was obtained by placing 10 ± 1 g of soil into a glass syringe with a layer of 3 cm of glass
153 wool inserted into the bottom. The syringe was inserted into a glass centrifuge tube and
154 centrifuged for 20 min at 2016 g to separate soil and soil pore water. The difference in density
155 of the polymer capsules and water is small so centrifugation would not be expected to affect
156 the recovery of the bifenthrin from the nano-treatments compared to the non nano-treatments
157 (Kah et al., 2016).

158 Earthworm samples were homogenized for 5 minutes using a LabGen Series 7 homogenizer
159 with 5 mL of acetonitrile. The suspension was transferred, with rinsing using an additional 5
160 ml of acetonitrile, to a glass vial. The extracts were centrifuged for 20 min at 2016 g. The
161 samples were then filtered using $0.45 \mu\text{m}$ nylon filters and a 2 mL aliquot of the supernatant
162 was taken for further analysis.

163 Soil and earthworm extracts and pore water were analysed using High-performance Liquid
164 Chromatography (HPLC; Perkin Elmer, Flexar) coupled with photodiode array detection. More
165 detail on the methods used are provided in the Supporting Information. The limits of detection
166 and quantification were 1.2 and 3.7 ng mL^{-1} for the analytical grade, 1.5 and 4.7 ng mL^{-1} for
167 the conventional formulation, 1.9 and 5.9 ng mL^{-1} for nano A and 2.1 and 6.5 ng mL^{-1} for nano
168 B. Recoveries for analytical method ranged from 90-107% for water, 88-103% for soil, 84-

169 102% for *E. fetida* and 90-107% for *L. terrestris* and recoveries of the filtration method ranged
170 from 87-100% (see supporting information).

171 *Data analysis*

172 Determination of sorption coefficient, k_d

173 Sorption coefficient, k_d values were calculated at each time point (Equation 1) where: C_{water}
174 and C_{soil} are the concentrations of bifenthrin in soil pore water ($\mu\text{g mL}^{-1}$) and soil ($\mu\text{g g}^{-1}$),
175 MWHC is the maximum water holding capacity of the soil (%), and %water is the moisture
176 content of the soil (%). Averages of k_d -values were then determined.

$$177 \quad k_d = \frac{C_{soil}}{C_{water} * \left(\frac{\%water}{MWHC} \right)} - 1 \quad (1)$$

178 Kinetic modelling

179 We wanted to evaluate whether data on the uptake and depuration characteristics could be
180 used to inform the uptake and depuration of bifenthrin resulting from exposure to Capture LFR
181 and the two nano formulations. Three models were explored with increasing complexity. Data
182 from the analytical grade bifenthrin treatment was always used to parameterise the models.

183 Model 1 was the first order one compartment toxicokinetic model outlined by Ashauer *et al.*
184 (2010) (Equation 2).

$$185 \quad \frac{dC_{organism}}{dt} = k_{in} * C_{water}(t) - k_{out} * C_{organism}(t) \quad (2)$$

186 Where: $C_{organism}$ is the internal concentration ($\mu\text{g g}^{-1}$); C_{water} is the concentration in the pore
187 water ($\mu\text{g mL}^{-1}$); and k_{in} and k_{out} are the uptake rate constant ($\text{mL g}^{-1} \text{h}^{-1}$) and the depuration
188 rate constant (h^{-1}), respectively.

189 Model 2 was designed for estimating uptake of an active ingredient from a nonencapsulated
190 treatment. This model is an adaptation of Model 1 modified to account for the release of
191 bifenthrin from the polymer capsules into the soil pore water (Equation 3).

$$192 \quad \frac{dC_{organism}}{dt} = k_{in} * C_{water_2}(t) - k_{out} * C_{organism}(t) \quad (3.1)$$

193 with

$$194 \frac{dC_{water_2}}{dt} = (C_{water}(t) - C_{water_2}(t)) * k_r \quad (3.2)$$

195 Where: C_{water_2} is the concentration of the compound in the pore water released from the
196 nanoformulation ($\mu\text{g g}^{-1}$) and k_r is the release rate of the nanoformulation (h^{-1}). The release
197 rate can be calculated by comparing the degradation rate of the bifenthrin a.i. with the
198 degradation rate of bifenthrin in the nanoencapsulated formulation (Kah et al., 2016). A full
199 description of the approach for estimating release rate is provided in Kah et al. (2016).

200 Model 3 was used for the distribution studies. This model extends either Model 1 or 2 to
201 account for the distribution of the compound in gut, skin and tissue (Equations 4).

$$202 C_{skin} = C_{organism}(t) * a \quad (4.1)$$

$$203 C_{tissue} = C_{organism}(t) * b \quad (4.2)$$

$$204 C_{gut} = C_{organism}(t) - C_{skin} - C_{tissue} \quad (4.3)$$

205 Where: C_{skin} , C_{gut} and C_{tissue} are the concentration of the compound in skin, gut and tissue
206 ($\mu\text{g g}^{-1}$), and a and b are distribution coefficients between the total internal concentration and
207 the skin, the total internal concentration and the tissue. The distribution coefficients are
208 obtained using studies on analytical grade a.i..

209 Statistical analysis

210 Statistical analysis was performed using SigmaPlot (Version 13.0; Systat Software, San Jose,
211 CA). Data were tested performing one-way - or two-way- Analysis of Variance (ANOVA) via
212 the Holm-Sidak pairwise comparison method with the Shapiro-Wilk test for normality of data
213 and the Brown-Forsythe test for equal variance of data. Modelling was conducted in
214 OpenModel V 2.4.2. (<http://openmodel.info/>) using the Runge-Kulta (4th Order) ordinary
215 differential equation method with Monte Carlo simulations to obtain the 95% confidence
216 interval and the Nash–Sutcliffe Efficiency calculation for goodness of fit indication. Nash–

217 Sutcliffe values (hereafter called Nash index) between 0 and 1 represent an acceptable fit of
218 the model to the data.

219 **Results and discussion**

220 Fate of bifenthrin in soil

221 Generally, throughout the uptake phase, there was a decrease in concentration of bifenthrin
222 in the soil and soil pore water which was associated with an increase in the concentration of
223 bifenthrin in the earthworms (Data are summarised in the Supporting information). At the end
224 of the uptake phase, 65-82% of bifenthrin was extractable and associated with the soil
225 particles, 16-33% had dissipated/degraded, around 1% was present in the pore water and <
226 1% was taken up by the earthworms (Figure 1 in the Supporting Information). Apparent half-
227 lives for bifenthrin in the different treatments increased in the order analytical grade bifenthrin
228 – Capture LFR – Nano B – Nano A. The observed DT_{50} for the analytical grade bifenthrin is at
229 the lower end of the values reported in the field for bifenthrin and is lower than reported in
230 laboratory studies (Pesticide Properties Database, 2017). Half-lives are also lower than those
231 obtained by Kah *et al.* (2016) in similar investigations into the differences in persistence of
232 analytical grade bifenthrin and the a.i. in Capture LFR and nanoencapsulated treatments,
233 although the order of half-lives is the same (Kah *et al.*, 2016). Release rates and associated
234 release times (RT_{50}) for the nanoformulations (Table 1) are lower than those found by Kah *et al.*
235 *et al.* (2016). Overall, these results indicate that even traditional formulations can affect the
236 persistence of an active ingredient but this impact is more enhanced in the nano-encapsulated
237 treatments, possibly due to the nanocapsules ‘shielding’ the unreleased a.i. from the
238 degrading microbes.

239 Sorption coefficients (k_d), based on the soil and soil-pore water concentrations ranged from
240 154 to 585 L kg⁻¹ and increased significantly in the order Nano A = Nano B < Capture LFR <
241 analytical grade bifenthrin (Table 1; Two-Way ANOVA Holm-Sidak method $P < 0.001$).
242 Sorption coefficients are lower than previously reported for the a.i. which range from 882 to
243 6000 mL g⁻¹ in different soil types (Pesticides Property Database, 2017). Sorption coefficients

244 are also lower than those observed by Kah et al. for bifenthrin a.i. and bifenthrin in traditional
245 and nanoencapsulated formulations (Kah et al., 2016). The mismatch is possibly explained by
246 the fact that we derived k_d values based on pore water measurements, which is arguably more
247 realistic than the batch equilibrium approach employed in previous studies. The differences
248 might be explained by dissolved organic carbon in the pore water which may act as an
249 additional sink for the bifenthrin or due to differences in the nature of the organic carbon in the
250 soils used in the different studies.

251 The observations for Capture LFR demonstrate that even traditional co-formulants can affect
252 the distribution of the bifenthrin in soils although the effect is more enhanced in the nano-
253 encapsulated treatments. Other studies have explored the effects of formulation on pesticide
254 behaviour. In studies with chlorsulfuron, co-formulants reduced sorption (Foldenyi et al., 2013)
255 while studies with propyzamide (Khan and Brown, 2017) showed sorption to increase and
256 studies with triticonazole, cyprodinil, propetamphos and fludioxinil showed sorption to
257 increase (Beigel et al., 1998; Beigel and Barriuso, 2000; Garcia-Ortega et al., 2006; Pose-
258 Juan et al., 2011). The impacts of co-formulants therefore likely depend on the active
259 ingredient and the nature of the co-formulants used in a product. The observed reduction in
260 the k_d values for the nano-encapsulated materials is likely due to a combination of the co-
261 formulant effects and the fact that the polymer capsule shielded the bifenthrin from sorption
262 sites on the soil surface.

263 Uptake and depuration behaviour

264 *Uptake and depuration in E. fetida*

265 No mortality was recorded and the studies passed the validity criteria (based on earthworm
266 growth and mortality) according to the principles outlined in the OECD 317 (OECD, 2010).
267 Lower uptake and depuration was seen for bifenthrin in the analytical and Capture LFR
268 treatments compared to the two nanoformulation treatments (Figure 1). At the end of the
269 depuration phase, in the analytical grade and Capture LFR treatments, bifenthrin was still
270 detectable in the earthworms whereas for the two nanoformulation treatments, it was not

271 detectable. The pattern of uptake and depuration between the non-nano and nano treatments
272 was also different. The non-nano exposures were characterised by a steady uptake and
273 elimination of bifenthrin over time whereas in the nano treatments, an initial rapid period of
274 uptake or elimination was observed and this then tailed off (Figure 1).

275 The first order one-compartment model (Model 1) was successfully fitted to the data from the
276 analytical grade bifenthrin treatment (Nash index = 0.94) obtaining an uptake- and depuration
277 - rate constant of $0.222 \pm 0.009 \text{ mL g}^{-1} \text{ h}^{-1}$ and $0.0036 \pm 0.0002 \text{ h}^{-1}$. Use of the uptake and
278 depuration rates in the model to simulate the uptake and depuration bifenthrin from the
279 Capture LFR formulation worked well (Nash index = 0.68) but failed to acceptably simulate
280 the uptake of Nano A and Nano B (Nash index < -0.01) (Figure 1). The results for the Capture
281 modelling do, however, suggest that it may be possible to extrapolate from studies into the
282 uptake of analytical grade materials to estimate uptake of a.i.'s from traditional formulations.

283 Model 2, which incorporates the release rate of bifenthrin from the nanocapsule,
284 underestimated uptake and depuration of the bifenthrin from the two nano treatments (Nash
285 index < 0; Figure 1). Closer inspection of the simulation however revealed that this model
286 more accurately simulated the internal concentration at the end of the depuration phase. The
287 differences in kinetic patterns and model fits suggests that the nanoencapsulated bifenthrin
288 was accumulated *via* a different mechanism than in the analytical grade material and Capture
289 LFR treatments. As previous studies with earthworms have shown that other nanoparticles
290 accumulate in the earthworm gut rather than the actual tissue (Unrine et al., 2010; Waissi-
291 Leinonen 2012), we performed studies to explore whether there were any differences in the
292 distribution of the bifenthrin in earthworms exposed to the different treatments. Here, we not
293 only used *E. fetida* but also *L. terrestris* due to its larger size and consequent ease of handling.

294 *Distribution studies*

295 Concentrations of bifenthrin in *L. terrestris* were significantly greater than in *E. fetida* for all
296 treatments ($P < 0.001$). Within each species, uptake was significantly different between the
297 nano formulated and non-nano treatments, while within the non-nano and nano treatments no

298 significant difference in uptake was observed ($P < 0.001$). These differences in uptake might
299 be explained by differences in the way the two earthworm species process soil organic matter,
300 their ecological strategy and/or lipid content (Kelsey et al., 2005). *Eisenia fetida* is smaller than
301 *L. terrestris* and is an epigeic species living primarily at or near the soil surface and consumes
302 coarse particulate organic matter and surface litter. *L. terrestris* is an anecic species that lives
303 in deep burrows and comes to surface to feed on surface litter [Bouche, 1983]. Interestingly,
304 the interspecies difference in uptake that we see is the opposite to that observed in a similar
305 study using pharmaceuticals covering a range of physico-chemical properties (Carter et al.,
306 2016).

307 Significant differences were also seen in the depuration of bifenthrin by the two species ($P <$
308 0.001). For the analytical grade and Capture LFR treatments, *L. terrestris* still contained 57 -
309 59% of the accumulated bifenthrin after the 7 d depuration phase while concentrations in
310 *E. fetida* were significantly lower (43-47%; $P < 0.001$). Depuration of bifenthrin from the two
311 nano treatments was faster with *L. terrestris* containing 20-22% of accumulated bifenthrin after
312 7 d depuration while *E. fetida* contained only 10-13% of the accumulated mass. Bifenthrin from
313 the nanoformulations was eliminated 2.8 and 4 times more quickly than from the non-nano
314 treatments ($P < 0.001$).

315 In *L. terrestris*, following the uptake phase, concentrations of bifenthrin from the analytical
316 grade and Capture LFR treatments was significantly higher ($P < 0.01$) in the tissue compared
317 to the gut and skin which had similar ($P = 0.6$) bifenthrin concentrations (Figure 2). In contrast,
318 for the nanoformulation treatments significantly ($P < 0.001$) higher concentrations were
319 observed in the gut of the earthworms compared to the skin and tissue. Concentrations in skin
320 and tissue were also significantly different ($P < 0.001$). The concentration in the gut of the
321 nano- exposed animals was significantly higher ($P < 0.001$) than the non-nano exposed
322 earthworms even though the concentration of bifenthrin in the soil was the same. A significant
323 difference was also observed between the two nanoformulations ($P = 0.019$). This might
324 indicate that the earthworms are selectively consuming the polymer capsules and/or that the

325 capsules are becoming 'trapped' in the gut of the animal and are eliminated more slowly than
326 the bulk soil. This seems to be nanoformulation specific. While it is not known whether
327 earthworms are able to select finer material from coarser particles, there is a body of evidence
328 indicating that they do exhibit preferences for different food types (Curry and Schmidt, 2007).

329 At the end of the 7 d depuration phase, for the analytical grade and Capture LFR treatments,
330 highest concentrations of bifenthrin in *L. terrestris* were seen in the tissue while for the
331 nanoformulation treatments highest concentrations were seen in the gut (Figure 2). The
332 observation that nanopesticide treatments result in highest bifenthrin concentrations in the gut
333 are similar to findings from a previous study into the uptake and distribution of C60 and Au
334 nanoparticles into earthworms (Unrine et al., 2010; Waissi-Leinonen 2012; Petersen et al.,
335 2008, 2011). We found that gut associated bifenthrin was generally less eliminated *via* the gut
336 for the non-nanoformulation treatments compared to the nanoformulated treatments (Figure
337 3). Furthermore, a temporal shift in elimination of the nanoformulated bifenthrin occurred. Gut
338 associated elimination of bifenthrin was greatest for the non-nanoformulation treatments whilst
339 the earthworms were still in soil during the elimination phase of the experiment whilst
340 elimination for the nanoformulated treatments was greatest when the organisms were on the
341 filter paper after the elimination phase of the experiment.

342 Unfortunately, it was not practically possible to separate out the internal organs of *E. fetida*
343 from the tissue so we could only distinguish between bifenthrin in the skin and in tissue
344 combined with the gut (Figure 2). For the analytical grade and Capture LFR treatments,
345 concentrations in the gut + tissue and in the skin were significantly different after the uptake
346 phase (Figure 2; $P < 0.001$) with higher concentrations being seen in the skin. For the
347 nanoformulation treatments, concentrations of bifenthrin accumulated in the gut combined
348 with the internal organs were significantly higher than in the skin ($P < 0.001$). Following the
349 depuration phase, concentrations of bifenthrin in the gut + tissue and skin in all treatments
350 were similar (Figure 2) and not significantly different within all treatment ($P > 0.1$) except for

351 the analytical grade treatment ($P = 0.003$). As we were unable to fully characterise the
352 distribution of the a.i. in *E. fetida* our modelling efforts focused on the *L. terrestris* studies.

353 Model 1, and combinations of models 1 and 3 and models 2 and 3 were used to simulate
354 uptake and depuration in the different *L. terrestris* treatments (Figure 4). Model 1 performed
355 very well for estimating the concentrations of the a.i. in the whole organism for the analytical
356 grade and Capture LFR treatments (Nash Index > 0.90). Predictions for the nano treatments
357 were also good (Nash Index > 0.48) although the model overestimated whole organism
358 concentrations at the end of the depuration phase (Figure 4). When model 1 was combined
359 with model 3 to simulate distribution of the a.i. between gut- skin- and remaining-tissues, good
360 predictions were obtained for the analytical grade and Capture LFR treatments for all tissues
361 (Nash Index > 0.76) and for the skin in the two nano treatments (Nash Index > 0.39).
362 Underestimates of concentrations in the gut and overestimates of concentrations in the tissue
363 by a factor of 8-11 were obtained using a combination of models 1 and 3 for both nano
364 treatments (Nash Index < 0.05). The fact that the two models worked well for estimating
365 behaviour in the Capture treatment is encouraging and suggests that estimates of uptake and
366 distribution based on analytical grade material can be used to extrapolate to behaviours in
367 traditional formulations. The approach worked less well for the nanoformulations so we then
368 extended the modelling to factor in the effect of the release rate from the capsule.

369 Incorporation of the a.i. release from the nanocapsule into the modelling of the
370 nanoformulations (i.e. model 2 and model 3 were used) resulted in predictions that fitted the
371 whole organism data and the skin data well (Nash Index > 0.58). This approach
372 underestimated concentrations in the gut at the end of the uptake phase while predictions of
373 concentrations at the end of the depuration phase were close to the measured data. This is a
374 direct result of the model assumption that compound distribution between different tissues
375 (skin, gut and remaining tissue) is instantaneous and fixed by distribution factors and a
376 temporal change in gut clearance between nano and non-nano formulations (Figure 3).
377 Nonetheless, inclusion of the release rate resulted in better predictions, compared to the

378 approach not considering release, of internal tissue concentrations of the a.i. with
379 concentrations being a factor of 3.8 (nano A) and 5.1 (nano B) of measured data at the end of
380 the uptake phase and within a factor of 5.7 (nano A) to 7.5. (nano B) at the end of the
381 depuration phase. While the predictions were not perfect, these results indicate that to model
382 internal tissue exposure, which will likely represent the toxicologically important fraction of the
383 accumulate a.i., it is necessary to factor in the release rate of the a.i. from the nanocapsule
384 into the toxicokinetic modelling.

385 Implications for risk and a potential modelling approach

386 We have demonstrated that nanoencapsulation will affect the behaviour and uptake of
387 pesticides in soil. For bifenthrin, it will decrease the sorption of the active ingredient to soils,
388 increase the apparent persistence of the compound and alter the uptake behaviour of the
389 active ingredient into earthworms and the subsequent distribution. Consequently, the risk of
390 nanoencapsulated bifenthrin to earthworms will be different from a conventional product.
391 Whether nanoencapsulation increases, decreases or has no effect on risk is difficult to
392 establish at this stage. While nanoencapsulated bifenthrin is taken up more quickly by the
393 earthworms, from the *L. terrestris* studies, it appears that the majority of the bifenthrin taken up
394 is contained in the gut so the internalised concentration is lower than in earthworms exposed
395 to the analytical grade substance and a conventional formulation. If less active ingredient is
396 internalised, one would assume that less of the active ingredient will reach the site of toxic
397 action so the effects of the nanoformulation will be lower. However, nanoencapsulation also
398 increases the apparent persistence of the active ingredient which will lengthen the exposure
399 duration of the earthworms to the active ingredient in the nanoformulation compared to a
400 conventional ingredient. The increased efficacy of the nanoformulation compared to
401 conventional formulations could mean that application rates to field are decreased which will
402 also affect risk. The risks to birds and mammals feeding of earthworms could also be altered.
403 If a nanoformulation is applied at the same rate as a conventional product then the oral
404 exposure of these organisms will increase but differences in the bioaccessibility of

405 nanoencapsulated bifenthrin compared to free bifenthrin could be lower meaning less is
406 internalised. Again the duration of exposure will increase.

407 To answer some of these questions around the implications of changes in fate and uptake or
408 effects, a toxicokinetic toxicodynamic modelling approach is probably required (Ashauer and
409 Escher, 2010). In Figure 5, we present a conceptual model, based on our experimental
410 findings and investigations into the performance of the different toxicokinetic modelling
411 approaches, that could be used to model the toxicokinetics of a nanoencapsulated active
412 ingredient. The model assumes sorption to soil is instantaneous following release of pesticide
413 from the capsule and that it is the free (i.e. dissolved pore water) pesticide that is taken up into
414 the earthworm tissue – this assumption is supported by our distribution studies in *L. terrestris*
415 and the testing of Models 2 and 3. The internal concentration in the earthworm tissue over
416 time, needed for toxicokinetic toxicodynamic modelling of the effects, are then calculated
417 based on the release rate from the capsule, the soil-water distribution coefficient and the
418 uptake and depuration rates of the free active ingredient into/out of the earthworm. To estimate
419 oral exposure of birds and mammals, the mass concentration of the active ingredient in the
420 gut also needs to be considered and this is estimated based on the feeding rate of the
421 earthworm on whole soil and on the nanoparticles.

422 We believe that this conceptual model is a useful first step towards developing improved
423 environmental risk assessment approaches for estimating the uptake and effects of
424 nanoencapsulated pesticides in earthworms. The approach might also be applicable to other
425 materials (e.g. nanoencapsulated pharmaceuticals) and other organisms. In the future, we
426 recommend that the model be further parameterised for bifenthrin. We also recommend that
427 studies of the type reported here are done on a wider range of organisms using other pesticide
428 active ingredients with different persistence and physico-chemical properties contained in a
429 wider range of nanocarrier materials in order to evaluate the broader applicability of the model.

430 **Acknowledgements**

431 We would like to thank Dr. Darren Anderson from Vive Crop Protection Inc. who provided the
432 nanoencapsulated test materials. MFMA was funded by the Ministry of Higher Education of
433 Malaysia. AAs contribution to the study received support from the Innovate UK-funded VFETL
434 project.

435 **Data Availability**

436 The experimental data on which this manuscript is based can be obtained, on request, from
437 the corresponding author.

438 **References**

439 Ashauer R, Caravatti I, Hintermeister A, Escher BI. 2010. Bioaccumulation kinetics of organic
440 xenobiotic pollutants in the freshwater invertebrate *Gammarus Pulex* modelled with prediction
441 intervals. *Environ. Toxicol. Chem.* 29(7):1625 – 1636.

442 Ashauer R, Escher BI. 2010. Advantages of toxicokinetic and toxicodynamic modelling in
443 aquatic ecotoxicology and risk assessment. *J. Environ. Monit.* 12:2056 - 2061.

444 Beigel C, Barriuso E, Calvet R. 1998. Sorption of low levels of non-ionic and anionic
445 surfactants on soil: effects on sorption of triticonazole fungicide. *Pest. Sci.* 54:52-60.

446 Beigel, C. Barriuso, E. Influence of formulation on triticonazole solubilisation and sorption in a
447 soil-water system. *Pest. Manag. Sci.* **2000**, 56: 271-276.

448 Bouche A. 1983. The establishment of earthworm communities. In: Satchell JE. (ed)
449 Earthworm ecology: from Darwin to vermiculture. Chapman and Hall, London, pp 431–438

450 Carter LJ, Ryan JJ, Boxall ABA. 2016. Does Uptake of pharmaceuticals vary across
451 earthworm species? *Bull. Environ. Contam. Toxicol.* 97(3):316-322.

452 Curry JP, Schmidt O. 2007. The feeding ecology of earthworms: a review. *Pedobiologia*
453 50:463-477.

454 Dalby PR, Baker GH, Smith SE. 1996. Filter paper method to remove soil from earthworm
455 intestines and to standardise the water content of earthworm tissue. *Soil Biol. Biochem.* 28:
456 685 – 687.

457 Foldenyi R, Toth Z, Samu G, Ersek C. 2013. Study of sorption of two sulfonylurea type
458 herbicides and their additives on soils and soil components. *J. Environ. Sci. Health B.* 48:758-
459 766.

460 Garcia-Ortega S, Holliman PJ, Jones DL. 2006. Toxicology and fate of Pestanal and
461 commercial propetamphos formulations in river and estuarine sediment. *Sci. Tot. Environ.*
462 366:826-836.

463 Heikens A, Peijnenburg W, Hendriks AJ. 2001. Bioaccumulation of heavy metals in terrestrial
464 invertebrates. *Environ. Pollut.* 113:385 – 393.

465 Kah M, Beulke S, Tiede K, Hofmann T. 2013. Nanopesticides: state of knowledge,
466 environmental fate, and exposure modelling. *Crit. Rev. Environ. Sci. Technol.* 43:1823-1867.

467 Kah M, Weniger A-K, Hofmann T. 2016. Impacts of (nano)formulations on the fate of an
468 insecticide in soil and consequences for environmental exposure assessment. *Environ. Sci.*
469 *Technol.* 50:10960-10967.

470 Kah M. 2015. Nanopesticides and nanofertilizers: emerging contaminants or opportunities for
471 risk mitigation. *Frontiers Chem.* 3(64):1-6.

472 Kelsey JW, Colino A, White JC. 2005. Effect of species differences, pollutant concentration,
473 and residence time in soil on the bioaccumulation of 2,2-bis(p-chlorophenyl)-1,1-
474 dichloroethylene by three earthworm species. *Environ. Toxicol. Chem.* 24:703–708.

475 Kelsey JW, White JC. 2013. Effect of C60 fullerenes in the accumulation of weathered p,p'-
476 DDE by plant and earthworm species under single and multispecies conditions. *Environ.*
477 *Toxicol. Chem.* 32(5):1117-1123.

478 Khan MA, Brown CD. 2017. Influence of commercial formulation on the sorption and leaching
479 behaviour of propyzamide in soil. *Sci. Tot. Environ.* 578:158-166.

480 Kookana RS, Boxall ABA, Reeves PT, Ashauer R, Beulke S, Chaudhry Q, Cornelis G,
481 Fernandes TF, Gan J, Kah M, Lynch I, Ranville J, Sinclair C, Spurgeon D, Tiede K, Van Den
482 Brink PJ. 2014. Nanopesticides: Guiding principles for regulatory evaluation of environmental
483 risks. *J. Agric. Food Chem.* 62 (19):4227 – 4240.

484 Kwak JI, Youn-Joo A. 2005. Ecotoxicological Effects of Nanomaterials on Earthworms: A
485 Review. *Human Ecol. Risk Ass.* 21(6):1566-1575.

486 Langdon, CJ, Hodson ME, Arnold RE, Black S. 2005. Survival, Pb-uptake and behaviour of
487 three species of earthworm in Pb treated soils determined using an OECD-style toxicity test
488 and a soil avoidance test. *Environ. Pollut.* 138:368 – 375.

489 Li D, Fortner JD, Johnson DR, Chen C, Li Q, Alvarez PJJ. 2010. Bioaccumulation of ¹⁴C60
490 by the Earthworm *Eisenia fetida*. *Environ. Sci. Technol.* 44(23):9170-9175.

491 Matscheko N, Tysklind M, De Wit C, Bergek S, Andersson R, Sellstrom U. 2002. Application
492 of sewage sludge to arable land-soil concentrations of polybrominated diphenyl ethers and
493 polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls, and their accumulation in
494 earthworms. *Environ. Toxicol. Chem.* 21:2515 – 2525.

495 OECD. 2010. Guidelines for the Testing of Chemicals. Test No 317: Bioaccumulation in
496 Terrestrial Oligochaetes. OECD, Paris.

497 Petersen EJ, Huang Q, Weber JWJ. 2008. Bioaccumulation of Radio-Labeled Carbon
498 Nanotubes by *Eisenia foetida*. *Environ. Sci. Technol.* 42(8):3090-3095.

499 Petersen EJ, Pinto RA, Zheng L, Huang Q, Landrum PF, Weber WJ Jr. 2011. Effects of
500 Polyethyleneimine-Mediated Functionalization of Multi-Walled Carbon Nanotubes on
501 Earthworm Bioaccumulation and Sorption by soils. *Environ. Sci. Technol.* 45(8):3718-3724.

502 Pesticide Properties Database. Bifenthrin (Ref: FMC 54800). Accessed 3 August 2017 at
503 <http://sitem.herts.ac.uk/aeru/ppdb/en/Reports/78.htm>.

504 Pose-Juan E, Rial-Otero R, Paradelo M, Lopez-Periago JE. 2011. Influence of the adjuvants
505 in a commercial formulation of the fungicide 'switch' on the adsorption of their active
506 ingredients: cyprodinil and fludioxinil, on soils devoted to vineyard. *J. Hazard. Mater.* 193:288-
507 295.

508 Scott-Fordsmand JJ, Krogh PH, Schaefer M, Johansen A. 2008. The toxicity testing of double-
509 walled nanotubes-contaminated food to *Eisenia veneta* earthworms. *Ecotox. Environ. Safety*
510 71(3):616-619.

511 Spurgeon DJ, Hopkin SP. 1996. Risk assessment of the threat of secondary poisoning by
512 metals to predators of earthworms in the vicinity of a primary smelting works. *Sci. Total*
513 *Environ.* 187:167 – 183.

514 Unrine JM, Hunyadi SE, Tsyusko OV, Rao W, Shoults-Wilson WA, Bertsch PM. 2010.
515 Evidence for bioavailability of Au nanoparticles from soil and biodistribution within earthworms
516 (*Eisenia fetida*). *Environ. Sci. Technol.* 44:8308 – 8313.

517 Waissi-Leinonen GC, Petersen EJ, Pakarinen K, Akkanen J, Leppanen MT, Kukkonen, JVK.
518 2012. Toxicity of fullerene (C60) to sediment-dwelling invertebrate *Chironomus riparius* larvae.
519 *Environ. Toxicol. Chem.* 31(9):2108-2116.

520

521 Table 1: Sorption coefficients (k_d), dissipation half-lives (DT50), release half times (RT50),
 522 bioconcentration factors (BCFs) and rates for release, uptake and depuration (k) for the
 523 different bifenthrin treatments studied in the *Eisenia fetida* and *L. terrestris* studies.

Endpoint	Study	Unit	Bifenthrin	Capture LFR	Nano A	Nano B
k_d	Uptake and depuration	L Kg ⁻¹	550 ± 21	394 ± 15	186 ± 4	233 ± 17
k_d	Distribution study <i>E.fetida</i>	L Kg ⁻¹	494 ± 71	371 ± 47	154 ± 7	163 ± 9
k_d	Distribution study <i>L. terrestris</i>	L Kg ⁻¹	585 ± 92	488 ± 32	274 ± 42	251 ± 31
DT ₅₀	Uptake and depuration	d	25 - 27	33-35	49 - 50	38 - 40
Release rate (k_r)	Uptake and depuration	h ⁻¹	NA	NA	0.104 ± 0.008	0.182 ± 0.016
RT ₅₀	Uptake and depuration	d	NA	NA	6 - 7	3 - 4
k_{in} <i>E.fetida</i>	Uptake and depuration	LKg ⁻¹ h ⁻¹	0.222 ± 0.009	NA	NA	NA
k_{out} <i>E.fetida</i>	Uptake and depuration	h ⁻¹	0.0036 ± 0.0002	NA	NA	NA
BCF <i>E.fetida</i>	Uptake and depuration	-	61.7	NA	NA	NA
k_{in} <i>L. terrestris</i>	Uptake and depuration	LKg ⁻¹ h ⁻¹	0.7021 ± 0.0336	NA	NA	NA
k_{out} <i>L. terrestris</i>	Uptake and depuration	h ⁻¹	0.0033 ± 0.0003	NA	NA	NA
BCF <i>L. terrestris</i>	Uptake and depuration	-	212	NA	NA	NA

524 ¹ Determined from data presented here with additional data on soil concentrations over time in four other soils
 525 (unpublished).

526

Figure Legends

Figure 1: Measured (dots) and predicted (lines) total internal concentration of different formulations of bifenthrin in *E. fetida* over time using Model 1 (black) and Model 2 (grey). Dotted lines indicate the 95% confidence interval for the predictions. Model parameterisation was conducted with data from bifenthrin a.i..

Figure 2: Proportion of total internal concentration of different formulations of bifenthrin in earthworms after 10 d uptake and 7 d depuration as average \pm SD in relation to the internal concentration at the end of the uptake phase.

Figure 3: Proportion of the gut concentration at the end of the uptake phase of different formulations of bifenthrin recovered from *L. terrestris* faeces samples as average \pm SD.

Figure 4: Measurements (dots) and predictions from different model combinations (lines) of internal concentration of different formulations of bifenthrin in whole organism and different compartments of *L. terrestris* over time. Grey backgrounds indicate that predictions account for the released fraction of bifenthrin from the nanoformulation. Dotted lines indicate the 95% confidence intervals for the predictions.

Figure 5: Conceptual model for estimating residues of active ingredients contained in nanoencapsulated formulations in terrestrial invertebrates over time.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5.

