

This is a repository copy of Multisystemic therapy versus management as usual in the treatment of adolescent antisocial behaviour (START): A pragmatic, randomised controlled, superiority trial.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/126503/

Version: Accepted Version

## Article:

Fonagy, P, Butler, S, Cottrell, D orcid.org/0000-0001-8674-0955 et al. (12 more authors) (2018) Multisystemic therapy versus management as usual in the treatment of adolescent antisocial behaviour (START): A pragmatic, randomised controlled, superiority trial. The Lancet Psychiatry, 5 (2). pp. 119-133. ISSN 2215-0366

https://doi.org/10.1016/S2215-0366(18)30001-4

© 2018 Elsevier Ltd. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International http://creativecommons.org/licenses/by-nc-nd/4.0/

## Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

## **Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Table 5: Secondary outcomes: A. Young people's and parent's report on parenting skills and family functioning. B. Parents' report on family functioning.

A.

	Group and between- group significance (t-test)	APQ Problems of monitoring and supervision (YP) Mean (SD) [n]	APQ Problems of monitoring and supervision (P) Mean (SD) [n]	Loeber parental support score (P) Mean (SD) [n]	Level of Expressed Emotion (YP) Mean (SD) [n]
6 months-baseline	MST	8·4 (3·0) [n=341]	9·29 (3·33) [n=341]	44·44 (6·40) [n=337]	88·8 (20·0) [n=341]
	MAU	8·8 (2·8) [n=339]	9·37 (3·34) [n=339]	44·57 (6·04) [n=335]	89·1 (19·1) [n=339]
6-month follow-up	MST	7·7 (2·9) [n=292]	7·7 (3·2) [n=292]	47·7 (5·7) [n=288]	83·1 (18·9) [n=292]
	MAU	7·9 (2·9) [n=261]	8·5 (3·4) [n=268]	45·4 (6·6) [n=262]	85·3 (18·3) [n=268]
	Effect (95% CI)	-0.12 (-0.59  to  0.35)	-0.71 (-1.20  to  -0.22)	2.05 (1.09  to  3.01)	-2.58 (-5.32  to  0.16)
	p value	0.62	0.0039	<0.0001	0.065
12-month follow-up	MST	7·7 (3·1) [n=246]	7·8 (3·3) [n=248]	46·7 (6·4) [n=243]	81·4 (19·7) [n=248]
	MAU	7·9 (3·1) [n=233]	8·1 (3·3) [n=238]	45·5 (6·4) [n=230]	82·3 (17·3) [n=238]
	Difference (95% CI)	0.02 (-0.49  to  0.53)	-0·10 (-0·61 to 0·41)	0.88 (-0.14 to 1.90)	-0.82 (-3.72  to  2.08)
	p value	0.94	0.72	0.093	0.58
18-month follow-up	MST	7·6 (3·1) [n=235]	7·7 (3·2) [n=234]	46·0 (7·1) [n=231]	78·7 (19·3) [n=234]
	MAU	7·7 (3·0) [n=206]	7·7 (3·4) [n=217]	44·9 (6·8) [n=215]	79·9 (18·6) [n=217]
	Difference (95% CI)	0.14 (-0.39  to  0.67)	0·11 (-0·42 to 0·64)	0.76 (-0.30  to  1.82)	-0.98 (-3.96  to  2.00)
	p value	0.61	0.70	0.16	0.52

Data were obtained using the Alabama Parenting Questionnaire (APQ), Loeber Caregiver Questionnaire (Loeber), and Level of Expressed Emotion. MAU=management as usual. MST=Multisystemic Therapy. YP=completed by young person. P=completed by parent. Other APQ scales are reported in the appendix.

B.

	Group (n) and between-group significance (t-test)	FACES-IV family satisfaction Mean (SD)	FACES-IV cohesion Mean (SD)	FACES-IV family communication Mean (SD)	CTS Mean (SD)
6 months-baseline	MST (n=337)	27.94 (8.73)	53·36 (22·54)	34.24 (8.11)	8.90 (9.87)
	MAU (n=335)	28.24 (9.09)	53·52 (24·21)	34-22 (8-55)	8.77 (9.72)
6-month follow-up	MST (n=288)	33.7 (8.0)	61.3 (18.7)	37.4 (7.2)	7-1 (9-9)
	MAU (n=262)	30.0 (9.1)	55.7 (22.1)	35.0 (8.4)	8.0 (8.1)
	Effect (95% CI)	3.85 (2.60 to 5.10)	5·80 (2·49 to 9·11)	2.60 (1.48 to 3.72)	-1.13 (-2.85  to  0.59)
	p value	< 0.0001	0.00059	< 0.0001	0.20
12-month follow-up	MST (n=243)	32.6 (8.6)	59.5 (20.8)	37.3 (7.0)	6.1 (9.5)
	MAU (n=230)	30.5 (8.9)	56.5 (22.3)	36.2 (8.1)	6.6 (7.1)
	Difference (95% CI)	2·43 (1·10 to 3·76)	3·27 (-0·20 to 6·74)	0.99 (-0.21 to 2.19)	0·10 (-1·74 to 1·94)
	p value	0.00037	0.065	0-11	0-92
18-month follow-up	MST (n=231)	32.5 (8.2)	59.7 (20.2)	37.7 (6.9)	5.0 (8.7)
	MAU (n=215)	32.0 (9.5)	59.5 (21.5)	37.0 (8.1)	4.8 (5.5)
	Difference (95% CI)	0·13 (-1·24 to 1·50)	0.80 (-2.79 to 4.39)	0.59 (-0.64 to 1.82)	-0.06 (-1.94 to 1.82)
	p value	0.85	0.66	0.35	0.95

Data were obtained using the Family Adaptability and Cohesion Evaluation Scale (FACES-IV) and the Conflict Tactics Scale (CTS). MAU=management as usual. MST=Multisystemic Therapy.