

This is a repository copy of *Fragon: rapid high-resolution structure determination from ideal protein fragments*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/127271/>

Article:

Jenkins, Huw Thomas orcid.org/0000-0002-3302-6966 (2018) *Fragon: rapid high-resolution structure determination from ideal protein fragments*. *Acta crystallographica. Section D, Structural biology*. pp. 205-214. ISSN 2059-7983

<https://doi.org/10.1107/S2059798318002292>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Fragon: Rapid high-resolution structure determination from ideal protein fragments

Huw T. Jenkins^{a*}

^aYork Structural Biology Laboratory, Department of Chemistry, The University of York, Heslington, York, YO10 5DD, UK

Correspondence email: huw.jenkins@york.ac.uk




Figure 1 Test cases in the mixed α/β test set solved by *Fragon*. The fragment placed by *Phaser* is shown as yellow sticks.




Figure 2 Test cases in the all- β test set solved by *Fragon*. The fragment placed by *Phaser* is shown as yellow sticks.