

This is a repository copy of *Disruption of the GDP-mannose synthesis pathway in Streptomyces coelicolor results in antibiotic hyper-susceptible phenotypes.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/128230/>

Version: Accepted Version

---

**Article:**

Howlett, Robert, Anttonen, Katri, Read, Nicholas et al. (1 more author) (2018) Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor* results in antibiotic hyper-susceptible phenotypes. *Microbiology* (Reading, England). ISSN 1465-2080

<https://doi.org/10.1099/mic.0.000636>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Microbiology

## Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor* results in antibiotic hyper-susceptible phenotypes

--Manuscript Draft--

<b>Manuscript Number:</b>	MIC-D-18-00002R1
<b>Full Title:</b>	Disruption of the GDP-mannose synthesis pathway in <i>Streptomyces coelicolor</i> results in antibiotic hyper-susceptible phenotypes
<b>Article Type:</b>	Research Article
<b>Section/Category:</b>	Physiology and metabolism
<b>Corresponding Author:</b>	Margaret CM Smith, PhD University of Leeds Leeds, West Yorkshire UNITED KINGDOM
<b>First Author:</b>	Robert Howlett, PhD
<b>Order of Authors:</b>	Robert Howlett, PhD Katri Anttonen, PhD Nicholas Read, PhD Margaret CM Smith, PhD
<b>Abstract:</b>	<p>Actinomycete bacteria use polyprenol phosphate mannose as a lipid linked sugar donor for extra-cytoplasmic glycosyl transferases that transfer mannose to cell envelope polymers, including glycoproteins and glycolipids. We showed recently that strains of <i>Streptomyces coelicolor</i> with mutations in the <i>ppm1</i> gene, which encodes polyprenol phosphate mannose synthase, were both resistant to phage <math>\phi</math>C31 and have greatly increased susceptibility to antibiotics that mostly act on cell wall biogenesis. Here we show that mutations in the genes encoding enzymes that act upstream of Ppm1 in the polyprenol phosphate mannose synthesis pathway can also confer phage resistance and antibiotic hyper-susceptibility. GDP-mannose is a substrate for Ppm1 and is synthesized by GDP-mannose pyrophosphorylase (GMP;ManC) which uses GTP and mannose-1-phosphate as substrates. Phosphomannomutase (PMM;ManB) converts mannose-6-phosphate to mannose-1 phosphate. <i>S. coelicolor</i> strains with knocked down GMP activity or with a mutation in <i>sco3028</i> encoding PMM acquire phenotypes that resemble those of the <i>ppm1</i>- mutants i.e. <math>\phi</math>C31 resistant and susceptible to antibiotics. Differences in the phenotypes of the strains were observed, however. While the <i>ppm1</i>- strains have a small colony phenotype, the <i>sco3028::Tn5062</i> mutants had an extremely small colony phenotype indicative of an even greater growth defect. Moreover we were unable to generate a strain in which GMP activity encoded by <i>sco3039</i> and <i>sco4238</i> is completely knocked out, indicating that GMP is also an important enzyme for growth. Possibly GDP-mannose is at a metabolic branch point where it is modified to supply alternative nucleotide sugar donors.</p>

1 **Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor***  
2 **results in antibiotic hyper-susceptible phenotypes**

3 Robert Howlett<sup>a</sup>, Katri Anttonen<sup>b</sup>, Nicholas Read<sup>a</sup>, and Margaret C.M Smith<sup>a,b, #</sup>

4

5 <sup>a</sup> Department of Biology, University of York, York, United Kingdom;

6 <sup>b</sup> Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom;

7 #Address correspondence to Margaret C.M. Smith, [Maggie.smith@york.ac.uk](mailto:Maggie.smith@york.ac.uk)

8

9 Running Title: *S. coelicolor* strains disrupted in the synthesis of GDP-mannose

10

11 Abbreviations; Ppm1; polyprenol phosphate mannose synthase, PMM;

12 Phosphomannomutase, GMP; GDP-mannose pyrophosphorylase, PGM;

13 phosphoglucomutase, MESG; 2-amino-6-mercapto-7-methylpurine ribonucleoside,

14 PNP; purine nucleoside phosphorylase, SMMS; supplemented minimal medium

15 solid.

16

17

18

19 **Abstract**

20 Actinomycete bacteria use polyprenol phosphate mannose as a lipid linked sugar  
21 donor for extra-cytoplasmic glycosyl transferases that transfer mannose to cell  
22 envelope polymers, including glycoproteins and glycolipids. We showed recently that  
23 strains of *Streptomyces coelicolor* with mutations in the gene *ppm1* encoding  
24 polyprenol phosphate mannose synthase were both resistant to phage  $\phi$ C31 and  
25 have greatly increased susceptibility to antibiotics that mostly act on cell wall  
26 biogenesis. Here we show that mutations in the genes encoding enzymes that act  
27 upstream of Ppm1 in the polyprenol phosphate mannose synthesis pathway can also  
28 confer phage resistance and antibiotic hyper-susceptibility. GDP-mannose is a  
29 substrate for Ppm1 and is synthesized by GDP-mannose pyrophosphorylase  
30 (GMP;ManC) which uses GTP and mannose-1-phosphate as substrates.  
31 Phosphomannomutase (PMM; ManB) converts mannose-6-phosphate to mannose-1  
32 phosphate. *S. coelicolor* strains with knocked down GMP activity or with a mutation in  
33 *sco3028* encoding PMM acquire phenotypes that resemble those of the *ppm1*<sup>-</sup>  
34 mutants i.e.  $\phi$ C31 resistant and susceptible to antibiotics. Differences in the  
35 phenotypes of the strains were observed, however. While the *ppm1*<sup>-</sup> strains have a  
36 small colony phenotype, the *sco3028::Tn5062* mutants had an extremely small  
37 colony phenotype indicative of an even greater growth defect. Moreover we were  
38 unable to generate a strain in which GMP activity encoded by *sco3039* and *sco4238*  
39 is completely knocked out, indicating that GMP is also an important enzyme for  
40 growth. Possibly GDP-mannose is at a metabolic branch point where it is modified to  
41 supply alternative nucleotide sugar donors.

42

43 **Introduction**

44 *Streptomyces* spp. are prolific producers of secondary metabolites, many with potent  
45 antibiotic activity. In nature *Streptomyces* spp. produce antibiotics either to inhibit  
46 competitors thus providing the producer with a growth advantage or as signalling  
47 molecules in microbial communities [1, 2]. Either way *Streptomyces* bacteria are  
48 constantly exposed to antibiotics produced by other soil microorganisms and  
49 consequently have evolved resistance mechanisms [3]. As such *Streptomyces* spp.  
50 are a model system to study how the mechanisms of antibiotic resistance evolve in  
51 an environmental organism.

52

53 We recently showed that strains of *S. coelicolor* lacking the ability to synthesise  
54 polyprenol phosphate mannose due to mutations in polyprenol phosphate mannose  
55 synthase (Ppm1) were hyper-sensitive to multiple antibiotics (Howlett et al,  
56 submitted). We used RNA-seq and Raman spectroscopy to demonstrate that the  
57 strains had undergone changes to the membrane phospholipids, with possible  
58 subsequent changes to membrane functions. Polyprenol phosphate mannose  
59 synthase, Ppm1, transfers mannose from GDP-mannose to polyprenol phosphate  
60 (Fig. 1). Previously we demonstrated that the synthesis of polyprenol phosphate  
61 mannose was entirely dependent on membrane associated Ppm1 [4].

62

63 Polyprenol phosphate mannose is the mannose donor for extracytoplasmic glycosyl  
64 transferases. One of these is a protein mannosyl transferase (Pmt), which  
65 glycosylates periplasmic and membrane proteins in *Streptomyces* [4, 5]. Pmt  
66 defective strains also show increased antibiotic susceptibility compared to the parent  
67 strain, but to fewer antibiotics and to a lower level than the *ppm1* mutants (Howlett et

68 al, submitted). Loss of protein glycosylation is therefore likely to contribute in part to  
69 the antibiotic hyper susceptible phenotype of the *ppm1<sup>-</sup>* mutants. In addition both  
70 *ppm1<sup>-</sup>* strains and the *pmt* strains are resistant to the phage  $\phi$ C31, most likely  
71 through loss of the receptor, although the exact nature of the phage receptor is still  
72 unknown [5, 6].

73

74 Polyprenol phosphate mannose is likely to be a mannose donor for other cell  
75 envelope macromolecules with one of these likely to be phosphoinositol mannosides  
76 (PIMs) [7, 8]. In other Actinobacteria including *Mycobacterium* and *Corynebacterium*  
77 spp. PIMs are precursors for the synthesis of lipoarabinomannan and lipomannan [9],  
78 but neither of these polymers have been reported in *Streptomyces*. Ppm1 is an  
79 essential enzyme in mycobacteria and a *ppm1<sup>-</sup>* strain of *Corynebacterium* is growth  
80 retarded indicating the central role polyprenol phosphate mannose has in both  
81 organisms [10, 11]. The protein O-glycosylation pathway is present in most  
82 Actinobacteria and Pmt in Mtb has been shown to be important for virulence [12, 13].  
83 In *Streptomyces coelicolor* other putative glycosyl transferases are also likely to use  
84 polyprenol phosphate mannose as a sugar donor and some of these are described in  
85 Howlett et al (submitted).

86

87 The role of polyprenol phosphate mannose in antibiotic resistance and the pathway  
88 leading to its synthesis is addressed further in this paper (Fig. 1). D-mannose is  
89 either taken up from the medium and converted by hexokinase to D-mannose-6-  
90 phosphate or the latter can be produced from D-fructose-6-phosphate by  
91 phosphomannoisomerase (ManA). Phosphomannomutase (ManB: PMM) then  
92 converts D-mannose-6-phosphate is to D-mannose-1-phosphate which is a substrate

93 for GDP mannose pyrophosphorylase (ManC:GMP). In *Corynebacterium glutamicum*  
94 deletion of the *manC* homologue (NCgl0710) conferred retarded growth and loss of  
95 nearly all mannoglycans from the envelope [14]. This phenotype resembles that of  
96 the *ppm1* mutant of *C. glutamicum* [10] and suggests that the ManB, ManC pathway  
97 is responsible for the synthesis of GDP-mannose. We hypothesized that strains  
98 containing blocks in the pathway leading to the synthesis of GDP-mannose ought to  
99 be phenotypically similar to the *ppm1*<sup>-</sup> strains as they too will be deficient in  
100 polyprenol phosphate mannose. Here we analyzed the roles of three putative *manC*  
101 genes in the *S. coelicolor* genome and a *manB* gene. We show that both a GMP  
102 depleted strain and a strain lacking PMM do indeed have phenotypes reminiscent of  
103 the *ppm1*<sup>-</sup> mutants. The phenotype of the *S. coelicolor manB*<sup>-</sup> strains constructed  
104 here varied from that reported previously for a *manB*<sup>-</sup> strain [15, 16]. We conclude  
105 that GMP activity in *S. coelicolor* is provided by expression of two genes, *sco3039*  
106 and *sco4238*. Moreover both GMP and PMM activities are part of the same metabolic  
107 pathway leading to the synthesis of polyprenol phosphate mannose and ultimately to  
108 glycoprotein biosynthesis in *S. coelicolor*.

109

## 110 **Methods**

### 111 *DNA manipulations*

112 Chemically competent *E. coli* cells were prepared, stored and used in the  
113 transformation procedure as described previously [17]. Plasmid DNA extraction from  
114 *E. coli* was performed using a Spin Miniprep Kit following the protocol supplied by the  
115 manufacturer (QIAGEN). Cosmids were manipulated as described [18]. Restriction  
116 enzymes and T4 ligase were obtained from New England Biolabs (NEB) and used  
117 according to the manufacturer's instruction. Phusion<sup>®</sup> High-Fidelity DNA Polymerase

118 (NEB) was employed for PCR amplification. Primers used in the present study are  
119 listed (Table S1). In-fusion® HD cloning kit (Clontech) was used according to the  
120 protocol supplied by the manufacturer. DNA sequencing (Sanger) was outsourced to  
121 Source Bioscience.

#### 122 *Plasmid, cosmid and strain constructions*

123 A list of plasmids and cosmids used in this work is provided (Table 1). Plasmid  
124 pRH01 was produced by cloning of the PCR amplified product from primers RH11  
125 and RH12 and J1929 genomic DNA as template, into EcoRV digested vector  
126 pAV11b [19-21]. Plasmid pRH12 was produced by cloning the PCR product from  
127 primers RH91 and RH92 and J1929 template into NdeI digested pJ10257. Plasmids  
128 pRH11 and pRH14 were produced similar to pRH12 but using *E.coli* DH5  $\alpha$  genomic  
129 DNA as template and primer pairs RH93/RH94, and RH140/RH141, respectively.  
130 Expression plasmids for *sco3039* (pRH06) and *sco4238* (pRH07) were produced  
131 through the ligation of XhoI and NdeI digested PCR products from primers pairs  
132 RH71/RH72, and RH73/RH74, respectively, and *S. coelicolor* J1929 template DNA,  
133 into XhoI and NdeI digested pET21a fusing both ORFs to an inframe C-terminal  
134 hexa-histidine tag. All constructs were confirmed as correct through Sanger  
135 sequencing performed by Source Bioscience.

136 The apramycin resistance markers within the Tn5062 transposon of StD8A.2.D12  
137 and St1A8A.1.B09 were replaced with spectinomycin and hygromycin markers,  
138 respectively, using the REDIRECT methodology [18]. Cosmids were introduced into  
139 *S. coelicolor* J1929 by conjugation and resistant exconjugants were selected  
140 according to the marker on Tn5062 (apramycin, spectinomycin or hygromycin  
141 resistance). Those that had undergone a double cross over recombination event  
142 were identified initially as they lost the marker (kanamycin resistance) on the cosmid



143 vector backbone. Presence of the interrupted allele and loss of the wild type allele  
144 was confirmed by PCR and Southern blotting.

145

#### 146 *Phage sensitivity assays*

147 Plaque assays were performed as described [22]. Briefly Difco nutrient agar  
148 supplemented with 10 mM MgSO<sub>4</sub> and 8 mM Ca (NO<sub>3</sub>)<sub>2</sub> were inoculated with  
149 dilutions of  $\phi$ C31  $\Delta$ c25 (clear plaque) phage [23] and then overlaid with soft nutrient  
150 agar containing approximately  $1 \times 10^7$  spores of the desired test strain. The streak  
151 plate assay was performed using square 10 cm plates containing Difco nutrient agar  
152 (10 mM MgSO<sub>4</sub> and 8 mM Ca (NO<sub>3</sub>)<sub>2</sub>). One half of the plate was inoculated with 100  
153  $\mu$ l of  $\phi$ C31  $\Delta$ c25 (approx  $1 \times 10^8$  pfu/ml) and a single streak of the test spore  
154 preparation was inoculated across the plate beginning on the phage-free region.  
155 Plates were incubated at 30 °C.

156

#### 157 *Protein expression*

158 An overnight culture of *E. coli* BL21DE3 (pRH07) in LB containing ampicillin was  
159 grown at 37 °C and used to inoculate 2YT, which was grown to OD 0.6. IPTG (0.15  
160 mM) was then added to induce expression and the culture was further incubated (22  
161 °C for 22 hours). The bacteria were harvested by centrifugation and resuspended in  
162 binding buffer (30 ml; 20 mM Tris HCl pH 7.4, 0.5M NaCl, 30 mM imidazole) and  
163 sonicated. The cell lysate was cleared by centrifugation (4 °C, 5 minutes, 10,000g) to  
164 remove unlysed cells and debris and then the supernatant was loaded onto a HiTrap  
165 Ni<sup>2+</sup> affinity column (AKTA Purifier). After washing with 2 column volumes of binding  
166 buffer the bound protein was eluted with a gradient of increasing imidazole  
167 concentration using the elution buffer (20 mM Tris HCl pH 7.4, 0.5M NaCl, 500 mM

168 imidazole). Pooled fractions were then loaded onto a desalting column to remove  
169 imidazole and eluted in 20 mM Tris HCl pH 7.4, 0.5M NaCl. The protein was  
170 concentrated using Vivaspin (GE healthcare) spin columns to approximately 10  
171 mg/ml. Glycerol was added to a final concentration of 50% and aliquots were stored  
172 at -80 °C. Protein concentration was assayed using the BioRad protein assay  
173 solution and is based on the Bradford assay [24].

174

#### 175 *GDP mannose pyrophosphorylase assays*

176 Activity was measured by monitoring the release of pyrophosphate using the  
177 EnzChek® pyrophosphate assay kit (Thermofisher). Briefly the kit includes a  
178 pyrophosphatase that catalyses the conversion of the pyrophosphate released from  
179 the GMP activity to two equivalents of phosphate, which is then used as a substrate  
180 in a reaction with 2-amino-6-mercapto-7-methylpurine ribonucleoside (MESG) and  
181 purine nucleoside phosphorylase (PNP) to release ribose 1-phosphate and 2-amino-  
182 6-mercapto-7-methyl-purine. The latter compound was detected  
183 spectrophotometrically by absorbance at 360 nm. Assays were performed according  
184 to the manufacturers instructions except that they were scaled down to enable use of  
185 a 96 well plate reader (200  $\mu$ l assay volume per well). GMP activity rates were  
186 obtained using different nucleotides (1 mM ATP, GTP, CTP or dTTP) and sugars, (1  
187 mM mannose-1-phosphate or mannose-6-phosphate) as substrates. Initial rates  
188 were calculated and plotted against substrate concentration using SIGMAplot.

189

## 190 **Results**

### 191 *Identification of putative GDP-mannose pyrophosphorylases*

192 GDP-mannose, a substrate for Ppm1, is synthesized by GDP-mannose  
193 pyrophosphorylase (GMP) encoded by *manC* (Fig. 1). BLAST searches of the  
194 *Streptomyces coelicolor* genome with the characterised *Corynebacterium*  
195 *glutamicum* GMP/ManC (encoded by NCgl0710) and *Mycobacterium tuberculosis*  
196 GMP/ManC (encoded by Rv3264c) identified SCO1388, SCO3039 and SCO4238 as  
197 putative GMP candidates (Fig. 2) [14]. The nucleotidyl transferase domains of all  
198 three *Streptomyces* GMP/ManC candidates contain the GXGXRnK signature motif  
199 of phosphorylases, and variations on the F(V)EKP motif characteristic of the GMP  
200 active site (Fig. 2) [25, 26]. SCO3039 and SCO1388 have protein domains in addition  
201 to the nucleotidyl transferase domain; SCO1388 in particular appears to be a  
202 bifunctional enzyme with both GMP and phosphomannomutase (ManB) activity (Fig.  
203 2).

204

205 *SCO3039 and SCO4238 have overlapping functions*

206

207 The *S. coelicolor* genes *sco1388*, *sco3039* and *sco4238* were disrupted by allelic  
208 exchange with cosmids containing Tn5062 in the gene of interest to produce single  
209 insertion mutants RH501 (*sco1388::Tn5062*), SKA211 (*sco3039::Tn5062*) and  
210 SKA311 (*sco4238::Tn5062*). The cosmids were obtained from the transposon  
211 insertion cosmid library [27] (Table 1). Validated mutants were tested for  $\phi$ C31  
212 resistance as a potential indicator for a loss of GMP/ManC activity due to a lack of  
213 protein O-glycosylation [5, 6], but all of the mutants were still sensitive to phage  
214 infection. A mild increase in blue pigment production was seen in the mutant strains  
215 SKA311 (*sco4238::Tn5062*) and SKA211 (*sco3039::Tn5062*) when grown on  
216 supplemented minimal media (SMM) (not shown).

217  
218 Double mutants were created to assess whether there is redundancy in gene  
219 function between *sco3039*, *sco4238* and *sco1388*. The cosmid StD8A.2.D12<sup>spec</sup>  
220 (*sco4238::Tn5062<sup>spec</sup>*) was introduced into J1929 by conjugation to create the  
221 spectinomycin resistant *sco4238* insertion mutant, RH25. The cosmid StE34.1.G05  
222 (*sco3039::Tn5062*) was then introduced into RH25 by conjugation, selecting for  
223 apramycin resistance. Only 8 exconjugants from several hundred that were screened  
224 had the spectinomycin-resistant, apramycin-resistant, kanamycin-sensitive  
225 phenotype indicative of a *sco4238::Tn5062<sup>spec</sup>*, *sco3039::Tn5062* double mutant.  
226 However subsequent analysis by polymerase chain reaction (PCR) to amplify the  
227 genomic region containing *sco3039* showed that this gene was uninterrupted in all  
228 eight candidate double mutant strains and mutations must have occurred elsewhere  
229 to confer resistance to apramycin. Thus we were unable to create a simple double  
230 mutant containing *Tn5062* insertions in both *sco3039* and *sco4238*, suggesting that  
231 these genes share an important function for growth. Multiple *sco4238::Tn5062<sup>spec</sup>*,  
232 *sco1388::Tn5062* and *sco3039::Tn5062*, *sco1388::Tn5062<sup>hyg</sup>* double mutant strains  
233 were produced and confirmed through kanamycin sensitivity. The phenotypes of  
234 these strains were no different from the individual *sco4238::Tn5062* and  
235 *sco3039::Tn5062* mutants, SKA311 and SKA211, respectively. The product of  
236 *sco1388* therefore probably contributes little to the total GMP activity in *S. coelicolor*.

237  
238 We were able to create a strain containing both *sco4238::Tn5062<sup>spec</sup>* and  
239 *sco3039::Tn5062* insertions in the presence of a conditionally expressed *sco4238*.  
240 Plasmid pRH01, encoding *sco4238* under the control of the anhydrotetracycline  
241 (ATC) inducible promoter, *tcp830* [20], was introduced into RH25 to create strain

242 RH221 (*sco4238::Tn5062<sup>spec</sup>*, *tcp830-sco4238*, *hyg*). Conjugation of StE34.1.G05  
243 (*sco3039::Tn5062*) into RH221 in the presence of ATC resulted in multiple *spec<sup>R</sup>*,  
244 *apra<sup>R</sup>*, *hyg<sup>R</sup>*, *kan<sup>S</sup>* exconjugants (RH2213) that were subsequently confirmed as  
245 *sco4238::Tn5062<sup>spec</sup>*, *sco3039::Tn5062* double mutants through PCR. Surprisingly  
246 RH2213 could grow in the absence of ATC, an observation that was at odds with our  
247 inability to isolate the transposon double mutants in the absence of pRH01. Colony  
248 sizes of the RH2213 strains in the absence of ATC were indistinguishable from the  
249 wild type parent strain, J1929, but a significant increase in blue pigments were  
250 observed compared to the single mutants RH25 (*sco4238::Tn5062<sup>spec</sup>*) and SKA211  
251 (*sco3039::Tn5062*)(Fig. 2B). The *tcp830* promoter has been shown by others to be  
252 incompletely turned off in the absence of ATC and we propose that this is also the  
253 case in our experiments [19]. It seems likely that RH2213 grown in the absence of  
254 ATC has a depleted level of GMP compared to the parent strain and compared to  
255 RH2213 grown in the presence of ATC.

256

257

258 *Strains depleted in the putative GMPs SCO3039 and SCO4238 are hyper*  
259 *susceptible to antibiotics and partially resistant to  $\phi$ C31*

260

261 Ppm1 uses GDP-mannose as a substrate and we therefore hypothesized that  
262 inability to synthesize GDP-mannose, for example through GMP depletion, should  
263 result in a similar phenotype to those strains deficient in Ppm1. RH2213 isolates  
264 (*sco4239::Tn5062<sup>spec</sup>*, *sco3039::Tn5062*, pRH01 encoding inducible *sco4238*) were  
265 still able to support  $\phi$ C31 plaque formation but displayed resistance to  $\phi$ C31 on a  
266 streak assay in the absence of ATC (Fig 3A).

267

268 We then tested the putative GMP deficient strains for their susceptibilities to  
269 antibiotics, notably those to which the *ppm1* and *pmt* mutants were particularly  
270 sensitive. *S. coelicolor* strains SKA211 and RH25 containing Tn5062 insertions in  
271 *sco3039* and *sco4238*, respectively, had the same antibiotic resistances as the  
272 parent strain J1929. However RH2213, with depleted levels of GMP in the absence  
273 of ATC, was highly susceptible to antibiotics, strongly resembling the phenotypes of  
274 the *ppm1* mutants (Fig. 3B). The phenotypes of the GMP depleted mutants indicate  
275 that *sco3039* and *sco4238* provide the majority of the GMP activity in *S. coelicolor*.

276

277 *SCO4238 encodes a highly specific GDP-mannose pyrophosphorylase activity*

278

279 To confirm the phenotypes mentioned above were due to a depletion of GMP activity  
280 in RH2213, *sco4238* and *sco3039* were overexpressed in *E. coli* in order to assay  
281 GMP activity on purified proteins. Overproduced SCO4238 showed high GMP activity  
282 (Fig. 4). The enzyme was highly specific for GTP and D-mannose-1-phosphate  
283 substrates, with no or very low rates achieved with CTP, ATP and dTTP (not shown).  
284 Approximately 50% activity was observed with D-mannose-6-phosphate and GTP,  
285 with the Hill coefficient showing a loss of the cooperativity seen with D-mannose-1-  
286 phosphate. In *Mycobacterium tuberculosis* the essential enzyme, RmlA, catalyses  
287 the synthesis of dTDP-glucose, an intermediate in dTDP-rhamnose biosynthesis  
288 required for the integrity of the cell wall [28]. Given the apparent essentiality of GMP  
289 in *S. coelicolor* we tested whether SCO4238 had activity on glucose-1-phosphate in  
290 combination with any nucleotide, including dTTP but no activity was detected.

291 Attempts to obtain soluble, active SCO3039 from several overexpression constructs  
292 in *E. coli* failed (not shown).

293

294 *Strains with a mutation in the manB gene, sco3028 are also phenotypically similar to*  
295 *the ppm1 mutants*

296 Previous work has shown that SCO3028 is a dual functioning enzyme capable of  
297 phosphomannomutase (PMM, mannose-6-phosphate to mannose-1-phosphate) and  
298 phosphoglucomutase (PGM, glucose-6-phosphate to glucose-1-phosphate) activity  
299 [16]. The authors constructed a *manB* deletion mutant,  $\Delta manB$ , which had increased  
300 actinorhodin production and had lost chloramphenicol resistance but displayed  
301 apparently similar growth to the parent strain, M145 [15, 16]. Both phenotypes of the  
302  $\Delta manB$  strain were complemented when wild type *E. coli manB* but not *S. coelicolor*  
303 *pgm* gene (*sco7443*) were introduced. Thus PMM activity was shown to be solely  
304 responsible for an increase in chloramphenicol sensitivity and actinorhodin  
305 production in *S. coelicolor* M145. If SCO3028 is the sole PMM enzyme in *S.*  
306 *coelicolor* we would expect a similar phenotype in the *sco3028* mutant as we see for  
307 the GMP depleted strains. However, Yang *et al* did not detect increased susceptibility  
308 of their  $\Delta manB$  strain to vancomycin, bacitracin or ampicillin [16].

309

310 In order to assess the phenotype of an *sco3028* mutant in our  $\phi C31$  sensitive strain  
311 *S. coelicolor* J1929, a *pglY* derivative of M145 [29], the cosmid StE34.1.B03  
312 (*sco3028::Tn5062*) was introduced into J1929 by conjugation. Exconjugants that had  
313 undergone a double crossover (RHB42 strains, validated by PCR) were isolated at  
314 low frequency and had an extreme small (XS) colony phenotype, even smaller than  
315 the colony size seen in the *ppm1* mutant DT3017 (Fig. 5). The XS colony phenotype

316 in RHB42 could be fully restored to wild type through complementation with *S.*  
317 *coelicolor sco3028 (manB)*, and *Escherichia coli manB (cpsG)* as observed in strains  
318 RHB4212 and RHB4211, respectively. RHB42 containing *Escherichia coli pgm*,  
319 encoding phosphoglucomutase, was capable of partially restoring colony size  
320 (RHB4214), suggesting it is the loss of both PMM and PGM activity that had resulted  
321 in the XS colony phenotype in RHB42.

322

323 RHB42 was highly resistant to  $\phi$ C31 infection in a plaque assay, resembling phage  
324 resistance in the *pmt* and *ppm1* mutants (Fig. 5A). RHB42 was also highly  
325 susceptible to a number of cell wall acting antibiotics, as well as the RNA polymerase  
326 targeting antibiotic, rifampicin (Fig. 5B). Phage sensitivity and antibiotic resistance  
327 were restored to wild type in RHB4212 (*sco3028<sup>+</sup>*) and RHB4211 (*cpsG<sup>+</sup>*) but not  
328 RHB4214 (*pgm<sup>+</sup>*). No difference in chloramphenicol resistance between RHB42 and  
329 J1929 was observed. These phenotypes are consistent with SCO3028 being the  
330 primary PMM in *S. coelicolor* and in the same metabolic pathway that synthesizes  
331 polyprenol phosphate mannose.

332

333 An increase in pigment production was recorded in RHB42, similar to that seen  
334 previously following *sco3028* deletion [16] and that seen in the GMP depleted strain,  
335 RH2213 (Fig. 5C). The production of blue pigment was reduced to wild type level in  
336 RHB4212 (*sco3028<sup>+</sup>*) and RHB4211 (*cpsG<sup>+</sup>*) but not RHB4214 (*pgm<sup>+</sup>*) (Fig. 5). To  
337 further validate our observations (as they differ from those of Yang et al, [15, 16]), we  
338 created two more sets of *sco3028::Tn5062* mutants: First we used a different *Tn5062*  
339 insertion in J1929 using cosmid, STE34.2.D03, generating strain JD182 and second  
340 we generated derivatives of M145 containing the *Tn5062* insertions from both



341 STE34.1.B03 and STE34.2.D03 to generate strains MD202 and MB92, respectively.  
342 All three strains had an identical phenotype to RHB42 (Figs S1 and S2).

343

## 344 **Discussion**

345 Mannose is a component of cell envelope polymers including manolipids,  
346 phosphoinositol mannosides (PIMs) and glycoproteins in many bacteria [9, 30-32].  
347 Extracytoplasmic glycosyl transferases use polyprenol phosphate mannose as the  
348 lipid linked sugar donor in the biosynthesis of mannose containing polymers [4, 33].  
349 The synthesis of polyprenol phosphate mannose by Ppm1 is therefore an important  
350 activity and *ppm1* mutants are considerably less fit than the parent strains [10, 11]. In  
351 the case of *S. coelicolor*, *ppm1*<sup>-</sup> mutants have a small colony growth phenotype and  
352 are hyper-susceptible to multiple antibiotics, most of which inhibit cell wall biogenesis  
353 suggesting that these mutants are pleiotropically deficient in membrane and/or  
354 periplasmic function (Howlett et al, submitted). Mutants lacking Ppm1 or Pmt are also  
355 resistant to phage infection and we have proposed that  $\phi$ C31 uses a glycoprotein(s)  
356 as its receptor [5, 6]. We show here that depletion of enzymes in the mannose  
357 metabolism pathway prior to Ppm1 display a phenotype that resembles that of the  
358 *ppm1*<sup>-</sup> mutants. We conclude that synthesis of polyprenol phosphate mannose and  
359 its subsequent role as a mannose donor in the periplasm is required for a wild type  
360 antibiotic resistant phenotype.

361

362 Although the overall phenotypes of the GDP-mannose pyrophosphorylase (GMP)  
363 deficient and the *manB* mutant strains resembled the *ppm1*<sup>-</sup> strain there were some  
364 minor differences. We had difficulty in generating a GMP deficient strain. *S. coelicolor*  
365 has three candidate genes that could express GMP activity and we could only obtain

366 a double *sco3039*, *sco4238* mutant if *sco4238* was expressed conditionally using  
367 the anhydrotetracycline-inducible *tcp830* promoter. While this is not absolute proof  
368 that the GMP activity is essential in *S. coelicolor*, it would seem that some low level  
369 of activity, possibly that provided by the leakiness of the repressed *tcp830* promoter  
370 reading into an integrated copy of *sco4238*, is required for the simultaneous  
371 interruption of both *sco3039* and *sco4238* by Tn5062. Similarly the insertion in  
372 *sco3028* (*manB*) was obtained at very low frequency and the colonies were  
373 extremely small, indicative of a requirement for both phosphomannomutase (PMM)  
374 and phosphoglucomutase (PGM) activities encoded by this gene. If GMP activity is  
375 essential then we would also expect PMM activity to be essential, but there may be  
376 sufficient PMM activity from other closely related enzymes (such as other PGM  
377 paralogues; *sco7443* or *sco4916* a possible alternative phosphomannomutase) to  
378 allow growth. The more severe phenotypes of the GMP depletion mutant and the  
379 *manB* mutant also suggest that GDP-mannose could be located at a metabolic  
380 branch point i.e. GDP-mannose is required for polyprenol phosphate mannose  
381 synthesis but also perhaps for modification into other nucleotide sugars. For  
382 example, the *S. coelicolor* genome encodes a GDP-mannose dehydrogenase  
383 (SCO0382) that is predicted to make GDP-mannuronate, one of the building blocks  
384 in the synthesis of alginates in *Pseudomonas*. *Sco0382* lies within an operon *sco381*  
385 to *sco386* that has features of an extracellular polysaccharide biosynthesis gene  
386 cluster including a polyprenol dependent glycosyl transferase and various other  
387 membrane proteins.

388

389 Enzyme assays with purified SCO4238 showed it to be a monofunctional GMP  
390 (ManC) with a slim substrate tolerance similar to Rv3264 (previously miss-annotated

391 as *rmlA*) of *Mycobacterium* [28, 34] and in contrast to the more promiscuous ManC  
392 enzymes of *E. coli* and *P. furiosus* [35, 36]. In *Mycobacterium* and in *Corynebacterium*  
393 the ManC enzymes (Rv3264 and NCgl0710, respectively) provide essential supplies  
394 of GDP-mannose for phosphatidyl inositol mannoside (PIM) biosynthesis and  
395 lipoglycans [14].

396

397 The phenotype described here for RHB42 (*sco3028:Tn5062, manB*) has differences  
398 and similarities to a  $\Delta manB$  strain of *S. coelicolor* M145 that was described  
399 previously [15, 16]. A notable difference is the sensitivity to antibiotics of RHB42 as  
400 Yang *et al* did not detect an increase in susceptibility of their  $\Delta manB$  strain to  
401 vancomycin, bacitracin or ampicillin [16]. Whilst we cannot explain these differences  
402 in phenotypes, both RHB42 and the  $\Delta manB$  of Yang *et al* have increased pigment  
403 production. Pigment production is also greatly increased in the ManC deficient  
404 strains. The increase in pigment production could be indicative of the activation of  
405 several stress pathways or, as discussed by Yang *et al*, could be due to the increase  
406 in carbon flux through glycolysis as the pathway to GDP-mannose is blocked. The  
407 *ppm1<sup>-</sup>* strain DT3017 has a mild pigment overproduction phenotype (data not  
408 shown). Neither Yang *et al*, or Rajesh *et al* could test phage sensitivity in their  
409  $\Delta manB$  strain as they used a Pgl<sup>+</sup> strain of *S. coelicolor*, which confers  $\phi C31$   
410 resistance.

411

412 Mannose is used in other *Streptomyces spp* in the biosynthesis of antibiotics eg  
413 mannopeptimycins and amphotericin [37, 38]. ManB and ManC activities are required  
414 in *S. nodosus* for the glycosylation of amphotericin [38]. The identification of the  
415 *manC* genes and the construction of the *manC* deficient strain could be useful in

416 heterologous expression and combinatorial biosynthesis of several antibiotic  
417 pathways in *S. coelicolor*.

418

### 419 **Acknowledgements**

420 We are grateful to Professor Mervyn Bibb for providing plasmid pIJ10257 and to  
421 Professor Paul Dyson for the provision of mutagenized cosmids.

422

### 423 **Funding Information**

424 This work was funded by project grants\_BB/J016691 and BBS/B/05990 from the  
425 Biotechnology and Biological Sciences Research Council, UK.

426

### 427 **Conflicts of Interest and Ethics**

428 None

429

430

### 431 **References**

- 432 1. **Wang W, Ji J, Li X, Wang J, Li S et al.** Angucyclines as signals modulate the  
433 behaviors of *Streptomyces coelicolor*. *Proc Natl Acad Sci U S A* 2014;111(15):5688-  
434 5693.
- 435 2. **Nesme J, Simonet P.** The soil resistome: a critical review on antibiotic  
436 resistance origins, ecology and dissemination potential in telluric bacteria.  
437 *Environmental microbiology* 2015;17(4):913-930.
- 438 3. **D'Costa VM, McGrann KM, Hughes DW, Wright GD.** Sampling the antibiotic  
439 resistome. *Science* 2006;311(5759):374-377.
- 440 4. **Wehmeier S, Varghese AS, Gurcha SS, Tissot B, Panico M et al.**  
441 Glycosylation of the phosphate binding protein, PstS, in *Streptomyces coelicolor* by a  
442 pathway that resembles protein O-mannosylation in eukaryotes. *Mol Microbiol*  
443 2009;71(2):421-433.
- 444 5. **Cowlshaw DA, Smith MCM.** Glycosylation of a *Streptomyces coelicolor*  
445 A3(2) cell envelope protein is required for infection by bacteriophage  $\phi$ C31. *Mol*  
446 *Microbiol* 2001;41(3):601-610.
- 447 6. **Cowlshaw DA, Smith MCM.** A gene encoding a homologue of dolichol  
448 phosphate-b-D mannose synthase is required for infection of *Streptomyces coelicolor*  
449 A3(2) by phage  $\phi$ C31. *J Bacteriol* 2002;184(21):6081-6083.

- 450 7. **Hoischen C, Gura K, Luge C, Gumpert J.** Lipid and fatty acid composition of  
451 cytoplasmic membranes from *Streptomyces hygroscopicus* and its stable protoplast-  
452 type L form. *J Bacteriol* 1997;179(11):3430-3436.
- 453 8. **Sandoval-Calderon M, Geiger O, Guan Z, Barona-Gomez F, Sohlenkamp**  
454 **C.** A eukaryote-like cardiolipin synthase is present in *Streptomyces coelicolor* and in  
455 most actinobacteria. *J Biol Chem* 2009;284(26):17383-17390.
- 456 9. **Guerin ME, Kordulakova J, Alzari PM, Brennan PJ, Jackson M.** Molecular  
457 basis of phosphatidyl-myo-inositol mannoside biosynthesis and regulation in  
458 mycobacteria. *J Biol Chem* 2010;285(44):33577-33583.
- 459 10. **Gibson KJ, Eggeling L, Maughan WN, Krumbach K, Gurcha SS et al.**  
460 Disruption of Cg-Ppm1, a polyprenyl monophosphomannose synthase, and the  
461 generation of lipoglycan-less mutants in *Corynebacterium glutamicum*. *J Biol Chem*  
462 2003;278(42):40842-40850.
- 463 11. **Rana AK, Singh A, Gurcha SS, Cox LR, Bhatt A et al.** Ppm1-encoded  
464 polyprenyl monophosphomannose synthase activity is essential for lipoglycan  
465 synthesis and survival in mycobacteria. *PLoS one* 2012;7(10):e48211.
- 466 12. **Liu CF, Tonini L, Malaga W, Beau M, Stella A et al.** Bacterial protein-O-  
467 mannosylating enzyme is crucial for virulence of *Mycobacterium tuberculosis*. *Proc Natl*  
468 *Acad Sci USA* 2013;110(16):6560-6565.
- 469 13. **Mahne M, Tauch A, Puhler A, Kalinowski J.** The *Corynebacterium*  
470 *glutamicum* gene pmt encoding a glycosyltransferase related to eukaryotic protein-O-  
471 mannosyltransferases is essential for glycosylation of the resuscitation promoting  
472 factor (Rpf2) and other secreted proteins. *FEMS Microbiol Lett* 2006;259(2):226-233.
- 473 14. **Mishra AK, Krumbach K, Rittmann D, Batt SM, Lee OY et al.** Deletion of  
474 manC in *Corynebacterium glutamicum* results in a phospho-myo-inositol mannoside-  
475 and lipoglycan-deficient mutant. *Microbiology* 2012;158(Pt 7):1908-1917.
- 476 15. **Rajesh T, Song E, Lee BR, Park SH, Jeon JM et al.** Increased sensitivity to  
477 chloramphenicol by inactivation of *manB* in *Streptomyces coelicolor*. *Journal of*  
478 *microbiology and biotechnology* 2012;22(10):1324-1329.
- 479 16. **Yang YH, Song E, Park SH, Kim JN, Lee K et al.** Loss of  
480 phosphomannomutase activity enhances actinorhodin production in *Streptomyces*  
481 *coelicolor*. *Applied microbiology and biotechnology* 2010;86(5):1485-1492.
- 482 17. **Green RM, Sambrook J.** *Molecular Cloning: A laboratory Manual*, Fourth  
483 Edition ed. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press;  
484 2012.
- 485 18. **Gust B, Challis GL, Fowler K, Kieser T, Chater KF.** PCR-targeted  
486 *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis  
487 of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci U S A* 2003;100(4):1541-  
488 1546.
- 489 19. **Jyothikumar V, Klanbut K, Tiong J, Roxburgh JS, Hunter IS et al.**  
490 Cardiolipin synthase is required for *Streptomyces coelicolor* morphogenesis. *Mol*  
491 *Microbiol* 2012;84(1):181-197.
- 492 20. **Rodriguez-Garcia A, Combes P, Perez-Redondo R, Smith MCM.** Natural  
493 and synthetic tetracycline-inducible promoters for use in the antibiotic-producing  
494 bacteria *Streptomyces*. *Nucleic Acids Res* 2005;33(9):e87.
- 495 21. **Fayed B, Younger E, Taylor G, Smith MC.** A novel *Streptomyces* spp.  
496 integration vector derived from the *S. venezuelae* phage, SV1. *BMC biotechnology*  
497 2014;14:51.
- 498 22. **Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA.** *Practical*  
499 *Streptomyces Genetics*. Norwich: The John Innes Foundation; 2000.

- 500 23. **Sinclair RB, Bibb MJ.** The repressor gene (c) of the *Streptomyces* temperate  
501 phage  $\phi$ C31: nucleotide sequence, analysis and functional cloning. *Mol Gen Genet*  
502 1988;213(2-3):269-277.
- 503 24. **Bradford MM.** A rapid and sensitive method for the quantitation of microgram  
504 quantities of protein, utilising the principal of protein-dye binding. *Anal Biochem*  
505 1976;72:248-254.
- 506 25. **Pelissier MC, Lesley SA, Kuhn P, Bourne Y.** Structural insights into the  
507 catalytic mechanism of bacterial guanosine-diphospho-D-mannose  
508 pyrophosphorylase and its regulation by divalent ions. *J Biol Chem*  
509 2010;285(35):27468-27476.
- 510 26. **Sousa SA, Moreira LM, Leitao JH.** Functional analysis of the *Burkholderia*  
511 *cenocepacia* J2315 BceAJ protein with phosphomannose isomerase and GDP-D-  
512 mannose pyrophosphorylase activities. *Applied microbiology and biotechnology*  
513 2008;80(6):1015-1022.
- 514 27. **Fernandez-Martinez LT, Del Sol R, Evans MC, Fielding S, Herron PR et al.**  
515 A transposon insertion single-gene knockout library and new ordered cosmid library  
516 for the model organism *Streptomyces coelicolor* A3(2). *Antonie Van Leeuwenhoek*  
517 2011;99(3):515-522.
- 518 28. **Ma Y, Stern RJ, Scherman MS, Vissa VD, Yan W et al.** Drug targeting  
519 *Mycobacterium tuberculosis* cell wall synthesis: genetics of dTDP-rhamnose  
520 synthetic enzymes and development of a microtiter plate-based screen for inhibitors  
521 of conversion of dTDP-glucose to dTDP-rhamnose. *Antimicrob Agents Chemother*  
522 2001;45(5):1407-1416.
- 523 29. **Bedford DJ, Laity C, Buttner MJ.** Two genes involved in the phase-variable  
524  $\phi$ C31 resistance mechanism of *Streptomyces coelicolor* A3(2). *J Bacteriol*  
525 1995;177(16):4681-4689.
- 526 30. **Dobos KM, Khoo KH, Swiderek KM, Brennan PJ, Belisle JT.** Definition of  
527 the full extent of glycosylation of the 45-kilodalton glycoprotein of *Mycobacterium*  
528 *tuberculosis*. *J Bacteriol* 1996;178(9):2498-2506.
- 529 31. **Michell SL, Whelan AO, Wheeler PR, Panico M, Easton RL et al.** The  
530 MPB83 antigen from *Mycobacterium bovis* contains O-linked mannose and (1 $\rightarrow$ 3)-  
531 mannobiose moieties. *J Biol Chem* 2003;278(18):16423-16432.
- 532 32. **Mishra AK, Driessen NN, Appelmek BJ, Besra GS.** Lipoarabinomannan  
533 and related glycoconjugates: structure, biogenesis and role in *Mycobacterium*  
534 *tuberculosis* physiology and host-pathogen interaction. *FEMS Microbiol Rev*  
535 2011;35(6):1126-1157.
- 536 33. **Besra GS, Morehouse CB, Rittner CM, Waechter CJ, Brennan PJ.**  
537 Biosynthesis of mycobacterial lipoarabinomannan. *J Biol Chem* 1997;272(29):18460-  
538 18466.
- 539 34. **Ning B, Elbein AD.** Purification and properties of mycobacterial GDP-  
540 mannose pyrophosphorylase. *Arch Biochem Biophys* 1999;362(2):339-345.
- 541 35. **Mizanur RM, Pohl NL.** Phosphomannose isomerase/GDP-mannose  
542 pyrophosphorylase from *Pyrococcus furiosus*: a thermostable biocatalyst for the  
543 synthesis of guanidinediphosphate-activated and mannose-containing sugar  
544 nucleotides. *Organic & biomolecular chemistry* 2009;7(10):2135-2139.
- 545 36. **Yang YH, Kang YB, Lee KW, Lee TH, Park SS et al.** Characterization of  
546 GDP-mannose pyrophosphorylase from *Escherichia coli* O157 : H7 EDL933 and its  
547 broad substrate specificity. *J Mol Catal B-Enzym* 2005;37(1-6):1-8.
- 548 37. **Magarvey NA, Haltli B, He M, Greenstein M, Hucul JA.** Biosynthetic  
549 pathway for mannopeptimycins, lipoglycopeptide antibiotics active against drug-

550 resistant gram-positive pathogens. *Antimicrob Agents Chemother* 2006;50(6):2167-  
551 2177.  
552 38. **Nic Lochlainn L, Caffrey P.** Phosphomannose isomerase and  
553 phosphomannomutase gene disruptions in *Streptomyces nodosus*: impact on  
554 amphotericin biosynthesis and implications for glycosylation engineering. *Metabolic*  
555 *engineering* 2009;11(1):40-47.  
556 39. **Hong HJ, Hutchings MI, Hill LM, Buttner MJ.** The role of the novel Fem  
557 protein Vank in vancomycin resistance in *Streptomyces coelicolor*. *J Biol Chem*  
558 2005;280(13):13055-13061.  
559 40. **MacNeil DJ.** Characterization of a unique methyl-specific restriction system in  
560 *Streptomyces avermitilis*. *J Bacteriol* 1988;170(12):5607-5612.  
561

562

### 563 **Legends to Figures**

564 **Figure 1. The GDP-mannose biosynthesis pathway in *Streptomyces coelicolor*.**

565

566 **Figure 2. GDP-mannose pyrophosphorylases in *Streptomyces coelicolor* A.**

567 Domain structures for *S. coelicolor* genes with putative GDP-mannose  
568 phosphorylase activity. B. Pigment overproduction in *sco4238*, *sco3039* double  
569 mutants. Strain RH2213 (*sco4239::Tn5062spec*, *sco3039::Tn5062*, pRH01 encoding  
570 inducible *sco4238*) overproduced blue pigment on supplemented minimal medium  
571 solid (SMMS) agar in the absence of anhydrotetracycline (ATC) but not in the  
572 presence of 0.5  $\mu$ g/ml ATC.

573

574 **Figure 3. *Streptomyces coelicolor* strains depleted in GDP mannose  
575 pyrophosphorylase activity are partially resistant to  $\phi$ C31 and are hyper-**

576 **susceptible to some antibiotics** A. Spores of the indicated *S. coelicolor* strains  
577 were streaked from an area free from  $\phi$ C31 to an area inoculated with  $1 \times 10^7$  pfu  
578  $\phi$ C31 on Difco nutrient agar plates with or with the supplementation of 0.5  $\mu$ g/ml  
579 anhydrotetracycline (ATC). RH2213 (*sco4239::Tn5062<sup>spec</sup>*, *sco3039::Tn5062*, pRH01

580 encoding inducible *sco4238*) showed conditional phage resistance growing only in  
581 the absence of ATC. For comparison the phage resistant phenotype of the *ppm1*-  
582 mutant (DT3017), the parent strain (J1929) and the strains with single mutations in  
583 the *manC* candidate genes, *sco3039* and *sco4238*, (SKA211 and RH25,  
584 respectively). B. RH2213 showed increased susceptibility to antibiotics in the  
585 absence of ATC but not in the presence of ATC. This phenotype is comparable to the  
586 antibiotic hyper-susceptible phenotype of the *ppm1*- mutant (DT3017). Results show  
587 the mean diameter of the disc diffusion inhibition zones from at least 3 replicates.  
588 Antibiotics were all used at 4  $\mu$  g/disc with the exception of ampicillin that was used  
589 at 40  $\mu$  g/disc.

590

591 **Figure 4. Kinetics of GDP-mannose pyrophosphorylase activity of SCO4238.**

592 Dependence of initial velocity of SCO4238 GMP activity with increasing  
593 concentration of mannose-1-phosphate (A) and mannose-6-phosphate (B). C. Kinetic  
594 parameters  $K_m$ ,  $n_H$  and  $V_{max}$  for the two sugar phosphate substrates.

595

596 **Figure 5. Phenotypes of the *manB* (*sco3028::Tn5062*) mutants.** A. Extreme small

597 colony phenotype (left) and resistance to phage  $\phi$ C31 (right) in the RHB42  
598 (*sco3028::Tn5062*) mutant strains. The poor growth of RHB42 was complemented  
599 with an additional copy of *sco3028* (RHB4212) or the *E. coli* gene *cpsG* (*manB*  
600 homologue; RHB4211) but not *pgm* from *E. coli* (encoding phosphoglucomutase;  
601 RHB4214). The  $\phi$ C31 resistance in RHB42 reverted to phage sensitivity in the  
602 complemented strains RHB4211 and RHB4212. B. RHB42 has increased  
603 susceptibility to antibiotics resembling the phenotype of the *ppm1*- mutant (DT3017).  
604 Antibiotic susceptibility could be complemented with an additional copy of *sco3028*



605 and *E. coli cpsG* but not *pgm* from *E. coli*. C. Increased pigment production in RH42  
606 compared to the parental strain, J1929. This phenotype was complemented by  
607 additional copy of *sco3028* and *E. coli cpsG* but not *pgm* from *E. coli*.

608

609

610

611 Table 1. Bacteria, plasmids and cosmids

<b>Plasmid Name</b>	<b>Description</b>	<b>References</b>
pAVIIb	Integrating vector with <i>tcp830</i> promoter and <i>Tetris</i> cassette	[21]
pRH01	<i>sco4238</i> in pAV11b	This study
pDT16	<i>sco1423 (ppm1)</i> in pSET152	[6]
pDT10	<i>sco3154 (pmt)</i> in pSET152	[5]
pET21a	Overexpression vector containing HIS <sub>6</sub> -tag, T7 promoter	Merck Chemicals
pRH06	<i>sco3039</i> in pET21a	This study
pRH07	<i>sco4238</i> in pET21a	This study
pIJ10257	Integrating vector with constitutive promoter <i>ermEp*</i>	[39]
pRH11	<i>cpsG</i> in pIJ10257	This study
pRH12	<i>sco3028</i> in pIJ10257	This study
pRH14	<i>pgm</i> in pIJ10257	This study
<b>Cosmid</b>	<b>Description</b>	<b>References</b>
St1A8A.1.B09	SCO1388:: <i>Tn5062</i> at nt 572	[27]
StD8A.2.D12	SCO4238:: <i>Tn5062</i> at nt 69	[27]
StE34.1.G05	SCO3039:: <i>Tn5062</i> at nt 155	[27]
StE34.1.B03	SCO3028:: <i>Tn5062</i> at nt 590	[27]
StD8A.2.D12 <sup>spec</sup>	StD8A.2.D12 with <i>sco4238</i> :: <i>Tn5062</i> <sup>spec</sup>	This study
St1A8A.1.B09 <sup>hyg</sup>	St1A8A.1.B09 with <i>sco1388</i> :: <i>Tn5062</i> <sup>hyg</sup>	This study
<b><i>Streptomyces</i> strain</b>	<b>Genotype</b>	<b>References</b>
M145	Prototroph	[22]
J1929	<i>pglY</i> mutant	[29]
DT3017	<i>ppm1</i> <sup>E218V</sup> mutant	[6]
DT1020	<i>ppm1</i> <sup>H116D</sup> mutant	[6]
DT1029	<i>ppm1</i> <sup>S163L</sup> mutant	[6]
DT1025	<i>pmt</i> , frameshift from A121	[5]
DT2008	<i>pmt</i>	[5]

SKA211	<i>sco3039::Tn5062</i>	This study
SKA311	<i>sco4238::Tn5062</i>	This study
RH501	<i>sco1388::Tn5062</i>	This study
RH25	<i>sco4238::Tn5062<sup>spec</sup></i>	This study
RH221	<i>sco4238::Tn5062<sup>spec</sup>, pRH01</i>	This study
RH2213	<i>sco4238::Tn5062<sup>spec</sup> sco3039::Tn5062, pRH01</i>	This study
RHB42	<i>sco3039::Tn5062</i>	This study
RHB4211	<i>sco3028::Tn5062, pRH11</i>	This study
RHB4212	<i>sco3028::Tn5062, pRH12</i>	This study
RHB4214	<i>sco3028::Tn5062, pRH14</i>	This study
<b><i>E.coli</i> strain</b>	<b>Genotype</b>	<b>References</b>
ET12567(pUZ8002)	<i>dam- dcm- hsdS</i> , RP4 transfer genes	[40]
BL21 (DE3)	<i>lon, ompT, gal, hsdS</i> , □DE3	[17]
DH5α	F- Φ80 <i>lacZ</i> Δ M15 Δ ( <i>lacZYA-argF</i> ) U169 <i>recA1 endA1 hsdR17</i> (rk-, mk+) <i>phoA supE44 λ -thi-1 gyrA96 relA1</i>	[17]

612

613





Figure 1

A



B





Figure 2

A



B




Figure 3






Figure 4

A



B



C




Figure 5



M145 wt	MD202 <i>sco3028</i> <sup>-</sup>	MD202 <i>sco3028, cpsG</i> <sup>+</sup>	MD202 <i>sco3028, sco3028</i> <sup>+</sup>	MD202 <i>sco3028, pgm</i> <sup>+</sup>
	MB92 <i>sco3028</i> <sup>-</sup>	MB92 <i>sco3028, cpsG</i> <sup>+</sup>	MB92 <i>sco3028, sco3028</i> <sup>+</sup>	MB92 <i>sco3028, pgm</i> <sup>+</sup>
	JD182 <i>sco3028</i> <sup>-</sup>	JD182 <i>sco3028, cpsG</i> <sup>+</sup>	JD182 <i>sco3028, sco3028</i> <sup>+</sup>	JD182 <i>sco3028, pgm</i> <sup>+</sup>




Figure S1




Figure S2

## Supplementary Figure Legends. Howlett *et al.*

**Figure S1. Extreme small colony phenotype of various *manB*<sup>-</sup> strains of *S. coelicolor*.** MD202 is a derivative of *S. coelicolor* M145 containing *sco3028::Tn5062* at nt 590 derived from cosmid StE34.1.B03 (1, 2). MB92 and JD182 are derivatives of *S. coelicolor* M145 and J1929, respectively, containing *sco3028::Tn5062* at nt 576 derived from cosmid StE34.2.D03 (1, 2). In each case a transposon insertion in *sco3028*, (encoding ManB) confers an extreme small colony phenotype. The phenotype can be fully complemented by introduction of a wild type copy of *sco3028* (*sco3028*<sup>+</sup> in pRH12), *E. coli cpsG* (encoding ManB in pRH11) but not by *E. coli pgm* (encoding phosphoglucomutase, pRH14).

**Figure S2. Antibiotic hyper-susceptible phenotype of various *manB*<sup>-</sup> strains of *S. coelicolor*.** Strains are as described in the legend to Figure S2. For comparison the antibiotic susceptibility profiles of the parent strains J1929 and M145 and that of the *ppm1*<sup>-</sup> strain DT3017 are also shown.

### References

1. Bishop A, Fielding S, Dyson P, Herron P. Systematic insertional mutagenesis of a streptomycete genome: a link between osmoadaptation and antibiotic production. *Genome Res.* 2004;14(5):893-900.
2. Fernandez-Martinez LT, Del Sol R, Evans MC, Fielding S, Herron PR, Chandra G, et al. A transposon insertion single-gene knockout library and new ordered cosmid library for the model organism *Streptomyces coelicolor* A3(2). *Antonie Van Leeuwenhoek.* 2011;99(3):515-22.

Table S1. Oligonucleotides for Howlett, Read *et al*,

Primer	Sequence	Use
RH11	TCTAGAGACGTCGATATCATGACCGATCCGAACGCC GCGTCC	<i>sco4238</i> F Infusion; pRH01
RH12	CATGCATGATCAGATATCTCAGCGGCCGGACAGGG CCG	<i>sco4238</i> R Infusion; pRH01
RH71	GACGCCCATATGACAGAAGCGATCCTCCTG	<i>sco3039</i> (NdeI); pRH06
RH72	CGTTACAAGCTTCGTGTCCGGTGAGAAGCG	<i>sco3039</i> (HindIII); pRH06
RH73	GGCGCCCATATGACCGATCCGAACGCCGCGTC	<i>sco4238</i> (NdeI); pRH07
RH74	GTACCTCGAGGCGGCCGGACAGGGCCG	<i>sco4238</i> (XhoI); pRH07
RH91	ACAGGAGGCCCATATGGCTGCTGATCTGTGCAG	<i>sco3028</i> F Infusion; pRH12
RH92	ACTCGAGATCTCATATGTCACGCCGGATGATCGCC	<i>sco3028</i> R Infusion; pRH12
RH93	ACAGGAGGCCCATATGAAAAATTAACCTGCTTT	<i>cpsG(manB)</i> F Infusion; pRH11
RH94	ACTCGAGATCTCATATGTTACTCGTTCAGCAACG	<i>cpsG(manB)</i> R Infusion; pRH11
RH140	ACAGGAGGCCCATATGGCAATCCACAATCGTGC	<i>pgm</i> F Infusion; pRH14
RH141	ACTCGAGATCTCATATGTTACGCGTTTTTCAGAA	<i>pgm</i> R Infusion; pRH14