



UNIVERSITY OF LEEDS

This is a repository copy of *Developing ESCRT-III as a toolkit for bottom-up construction of eukaryote-like artificial cells*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/128587/>

Version: Accepted Version

Conference or Workshop Item:

Booth, A, Marklew, C, Ciani, B et al. (1 more author) (2017) Developing ESCRT-III as a toolkit for bottom-up construction of eukaryote-like artificial cells. In: 19th IUPAB Congress & 11th EBSA Congress, 16-20 Jul 2017, Edinburgh, Scotland.

<https://doi.org/10.1007/s00249-017-1222-x>

c) European Biophysical Societies' Association 2017. This is a post-peer-review, pre-copyedit version of an article published in European Biophysics Journal The final authenticated version is available online at: <https://doi.org/10.1007/s00249-017-1222-x>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Developing ESCRT-III as a toolkit for bottomup construction of eukaryote-like artificial cells

A. Booth¹, C. Marklew², B. Ciani², P. A. Beales¹

¹University of Leeds, UK; ²University of Sheffield, UK

The Endosomal Complex Required for Transport (ESCRT) is a ubiquitous class of proteins involved in most membrane remodeling processes in eukaryotic cells. Previous studies have demonstrated that ESCRTs can be reconstituted in artificial lipid vesicle systems, notably giant unilamellar vesicles (GUVs), where they are able to induce the formation of intraluminal vesicles (ILVs), encapsulating material from the extravesicular solution.[1] Our aim is to develop the ESCRT proteins as a toolkit to generate GUV-derived structures, with multiple compartments containing different chemical environments, enzymes or molecular probes. The resulting cell-like structures may find applications in synthetic biology, drug delivery, diagnostics and the development of nano-reactor technology.[2] Using confocal microscopy and flow cytometry, we have gained insights into how protein stoichiometry and membrane mechanics influence the size and number of the ILV compartments obtained. We will further develop the ESCRT toolkit using the ATP-ase Vps4 to induce multiple encapsulation events and introduce rationally designed ESCRT chimera proteins to further simplify the system.

[1] T. Wollert, J. Lippincroft-Schwartz, J. H. Hurley, *Nature*, 2009

[2] P. A. Beales, B. Ciani, *Phys. Chem. Chem. Phys.*, 2015