

This is a repository copy of *The changing role of ornamental horticulture in alien plant invasions*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/128636/>

Version: Accepted Version

---

**Article:**

van Kleunen, Mark, Essl, Franz, Pergl, Jan et al. (25 more authors) (2018) The changing role of ornamental horticulture in alien plant invasions. *Biological reviews*. pp. 1421-1437. ISSN 1469-185X

<https://doi.org/10.1111/brv.12402>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1 **The changing role of ornamental horticulture in alien plant**  
2 **invasions**

3  
4 Mark van Kleunen<sup>1,2,\*</sup>, Franz Essl<sup>3</sup>, Jan Pergl<sup>4</sup>, Giuseppe Brundu<sup>5</sup>, Marta  
5 Carboni<sup>6</sup>, Stefan Dullinger<sup>3</sup>, Regan Early<sup>7</sup>, Pablo González-Moreno<sup>8</sup>, Quentin J.  
6 Groom<sup>9</sup>, Philip E. Hulme<sup>10</sup>, Christoph Kueffer<sup>11,12</sup>, Ingolf Kühn<sup>13,14</sup>, Cristina  
7 Máguas<sup>15</sup>, Noëlie Maurel<sup>2</sup>, Ana Novoa<sup>4,12,16</sup>, Madalin Parepa<sup>17</sup>, Petr Pyšek<sup>4,18</sup>,  
8 Hanno Seebens<sup>19</sup>, Rob Tanner<sup>20</sup>, Julia Touza<sup>21</sup>, Laura Verbrugge<sup>22,23</sup>, Ewald  
9 Weber<sup>24</sup>, Wayne Dawson<sup>25</sup>, Holger Kreft<sup>26</sup>, Patrick Weigelt<sup>26</sup>, Marten Winter<sup>14</sup>,  
10 Günther Klöner<sup>3</sup>, Matthew V. Talluto<sup>6</sup> & Katharina Dehnen-Schmutz<sup>27</sup>

11  
12 <sup>1</sup>*Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation,*  
13 *Taizhou University, Taizhou 318000, China*

14 <sup>2</sup>*Ecology, Department of Biology, University of Konstanz, Universitätsstrasse 10, D-78457*  
15 *Konstanz, Germany*

16 <sup>3</sup>*Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030*  
17 *Vienna, Austria*

18 <sup>4</sup>*Institute of Botany, Department of Invasion Ecology, The Czech Academy of Sciences, CZ-*  
19 *252 43 Průhonice, Czech Republic*

20 <sup>5</sup>*Department of Agriculture, University of Sassari, Viale Italia 39, 07100 Sassari, Italy*

21 <sup>6</sup>*Université Grenoble Alpes, CNRS, LECA, Laboratoire d'Écologie Alpine, F-38000*  
22 *Grenoble, France*

23 <sup>7</sup>*Centre for Ecology and Conservation, University of Exeter, Penryn Campus, UK*

24 <sup>8</sup>*CABI, Bakeham Lane, Egham TW20 9TY, UK*

25 <sup>9</sup>*Botanical Garden Meise, Bouchout Domain, Nieuwelaan 38, 1860 Meise, Belgium*

26 <sup>10</sup>*Bio-Protection Research Centre, Lincoln University, Lincoln 7648, Canterbury, New*  
27 *Zealand*

28 <sup>11</sup>*Institute of Integrative Biology, ETH Zurich, Universitätsstrasse 16, 8092 Zurich,*  
29 *Switzerland*

30 <sup>12</sup>*Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University,*  
31 *Matieland 7602, South Africa*

32 <sup>13</sup>*Helmholtz Centre for Environmental Research – UFZ, Dept. Community Ecology, Theodor-*  
33 *Lieser-Str. 4, 06120 Halle, Germany*

34 <sup>14</sup>*German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher*  
35 *Platz 5e, 04103 Leipzig, Germany*

36 <sup>15</sup>*Centre for Ecology, Evolution and Environmental Changes (cE3c), Faculty of Sciences,*  
37 *University of Lisbon, Campo Grande, 1749-016 Lisboa, Portugal*

38 <sup>16</sup>*Invasive Species Programme, South African National Biodiversity Institute, Kirstenbosch*  
39 *Research Centre, South Africa*

40 <sup>17</sup>*Institute of Evolution & Ecology, University of Tübingen, Auf der Morgenstelle 5, 72076*  
41 *Tübingen, Germany*

42 <sup>18</sup>*Department of Ecology, Faculty of Science, Charles University, Viničná 7, CZ-128 44*  
43 *Prague, Czech Republic*

44 <sup>19</sup>*Senckenberg Biodiversity and Climate Research Centre, Georg-Voigt-Straße 14-16, 60325*  
45 *Frankfurt, Germany*

46 <sup>20</sup>*European and Mediterranean Plant Protection Organization, 21 boulevard Richard Lenoir*  
47 *75011, Paris, France*

48 <sup>21</sup>*Environment Department, University of York, Wentworth Way, Heslington, YO10 5NG,*  
49 *York, UK*

50 <sup>22</sup>*Institute for Science in Society, Radboud University, PO Box 9010, 6500 GL Nijmegen, The*  
51 *Netherlands*

52 <sup>23</sup>*Netherlands Centre of Expertise for Exotic Species, Nijmegen, The Netherlands*

53 <sup>24</sup>*Biodiversity Research, University of Potsdam, Maulbeerallee 1, Potsdam D-14469,*  
54 *Germany*

55 <sup>25</sup>*Department of Biosciences, Durham University, South Road, Durham DH1 2LF, UK*

56 <sup>26</sup>*Biodiversity, Macroecology & Biogeography, University of Goettingen, Büsngenweg 1,*  
57 *37077 Göttingen, Germany*

58 <sup>27</sup>*Centre for Agroecology, Water and Resilience, Coventry University, Ryton Gardens,*  
59 *Coventry, CV8 3LG, UK*

60

61 **Running head:** Horticulture and plant invasions

62

63 \* Author for correspondence (E-mail: mark.vankleunen@uni-konstanz.de; Tel.: +49 7531 88  
64 2997).

65

66 **ABSTRACT**

67 The number of alien plants escaping from cultivation into native ecosystems is increasing  
68 steadily. We provide an overview of the historical, contemporary and potential future roles of  
69 ornamental horticulture in plant invasions. We show that currently at least 75% and 93% of  
70 the global naturalised alien flora is grown in domestic and botanical gardens, respectively.  
71 Species grown in gardens also have a larger naturalised range than those that are not. After  
72 the Middle Ages, particularly in the 18<sup>th</sup> and 19<sup>th</sup> centuries, a global trade network in plants

73 emerged. Since then, cultivated alien species also started to appear in the wild more  
74 frequently than non-cultivated aliens globally, particularly during the 19<sup>th</sup> century.  
75 Horticulture still plays a prominent role in current plant introduction, and the monetary value  
76 of live-plant imports in different parts of the world is steadily increasing. Historically,  
77 botanical gardens – an important component of horticulture – played a major role in  
78 displaying, cultivating and distributing new plant discoveries. While the role of botanical  
79 gardens in the horticultural supply chain has declined, they are still a significant link, with  
80 one-third of institutions involved in retail-plant sales and horticultural research. However,  
81 botanical gardens have also become more dependent on commercial nurseries as plant  
82 sources, particularly in North America. Plants selected for ornamental purposes are not a  
83 random selection of the global flora, and some of the plant characteristics promoted through  
84 horticulture, such as fast growth, also promote invasion. Efforts to breed non-invasive plant  
85 cultivars are still rare. Socio-economical, technological, and environmental changes will lead  
86 to novel patterns of plant introductions and invasion opportunities for the species that are  
87 already cultivated. We describe the role that horticulture could play in mediating these  
88 changes. We identify current research challenges, and call for more research efforts on the  
89 past and current role of horticulture in plant invasions. This is required to develop science-  
90 based regulatory frameworks to prevent further plant invasions.

91

92 *Key words:* botanical gardens, climate change, horticulture, naturalised plants, ornamental  
93 plants, pathways, plant invasions, plant nurseries, trade, weeds.

94

## 95 CONTENTS

96 I. Introduction ..... 5

97 II. Contemporary gardens and the naturalised alien flora of the world..... 7

98	III. The history of ornamental horticulture and implications for current plant invasions .....	9
99	(1) Garden-plant introductions .....	9
100	(2) Historical garden-fashion trends .....	15
101	IV. The recent role of horticulture in plant invasions .....	16
102	(1) Global patterns, changing dynamics and likely future trends.....	16
103	(2) Modern garden-fashion trends .....	19
104	(3) Horticultural selection favours traits related to invasiveness .....	20
105	V. The next generation of invading alien horticultural plants.....	22
106	(1) New pathways and horticultural practices .....	22
107	(2) Climate change .....	23
108	VI. Research opportunities and needs .....	25
109	VII. Conclusions.....	28
110	VIII. Acknowledgements .....	29
111	IX. References .....	30
112	X. Supporting information	

113

114 **I. INTRODUCTION**

115 With increasing globalisation, many plant species have been introduced beyond their natural  
116 ranges, and some of these have established and sustain persistent populations without human  
117 assistance (van Kleunen *et al.*, 2015; Pyšek *et al.*, 2017). Most of these alien species (*sensu*  
118 Richardson *et al.*, 2000) have comparatively small naturalised ranges (Pyšek *et al.*, 2017) and  
119 do not cause major ecological or economic damage. Some alien species, however, have  
120 become invasive (*sensu* Richardson *et al.*, 2000), impact upon native species, and can result  
121 in a significant burden on global economies, ecosystem services and public health (Pimentel,  
122 Zuniga & Morrison, 2005; Vilà *et al.*, 2011; Pyšek *et al.*, 2012*b*). Alien species introductions

123 have sometimes occurred unintentionally through various pathways (e.g. as seed  
124 contaminants), but most invasive alien plants have been introduced intentionally, particularly  
125 for cultivation as ornamentals in public and private gardens (Hulme *et al.*, 2008; Pyšek,  
126 Jarošík & Pergl, 2011).

127 Alien plant invasions have been facilitated by an increase in species traded and trade  
128 volumes, complexity of the trade network, improved long-distance connections, and new  
129 ways of trading (Humair *et al.*, 2015; Pergl *et al.*, 2017). The horticultural introduction  
130 pathway is characterised by a wide range of supply-chain actors (Fig. 1; also see Drew,  
131 Anderson & Andow, 2010; Hulme *et al.*, 2018), whose roles have changed over time  
132 (Daehler, 2008). Some of the first actors were professional ‘plant hunters’ – individuals who  
133 collected seeds, bulbs, roots and tubers of wild species for cultivation and trade. Although the  
134 heydays of plant hunting were in the 18<sup>th</sup> and 19<sup>th</sup> century, such practices continue today  
135 (Ward, 2004). Many of the species collected by plant hunters are not grown easily or are not  
136 chosen by breeders and propagators, limiting the eventual size of the cultivated species pool  
137 (Fig. 1). Through selection and hybridisation, however, breeders also create novel ornamental  
138 cultivars and species, increasing the gene pool for cultivation (Fig. 1). The availability of  
139 plant species through wholesalers and retailers largely determines the alien species that are  
140 cultivated in botanical gardens, public green spaces and domestic gardens, from which some  
141 of these alien species may escape into the wild and become invasive. While certain native  
142 species show similar behaviour to invasive alien species, we use the term ‘invasive’  
143 exclusively to refer to species that spread outside their native range through human  
144 intervention (Richardson *et al.*, 2000).

145 To interpret current trends and to predict likely future developments, we need a better  
146 understanding of the number and diversity of alien plants grown in gardens. Furthermore, we  
147 also need to know their introduction history and the species characteristics that promote both

148 their horticultural usage and potential invasion success. Therefore, we here integrate  
149 information from invasion biology and horticulture to provide a broad overview of the role of  
150 ornamental horticulture in alien plant invasions. We do this by (i) using a scheme describing  
151 the pathways and processes involved in ornamental plant invasions (Fig. 1; also see Drew *et*  
152 *al.*, 2010), (ii) covering a wide range of relevant issues, such as introduction dynamics,  
153 garden fashions and plant traits promoted by horticulture, from both historical and  
154 contemporary perspectives, (iii) discussing the potential future role of horticulture, and (iv)  
155 highlighting research needs.

156

## 157 **II. CONTEMPORARY GARDENS AND THE NATURALISED ALIEN FLORA OF** 158 **THE WORLD**

159 Regional analyses of alien naturalised floras have shown that usually more than half of these  
160 species were introduced for ornamental horticulture purposes (e.g. Germany: Kühn & Klotz,  
161 2002; Czech Republic: Pyšek *et al.*, 2012a; Britain: Clement & Foster, 1994; USA: Mack &  
162 Erneberg, 2002; Australia: Groves, 1998; South Africa: Faulkner *et al.*, 2016). Furthermore, a  
163 comparison of the frequency of invasive species across the world reveals that most have  
164 originated from ornamental horticulture (Hulme *et al.*, 2018). However, a global analysis of  
165 naturalised alien plants is still missing. In order to obtain a benchmark estimate of the  
166 proportion of naturalised species that have been introduced as garden plants globally, we  
167 compared the naturalised alien flora and the cultivated garden flora. The recently compiled  
168 Global Naturalized Alien Flora (GloNAF) database revealed that more than 13,000 vascular  
169 plant species have become naturalised somewhere in the world (van Kleunen *et al.*, 2015;  
170 Pyšek *et al.*, 2017). The number of plant species grown in domestic gardens, public green  
171 spaces and botanical gardens is much larger but precise numbers are yet unknown  
172 (Khoshbakht & Hammer, 2008). In order to obtain a minimum estimate of the size of the



173 global domestic garden flora, we extracted the lists of species in Dave's Garden PlantFiles  
174 (<http://davesgarden.com/guides/pf/>, accessed 23 March 2016) and in the Plant Information  
175 Online database (<https://plantinfo.umn.edu/>, accessed 22 November 2017). Furthermore, to  
176 obtain a minimum estimate of the number of species planted in botanical gardens, we  
177 extracted the list of species in the PlantSearch database of Botanic Gardens Conservation  
178 International ([http://www.bgci.org/plant\\_search.php](http://www.bgci.org/plant_search.php), accessed 25 May 2016), which includes  
179 species accessions of 1,144 botanical institutions worldwide. All species names were  
180 taxonomically harmonised using The Plant List (version 1.1; <http://www.theplantlist.org/>,  
181 accessed in December 2017), which also provided us with an estimate of the number of  
182 species in the global vascular plant flora. Ornamental cultivars that could not be assigned to  
183 species were not considered as they are not included in The Plant List.

184         At least 51% of all known species of vascular plants worldwide (337,137) are grown  
185 in domestic (70,108) or botanical gardens (162,846; Fig. 2). Most of the species grown in  
186 domestic gardens are also grown in botanical gardens (88%; Fig. 2), and it is likely that most,  
187 if not all species grown in public green spaces, for which we have no estimates, are also  
188 grown in domestic or botanical gardens (Mayer *et al.*, 2017). Although not all species in these  
189 gardens are cultivated for decorative purposes, and not all of them are cultivated outside their  
190 native ranges, these large numbers of garden species suggest that ornamental horticulture is  
191 the major pathway of alien plant introduction. Thus, it is not surprising that at least 75% and  
192 93%, respectively, of the naturalised alien plants worldwide are grown in domestic and  
193 botanical gardens (Fig. 2). Moreover, among the naturalised species, those grown in domestic  
194 or botanical gardens are also naturalised in more regions around the globe (Fig. 3).  
195 Furthermore, Hulme (2011) showed for the 450 invasive alien plant species listed in Weber  
196 (2003) that the number of regions in which each of these species is invasive is positively  
197 correlated with their frequency in botanical garden collections worldwide. Some of these

198 species may also have been introduced via additional pathways (e.g. agriculture or forestry).  
199 For example, *Robinia pseudoacacia* has been introduced as ornamental plant, forestry tree  
200 and nectar source, and for soil stabilization (Vítková *et al.*, 2017). Particularly, during the so-  
201 called utilitarian phase of the history of global weed movement (Mack & Lonsdale, 2001),  
202 the chances of becoming invasive may be high. So, while other deliberate introduction  
203 pathways are also important, there is strong evidence that ornamental horticulture remains a  
204 major contributor to plant invasions (Mack & Erneberg, 2002; Dehnen-Schmutz *et al.*, 2007;  
205 Hanspach *et al.*, 2008; Lambdon *et al.*, 2008; Hulme, 2011, Pyšek *et al.*, 2011; Pergl *et al.*,  
206 2016; Saul *et al.*, 2017; Hulme *et al.*, 2018).

207

### 208 **III. THE HISTORY OF ORNAMENTAL HORTICULTURE AND IMPLICATIONS** 209 **FOR CURRENT PLANT INVASIONS**

#### 210 **(1) Garden-plant introductions**

211 Archaeological evidence has revealed that plant species were transported by modern humans  
212 when humans expanded their range from the Late Pleistocene onwards (Bolvin *et al.*, 2016).  
213 Most of these alien species were used as food crops or as medicinal plants. It has also been  
214 speculated that Pleistocene people, and even Neanderthals, used ornamental flowers in burial  
215 sites (Leroi-Gourham, 1975). However, these claims are very controversial (Fiacconi &  
216 Hunt, 1995) and there is no evidence that these ornamentals were alien species. In the  
217 Americas, there is evidence for the existence of intensive trade of agricultural crops between  
218 areas in current Mexico and the coastal areas of Peru approximately 3000 years ago  
219 (Manrique, 2010). Around the same time, regions in current Panama had established a trade  
220 of plants with regions in current Ecuador, Colombia, Guatemala or Mexico (Sánchez, 1997).  
221 To what extent these traded plants included ornamentals remains unknown.

222            Since pre-Roman times, and increasingly with the Romans and in the Middle Ages,  
223 plant species were transported across Europe. In particular, Mediterranean plants were carried  
224 to other parts of Europe, and occasionally plants from more distant regions, such as Central  
225 and East Asia, were introduced to Europe (e.g. Jacomet & Kreuz, 1999; Campbell-Culver,  
226 2001). In their colonisation of Pacific islands, Polynesians introduced several crop and fibre  
227 species to Hawaii and later New Zealand (Cox & Barnack, 1991; Roullier *et al.*, 2013). From  
228 China, there is evidence of the early use of alien plants during the Han-Dynasty, where the  
229 new long-distance trade network of the ‘silk road’ was used to introduce ornamental alien  
230 plants for the extensive park created by Emperor Wu-Ti (140–89 BC; Hill, 1915; Keller,  
231 1994). In pre-Columbian Mexico, there were already gardens, such as that of the Acolhua  
232 king Netzahualcōyotl (1402–1472) and those of the Aztec kings Moctezuma I (1390–1469)  
233 and Moctezuma II (1465–1520), with plants collected in Mexico and elsewhere in the  
234 Americas (Hill, 1915; Sánchez, 1997). For other parts of the world, little or no information is  
235 available on such historical plant introductions.

236            It is known that roses were cultivated and traded as early as in the times of the ancient  
237 Romans, Greeks and Phoenicians (Harkness, 2003). For the medieval period, there are  
238 documents that detail the plants grown in the gardens of monasteries and castles. An example  
239 is Walafried Strabo’s *Liber de cultura hortorum*, published around the year 840 and  
240 describing 24 garden herbs. Although most of the species listed in these works were used as  
241 spices or as medicinal plants, some also had symbolic value and were appreciated as  
242 ornamentals (e.g. roses, lavender and poppies). Certain alien plant species introduced to  
243 medieval European castle gardens still persist as naturalised species in the areas around these  
244 castles today (e.g. *Erysimum cheiri*; Dehnen-Schmutz, 2004).

245            After the Middle Ages, global exploration by European nations expanded rapidly, the  
246 intercontinental exchange of species gained momentum, and eventually a truly global

247 network of plant species trade and exchange emerged (Mack, 2000). The explorers and plant  
248 hunters sent out by the different European countries in the 15<sup>th</sup> and 16<sup>th</sup> century were  
249 instructed to collect (economically) interesting plants (e.g. Stöcklin, Schaub & Ojala, 2003).  
250 Driven by the discoveries of new lands and the growing demands of private collectors,  
251 nurseries and botanical gardens for botanical novelties, plant hunting became a recognized  
252 occupation in Europe during the mid-16<sup>th</sup> century (Janick, 2007). In the 17<sup>th</sup> century, John  
253 Tradescant the elder and his son were among the first Europeans to explore the floras of the  
254 Middle East and Russia, and later North America (Reichard & White, 2001). They collected  
255 for example *Rhus typhina*, *Tradescantia virginiana* and *Liriodendron tulipifera* (Musgrave,  
256 Gardner & Musgrave, 1999), species that are now widely naturalised in different parts of the  
257 world. During the 18<sup>th</sup> and 19<sup>th</sup> centuries, many plant hunters collected plants for botanical  
258 institutions such as the Royal Botanical Gardens, Kew in the UK, the Leiden Hortus  
259 Botanicus in the Netherlands and the Jardin du Roi in France (Whittle, 1970), and for clubs of  
260 plant enthusiasts such as Der Esslinger Botanische Reiseverein in Germany (Wörz, 2016).  
261 During this period, plant exploration became very popular. For example, by the 18<sup>th</sup> century  
262 almost 9,000 ornamental plant species from all over the world were introduced to the British  
263 Isles (Clement & Foster, 1994). Many of the ornamental species currently naturalised in  
264 Europe were introduced in this period (e.g. Maurel *et al.*, 2016).

265         Similarly, many new ornamentals were introduced to North America from the 18<sup>th</sup> to  
266 the 20<sup>th</sup> centuries from plant-collection expeditions in Eastern and Central Asia, North Africa  
267 and the Middle East (Stoner & Hummer, 2007). During the first expedition of this kind  
268 funded by the federal government of the USA, Robert Fortune (1812–1880) introduced  
269 species of *Chrysanthemum*, *Paeonia* and *Rhododendron* (azaleas) as ornamentals into the  
270 USA (Musgrave *et al.*, 1999). Another noteworthy plant hunter was Ernest Henry Wilson  
271 (1876–1930), who introduced >2,000 plant species from Asia to Europe and North America.

272 Some of these species, such as *Lonicera maackii* and *Pyrus calleryana* (Farrington, 1931),  
273 are now widely naturalised in North America (<http://bonap.org/>). Taken together, the efforts  
274 of plant hunters brought many new species to botanical gardens and private collections, and  
275 fuelled the horticultural trade from the 16<sup>th</sup> until the early 20<sup>th</sup> century.

276 Governments also played active roles in alien plant introductions. For example, US  
277 President John Quincy Adams (1767–1829) requested all US consuls to forward rare seeds to  
278 Washington for distribution (Hodge & Erlanson, 1956). In 1839, the US Congress  
279 appropriated \$1000 for the handling and distribution of seeds of introduced alien plants, and  
280 the United States Department of Agriculture (USDA) created in 1898 the Office of Foreign  
281 Plant Introductions with the aim of building up new plant industries (Fairchild, 1898; Hodge  
282 & Erlanson, 1956). Until the end of World War II, the USDA office introduced  
283 approximately 250,000 accessions (i.e. species and varieties combined), and coordinated the  
284 initial propagation, testing and distribution of the plants (Hodge & Erlanson, 1956). Most of  
285 these plants were introduced for agricultural purposes, but they also included species for  
286 ornamental horticulture (Fairchild, 1898; Dorsett, 1917). Similarly, government agencies  
287 were responsible for the introduction of alien plant species in countries like Australia (Cook  
288 & Dias, 2006) and New Zealand (Kirkland & Berg 1997).

289 Ornamental alien plants were not only introduced to the home countries of the  
290 predominantly European plant hunters, but plants native to Europe were also introduced into,  
291 and exchanged among the colonies. An important role in this exchange was played by the  
292 acclimatisation societies, which arose in Europe and its colonies during the 19<sup>th</sup> century.  
293 Initially, the acclimatisation societies were fuelled by interest in novel flora and fauna from  
294 the colonies for introduction into European gardens and zoos (Dunlap, 1997). Later, the focus  
295 changed to transplanting the biotic landscape from the mother country into the colonies and  
296 the exchange of ornamental and crop species among colonies (di Castri, 1989; Osborne,

297 2001). Subsidies and free transport of explorers, plants and animals on cargo ships to and  
298 from the colonies was offered by supporting governments (Grove, 1995). Many crops but  
299 also ornamentals were transported this way, including bamboos and species of *Araucaria*,  
300 *Acacia* and *Camellia* (Bennett, 1870). Soon after their foundation, popularity of the  
301 acclimatisation societies waned due to growing concerns for the preservation of indigenous  
302 biota (Dunlap, 1997). Twenty years after their rapid appearance, most acclimatisation  
303 societies had been dissolved, and the few remaining ones started to focus on reintroduction of  
304 threatened native species.

305         While botanical gardens were used as showcases by the acclimatisation societies in  
306 the second half of the 19<sup>th</sup> century, their role in introducing and cultivating alien plants  
307 started much earlier and continues today. Particularly, during the 17<sup>th</sup> and 18<sup>th</sup> century,  
308 botanical gardens were part of the colonial infrastructure that facilitated the distribution of  
309 useful plants around the world (Hulme, 2011). Between 1750 and 1850, the first botanical  
310 gardens were founded in all non-European continents (with the exception of Antarctica):  
311 Bartram's Garden (1728) in North America, the Calcutta Botanic Garden (1786) in Asia, the  
312 Sydney Gardens (1788) in Australia, the Rio de Janeiro Botanical Garden (1808) in South  
313 America, and Cape Town Botanic Garden (1848) in Africa (Hill, 1915). Botanical gardens  
314 were also instrumental in the collation, evaluation and dissemination of new discoveries of  
315 foods, agricultural products and ornamentals, generally sponsored by governments and  
316 commercial enterprises (e.g. Diagre-Vanderpelen, 2011). Unsurprisingly, many of the  
317 currently naturalised and invasive alien plant species were first planted in botanical gardens.  
318 For example, in Europe, *Solidago canadensis* and *S. gigantea* were first planted in Paris and  
319 London, respectively (Wagenitz, 1964; Weber, 1998), and *Agave americana* was first planted  
320 in the Padua Botanical Garden (Italy; [http://www.ortobotanicopd.it/en/piante-introdotte-](http://www.ortobotanicopd.it/en/piante-introdotte-italia-dallorto-botanico)  
321 [italia-dallorto-botanico](http://www.ortobotanicopd.it/en/piante-introdotte-italia-dallorto-botanico); accessed 23 March 2017). Many of the species introduced to

322 botanical gardens may first have been distributed to other gardens and public green spaces  
323 before they escaped into the wild. However, some alien species escaped directly from  
324 botanical gardens (Harris, 2002; Sukopp, 2006), including several listed among the worst  
325 aliens worldwide (Hulme, 2011).

326         With the emergence and intensification of the global network of ornamental plant  
327 species trade after the Middle Ages, it is not surprising that the rate at which new alien  
328 species established in the wild increased dramatically (Seebens *et al.*, 2017). Some of these  
329 species were not introduced intentionally for their economic and ornamental value, but were  
330 accidentally transported with other cargo or in ballast soil (e.g. Brown, 1878; Hulme *et al.*,  
331 2008). The exact role of ornamental horticulture in the temporal dynamics of naturalisation  
332 events is therefore difficult to quantify. To gain some insights, we used the database of  
333 Seebens *et al.* (2017) on first-record rates of established alien plants in combination with data  
334 on their cultivation in domestic (data from Dave’s Garden PlantFiles and the Plant  
335 Information Online database) and botanical (data from Botanic Gardens Conservation  
336 International PlantSearch database) gardens. The first-record rate in the 19<sup>th</sup> century  
337 increased faster for species that are now cultivated in gardens, particularly in botanical  
338 gardens, than for species not known to be cultivated (Fig. 4). This suggests that species  
339 introduced for horticultural purposes naturalised earlier than alien species introduced by other  
340 pathways. However, while the first-record rates of species grown in domestic gardens only  
341 and species not known to be cultivated are still increasing rapidly, the first-record rate  
342 appears to slow down for species grown in botanical gardens (Fig. 4). Possibly, this is partly  
343 a consequence of the increasing awareness about invasive plants among botanical gardens  
344 and their stronger focus on native plants in recent times (Hulme, 2015).

345

346 **(2) Historical garden-fashion trends**

347 Changing garden and landscaping fashions impact on plant introductions and subsequent  
348 invasions through floral design, style elements and layouts of gardens, parks and other green  
349 spaces, as well as through the choice of plants they promote (e.g. Müller & Sukopp, 2016).  
350 Historic fashion trends were not only driven by demand but also by the chronological order in  
351 which plants from different parts of the world became available. For example, with the  
352 discovery of the New World, novel ornamental plants were introduced into European  
353 horticulture as early as the 16<sup>th</sup> century, many of which are still common in today's gardens –  
354 e.g. *Helianthus* spp., *Amaranthus caudatus* and *Mirabilis jalapa*. Increased trade with the  
355 Orient also opened the door to plants from Asia (e.g. *Hemerocallis* spp.) into Europe. While  
356 most of these species are herbaceous, the development of landscape gardens and arboreta in  
357 the 18<sup>th</sup> and 19<sup>th</sup> centuries marked the start of the widespread introduction of ornamental trees  
358 to Europe (see e.g. Goeze, 1916). Landscape gardens were characterised by the opening up of  
359 gardens into a wider landscape accompanied by careful positioning of artificial lakes, trees  
360 and hedges. Many alien trees introduced to create such gardens still characterise urban parks  
361 today, and some of them – such as the North American species *Acer negundo*, *Robinia*  
362 *pseudoacacia*, *Pinus strobus*, *Prunus serotina* and *Quercus rubra* – have also become  
363 naturalised in Europe and elsewhere (Brundu & Richardson, 2016; Richardson & Rejmánek,  
364 2011; Campagnaro, Brundu & Sitzia, 2017).

365         The second half of the 19<sup>th</sup> century saw the development of ecologically and  
366 biogeographically focused plantings that aimed to recreate representative examples of  
367 specific vegetation types from around the world (Woudstra, 2003). This period also saw a  
368 broadening interest in different growth forms besides plantings of woody species, with an  
369 increasing representation of perennial forbs and later also grasses. Specific habitats such as  
370 rockeries, bogs and woodlands were created in gardens to accommodate high plant diversity.



371 Plant recommendations for these habitats in Britain were provided by William Robinson with  
372 his influential book *The wild garden or, our groves and shrubberies made beautiful by the*  
373 *naturalization of hardy alien plants* (Robinson, 1870). The trend of using hardy perennial  
374 plants continued into the 20<sup>th</sup> century, first driven by the desire to create *Colour in the flower*  
375 *garden* as Gertrude Jekyll (1908) titled her influential book. It was also influenced by the  
376 ornamental plant breeder Karl Foerster (1874–1970), one of the first to promote the use of  
377 grasses as ornamentals in Germany (Hottenträger, 1992). These are just a few of the  
378 individuals that influenced garden fashions in Europe. Examples of influential people in the  
379 Americas are Andrew Jackson Downing (1815–1852) and Frederick Law Olmsted (1822–  
380 1903), who both preached the English or natural style of landscape gardening, and more  
381 recently Thomas Church (1902–1978), who designed the ‘California Style’ of garden  
382 landscapes (<https://www.gardenvisit.com>, accessed 28 November 2017). The consequences  
383 of these different ‘garden fashions’ initiated by these people on plant invasions in different  
384 regions of the world still need more research.

385

#### 386 **IV. THE RECENT ROLE OF HORTICULTURE IN PLANT INVASIONS**

##### 387 **(1) Global patterns, changing dynamics and likely future trends**

388 Horticulture continues to play a prominent role in alien plant introductions (Reichard &  
389 White, 2001; Bradley *et al.*, 2011; Humair *et al.*, 2015). This is confirmed by analyses of the  
390 monetary value of live-plant imports in different parts of the world, which show a steady  
391 increase in live-plant imports in Europe and North America (Fig. 5). This may, however, not  
392 necessarily translate into a higher diversity of species traded, as such trade statistics do not  
393 specify the number of species traded, and include non-ornamental plants. Live-plant imports  
394 in South and Central Asia are rising at an increasing rate, and, while imports to East Asia  
395 appear to have undergone a rise and fall at the end of the 1990s, imports are increasing once

396 again (Fig. 5). Understanding who is involved in horticulture in these regions would help  
397 invasive-plant management plans to be targeted to the appropriate audience.

398         The most data on the role of ornamental horticulture in plant invasions are available  
399 for Europe and North America. However, horticulture was recently identified as a strong  
400 driver of invasions in Argentina (Giorgis & Tecco, 2014), Brazil (Zenni, 2014), and Puerto  
401 Rico and the Virgin Islands (Rojas-Sandoval & Acevedo-Rodríguez, 2014). This is despite  
402 slow growth of live-plant imports to the Caribbean, Central and South America (Fig. 5).  
403 Furthermore, while gardening is a popular hobby in North America, Australasia and Europe  
404 (Bradbury, 1995; Crespo *et al.*, 1996; Soga, Gaston & Yamaura, 2017), information on the  
405 prevalence of recreational gardening outside these regions is harder to find. In Japan, one in  
406 four people gardens daily, and at least five studies have assessed the effect of gardening on  
407 mental health in Asia (Soga *et al.*, 2017), suggesting public interest in this hobby.

408         The establishment of botanical gardens was historically driven by the needs of  
409 economic botany and ornamental horticulture. This role has decreased with the increasing  
410 importance of many botanical gardens in global plant conservation (Havens *et al.*, 2006).  
411 Currently, private and public sector breeding programs play major roles in the release of alien  
412 plants through the ornamental nursery supply-chain. The role of botanical gardens in the  
413 ornamental nursery supply-chain, however, is not negligible (Fig. 1; Hulme 2011, 2015). An  
414 analysis of the Botanic Garden Conservation International (BGCI) Garden Search database  
415 ([http://www.bgci.org/garden\\_search.php](http://www.bgci.org/garden_search.php), accessed on 1 November 2016) shows that  
416 approximately one-third of botanical gardens worldwide are involved in retail-plant sales,  
417 particularly in developing countries (Fig. 6). Similarly, approximately one-third of botanical  
418 gardens undertake horticultural research and around 10% are involved in plant breeding (Fig.  
419 6). In both cases, the levels of participation in this research seem particularly high in Asia,  
420 and low in North America ( $\chi^2=28.02$  and  $26.03$ ,  $df=5$ ,  $P < 0.0001$ , respectively).

421 Nevertheless, North American botanical gardens play a leading role in using their living  
422 collections of alien ornamentals as a basis for commercial breeding and marketing (Pooler,  
423 2001; Kintgen, Krishnan & Hayward, 2013; Ault & Thomas, 2014).

424 The participation of botanical gardens in plant exploration varies among continents  
425 ( $\chi^2=48.02$ ,  $df=5$ ,  $P < 0.0001$ ), and is most important in continents with many developing  
426 countries, Asia in particular (Fig. 6). While much of this exploration advances the knowledge  
427 of the native flora, it also highlights a potential route for new ornamental plants to enter the  
428 global horticulture market. The combination of a rapid growth in numbers and importance of  
429 botanical gardens in Asia (Hulme, 2015), an increased emphasis on horticulture and breeding  
430 research in these institutions and a significant role of retail-plant sales suggest that Asia will  
431 contribute to increasing global trade in ornamental plants in the future. This is certainly the  
432 philosophy and expectation of botanical gardens in China (Zhao & Zhang, 2003). Given the  
433 increasing evidence that alien plants from Asia are particularly successful invaders elsewhere  
434 in the world (Lambdon *et al.*, 2008; Fridley & Sax, 2014; van Kleunen *et al.*, 2015), we can  
435 expect even more horticulture-driven plant invasions from Asia in the future.

436 With already a significant proportion of the global flora in cultivation (Fig. 2) and  
437 increased availability of plant propagules through other sources, wild collection has probably  
438 decreased in the last decades. It is likely to decrease further due to global restrictions on  
439 collecting wild plants imposed by the Nagoya Protocol on access and benefit-sharing of the  
440 Convention of Biological Diversity (2011; <https://www.cbd.int/abs/>). This means that home  
441 gardens and plantings in public green spaces will rely on nurseries, but also that botanical  
442 gardens will have to maintain or expand their collections using commercially bought plant  
443 material or through exchange with other botanical gardens. To obtain an impression of the  
444 importance of different plant sources for current botanical garden collections, we sent a  
445 questionnaire to botanical gardens around the globe (Appendix 1). Of the 161 respondents,

446 37%, 29% and 27% indicated that their major sources of plants are commercial nurseries,  
447 other botanical gardens and collections from the wild, respectively (Fig. 7). Commercial  
448 nurseries were particularly important sources for North American botanical gardens, whereas  
449 other botanical gardens were particularly important sources for European botanical gardens  
450 (Fig. 7). The latter might reflect that many European botanical gardens produce an Index  
451 Seminum (i.e. seed catalogue) of the species available for exchange.

452

## 453 **(2) Modern garden-fashion trends**

454 Since the 1990s, there has been a resurgence in cultivating herbaceous perennials, frequently  
455 prairie species from North America, in more naturalistic plantings. This is motivated by the  
456 ease and low costs of management and by an increased interest in species-rich gardens  
457 (Hitchmough & Woudstra, 1999). These plantings often combine native and alien species that  
458 originate from different continents but belong to the same habitat type (e.g. prairies).  
459 Regarding other more recent gardening fashions, few formal studies exist that document  
460 them, and even fewer link them to plant invasions (e.g. Dehnen-Schmutz, 2011; Humair,  
461 Kueffer & Siegrist, 2014a; Pergl *et al.*, 2016). For example, although the surge in invasive  
462 aquatic plants is most likely the result of increasing interest in water gardening since the  
463 middle of the 20<sup>th</sup> century, robust data are hard to find (Maki & Galatowitsch, 2004). Other  
464 recent fashions are ‘jungle’ and desert gardens, living walls, and guerrilla gardening (i.e.  
465 gardening on land not owned by the gardener), all of which depend on and promote their own  
466 selection of mainly alien plants (Dunnett & Kingsbury, 2008; Reynolds, 2014). There is also  
467 a rising interest in increasing the services provided by urban vegetation, such as food  
468 production (Smardon, 1988), and therefore an increasing number of urban parks include  
469 ornamental aliens that are edible (Viljoen, Bohn & Howe, 2005). In addition to the fashion  
470 trends that mainly use alien plants, there is also an increasing interest in gardening with

471 native species (e.g. Kruckeberg, 2001; Shaw, Miller & Wescott, 2017). This is likely due to  
472 awareness of biological invasions but also because people want to have gardens that promote  
473 diversity and wildlife, and are less labour intensive.

474

### 475 **(3) Horticultural selection favours traits related to invasiveness**

476 The horticultural industry identifies particularly prized species, varieties or cultivars through  
477 specific accolades, e.g. Awards of Garden Merit (Great Britain), Mérites de Courson  
478 (France), All-America Selection Winners (USA), Gold Medal Plant (Pennsylvania). Such  
479 accolades are an important marketing strategy to promote specific plants, and are an  
480 important aspiration for many ornamental plant breeders. While the criteria differ for  
481 individual accolades, in general the plants must be excellent for garden use, exhibit  
482 consistently good performance in different garden environments and climates, should be easy  
483 to grow, and should not be particularly susceptible to insect pests or pathogens (Hulme,  
484 2011). Such characteristics, together with the higher market frequency of these species may  
485 have contributed to the high propensity of award-winning plants to become invasive (Hulme,  
486 2015).

487         There are several plant characteristics that might promote both horticultural use and  
488 invasion. Environmental matching is an obvious criterion when considering a species for  
489 horticulture (Reichard, 2011), and at the same time is also important for naturalisation and  
490 invasiveness (Richardson & Pyšek, 2012). For example, in Germany – a temperate region  
491 with winter frost – hardier species are planted more frequently (Maurel *et al.*, 2016) and have  
492 a higher probability of naturalisation (Hanspach *et al.*, 2008; Maurel *et al.*, 2016) than less  
493 hardy species. Horticultural usage should also be favoured by ease of propagation (Mack,  
494 2005; Reichard, 2011), and alien species with rapid and profuse seedling emergence are also  
495 more likely to naturalise (van Kleunen & Johnson, 2007). Similarly, fast vegetative growth is

496 promoted by the horticultural industry (Reichard, 2011), and also promotes invasiveness of  
497 plants (Dawson, Fischer & van Kleunen, 2011; Grotkopp, Erskine-Ogden, & Rejmánek,  
498 2010). Furthermore, early-flowering species and genotypes often have a long flowering  
499 period or have repeated bouts of flowering (Mack, 2005) and can be sold sooner or for a  
500 longer time, thus increasing profit (Reichard, 2011). At the same time, a longer flowering  
501 period has also been found to be associated with invasiveness (Lloret *et al.*, 2005; Gallagher,  
502 Randall & Leishman, 2015). So, horticulture may facilitate plant invasions by screening  
503 species and genotypes of ornamental value based on traits that inadvertently promote spread  
504 (Drew *et al.*, 2010; Knapp *et al.*, 2012).

505         Although horticulture seems to foster plant invasions overall by filtering species based  
506 on characteristics that increase their success inside and outside of gardens, this is not  
507 systematically the case. In some taxonomic groups, the most valued species are actually the  
508 ones with traits that make them less successful outside of gardens. For example, among cacti,  
509 slow-growing species are usually favoured by gardeners (Novoa *et al.*, 2017), and they  
510 should be less likely to naturalise and become invasive (Novoa *et al.*, 2015b). For orchids,  
511 which are strongly underrepresented in the global naturalised flora (Pyšek *et al.*, 2017), some  
512 hobby growers are willing to pay more for species that are rare in trade and most likely  
513 difficult to cultivate (Hinsley, Verissimo & Roberts, 2015). Furthermore, many ornamental  
514 cultivars have showy flowers that are sterile (e.g. in roses; Debener *et al.*, 2001), which  
515 diminishes their invasion potential. Thus, there is potential to select ornamental species or  
516 breed cultigens that are less likely to become invasive.

517         To date there has been very limited involvement of plant breeders in reducing  
518 invasion risk of ornamental plants (e.g. Burt *et al.*, 2007; Novoa *et al.*, 2015a). Anderson,  
519 Gomez & Galatowitsch (2006) proposed 10 traits to reduce invasiveness while retaining  
520 commercial value of ornamentals: reduced genetic variation in propagules, slowed growth

521 rates, non-flowering, elimination of asexual propagules, lack of pollinator rewards, non-  
522 dehiscing fruits (to prevent seed dispersal), lack of edible fruit flesh, lack of seed  
523 germination, sterility and programmed death prior to seed production. So far, most effort in  
524 producing non-invasive cultivars has focussed on reduced fecundity (e.g. Freyre *et al.*, 2016).  
525 Unfortunately, for perennial species, even relatively low levels of seed production may be  
526 sufficient for plant invasions (Knight, Havens & Vitt, 2011). Furthermore, traits such as seed  
527 sterility and dwarfism, bred into cultivars to reduce invasion potential, may revert back to  
528 their original states (Brand, Lehrer & Lubell, 2012). Perhaps the way forward is for  
529 horticultural accolades to recognise the risk of invasiveness more formally and at least  
530 account for this in field trials and subsequent selection of award-winning taxa.

531

## 532 **V. THE NEXT GENERATION OF INVADING ALIEN HORTICULTURAL PLANTS**

### 533 **(1) New pathways and horticultural practices**

534 A major future challenge might be that social, technological and environmental changes will  
535 lead to fundamentally novel patterns of plant introductions resulting in invasion risks by new  
536 types of plants for which past invasions give only partial guidance (Kueffer, 2010). Through  
537 internet trade, a much broader range of taxa from many more source regions becomes  
538 available for buyers worldwide (Humair *et al.*, 2015). Many of these new species might  
539 initially be traded in low numbers, but marketing, promotion by celebrity gardeners, and  
540 popularity in social media of specialised gardening groups can result in sudden interest in a  
541 new plant species. One example is the recent rise in trade and illegal import into Europe of  
542 *Lycium barbarum*, the shrub that produces the putative ‘superfood’ goji berry (Giltrap, Eyre  
543 & Reed, 2009) and is widely naturalised in Europe ([http://www.europe-](http://www.europe-aliens.org/speciesFactsheet.do?speciesId=20401#)  
544 [aliens.org/speciesFactsheet.do?speciesId=20401#](http://www.europe-aliens.org/speciesFactsheet.do?speciesId=20401#), accessed on 13 July 2017). Unsurprisingly,  
545 horticulturalists are continually searching for new plants with ‘unique’ features to be sold.

546 Seaton, Bettin & Grüneberg (2014 ) for instance wrote that “Introduction of new plants is  
547 critical to the survival and profitability of the horticultural industries” in their article on how  
548 to find new plant species in the world’s existing plant diversity. Furthermore, new molecular-  
549 based breeding technologies have reached the horticultural industry (e.g. Chandler &  
550 Brugliera, 2011; Xiong, Ding & Li, 2015). One primary target of current breeding efforts is  
551 to increase resistance to diseases and herbivores, which could then also increase invasiveness  
552 of some cultivars.

553

## 554 **(2) Climate change**

555 Environmental changes, such as atmospheric nitrogen deposition, habitat fragmentation and  
556 disturbance due to land-use change, have contributed to plant invasions and are likely to do  
557 so in the future (Bradley *et al.*, 2010; Sheppard, Burns & Stanley, 2014; Dullinger *et al.*,  
558 2017; Liu *et al.*, 2017). In addition, it is commonly expected that climate change will increase  
559 plant invasions globally, although its impacts may vary considerably among geographic areas  
560 and species (Lambdon *et al.*, 2008; Hulme, 2009; Bradley *et al.*, 2010; Seebens *et al.*, 2015;  
561 Early *et al.*, 2016; Dullinger *et al.*, 2017). This expectation is mainly based on the anticipated  
562 destabilisation of resident native plant communities caused by an emerging disequilibrium  
563 with climatic conditions (Svenning & Sandel, 2013) and by increased frequencies of extreme  
564 events, such as droughts, hurricanes and heat waves (Diez *et al.*, 2012). Both will likely  
565 decrease the biotic resistance of resident vegetation against the establishment and spread of  
566 alien species (e.g. Eschtruth & Battles, 2009; Early *et al.*, 2016; Haeuser, Dawson & van  
567 Kleunen, 2017).

568         Although climatic suitability is an important criterion in horticulture, many  
569 ornamental species are grown beyond the climatic conditions they would be able to tolerate in  
570 the wild (Van der Veken *et al.*, 2008). A warming climate potentially increases the match  
571 between current cultivation areas and suitable climatic conditions, especially in temperate



572 regions where many garden plants have been introduced from warmer parts of the world  
573 (Niinimets & Peñuelas, 2008; Bradley *et al.*, 2011; Dullinger *et al.*, 2017). Cultivated  
574 ornamental plants will have a ‘head start’ (Van der Veken *et al.*, 2008) allowing them to  
575 colonise newly suitable areas long before other range-shifting species arrive. This head-start  
576 advantage may become even more important in the coming decades. First, adaptation of  
577 gardeners’ demands to anticipate changes in regional climates could improve the climatic  
578 match of newly planted species. Demand for drought-tolerant ornamental species is already  
579 growing in the USA in response to forecasted drier conditions (Bradley *et al.*, 2011). Second,  
580 rising urbanisation all around the world will lead to an increased concentration of demand for  
581 ornamental plants in metropolitan areas. These areas usually have higher temperatures than  
582 the surrounding rural areas (i.e. the urban heat-island effect). Consequently, warm-adapted  
583 garden plants will have the chance to establish naturalised populations in cities, which may  
584 facilitate their spread into the surrounding landscapes (e.g. Essl, 2007; but see Botham *et al.*,  
585 2009).

586         A warming climate may also foster the establishment of ornamental plants in those  
587 ecosystems that have so far been less affected by biological invasions. Mountains, for  
588 example, have few invasive species so far due to climatic constraints and low human  
589 population densities, and hence low propagule pressure (Pauchard *et al.*, 2016). Indeed, the  
590 few alien species currently found in mountains are mostly lowland generalists able to cope  
591 with the cold climate (Alexander *et al.*, 2011). However, climate warming, in combination  
592 with changing land use and increased tourism, will potentially relax these constraints and  
593 increase invasion risks at higher elevations (Pyšek *et al.*, 2011; Petitpierre *et al.*, 2016;  
594 Dainese *et al.*, 2017). Specifically, ornamental plants currently cultivated in mountain  
595 villages and resorts will have a head start under a warming climate and profit from greater  
596 propagule availability with increasing human population (Pauchard *et al.*, 2009). Further, in

597 order to satisfy the growing demands of tourism, nurseries selling into mountainous regions  
598 are also likely to increase the supply of garden plants pre-adapted to mountain conditions, i.e.  
599 originating from other alpine environments around the world (Kueffer *et al.*, 2013; Alexander  
600 *et al.*, 2017). The threat posed to mountains by escaping ornamental plants will thus probably  
601 increase in the future because of globalisation and climate change.

602

## 603 **VI. RESEARCH OPPORTUNITIES AND NEEDS**

604 To address new research frontiers identified in this overview, we provide an agenda of  
605 pressing research challenges that lie ahead in order to foster our understanding of the role of  
606 horticulture in plant invasions (Table 1). One overarching scientific challenge is advancing  
607 our understanding of how different practices, related features and characteristics of  
608 horticulture, and processes and impacts of plant invasions are linked to one another (Fig. 1).  
609 This will benefit greatly from an interdisciplinary scientific approach that jointly considers  
610 the human dimensions (e.g. behaviour, preferences, governance, culture), and their  
611 interactions with the biophysical environment. Addressing this topic in well-circumscribed  
612 study systems may be an appropriate way forward. *Inter alia* this can be achieved by  
613 focussing research questions on specific geographical regions or by focusing on subsets of  
614 ornamental species (e.g. certain families, or species with certain traits). This general research  
615 background can be broken down into eight specific research challenges (Table 1).

616 **Topic 1: an improved understanding of the origins of ornamental alien species**  
617 **and the means by which they arrive and are distributed.** Here, it is important to go  
618 beyond analyses on where from and by which pathway the most successful (most frequent)  
619 species, or those with the highest impacts arrived. It is crucial to take into account the species  
620 pool in the area of their origin and the trade pattern and volume to disentangle the effect of  
621 propagule pressure ('transport mass effect') from other factors related to invasion success or

622 impact. In this light, it is also important to know how species are distributed through new  
623 ways of trading or social networks. For example, how important is garden-plant exchange  
624 among relatives and friends (Verbrugge *et al.*, 2014)? In addition, there might be certain plant  
625 traits associated with specific origins and pathways.

626 **Topic 2: knowledge of temporal trends and fashions related to import and the**  
627 **consequences for invasion success and impact.** For example, are species that were  
628 introduced earlier more likely to be invasive now because they have had more time to  
629 become invasive or because plant hunters initially introduced plant species that could be  
630 cultivated easily and thus are better pre-adapted and more competitive? How do changes in  
631 breeding, fashions, and cultivation patterns affect plant invasions and impacts?

632 **Topic 3: improve understanding of the drivers of horticulture-related plant**  
633 **invasions including the identification of future invaders.** For example, what are the roles  
634 of changing trade partners and consequently trade patterns, plant traits and environmental  
635 conditions in invasion success, and how can the different drivers be ranked in importance?  
636 This, to some degree, is different from, but can be dependent on, origins and pathways.

637 **Topic 4: forecasting whether global environmental change will influence the**  
638 **naturalisation of ornamental species that were not a problem in the past.** Emerging  
639 patterns in global environmental change, like for example increased landscape fragmentation  
640 and climate change impacts, might differ among regions and among habitats (i.e. some  
641 combinations of these changes may synergistically promote invasions, while other  
642 combinations may inhibit invasions). Moreover, some of the solutions proposed to help  
643 native species survive might also affect plant invasions. For example, the creation of habitat  
644 corridors to promote dispersal and migration of native species in the light of habitat  
645 fragmentation and climate change may also benefit invasive alien species (Procheş *et al.*,

646 2005). However, it is not known whether these corridors provide appropriate dispersal habitat  
647 for many ornamental alien species.

648 **Topic 5: a much better understanding of the current and future impacts of**  
649 **horticulture-related plant invasions.** For instance, what are the impacts of horticultural  
650 invaders on biodiversity, human livelihoods, and ecosystem services provision, including  
651 cultural ecosystem services; and where do they occur?

652 **Topic 6: evaluation and development of tools for detecting, managing and**  
653 **monitoring of horticulture-driven plant invasions.** Based on evaluations of current early-  
654 detection programs, this should involve developing best practices for comprehensive early-  
655 detection programs for colonising and spreading alien horticultural species. This should  
656 consider how effective monitoring and prevention strategies can be implemented, and which  
657 management methods would be most efficient and effective.

658 **Topic 7: legal regulations that permit a thriving industry with a low risk of plant**  
659 **invasions.** First, one would need to review the existing regulatory frameworks (Hulme *et al.*,  
660 2018), identify gaps, address the demands of nature conservation to prevent the spread of  
661 ornamental species, and investigate how to promote the success of novel schemes (e.g.  
662 assurance schemes) in the industry that can incentivise behavioural changes. Given the  
663 diversity of stakeholders, this needs to be done sensitively to gain support from a diverse  
664 community. Importantly, sufficient long-term funding should be made available for  
665 monitoring by regulatory agents and land managers.

666 **Topic 8: public awareness and building partnerships with stakeholders.** Finally,  
667 we need to inform, educate and convince the public to promote native or benign alien plants  
668 as ornamentals rather than detrimental ones. Public awareness campaigns need to be  
669 underpinned by research on the role of cultural and social values in processes leading to new  
670 introductions. In addition to raising awareness, we need to build long-term, enduring

671 partnerships with stakeholders, such as the plant industry, gardeners and the public (Humair,  
672 Siegrist & Kueffer, 2014*b*). They harness important knowledge about how to regulate trade  
673 and inform the involved actors. Moreover, they are also interested in avoiding unregulated  
674 trade that leads to the introduction of new plant diseases and pests.

675

## 676 **VII. CONCLUSIONS**

677 (1) It is clear that ornamental horticulture is the major introduction pathway of naturalised  
678 and invasive alien plants (Figs 2 and 3). Therefore, a better knowledge and understanding of  
679 the ornamental plant supply chain (Fig. 1) and historical changes therein might help us  
680 predict the potential next generation of plant invaders.

681 (2) The efforts of plant hunters brought many new species to botanical gardens and private  
682 collections, and fuelled the horticultural trade. Species that came in through this horticultural  
683 pathway naturalised earlier than alien species introduced by other pathways (Fig. 4).

684 (3) Garden fashions, and the plant species promoted by them, have changed in the last  
685 centuries, and differ among regions. However, the consequences of the different garden  
686 fashions on plant invasions still need more research.

687 (4) The horticultural industry continues to play a prominent role in alien plant introductions,  
688 as is evident from the high monetary value of the live-plant import market in different parts  
689 of the world (Fig. 5). Botanical gardens still play an important role in horticultural activities  
690 (Fig. 6), but their collections have become more dependent on commercial nurseries and  
691 exchange among botanical collections than on wild collection (Fig. 7).

692 (5) Some of the species traits promoted by horticulture, such as fast growth, are also likely to  
693 promote invasiveness. On the other hand, there is great potential to breed non-invasive  
694 ideotypes of ornamental plants, but the efforts of the horticultural industry in this regard are  
695 still very limited.

696 (6) A major future challenge is that social and technological changes, such as internet trade  
697 and molecular genetic breeding techniques, will lead to fundamentally novel patterns of plant  
698 introductions. In addition, environmental change, and climate change in particular, is likely to  
699 change the invasion opportunities of the ornamental species that have already been  
700 introduced.

701 (7) There is a need for analysis of current and future invasion risks for ornamental species in  
702 many regions of the world (Mayer *et al.*, 2017). Ecological and socio-economic impact-  
703 categorisation frameworks such as EICAT (Blackburn *et al.*, 2014) and SEICAT (Bacher *et*  
704 *al.*, 2017), as well as global lists of currently widely naturalised species (Pyšek *et al.*, 2017)  
705 will be very useful in this regard.

706 (8) There are still many open questions on the role of horticulture in plant invasions (Table  
707 1). Therefore, more intensive research efforts on the role of horticulture are urgently needed  
708 to develop science-based regulatory frameworks that help to prevent further plant invasions.  
709

## 710 **VIII. ACKNOWLEDGEMENTS**

711 We thank the COST Action TD1209 ‘Alien Challenge’ for funding the workshop that was at  
712 the basis of this paper. M.v.K., F.E., M.C., S.D., G.K. thank the ERA-Net BiodivERsA, with  
713 the national funders ANR (French National Research Agency), DFG (German Research  
714 Foundation; to M.v.K. and W.D.) and FWF (Austrian Science Fund; to S.D. and F.E.), part of  
715 the 2012-2013 BiodivERsA call for research proposals. M.v.K., W.D. (both KL 1866/9-1)  
716 and H.S. (SE 1891/2-1) acknowledge funding by the German Research Foundation. J.P., A.N.  
717 and P.P. are supported by grants (DG16P02M041, MSMT CR), Centre of Excellence  
718 PLADIAS, no. 14-15414S (Czech Science Foundation) and long-term research development  
719 project RVO 67985939 (The Czech Academy of Sciences). P.P. acknowledges funding by  
720 Praemium Academiae award from The Czech Academy of Sciences.

721

722 **IX. REFERENCES**

723 ALEXANDER, J. M., KUEFFER, C., DAEHLER, C. C., EDWARDS, P. J., PAUCHARD, A., SEIPEL, T.

724 & THE MIREN CONSORTIUM. (2011). Assembly of nonnative floras along elevational

725 gradients explained by directional ecological filtering. *Proceedings of the National*

726 *Academy of Sciences U.S.A.* **108**, 656–661.

727 ALEXANDER, J. M., LEMBRECHTS, J. J., CAVIERES, A. L., DAEHLER, C. C., HAIDER, S.,

728 KUEFFER, C., LIU, G., MCDOUGALL, K., MILBAU, A., PAUCHARD, A., REW, L. J. & SEIPEL

729 T. (2017). Plant invasions into mountains and alpine ecosystems: current status and future

730 challenges. *Alpine Botany* **126**, 89–103.

731 ANDERSON, N. O., GOMEZ, N. & GALATOWITSCH, S. M. (2006). A non-invasive crop ideotype

732 to reduce invasive potential. *Euphytica* **148**, 185–202.

733 AULT, J., & THOMAS, C. (2014). Plant propagation for the breeding program at Chicago

734 botanical garden©. *Acta Horticulturae* **1055**, 265–272.

735 BACHER, S., BLACKBURN, T. M., ESSL, F., GENOVESI, P., HEIKKILÄ, J., JESCHKE, J. M., JONES,

736 G., KELLER, R., KENIS, M., KUEFFER, C., MARTINOU, A. F., NENTWIG, W., PERGL, J.,

737 PYŠEK, P., RABITSCH, W., *ET AL.* (2017). Socio-economic impact classification of alien taxa

738 (SEICAT). *Methods in Ecology and Evolution*, DOI: 10.1111/2041-210X.12844.

739 BENNETT, A. W. (1870). Acclimatization of foreign trees and plants. *The American Naturalist*

740 **5**, 528–534.

741 BLACKBURN, T. M., ESSL, F., EVANS, T, HULME, P. E., JESCHKE, J. M., KÜHN, I., KUMSCHICK,

742 S., MARKOVÁ, Z., MRUGALA, A, NENTWIG, W., PERGL, J., PYŠEK, P., RABITSCH, W.,

743 RICCIARDI, A., RICHARDSON, D. M., *ET AL.* (2014). A unified classification of alien species

744 based on the magnitude of their environmental impacts. *PLoS Biology* **12**, e1001850.

745 BOLVIN, N. L., ZEDER, M. A., FULLER, D. Q., CROWTHER, A., LARSON, G., ERLANDSON, J. M.,  
746 DENHAM, T. & PETRAGLIA, M. D. (2016). Ecological consequences of human niche  
747 construction: Examining long-term anthropogenic shaping of global species distributions.  
748 *Proceedings of the National Academy of Sciences USA* **113**, 6388–6396.

749 BOTHAM, M. S., ROTHERY, P., HULME, P. E., HILL, M. O., PRESTON, C. D. & ROY, D. B.  
750 (2009). Do urban areas act as foci for the spread of alien plant species? An assessment of  
751 temporal trends in the UK. *Diversity and Distributions* **15**, 338–345.

752 BRADBURY, M. (ED.) (1995). *A history of the garden in New Zealand*. Viking, Auckland.

753 BRADLEY, B. A., BLUMENTHAL, D. M., EARLY, R., GROSHOLZ, E. D., LAWLER, J. J., MILLER,  
754 L. P., SORTE, C. J. B., D'ANTONIO, C. M., DIEZ, J. M., DUKES, J. S., IBANEZ, I. & OLDEN, J.  
755 D. (2011). Global change, global trade, and the next wave of plant invasions. *Frontiers in*  
756 *Ecology and the Environment* **10**, 20–28.

757 BRADLEY, B. A., BLUMENTHAL, D. M., WILCOVE, D. S. & ZISKA, L. H. (2010). Predicting  
758 plant invasions in an era of global change. *Trends in Ecology and Evolution* **25**, 310–318.

759 BRAND, M. H., LEHRER, J. M. & LUBELL, J. D. (2012). Fecundity of Japanese barberry  
760 (*Berberis thunbergii*) cultivars and their ability to invade a deciduous woodland. *Invasive*  
761 *Plant Science and Management* **5**, 464–476.

762 BROWN, A. (1878). Plants introduced with ballast on made land. *Bulletin of the Torrey*  
763 *Botanical Club* **6**, 255–258.

764 BRUNDU, G. & RICHARDSON, D. M. (2016). Planted forests and invasive alien trees in Europe:  
765 a code for managing existing and future plantings to mitigate the risk of negative impacts  
766 from invasions. *NeoBiota* **30**, 5–47.

767 BRUMMIT, R. K. (2001). *World geographical scheme for recording plant distributions*. 2<sup>nd</sup>  
768 *edition*. Hunt Institute for Botanical Documentation, Pittsburgh.



769 BURT, J. W., MUIR, A. A., PIOVIA-SCOTT, J., VEBLEN, K. E., CHANG, A. L., GROSSMAN, J. D.  
770 & WEISKEL, H. W. (2007). Preventing horticultural introductions of invasive plants:  
771 potential efficacy of voluntary initiatives. *Biological Invasions* **9**, 909–923.

772 CAMPAGNARO, T., BRUNDU, G. & SITZIA, T. (2017). Five major invasive alien tree species in  
773 European Union forest habitat types of the Alpine and Continental biogeographical  
774 regions. *Journal for Nature Conservation*, doi: 10.1016/j.jnc.2017.07.007.

775 CAMPBELL-CULVER, M. (2001). *The origin of plants: the people and plants that have shaped*  
776 *Britain's garden history since the year 1000*. Headline Book Publishing, London.

777 CHANDLER, S. F. & BRUGLIERA, F. (2011). Genetic modification in floriculture.  
778 *Biotechnology letters* **33**, 207–214.

779 CIESIN, IFPRI, WORLD BANK & CIAT (2011). *Global Rural-Urban Mapping Project,*  
780 *version 1 (GRUMPv1): land and geographic unit area grids*. Center for International  
781 Earth Science Information Network (CIESIN) of Columbia University, International Food  
782 Policy Research Institute (IFPRI), the World Bank, Centro Internacional de Agricultura  
783 Tropical (CIAT).

784 CLEMENT, E. J. & FOSTER, M. C. (1994). *Alien plants of the British Isles*. Botanical Society of  
785 the British Isles, London.

786 CONVENTION ON BIOLOGICAL DIVERSITY (2011). *Nagoya protocol on access to genetic*  
787 *resources and the fair and equitable sharing of benefits arising from their utilization to the*  
788 *Convention on Biological Diversity*. Convention on Biological Diversity. United Nations.

789 COOK, G. D. & DIAS, L. (2006). It was no accident: deliberate plant introductions by  
790 Australian government agencies during the 20<sup>th</sup> century. *Australian Journal of Botany* **54**,  
791 601–625.

792 COX, P. A. & BANACK, S. A. (1991). *Islands, plants, and Polynesians: an introduction to*  
793 *Polynesian ethnobotany*. Dioscorides Press Press, Portland.

794 CRESPO, C. J., KETEYIAN, S. J., HEATH, G. W. & SEMPOS, C. T. (1996). Leisure-time physical  
795 activity among US adults: results from the third National Health and Nutrition  
796 Examination Survey. *Archives of Internal Medicine* **156**, 93–98.

797 DAEHLER, C. C. (2008). Invasive plant problems in the Hawaiian Islands and beyond: insights  
798 from history and psychology. In *Plant invasions: human perception, ecological impacts*  
799 *and management* (eds B. TOKARSKA-GUZIK, J. H. BROCK, G. BRUNDU, L. CHILD, C. C.  
800 DAEHLER & P. PYŠEK), pp. 3–20. Backhuys Publishers, Leiden, The Netherlands.

801 DAINESE, M., AIKIO, S., HULME, P. E., BERTOLLI, A., PROSSER, F. & MARINI, L. (2017).  
802 Human disturbance and upward expansion of plants in a warming climate. *Nature Climate*  
803 *Change* DOI:10.1038/nclimate3337.

804 DAWSON, W., FISCHER, M & VAN KLEUNEN, M. (2011). Maximum relative growth rate of  
805 common UK plant species is positively associated with their global invasiveness. *Global*  
806 *Ecology and Biogeography* **20**, 299–306.

807 DEBENER, T., VON MALEK, B., MATTIESCH, L. & KAUFMANN, H. (2001). Genetic and  
808 molecular analysis of important characters in roses. *Acta Horticulturae* **547**, 45–49.

809 DEHNEN-SCHMUTZ, K. (2004). Alien species reflecting history: medieval castles in Germany.  
810 *Diversity and Distributions* **10**, 147–151.

811 DEHNEN-SCHMUTZ, K. (2011). Determining non-invasiveness in ornamental plants to build  
812 green lists. *Journal of Applied Ecology* **48**, 1374–1380.

813 DEHNEN-SCHMUTZ, K., TOUZA, J., PERRINGS, C. & WILLIAMSON, M. (2007). A century of the  
814 ornamental plant trade and its impact on invasion success. *Diversity and Distributions* **13**,  
815 527–534.

816 DI CASTRI, F. (1989). History of biological invasions with special emphasis on the Old World.  
817 In *Biological invasions: a global perspective* (eds J. A. Drake, H. A. Mooney, F. di Castri,

818 R. H. Groves, F. J. Kruger, M. Rejmánek and M. Williamson.), pp. 1–30. John Wiley and  
819 Sons, Chichester.

820 DIAGRE-VANDERPELEN, D. (2011). *The Botanical garden of Brussels (1826–1912): reflection*  
821 *of a changing nation*. Botanical Garden Meise, Sc. Coll. U.L.B., Belgium.

822 DIEZ, J. M., D'ANTONIO, C. M., DUKES, J. S., GROSHOLZ, E. D., OLDEN, J. D., SORTE, C. J. B.,  
823 BLUMENTHAL, D. M., BRADLEY, B. A., EARLY, R., IBANEZ, I., JONES, S. J., LAWLER, J. J. &  
824 MILLER, L. P. (2012). Will extreme climatic events facilitate biological invasions.  
825 *Frontiers in Ecology and the Environment* **10**, 249–257.

826 DORSETT, P. H. (1917). The plant-introduction gardens of the Department of Agriculture.  
827 *Yearbook of the United States Department of Agriculture* **1916**, 135–144.

828 DREW, J., ANDERSON, N. & ANDOW, D. (2010). Conundrums of a complex vector for invasive  
829 species control: a detailed examination of the horticultural industry. *Biological Invasions*  
830 **12**, 2837–2851.

831 DULLINGER, I., WESSELY, J., BOSSDORF, O., DAWSON, W., ESSL, F., GATTRINGER, A.,  
832 KLONNER, G., KREFT, H., KUTTNER, M., MOSER, D., PERGL, J., PYŠEK, P., THULLER, W.,  
833 VAN KLEUNEN, M., WEIGELT, P., *ET AL.* (2017). Climate change will increase the  
834 naturalization risk from garden plants in Europe. *Global Ecology and Biogeography* **26**,  
835 43–53.

836 DUNLAP, T. R. (1997). Remaking the land: the acclimatization movement and Anglo ideas of  
837 nature. *Journal of World History* **8**, 303–319.

838 DUNNETT, N. & KINGSBURY, N. (2008). *Planting green roofs and living walls*. Timber Press,  
839 Portland, USA.

840 EARLY, R., BRADLEY, B. A., DUKES, J. S., LAWLER, J. J., OLDEN, J. D., BLUMENTHAL, D. M.,  
841 GONZALEZ, P., GROSHOLZ, E. D., IBANEZ, I., MILLER, L. P., SORTE, C. J. B. & TATEM, A. J.

842 (2016). Global threats from invasive alien species in the twenty-first century and national  
843 response capacities. *Nature Communications* **7**, 12485.

844 ESCHTRUTH, A. K. & BATTLES, J. J. (2009). Assessing the relative importance of disturbance,  
845 herbivory, diversity, and propagule pressure in exotic plant invasions. *Ecological*  
846 *Monographs* **79**, 265–280.

847 ESSL, F. (2007). From ornamental to detrimental? The incipient invasion of Central Europe  
848 by *Paulownia tomentosa*. *Preslia* **79**, 377–389.

849 FAIRCHILD, D. G. (1898). *Systematic plant introduction: its purposes and methods*.  
850 Government Printing Office, Washington.

851 FARRINGTON, E. I. (1931). *Ernest H. Wilson plant hunter. With a list of his most important*  
852 *introductions and where to get them*. The Stratford Company, Boston, Massachusetts.

853 FAULKNER, K. T., ROBERTSON, M. P., ROUGET, M. & WILSON, J. R. U. (2016). Understanding  
854 and managing the introduction pathways of alien taxa: South Africa as a case study.  
855 *Biological Invasions* **18**, 73–87.

856 FIACCONI, M. & HUNT, C. O. (2015). Pollen taphonomy at Shanidar Cave (Kurdish Iraq): an  
857 initial evaluation. *Review of Palaeobotany and Palynology* **223**, 87–93.

858 FREYRE, R., DENG, Z., KNOX, G. W., MONTALVO, S. & ZAYAS, V. (2016). Fruitless *Ruellia*  
859 *simplex* R12-2-1 (Mayan Compact Purple). *HortScience* **51**, 1057–1061.

860 FRIDLEY, J. D. & SAX, D. F. (2014). The imbalance of nature: revisiting a Darwinian  
861 framework for invasion biology. *Global Ecology and Biogeography* **23**, 1157–1166.

862 GALLAGHER, R. V., RANDALL, R. P. & LEISHMAN, M. R. (2015). Trait differences between  
863 naturalized and invasive plant species independent of residence time and phylogeny.  
864 *Conservation Biology* **29**, 360–369.

865 GILTRAP, N., EYRE, D. & REED, P. (2009). Internet sales of plants for planting – an increasing  
866 trend and threat? *EPPO Bulletin* **39**, 168–170.

867 GIORGIS, M. A. & TECCO, P. A. (2014). Invasive alien trees and shrubs in Córdoba province  
868 (Argentina): a contribution to the systematization of global bases. *Boletín de la Sociedad*  
869 *Argentina de Botánica* **49**, 581–603.

870 GOEZE, E. (1916). Liste der seit dem 16. Jahrhundert eingeführten Bäume und Sträucher.  
871 *Mitteilungen der Deutschen Dendrologischen Gesellschaft* **25**, 129–201 (in German).

872 GROTKOPP, E., ERSKINE-OGDEN, J. & REJMÁNEK, M. (2010). Assessing potential invasiveness  
873 of woody horticultural plant species using seedling growth rate traits. *Journal of Applied*  
874 *Ecology* **47**, 1320–1328.

875 GROVE, R. (1995). *Green imperialism: colonial expansion, tropical island Edens and the*  
876 *origins of environmentalism, 1600-1860*. Cambridge University Press, New York.

877 GROVES, R. H. (1998). Recent incursions of weeds to Australia 1971–1995. *CRC for Weed*  
878 *Management Systems technical series*, **3**, 1–74.

879 HANSPACH, J., KÜHN, I., PYŠEK, P., BOOS, E. & KLOTZ, S. (2008). Correlates of naturalization  
880 and occupancy of introduced ornamentals in Germany. *Perspectives in Plant Ecology*  
881 *Evolution and Systematics* **10**, 241–250.

882 HARKNESS, P. (2003). *The rose: an illustrated history*. Firefly Books.

883 HARRIS, S. A. (2002). Introduction of Oxford Ragwort, *Senecio squalidus* L. (Asteraceae), to  
884 the United Kingdom". *Watsonia* **24**, 31–43.

885 HAEUSER, E., DAWSON, W. & VAN KLEUNEN, M. (2017). The effects of climate warming and  
886 disturbance on the colonization potential of ornamental alien plant species. *Journal of*  
887 *Ecology* **105**, 1698–1708.

888 HAVENS, K., VITT, P., MAUNDER, M., GUERRANT JR., E. O. & DIXON, K. (2006). *Ex situ* plant  
889 conservation and beyond. *BioScience* **56**, 525–583.

890 HILL, A. W. (1915). The history and functions of botanic gardens. *Annals of the Missouri*  
891 *Botanical Garden* **2**, 185-240.

892 HINSLEY, A., VERISSIMO, D. & ROBERTS, D. L. (2015). Heterogeneity in consumer  
893 preferences for orchids in international trade and the potential for the use of market  
894 research methods to study demand for wildlife. *Biological Conservation* **190**, 80–86.

895 HITCHMOUGH, J & WOULDSTRA, J. (1999). The ecology of exotic herbaceous perennials grown  
896 in managed, native grassy vegetation in urban landscapes. *Landscape and Urban Planning*  
897 **45**, 107–121.

898 HODGE, W. H. & ERLANSON, C. O. (1956). Federal plant introduction – a review. *Economic*  
899 *Botany* **10**, 299–334.

900 HOTTENTRÄGER, G. (1992). New flowers-new gardens. *The Journal of Garden History* **12**,  
901 207–227.

902 HULME, P. E. (2009). Relative roles of life-form, land use and climate in recent dynamics of  
903 alien plant distributions in the British Isles. *Weed Research* **49**, 19–28.

904 HULME, P. E. (2011). Addressing the threat to biodiversity from botanical gardens. *Trends in*  
905 *Ecology and Evolution* **26**, 168–174.

906 HULME, P. E. (2015). Resolving whether botanical gardens are on the road to conservation or  
907 a pathway for plant invasions. *Conservation Biology* **29**, 816–824.

908 HULME, P. E., BACHER, S., KENIS, M., KLOTZ, S., KÜHN, I., MINCHIN, D., NENTWIG, W.,  
909 OLENIN, S., PANOV, V., PERGL, J., PYŠEK, P., ROQUES, A., SOL, D., SOLARZ, W. & VILÀ, M.  
910 (2008). Grasping at the routes of biological invasions: a framework for integrating  
911 pathways into policy. *Journal of Applied Ecology* **45**, 403–414.

912 HULME, P. E., BRUNDU, G., CARBONI, M., DEHNEN-SCHMUTZ, K., DULLINGER, S., EARLY, R.,  
913 ESSL, F., GONZÁLEZ-MORENO, P., GROOM, Q. J., KUEFFER, C., KÜHN, I., MAUREL, N.,  
914 NOVOA, A., PERGL, J., PYŠEK, P., *ET AL.* (2018). Integrating invasive species policies across  
915 ornamental horticulture supply-chains to prevent plant invasions. *Journal of Applied*  
916 *Ecology* **55**, 92–98.

- 917 HUMAIR, F., HUMAIR, L., KUHN, F. & KUEFFER, C. (2015). E-commerce trade in invasive  
918 plants. *Conservation Biology* **29**, 1658–1665.
- 919 HUMAIR, F., KUEFFER, C. & SIEGRIST, M. (2014a). Are non-native plants perceived to be  
920 more risky? Factors influencing horticulturists' risk perceptions of ornamental plant  
921 species. *PLoS ONE* **9**, e102121.
- 922 HUMAIR, F., SIEGRIST, M. & KUEFFER, C. (2014b). Working with the horticultural industry to  
923 limit invasion risks: the Swiss experience. *EPPO Bulletin* **44**, 232–238.
- 924 JACOMET, S. & KREUZ, A. (1999). Archäobotanik. Ulmer, Stuttgart, Germany.
- 925 JANICK, J. (2007). Plant exploration: from Queen Hatshepsut to Sir Joseph Banks.  
926 *Horticultural Science* **42**, 191–196.
- 927 JEKYLL, G. (1908). *Colour in the flower garden*. Country Life, London, UK.
- 928 KELLER, H. (1994). *Kleine Geschichte der Gartenkunst*. Blackwell Wissenschafts-Verlag,  
929 Berlin, Germany (in German).
- 930 KHOSHBAKHT, K. & HAMMER, K. (2008). How many plant species are cultivated. *Genetic*  
931 *Resources and Crop Evolution* **55**, 925–928.
- 932 KINTGEN, M., KRISHNAN, S. & HAYWARD, P. (2013). Plant Select® a brief overview, history  
933 and future of a plant introduction program. *Acta Horticulturae* **1000**, 585–589.
- 934 KIRKLAND, A. & BERG, P. (1997). *A century of state-honed enterprise: 100 years of state*  
935 *plantation forestry in New Zealand*. Profile Books, Masterton.
- 936 KNAPP, S., DINSMORE, L., FISSORE, C., HOBBIE, S. E., JAKOBSDOTTIR, I., KATTGE, J., KING, J.  
937 Y., KLOTZ, S., MCFADDEN, J. P. & CAVENDER-BARES, J. (2012). Phylogenetic and  
938 functional characteristics of household yard floras and their changes along an urbanization  
939 gradient. *Ecology* **93**, 83–98.
- 940 KNAPP, S. & KÜHN, I. (2012). Origin matters: widely distributed native and non-native  
941 species benefit from different functional traits. *Ecology Letters* **15**, 696–703.

942 KNIGHT, T. M., HAVENS, K. & VITT, P. (2011). Will the use of less fecund cultivars reduce the  
943 invasiveness of perennial plants? *Bioscience* **61**, 816–822.

944 KRUCKEBERG, A. R. (2001). *Gardening with native plants, 2nd edition*. University of  
945 Washington Press, Seattle, USA.

946 KUEFFER, C. (2010). Transdisciplinary research is needed to predict plant invasions in an era  
947 of global change. *Trends in Ecology and Evolution* **20**, 619–620.

948 KUEFFER, C., MCDUGALL, K., ALEXANDER, J., DAEHLER, C., EDWARDS, P., HAIDER, S.,  
949 MILBAU, A., PARKS, C., PAUCHARD, A., RESHI, Z. A., REW, L. J., SCHRODER, M. & SEIPEL,  
950 T. (2013). Plant invasions into mountain protected areas: assessment, prevention and  
951 control at multiple spatial scales. In *Plant invasions in protected areas: patterns, problems*  
952 *and challenges* (eds L. C. FOXCROFT, P. PYŠEK, D. M. RICHARDSON & P. GENOVESI.), pp.  
953 89–113. Springer, Dordrecht, The Netherlands.

954 KÜHN, I. & KLOTZ, S. (2002). Floristischer Status und gebietsfremde Arten. *Schriftenreihe*  
955 *Vegetationskunde* **38**, 47–56 (in German).

956 LAMBDon, P. W., PYŠEK, P., BASNOU, C., HEJDA, M., ARIANOUTSOU, M., ESSL, F., JAROŠÍK,  
957 V., PERGL, J., WINTER, M., ANASTASIU, P., ANDRIOPOULOS, P., BAZOS, I., BRUNDU, G.,  
958 CELESTI-GRAPow, L., CHASSOT, P., *ET AL.* (2008). Alien flora of Europe: species diversity,  
959 temporal trends, geographical patterns and research needs. *Preslia* **80**, 101–149.

960 LEROI-GOURHAN, A. (1975). The flowers found with Shanidar IV, a Neanderthal burial in  
961 Iraq. *Science* **190**, 562–564.

962 LIU, Y., ODUOR, A. M. O., ZHANG, Z., MANEA, A., TOOTH, I. M., LEISHMAN, M. R., XU, X. &  
963 VAN KLEUNEN, M. (2017). Do invasive alien plants benefit more from global  
964 environmental change than native plants? *Global Change Biology* **23**, 3363–3370.



965 LLORET, F., MÉDAIL, F., BRUNDU, G., CAMARDA, I., MORAGUES, E., RITA, J., LAMBTON, P. &  
966 HULME, P. E. (2005). Species attributes and invasion success by alien plants in  
967 Mediterranean islands. *Journal of Ecology* **93**, 512–520.

968 MACK, R. N. (2000). Cultivation fosters plant naturalization by reducing environmental  
969 stochasticity. *Biological Invasions* **2**, 111–122.

970 MACK, R. N. (2005). Predicting the identity of plant invaders: future contributions from  
971 horticulture. *HortScience* **40**, 1168–1174.

972 MACK, R. N. & ERNEBERG, M. (2002). The United States naturalized flora: largely the product  
973 of deliberate introductions. *Annals of the Missouri Botanical Garden* **89**, 176–189.

974 MACK, R. N. & LONSDALE, W. M. (2001). Humans as global plant dispersers: getting more  
975 than we bargained for: current introductions of species for aesthetic purposes present the  
976 largest single challenge for predicting which plant immigrants will become future pests.  
977 *BioScience* **51**, 95–102.

978 MAKI, K. & GALATOWITSCH, S. (2004). Movement of invasive aquatic plants into Minnesota  
979 (USA) through horticultural trade. *Biological Conservation* **118**, 389–396.

980 MANRIQUE, L. R. (2010). Intercambio y difusión de plantas agrícolas entre el nuevo y el viejo  
981 mundo. *Revista de la Ofil* **21**, 89–91 (in Spanish).

982 MAUREL, N., HANSPACH, J., KÜHN, I., PYŠEK, P. & VAN KLEUNEN, M. (2016). Introduction  
983 bias affects relationships between the characteristics of ornamental alien plants and their  
984 naturalization success. *Global Ecology and Biogeography* **25**, 1500–1509.

985 MAYER, K., HAEUSER, E., DAWSON, W., ESSL, F., KREFT, H., PERGL, J., PYŠEK, P., WEIGELT,  
986 P., WINTER, M., LENZNER, B. & VAN KLEUNEN, M. (2017). Current and future local  
987 naturalization potential of ornamental species planted in public green spaces and private  
988 gardens. *Biological invasions* **19**, 3613–3627.

989 MÜLLER, N. & SUKOPP, H. (2016). Influence of different landscape design styles on plant  
990 invasions in Central Europe. *Landscape and Ecological Engineering* **12**, 151–169.

991 MUSGRAVE, T., GARDNER, C. & MUSGRAVE, W. (1999). *The plant hunters two hundred years*  
992 *of adventure and discovery*. Ward Lock, London, UK.

993 NIINIMENTS, Ü. & PEÑUELAS, J. (2008). Gardening and urban landscaping: significant players  
994 in global change. *Trends in Plant Science* **13**, 60–65.

995 NOVOA, A., KAPLAN, H., KUMSCHICK, S., WILSON, J. R. U. & RICHARDSON, D. M. (2015a).  
996 Soft touch or heavy hand? Legislative approaches for preventing invasions: insights from  
997 cacti in South Africa. *Invasive Plant Science and Management* **8**, 307–316.

998 NOVOA, A., LE ROUX, J. J., ROBERTSON, M. P., WILSON, J. R. U. & RICHARDSON, D. M.  
999 (2015b). Introduced and invasive cactus species: a global review. *AoB Plants* **7**, 14.

1000 NOVOA, A., ROUX, J. J., RICHARDSON, D. M. & WILSON, J. R. U. (2017). Level of  
1001 environmental threat posed by horticultural trade in Cactaceae. *Conservation Biology*  
1002 DOI:10.1111/cobi.12892.

1003 OSBORNE, M. A. (2001). Nature and empire: science and the colonial enterprise. *Osiris* **15**,  
1004 135–151.

1005 PAUCHARD, A., KUEFFER, C., DIETZ, H., DAEHLER, C. C., ALEXANDER, J., EDWARDS, P. J.,  
1006 ARÉVALO, J. R., CAVIERES, L. A., GUISAN, A., HAIDER, S., JAKOBS, G., MCDOUGALL, K.,  
1007 MILLAR, C. I., NAYLOR, B. J., PARKS, C. G., *ET AL.* (2009). Ain't no mountain high enough:  
1008 plant invasions reaching new elevations. *Frontiers in Ecology and the Environment* **7**,  
1009 479–486.

1010 PAUCHARD, A., MILBAU, A., ALBIHN, A., ALEXANDER, J., NUN, M. A., DAEHLER, C.,  
1011 ENGLUND, G., ESSL, F., EVENGARD, B., GREENWOOD, G. B., HAIDER, S., LENOIR, J.,  
1012 MCDOUGALL, K., MUTHS, E., NUNEZ, M. A., *ET AL.* (2016). Non-native and native

1013 organisms moving into high elevation and high latitude ecosystems in an era of climate  
1014 change: new challenges for ecology and conservation. *Biological Invasions* **18**, 345–353.

1015 PERGL, J., PYŠEK, P., BACHER, S., ESSL, F., GENOVESI, P., HARROWER, C. A., HULME, P. E.,  
1016 JESCHKE, J. M., KENIS, M., KÜHN, I., PERGLOVÁ, I., RABITSCH, W., ROQUES, A., ROY, D.  
1017 B., ROY, H. E., *ET AL.* (2017). Troubling travellers: are ecologically harmful alien species  
1018 associated with particular introduction pathways? *NeoBiota* **32**, 1–20.

1019 PERGL, J., SÁDLO, J., PETŘÍK, P., DANIHELKA, J., CHRTEK JR., J., HEJDA, M., MORAVCOVÁ, L.,  
1020 PERGLOVÁ, I., ŠTAJEROVÁ, K. & PYŠEK, P. (2016). Dark side of the fence: ornamental  
1021 plants as a source for spontaneous flora of the Czech Republic. *Preslia* **88**, 163–184.

1022 PETITPIERRE, B., MCDUGALL, K., SEIPEL, T., BROENNIMANN, O., GUISAN, A. & KUEFFER, C.  
1023 (2016). Will climate change increase the risk of plant invasions into mountains?  
1024 *Ecological Applications* **26**, 530–544.

1025 PIMENTEL, D., ZUNIGA, R. & MORRISON, D. (2005). Update on the environmental and  
1026 economic costs associated with alien-invasive species in the United States. *Ecological*  
1027 *Economics* **52**, 273–288.

1028 POOLER, M. R. (2001). Plant breeding at the US National Arboretum: Selection, evaluation,  
1029 and release of new cultivars. *Horttechnology* **11**, 365–367.

1030 PROCHEŞ, Ş., WILSON, J. R. U., VELDTMAN, R., KALWIJ, J. M., RICHARDSON, D. M. & CHOWN,  
1031 S. L. (2005). Landscape corridors: possible dangers? *Science* **310**, 781–782.

1032 PYŠEK, P., DANIHELKA, J., SÁDLO, J., CHRTEK, J. JR, CHYTRÝ, M., JAROŠÍK, V., KAPLAN, Z.,  
1033 KRAHULEC, F., MORAVCOVÁ, L., PERGL, J., ŠTAJEROVÁ, K. & TICHÝ, L. (2012*a*).  
1034 Catalogue of alien plants of the Czech Republic (2nd edn): checklist update, taxonomic  
1035 diversity and invasion patterns. *Preslia* **84**, 155–255.

1036 PYŠEK, P., JAROŠÍK, V., HULME, P. E., PERGL, J., HEJDA, M., SCHAFFNER, U. & VILÀ, M.  
1037 (2012*b*). A global assessment of invasive plant impacts on resident species, communities

1038 and ecosystems: the interaction of impact measures, invading species' traits and  
1039 environment. *Global Change Biology* **18**, 1725–1737.

1040 PYŠEK, P., JAROŠÍK, V. & PERGL, J. (2011). Alien plants introduced by different pathways  
1041 differ in invasion success: unintentional introductions as greater threat to natural areas?  
1042 *PLoS ONE* **6**, e24890.

1043 PYŠEK, P., JAROŠÍK, V., PERGL, J. & WILD, J. (2011). Colonization of high altitudes by alien  
1044 plants over the last two centuries. *Proceedings of the National Academy of Sciences USA*  
1045 **108**, 439–440.

1046 PYŠEK, P., PERGL, J., ESSL, F., LENZNER, B., DAWSON, W., KREFT, H., WEIGELT, P., WINTER,  
1047 M., KARTESZ, J., NISHINO, M., ANTONOVA, L. A., BARCELONA, J. F., CABEZAS, F. J.,  
1048 CÁRDENAS, D., CÁRDENAS-TORO, J., *ET AL.* (2017). Naturalized and invasive alien flora of  
1049 the world: species diversity, taxonomic and phylogenetic patterns, geographic distribution  
1050 and global hotspots of plant invasion. *Preslia* **89**, 203–274.

1051 REICHARD, S. H. (2011). Horticulture. In *Encyclopedia of Biological Invasions* (eds D.  
1052 SIMBERLOFF & M. REJMÁNEK.), pp. 336–342. University of California Press, USA.

1053 REICHARD, S. H. & WHITE, P. (2001). Horticulture as a pathway of invasive plant  
1054 introductions in the United States. *Bioscience* **51**, 103–113.

1055 REYNOLDS, R. (2014). *On guerrilla gardening: a handbook for gardening without*  
1056 *boundaries*. Bloomsbury Publishing, London, UK.

1057 RICHARDSON, D. M. & PYŠEK, P. (2012). Naturalization of introduced plants: ecological  
1058 drivers of biogeographical patterns. *New Phytologist* **196**, 383–396.

1059 RICHARDSON, D. M., PYŠEK, P., REJMÁNEK, M., BARBOUR, M. G., PANETTA, F. D. & WEST, C.  
1060 J. (2000). Naturalization and invasion of alien plants: concepts and definitions. *Diversity*  
1061 *and Distributions* **6**, 93–107.

- 1062 RICHARDSON, D. M. & REJMÁNEK, M. (2011). Trees and shrubs as invasive alien species – a  
1063 global review. *Diversity and Distributions* **17**, 788–809.
- 1064 ROBINSON, W. (1870). *The wild garden*. John Murray, London, UK.
- 1065 ROJAS-SANDOVAL, J. & ACEVEDO-RODRÍGUEZ, P. (2014). Naturalization and invasion of alien  
1066 plants in Puerto Rico and the Virgin Islands. *Biological Invasions* **17**, 149–163.
- 1067 ROULLIER, C., BENOIT, L., MCKEY, D. B. & LEBOT, V. (2013) Historical collections reveal  
1068 patterns of diffusion of sweet potato in Oceania obscured by modern plant movements and  
1069 recombination. *Proceedings of the National Academy of Sciences U.S.A.* **110**, 2205–2210.
- 1070 SÁNCHEZ, A. L. (1997). Los jardines botánicos neotropicales y el intercambio de plantas:  
1071 pasado, presente y futuro. *Monografías del Real Jardín Botánico de Córdoba* **5**, 75–84  
1072 (inSpanish).
- 1073 SAUL, W.-C., ROY, H. E., BOOY, O., CARNEVALI, L., CHEN, H.-J., GENOVESI, P., HARROWER,  
1074 C. A., HULME, P. E., PAGAD, S., PERGL, J. & JESCHKE, J. M. (2017). Assessing patterns in  
1075 introduction pathways of alien species by linking major invasion data bases. *Journal of*  
1076 *Applied Ecology* **54**, 657–669.
- 1077 SEATON, K., BETTIN, A. & GRÜNEBERG, H. (2014). New ornamental plants for horticulture. In  
1078 *Horticulture: plants for people and places* (eds G. R. DIXON & D. E. ALDOUS.), pp. 435–  
1079 463. Springer, Furth, Germany.
- 1080 SEEBENS, H., BLACKBURN, T. M., DYER, E. E., GENOVESI, P., HULME, P. E., JESCHKE, J. M.,  
1081 PAGAD, S., PYŠEK, P., WINTER, M., ARIANOUTSOU, M., BACHER, S., BLASIUS, B., BRUNDU,  
1082 G., CAPINHA, C., CELESTI-GRAPOW, L., *ET AL.* (2017). No saturation in the accumulation of  
1083 alien species worldwide. *Nature Communications* **8**, 14435.
- 1084 SEEBENS, H., ESSL, F., DAWSON, W., FUENTES, N., MOSER, D., PERGL, J., PYŠEK, P., VAN  
1085 KLEUNEN, M., WEBER, E., WINTER, M. & BLASIUS, B. (2015). Global trade will accelerate

1086 plant invasions in emerging economies under climate change. *Global Change Biology* **21**,  
1087 4128–4140.

1088 SHAW, A., MILLER, K. K. & WESCOTT, G. (2017). Australian native gardens: Is there scope for  
1089 a community shift? *Landscape and Urban Planning* **157**, 322–330.

1090 SHEPPARD, C., BURNS, B. & STANLEY, M. (2014). Predicting plant invasions under climate  
1091 change: are species distribution models validated by field trials? *Global Change Biology*  
1092 **20**, 2800–2814.

1093 SMARDON, R. C. (1988). Perception and aesthetics of the urban environment: review of the  
1094 role of vegetation. *Landscape and Urban Planning* **15**, 85–106.

1095 SOGA, M., GASTON, K. J. & YAMAURA, Y. (2017). Gardening is beneficial for health: a meta-  
1096 analysis. *Preventive Medicine Reports* **5**, 92–99.

1097 STÖCKLIN, J., SCHAUB, P. & OJALA, O. (2003). Häufigkeit und Ausbreitungsdynamik von  
1098 Neophyten in der Region Basel: Anlass zur Besorgnis oder Bereicherung? *Bauhinia* **17**,  
1099 11–23 (in German).

1100 STONER, A. & HUMMER, K. (2007). 19th and 20th century plant hunters. *Horticultural*  
1101 *Science* **42**, 197–199.

1102 STRABO W. (c. 840). *De cultura hortorum Über den Gartenbau*. Reclam, Stuttgart, German  
1103 (in Latin and German)

1104 SUKOPP, H. (2006). Botanische Gärten und die Berliner Flora. *Willdenowa* **36**, 115–125 (in  
1105 German).

1106 SVENNING, J.C. & SANDEL, B. (2013). Disequilibrium vegetation dynamics under future  
1107 climate change. *American Journal of Botany* **100**, 1266–1286.

1108 VAN DER VEKEN, S., HERMY, M., VELLEND, M., KNAPEN, A. & VERHEYEN, K. (2008). Garden  
1109 plants get a head start on climate change. *Frontiers in Ecology and the Environment* **6**,  
1110 212–216.

- 1111 VAN KLEUNEN, M., DAWSON, W., ESSL, F., PERGL, J., WINTER, M., WEBER, E., KREFT, H.,  
1112 WEIGELT, P., KARTESZ, J., NISHINO, M., ANTONOVA, L. A., BARCELONA, J. F., CABEZAS, F.  
1113 J., CÁRDENAS, D., CÁRDENAS-TORO, J., *ET AL.* (2015). Global exchange and accumulation  
1114 of non-native plants. *Nature* **525**, 100–103.
- 1115 VAN KLEUNEN, M. & JOHNSON, S. D. (2007). South African Iridaceae with rapid and profuse  
1116 seedling emergence are more likely to become naturalized in other regions. *Journal of*  
1117 *Ecology* **95**, 674–681.
- 1118 VERBRUGGE, L. N. H., LEUVEN, R. S. E. W., VAN VALKENBURG, J. L. C. H. & VAN DEN BORN,  
1119 R. J. G. (2014). Evaluating stakeholder awareness and involvement in risk prevention of  
1120 aquatic invasive plant species by a national code of conduct. *Aquatic Invasions* **9**, 369–  
1121 381.
- 1122 VILÀ, M., ESPINAR, J. L., HEJDA, M., HULME, P. E., JAROŠÍK, V., MARON, J. L., PERGL, J.,  
1123 SCHAFFNER, U., SUN, Y. & PYŠEK, P. (2011). Ecological impacts of invasive alien plants: a  
1124 meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters*  
1125 **14**, 702–708.
- 1126 VILJOEN, A., BOHN, K. & HOWE, J. (EDS.) (2012). *Continuous productive urban landscapes*.  
1127 Architectural Press, Amsterdam.
- 1128 VÍTKOVÁ, M., MÜLLEROVÁ, J., SÁDLO, J., PERGL, J. & PYŠEK, P. (2017). Black locust  
1129 (*Robinia pseudoacacia*) beloved and despised: a story of an invasive tree. *Forest Ecology*  
1130 *and Management* **384**, 287–302.
- 1131 WAGENITZ, G. (1964). *Solidago* L. In *Illustrierte Flora von Mitteleuropa* (ed G. HEGI), pp.  
1132 16–29. Calr Hanser, Munich, Germany (in German).
- 1133 WARD, B. J. (2004). *The plant hunter's garden. The new explorers and their discoveries*.  
1134 Timber Press, Portland, USA.

- 1135 WEBER, E. (1998). The dynamics of plant invasions. A case study of three exotic goldenrod  
1136 species (*Solidago* L.) in Europe. *Journal of Biogeography* **25**, 147–154.
- 1137 WEBER, E. (2003). *Invasive plant species of the world: a reference guide to environmental*  
1138 *weeds*. CABI Publishing, Wallingford, UK.
- 1139 WHITTLE, T. (1970). *The plant hunters*. Heinemann, London, UK.
- 1140 WÖRZ, A. (2016). *Der Esslinger Botanische Reiseverein 1825–1845*. Logos Verlag, Berlin,  
1141 Germany (in German).
- 1142 WOULDSTRA, J. (2003). The changing nature of ecology: a history of ecological planting  
1143 (1800–1980). In *The dynamic landscape* (eds N. DUNNETT & J. HITCHMOUGH) pp. 23–57.  
1144 Spon Press, London, UK.
- 1145 XIONG, J.-S., DING, J. & LI, Y. (2015). Genome-editing technologies and their potential  
1146 application in horticultural crop breeding. *Horticulture Research* **2**, 15019.
- 1147 ZENNI, R. D. (2014). Analysis of introduction history of invasive plants in Brazil reveals  
1148 patterns of association between biogeographical origin and reason for introduction. *Austral*  
1149 *Ecology* **39**, 401–407.
- 1150 ZHAO, L. J. & ZHANG, D. L. (2003). Ornamental plant resources from China. *Acta*  
1151 *Horticulturae* **620**, 365–375.

1152

## 1153 X. SUPPORTING INFORMATION

1154 Additional supporting information may be found in the online version of this article.

1155 **Appendix S1.** The questionnaire sent to botanical gardens.



1156 Table 1. Eight key research topics proposed for studying horticulture and plant invasions,  
 1157 associated priority research questions, and the required data and methods.

#	Research topics	Priority questions	Required data and methods
1	Origins of ornamentals and routes of introduction and distribution	Why are new species being introduced? How are they selected? From where do they come? What is the import volume? How are introduced species distributed?	Qualitative and quantitative data on species introductions from the horticultural trade, customs duties, sales volume
2	Temporal dimensions, predicting new developments and emerging trends on horticultural trade and plant invasion	What will the future trends in horticulture be? Which species will be next to become invasive? How did and how will horticultural invaders change (fashions, traits, trade volume)?	Questionnaire to horticultural experts, qualitative and quantitative data and approaches from different scientific domains, phenomenological and mechanistic models
3	Identifying the drivers of horticulture-related plant invasions, identifying future invaders from the horticultural trade	How does trade volume and planting frequency affect invasiveness of horticultural species? How does this depend on habitat characteristics, species traits, and global change (habitat loss, land-use change, climate warming)?	Measuring propagule pressure, assessing ability to become naturalised by experimental means
4	Interactions with other features of global change: climate, land-use, urbanisation, eutrophication, habitat loss and fragmentation	How will global environmental change interact with horticulture on plant invasions?	Quantitative models on the current and future interactions of horticulture and other environmental changes
5	Assessing and predicting impacts of alien plants introduced by horticulture	What are the current impacts of alien plants introduced by horticulture? What will be the impacts of current and future ornamental plants?	Qualitative and quantitative data and approaches from different scientific domains, phenomenological and mechanistic models
6	Management: tools, effectiveness, monitoring and implementation	Do we have enough expertise to detect, monitor and manage invasive alien species introduced by horticulture? How can the	Data and models on monitoring and management measures, implementation, analysing and improving management efficiency

relevant methods be improved? Are efficient management and methods species and site specific or can generalisations be made?

7 Legal frameworks

Are current legal frameworks for combating invaders from the horticultural trade sufficient and effective? What roles do voluntary codes of conduct have?

Analyses of the coverage, implementation and effectiveness of current legislation, assessment of different legal tools

8 Raising public awareness, stakeholder partnerships, capacity building and promoting non-invasive species/cultivars

Are people sufficiently informed about invaders? How can communication tools be adapted to maximise the number of people reached? Who are the key people to reach? How to build mutually beneficial partnerships?

Qualitative and quantitative surveys and questionnaires of gardeners, authorities, and managers of invasive species

---

1158

1159

1160 **Fig. 1.** The main pools (boxes) and flows (arrows) of species introduced for ornamental  
1161 purposes, and the actors and processes involved. The width of the different species pools  
1162 illustrate differences in their sizes: the cultivated species pool represents a subset of the wild  
1163 species pool, and the escaped species pool is a subset of the cultivated species pool. Note that  
1164 although we do not include arrows from breeders and propagators, and from wholesalers and  
1165 retailers to the escaped species pool, alien plants may also escape at those stages of the  
1166 supply chain. The dashed arrow indicates that the escaped alien species become part of the  
1167 wild species pool, and thus that in certain regions alien species might subsequently be  
1168 collected again for ornamental purposes. Across the different horticultural and ornamental  
1169 trade stages, the size of the cultivated species pool changes; some of the species collected by  
1170 plant hunters will not be used by breeders and propagators, but the latter will through  
1171 breeding and hybridisation create new taxa, and some of the species offered by the nursery  
1172 trade network of wholesalers and retailers will not be sold and planted. The thin arrows from  
1173 plant hunters to botanical gardens and domestic gardens, indicate that some species planted in  
1174 these gardens were collected in the wild, and by-passed the commercial ornamental plant  
1175 industry. The looped arrow for botanical gardens indicates the exchange of seeds/plants  
1176 among botanical gardens and the looped arrow for domestic gardens indicates the exchange  
1177 of seeds/plants among hobby gardeners. Public spaces include both public green spaces (e.g.  
1178 city parks) and infrastructure (e.g. road-side plantings). For similar diagrams, see Drew *et al.*  
1179 (2010) and Hulme *et al.* (2018).

1180

1181 **Fig. 2.** Venn diagram illustrating that most of the species that have become naturalised  
1182 somewhere in the world are grown in private gardens and in botanical gardens. A circle  
1183 illustrating the size of the global vascular plant flora has been added for comparison. Data on  
1184 the global naturalised flora were extracted from the Global Naturalized Alien Flora database

1185 (GloNAF version 1.1; van Kleunen *et al.*, 2015). Data on species grown in private gardens  
1186 were extracted from Dave’s Garden PlantFiles (<http://davesgarden.com/guides/pf/>) and the  
1187 Plant Information Online database (<https://plantinfo.umn.edu/>). Data on species grown in  
1188 botanical gardens were extracted from the PlantSearch database of Botanic Gardens  
1189 Conservation International (BGCI; [http://www.bgci.org/plant\\_search.php](http://www.bgci.org/plant_search.php)). All species names  
1190 were standardised according to The Plant List (<http://www.theplantlist.org/>), which also  
1191 provided the number for the size of the global vascular plant flora.

1192

1193 **Fig. 3.** Among naturalised species, those grown in domestic or botanical gardens have  
1194 become naturalised in more regions around the globe than species not known to be grown  
1195 (labelled ‘No’ on figure) in gardens (Kruskal-Wallis  $\chi^2 = 1379.8$ ,  $df = 3$ ,  $P < 0.001$ ). Data  
1196 were taken from the Global Naturalized Alien Flora database (version 1.1; van Kleunen *et al.*,  
1197 2015), Dave’s Garden PlantFiles (<http://davesgarden.com/guides/pf/>), the Plant Information  
1198 Online database (<https://plantinfo.umn.edu/>) and PlantSearch of Botanic Gardens  
1199 Conservation International ([http://www.bgci.org/plant\\_search.php](http://www.bgci.org/plant_search.php)).

1200

1201 **Fig. 4.** (A) Absolute and (B) normalised first-record rates for naturalised species that are not  
1202 known to be planted in gardens, and that are planted in domestic gardens (Dave’s Garden  
1203 PlantFiles, <http://davesgarden.com/guides/pf/>; the Plant Information Online database,  
1204 <https://plantinfo.umn.edu/>), botanical gardens (PlantSearch of Botanic Gardens Conservation  
1205 International, [http://www.bgci.org/plant\\_search.php](http://www.bgci.org/plant_search.php)) or both. The data on first-record rates  
1206 were taken from Seebens *et al.* (2017). First-record rates are defined as the number of first  
1207 records of alien species per ten-year period. As the first-record rates for naturalised species  
1208 that are only known to occur in domestic gardens or in no garden at all were very low, the  
1209 inset of A zooms in on those species. In B, the data were normalised by setting the highest

1210 first-record rate of each group equal to 1, and changing the other values proportionally. The  
1211 trends in B are indicated by running medians (lines).

1212

1213 **Fig. 5.** (A) The import value (US\$) of live plants to each country averaged for the period  
1214 2001–2010, and expressed per person. Plant import data were extracted from the United  
1215 Nations Commodity Trade Statistics database (Comtrade; <http://comtrade.un.org>), and  
1216 included commodity codes 0601 (bulbs and seeds) and 0602 (other live plants). Human  
1217 population data were taken from CIESIN *et al.* (2011). Values are presented as 20%  
1218 quantiles. (B) The increase in the imports of live plants expressed relative to the region with  
1219 the greatest increase, Europe. Rates of increase were calculated as the area under the trend  
1220 curve, and for East Asia was calculated from 2005 to 2015 due to the decrease in plant  
1221 imports that occurred prior to that. (C, D) Change in import value (US\$) of live plants (from  
1222 1995 to 2015, reliable plant import data were not available before 1995), for the highest four  
1223 (C) and lowest five (D) importing regions shown in B. Colours correspond to the legend in B.  
1224 As the rates of increase for Africa and Western Asia were identical, we distinguish Africa  
1225 with white stippling on the map in panel B, and a dashed line on the graph in panel D. Import  
1226 values were summed across all countries in a region, and regions were defined according to  
1227 sub-continent and similarity among import trends. Import values and trends were very similar  
1228 for some geographically disjunct regions, and so values were aggregated to reduce the  
1229 number of lines and maximise colour differences: for Central-South America and Africa  
1230 Pearson's  $r=0.81$ ,  $P<0.00001$ ,  $df=19$ ; the combined import values for Central-north Asia,  
1231 south and south-east Asia, and Oceania were grouped as they were relatively low.

1232


1233 **Fig. 6.** Proportion of 947 botanical gardens across six continents that participate in retail plant  
1234 sales, horticulture or plant breeding research, or undertake plant explorations. Data from

1235 Botanic Garden Conservation International Garden Search  
1236 ([www.bgci.org/garden\\_search.php](http://www.bgci.org/garden_search.php); accessed on 1 November 2016).

1237

1238 **Fig. 7.** Main sources of plants in botanical gardens, based on a questionnaire to which 161  
1239 botanical gardens responded. Six of the botanical gardens indicated two sources as the main  
1240 ones; these were assigned to both sources. The botanical gardens were grouped according to  
1241 continent (TDWG continent; Brummitt, 2001).

1242 **FIGURE 1**




1243

1244

1245

1246 **FIGURE 2**




1247

1248

1249






1251

1252


1253 **FIGURE 4**



1254

1255

1256




1258

1259

1260

1261 **FIGURE 6**

1262




1263

1264

1265 **FIGURE 7**

1266



1267

1268