



**UNIVERSITY OF LEEDS**

This is a repository copy of *Advanced Radiotherapy Technologies Network in the UK (ART-NET) - focus on lung cancer*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/128682/>

Version: Accepted Version

---

**Proceedings Paper:**

Brown, S, Van Herk, M, Chuter, R et al. (18 more authors) (2018) *Advanced Radiotherapy Technologies Network in the UK (ART-NET) - focus on lung cancer*. In: *Lung Cancer. 16th Annual British Thoracic Oncology Group Conference 2018, 24-26 Jan 2018, Dublin, Ireland*. Elsevier , S60-S60.

[https://doi.org/10.1016/S0169-5002\(18\)30166-1](https://doi.org/10.1016/S0169-5002(18)30166-1)

---

(c) 2018, Elsevier Ireland Ltd. Published by Elsevier Inc. This manuscript version is made available under the CC BY-NC-ND 4.0 license  
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Introduction:** Lung cancer is the most frequently diagnosed cancer worldwide. Whilst the role of curative-intent radiotherapy is widely accepted, the thorax remains a challenging anatomical site to irradiate. New advanced radiotherapy technologies including stereotactic ablative radiotherapy (SABR), magnetic resonance image-guided radiotherapy (MR-Linac) and proton-beam therapy (PBT) have potential to yield substantial clinical benefits. However there are significant challenges for development, assessment and rational implementation within the NHS.

**Methods:** ART-NET is funded by a Cancer Research UK accelerator award. The goal of ART-NET is to accelerate clinical translation of SABR, MR-Linac and PBT, around the unifying theme of hypo-fractionated radiotherapy, and to provide an evidence-based rationale across a range of tumour types. This is being conducted across 5 UK centres and will develop and disseminate expertise in medical physics required to lead the development of planning and image-guided solutions for these technologies.

**Results:** ART-NET will implement methodologies for reliable treatment plan optimization on MR-images, acquired directly prior to or during treatment, enabling the clinical benefit of MR-guided SABR and MR-Linac to be fully exploited. ART-NET will also co-ordinate methodological developments in design and conduct of clinical trials to streamline assessments of new health technologies in specific tumours. An example is the development of a traffic light system to facilitate online radiographer review of CBCTs in patients undergoing lung SABR to assess organ motion and anatomical changes. This will inform the need for plan adaptation nationally.

**Conclusion:** A key aim of ART-NET is to generate national treatment protocols and disseminate them across the UK, improving and harmonising practice. Additionally, involvement of medical physicists, research radiographers and clinical research fellows in ART-NET will develop a skilled workforce that will be a huge asset for future radiotherapy research in the UK. Ultimately this will improve patient access to state-of-the-art radiotherapy and increase recruitment into clinical trials.