



UNIVERSITY OF LEEDS

This is a repository copy of *Health-related quality of life after treatment for bladder cancer in England*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/130368/>

Version: Supplemental Material

Article:

Mason, SJ, Downing, A orcid.org/0000-0002-0335-7801, Wright, P orcid.org/0000-0001-6129-4143 et al. (6 more authors) (2018) Health-related quality of life after treatment for bladder cancer in England. *British Journal of Cancer*, 118 (11). pp. 1518-1528. ISSN 0007-0920

<https://doi.org/10.1038/s41416-018-0084-z>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Supplementary Table 2. Comparison of Demographics and Clinical Characteristics Between Respondents and Nonrespondents						
Demographic or clinical characteristic	Non-respondents		Respondents		Total	Response Rate (%)
	N	%	N	%		
Sex						
Male	413	71.5%	508	75.5%	921	55.2
Female	165	28.5%	165	24.5%	330	50.0
Total	578	100.0%	673	100.0%	1251	53.8
Age						
<55	44	7.6%	47	7.0%	91	51.6
55-64	91	15.7%	128	19.0%	219	58.4
65-74	164	28.4%	252	37.4%	416	60.6
75-84	198	34.3%	194	28.8%	392	49.5
85+	81	14.0%	52	7.7%	133	39.1
Total	578	100.0%	673	100.0%	1251	53.8
Stage						
1	42	7.3%	73	10.8%	115	63.5
2	15	2.6%	23	3.4%	38	60.5
3	6	1.0%	3	0.4%	9	33.3
4	3	0.5%	6	0.9%	9	66.7
Unknown	512	88.6%	568	84.4%	1080	52.6
Total	578	100%	673	100.0%	1251	53.8

NOTE. Based on cancer registry data available for all individuals