



This is a repository copy of *Progressive ductile shearing during till accretion within the deforming bed of a palaeo-ice stream*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/132079/>

Version: Accepted Version

---

**Article:**

Phillips, E., Spagnolo, M., Pilmer, A. et al. (4 more authors) (2018) Progressive ductile shearing during till accretion within the deforming bed of a palaeo-ice stream. *Quaternary Science Reviews*, 193. pp. 1-23. ISSN 0277-3791

<https://doi.org/10.1016/j.quascirev.2018.06.009>

---

Article available under the terms of the CC-BY-NC-ND licence  
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Progressive ductile shearing during till accretion within the deforming bed of a palaeo-ice stream

Emrys Phillips<sup>1\*</sup>, Matteo Spagnolo<sup>2</sup>, Alasdair C.J. Pilmer<sup>2</sup>, Brice R. Rea<sup>2</sup>, Jan A. Piotrowski<sup>3</sup>, Jeremy C. Ely<sup>4</sup> and Simon Carr<sup>5</sup>

<sup>1</sup> British Geological Survey, The Lyell Centre, Research Avenue South, Edinburgh EH14 4AP, UK

<sup>2</sup> Department of Geography and the Environment, School of Geosciences, University of Aberdeen, Aberdeen AB24 3UF, UK

<sup>3</sup> Department of Geoscience, Aarhus University, Denmark

<sup>4</sup> Department of Geography, University of Sheffield, Sheffield S10 2TN, UK

<sup>5</sup> School of Geography, Queen Mary, University of London, Mile End Road, London E1 4NS, UK

\* Corresponding author: e-mail - [erp@bgs.ac.uk](mailto:erp@bgs.ac.uk)

**Keywords:** micromorphology; glacier bed deformation; ductile shearing; foliation development; palaeo-ice stream

## Abstract

This paper presents the results of a detailed microstructural study of a thick till formed beneath the Weichselian (Devensian) Odra palaeo-ice stream, west of Środa Wielkopolska, Poland. This SE-flowing ice stream was one of a number of corridors of faster flowing ice which drained the Scandinavian Ice Sheet in the Baltic region. Macroscopically, the massive, laterally extensive till which formed the bed of this ice stream lacks any obvious evidence of glaciotectonism (thrusting, folding). However, microscale analysis reveals that bed deformation was dominated by foliation development, recording progressive ductile shearing within a subhorizontal subglacial shear zone. Five successive generations of clast microfabric (S1 to S5) have been identified defining a set of up-ice and down-ice dipping Riedel shears, as well as a subhorizontal shear foliation coplanar to the ice-bed interface. Cross-cutting relationships between the shear fabrics record temporal changes in the style of deformation during this progressive shear event. Kinematic indicators (S-C and ECC-type fabrics) within the till indicate a consistent SE-directed shear sense, in agreement with the regional ice flow pattern. A model of bed deformation involving incremental progressive simple shear during till accretion is proposed. The relative age of this deformation was diachronous becoming progressively younger upwards, compatible with subglacial shearing having accompanied till accretion at the top of the deforming bed. Variation in the relative intensity of the microfabrics

31 records changes in the magnitude of the cumulative strain imposed on the till and the degree of  
32 coupling between the ice and underlying bed during fast ice flow.

## 33 **1. Introduction**

34 Ice streams play an important role in regulating the behaviour of modern ice sheets (e.g. Antarctica,  
35 Bamber *et al.*, 2000) and take the form of corridors of fast flowing ice bounded by ice flowing up to  
36 an order of magnitude slower (Stokes and Clark, 2001; Bennett, 2003). However the factors  
37 controlling fast ice flow are incompletely understood. Published studies of modern and ancient ice  
38 stream beds have led to two possible explanations governing ice stream flow: (i) basal sliding  
39 facilitated by elevated water pressures at the ice-bed interface with the ice stream effectively  
40 becoming decoupled from the underlying sediments (e.g. Alley, 1989; Piotrowski and Tulaczyk, 1999)  
41 or hard bedrock substrate (Margold *et al.*, 2015); and (ii) basal motion accommodated by  
42 deformation of either a thick (several metres) or thin (centimetres to decimetres) layer of 'soft'  
43 sediments (till) (e.g. Alley *et al.*, 1986, 1987a, b; Boulton and Hindmarsh, 1987; Clarke, 1987;  
44 Humphrey *et al.*, 1993; Boulton *et al.*, 2001). However, in reality these two processes are not  
45 mutually exclusive and may periodically "switch" to form the dominant movement mechanism of ice  
46 stream movement depending upon the water content and/or pressure within the bed.  
47 Understanding these processes has fundamental implications for our understanding of subglacial  
48 sediment erosion, transport and deposition. Furthermore a greater understanding of the subglacial  
49 environment of ice streams may also elucidate controls on ice streaming such as basal thermal  
50 regime (Hindmarsh, 2009) and/or subglacial hydrology (Kyrke-Smith *et al.*, 2015), leading to the  
51 development of more sophisticated and robust models of ice stream flow dynamics and, ultimately,  
52 ice sheet mass balance and sea-level change.

53 The recognition of a characteristic suite of subglacial landforms (including megascale glacial  
54 lineations) formed beneath palaeo-ice streams (e.g. Dyke and Morris, 1988; Hodgson, 1994;  
55 Patterson, 1997, 1998; Clark and Stokes, 2001, 2002, 2003) has enabled the establishment of a set of  
56 criteria for identifying the presence and areal extent of these ancient ice streams (Stokes and Clark,  
57 1999). These criteria have been, at least partially, validated by observations of the subglacial  
58 landscape beneath contemporary Antarctic ice-streams (King *et al.*, 2009; Bingham *et al.*, 2017). The  
59 exposed beds of palaeo-ice streams provide an ideal laboratory to investigate the sedimentary and  
60 structural processes occurring beneath fast flowing ice. However, on a macroscale the sediments  
61 (tills) deposited beneath many palaeo-ice streams are massive, lacking any visible signs of  
62 stratification and/or glaciectonic deformation (see Evans, 2018 and references therein). As a  
63 consequence micromorphology is increasingly being used as a primary tool for the analysis of these

64 and other subglacial sediments (tills) (see Menzies and Maltman, 1992; van der Meer, 1979, 1987;  
65 Menzies *et al.*, 1997; Khatwa and Tulaczyk, 2001; van der Meer *et al.*, 2003; Hiemstra *et al.*, 2005;  
66 Baroni and Fasano, 2006; Larsen *et al.*, 2006, 2007; Phillips *et al.*, 2007, 2011, 2013, 2018; Narloch *et al.*,  
67 *et al.*, 2012; Neudorf *et al.*, 2013; Gehrmann *et al.*, 2017; Evans, 2018). This technique can provide far  
68 greater detail on the depositional and deformation histories recorded by these sediments than can  
69 be obtained from macroscale studies alone; for example, unravelling the often complex deformation  
70 histories recorded by glacial sequences (van der Meer, 1993; Phillips and Auton, 2000; van der  
71 Wateren *et al.*, 2000; Menzies, 2000; Phillips *et al.*, 2007; Lee and Phillips, 2008; Vaughan-Hirsch *et al.*,  
72 *et al.*, 2013; Narloch *et al.*, 2012, 2013) and the role played by pressurised meltwater during their  
73 deformation (Hiemstra and van der Meer, 1997; Phillips and Merritt, 2008; van der Meer *et al.*,  
74 2009; Denis *et al.*, 2010; Phillips *et al.*, 2013; 2018; Narloch *et al.*, 2012, 2013).

75 This paper presents the results of a detailed micromorphological study of the thick till  
76 sequence laid down by the Weichselian (Devensian) Odra palaeo-ice stream as it flowed SE across  
77 Wielkopolska Lowlands of Poland (Fig. 1). The study area is located near Poznań, in a region  
78 dominated by NW-SE-trending subglacial landforms (megascala lineations) interpreted as having  
79 been formed during fast ice flow (Przybylski, 2008; Spagnolo *et al.*, 2016). Thin sections are used to  
80 investigate the strain signature imparted by this palaeo-ice stream on the laterally extensive till  
81 formed within its bed. The results of this detailed microstructural study have been used to  
82 investigate the nature of deformation and in particular foliation development during progressive  
83 ductile simple shear within an evolving subhorizontal subglacial shear zone. Spatial variations in the  
84 relative intensity of the microfabrics are interpreted as recording changes in the magnitude of the  
85 cumulative strain imposed on the till, potentially reflecting the degree of ice-bed coupling during fast  
86 ice flow.

## 87 **2. Location of study area and geological setting**

88 During the Weichselian (Devensian) glaciation much of the Baltic region was covered by the  
89 Scandinavian Ice Sheet. This ice sheet was drained by a series of ice streams, including the Odra  
90 palaeo-ice stream (OPIS) which flowed SE across the Wielkopolska Lowland region of western Poland  
91 (Przybylski, 2008; Spagnolo *et al.*, 2016). In this region, the bed of the Odra palaeo-ice stream (over  
92 1000 km<sup>2</sup>) is characterised by a suite of well-preserved NW-SE-trending megascala glacial lineations  
93 (MSGSL) underlain by a thick (c. 30 m) sequence of Quaternary sediments. This study is focused on  
94 the bed of the OPIS to the west of the town of Środa Wielkopolska, approximately 30 km southeast  
95 of Poznań (Fig. 1a, b), close to the c. 21 ka Leszno phase ice margin (Kozarski, 1988; Przybylski, 2008;  
96 Marks, 2012). The geomorphology of the study area (c. 180 km<sup>2</sup>) is dominated by a suite of elongate

97 (>16 km long), low-relief (2-4 m high) MSGL with a crest-to-crest spacing of 600-800 m (Fig. 1c). It is  
98 possible that these landforms were originally much longer (Przybylski, 2008) as they have been  
99 locally truncated by glacial erosion, as well as the extensive urbanisation of the region which has  
100 locally overprinted/strongly modified this subglacial landscape. Although locally modified the  
101 morphology of these subglacial landforms are comparable to MSGL described from other palaeo-ice  
102 stream settings worldwide (Spagnolo *et al.*, 2014).

103 The sediments making up the bed of the OPIS are in general poorly exposed and detailed  
104 sedimentological analysis of these deposits has relied upon trenches excavated at key positions  
105 across the MSGL's (Fig. 1c; see below). The trenches reveal that these subglacial landforms are  
106 composed of a homogeneous, matrix-supported, yellow coloured silty-sandy diamicton (Fig. 1d)  
107 containing rare gravel (2-64 mm) and extremely rare cobble (>64 mm) sized clasts (Spagnolo *et al.*,  
108 2016). The massive, laterally extensive diamicton (interpreted as a subglacial traction till; *sensu*  
109 Evans *et al.*, 2006) lacks any obvious macroscale evidence of glaciotectonism (e.g. thrusting, folding...  
110 etc.) and no other sedimentary units have been recognised. Fine gravel clasts (2-4 mm) contained  
111 within the diamicton are composed of a range of sedimentary and crystalline rock fragments,  
112 including Palaeozoic limestones derived from the Baltic Basin as well as metamorphic and igneous  
113 rocks from Scandinavia, indicating that this deposit contains a significant far-travelled component  
114 (Spagnolo *et al.*, 2016). The diamicton is relatively unaltered, exhibiting only very minor to rare  
115 macroscopic evidence of calcification, typically occurring in patches of <200 cm<sup>2</sup>. Clast a-axis  
116 macrofabric data published by Spagnolo *et al.* (2016) are remarkably uniform (vertically and  
117 laterally) across the bed of the OPIS within the study area. These shallow dipping macrofabrics are  
118 orientated NW-SE, concordant with the long axes of the MSGL and parallel to the regional ice flow  
119 direction.

### 120 **3. Methods**

121 Detailed analysis of the till forming the bed of the OPIS has focussed on 10 sites located on the crests  
122 (A, B, C, D, E, K, T; Fig. 1c) and flanks (X, Y, Z; Fig. 1c) of three of the mega-scale lineations. A trench  
123 (6-10 m long, 2-3 m wide and 3-5 m deep) was excavated at each site (Fig. 1d) and the samples for  
124 thin section preparation collected using standard Kubiena tins. Prior to sampling, the temporarily  
125 exposed sections were logged, photographed and described in detail with particular emphasis being  
126 placed on recording the macroscale variation in lithology and structure of the till. The samples were  
127 collected in a vertical profile (e.g. C1M highest to C6M lowest) below the base of the modern soil  
128 and with a 20 cm spacing between each Kubiena tin (Fig. 1d). This approach was adopted to provide  
129 detailed information on the range of microstructures developed at different

130 stratigraphical/structural levels within the till. The Kubiena tins were either cut or pushed into the  
131 face in order to limit sample disturbance. The position of the sample within the sequence, its  
132 orientation relative to magnetic north, depth and way-up were marked on the outside of the tin  
133 during collection.

134 Sample preparation was carried out at Royal Holloway, University of London, using the  
135 methods outlined by Palmer (2005). Large format (10 x 8 cm), orientated (parallel to the long axis of  
136 the MSGL and former ice flow direction) thin sections were taken from the centre of each of the  
137 resin impregnated samples, avoiding artefacts associated with sample collection. The thin sections  
138 were described using a Zeiss petrological microscope revealing that the composition, texture and  
139 structure of the diamicton are uniform across the study area. The detailed microscale study focused  
140 upon site C, located on the crest of a prominent NW-SE-trending MSGL, as well as three samples  
141 from sites X, Y and Z, which provide a traverse across the flank of one of these landforms (Fig. 1c).  
142 The location of site C on the crest of the MSGL means that the thin sections represent a vertical  
143 section through the landform and therefore provide a valuable insight into the processes which may  
144 have occurred during the formation of this landform. The terminology used to describe the various  
145 microtextures developed within these sediments follows that proposed by van der Meer (1987,  
146 1993) and Menzies (2000) with modifications. Microstructural maps and quantitative data for the  
147 clast microfabrics (Figs. 2 to 7) developed within the till were obtained using the methodology of  
148 Phillips *et al.* (2011) (also see Vaughan-Hirsh *et al.*, 2013; Neudorf *et al.*, 2013; Gehrmann *et al.*,  
149 2017; Phillips *et al.*, 2013, 2018; Brumme, 2015). During this process the relationships between  
150 successive generations of clast microfabrics (S1 oldest to Sn youngest) and other microstructures  
151 (e.g. plasmic fabrics, turbate structures, folds, faults, shears..etc) present within the diamicton are  
152 determined, allowing a detailed relative chronology of fabric development to be established,  
153 enabling the investigation of the complex polyphase deformation histories recorded in these  
154 deposits (see Phillips *et al.*, 2011 for details of this process). Each thin section was divided into 16  
155 rectangular areas (A to P on Figs. 2 to 7) and the orientation of the long axes of the clasts (skeleton  
156 grains) plotted on a series of rose diagrams and the eigenvalues (E1, E2) calculated for each area  
157 using the commercial software package StereoStat by RockWare™ (see Fig. 2 to 7). In order to  
158 assess the effects of grain size on clast microfabric development within tills, the long axis data were  
159 divided into three sets: (i) grains < 0.25 mm in length (fine-sand and below); (ii) grains between 0.25  
160 to 0.5 mm in size (medium-sand); and (iii) grains over > 0.5 mm in length (coarse-sand and above).  
161 The resultant data sets were plotted on a series of histograms and rose diagrams to highlight any  
162 variation in clast size versus long axis orientation.

163 In conjunction with the manual microstructural mapping methodology (Phillips *et al.*, 2011)  
164 an automated approach using ArcGIS line density tools was conducted on selected thin sections  
165 (X1M, Y1M, Z1M) to provide a robust, objective interpretation of the clast microfabrics. These  
166 automated tools allow the calculation of the magnitude of long axes per unit area within a thin  
167 section and was applied to each of the clast microfabrics. The variation in density ( $\text{mm}^2$ ) of the clasts  
168 defining each microfabric was calculated with the resulting output raster files providing a map of  
169 relative intensity of clast microfabric for each thin-section (Fig. 8).

## 170 **4. Results of the micromorphological and microstructural analysis**

### 171 **4.1. Composition and provenance of the till**

172 In thin section (C1M, C2M, C3M, C4M, C5M, C6M) the till is massive, lacking any obvious  
173 stratification (e.g. bedding) or other primary structure. It is composed of fine- to medium-grained,  
174 open-packed, matrix-supported, silty sand (Figs. 2-7) containing scattered, angular to well-rounded  
175 granule, to small pebble-sized rock fragments composed of sedimentary rocks (siltstone, sandstone,  
176 mudstone, indurated quartz-arenite, bioclastic limestone, micritic limestone), igneous (biotite-  
177 granite, muscovite-granite, alkali granite, micrographic intergrowth, altered volcanic rocks) and  
178 metamorphic rocks (amphibolite, biotite-schist) (Table 1). Angular to subangular, coarse-silt to sand-  
179 sized grains within the till matrix are composed of monocrystalline quartz and feldspar (plagioclase,  
180 K-feldspar). The compositional data support the conclusion of Spagnolo *et al.* (2016) that the till was  
181 laid down by ice advancing from the NW and contains far-travelled material derived from Palaeozoic  
182 sedimentary sequences within the Baltic Basin and crystalline basement rocks from Scandinavia;  
183 similar till compositions have also been reported from Germany and Denmark (Piotrowski, 1994a, b;  
184 Kjær *et al.*, 2003).

185 The thin sections reveal that the till is compositionally, essentially homogenous with only a  
186 slight increase in the proportion of limestone and fine carbonate grains downwards through the  
187 sequence (Table 1). This increase in detrital carbonate is accompanied by the appearance of small,  
188 irregular patches of a micritic carbonate cement which appears to replace the clay within the matrix  
189 (Figs. 9a, b, c; red areas on Figs. 4 to 7). Small, rounded to irregular voids and fractures within the till  
190 are lined or filled by massive to very finely laminated, highly birefringent clay (Figs. 9d, e, f). These  
191 clay-filled features form between 5 and 15% (visual estimate) of the matrix, and locally (e.g. C1M)  
192 define a weakly developed subhorizontal "foliation". The dark orange-brown clay is petrographically  
193 similar to clay cutan within soils, suggesting that it was deposited by water flowing through the till  
194 matrix, with the laminated nature of these fines recording several phases of fluid flow.

## 195 **4.2. Microstructures developed in response to subglacial deformation**

196 Microstructural analysis of the thin sections (C1M to C6M; Figs. 2 to 7, respectively) has revealed  
197 that the tills possess five successive generations of clast microfabric (S1 to S5) defined by the  
198 preferred shape alignment of elongate coarse silt to sand-grade clasts. The relative intensity of these  
199 microfabrics varies across the thin section, reflecting the heterogeneous nature of shearing within  
200 the glacier bed. The spacing of the microfabric domains is controlled by the overall grain size of the  
201 diamicton and occurrence of coarse-sand to small pebble-sized clasts which acted as rigid bodies  
202 during deformation. Although the results of detailed microfabric analysis described below focus  
203 upon site C, comparable fabric geometries have been observed in the thin sections from sites X, Y  
204 and Z (Fig. 8).

205 The earliest microfabric is a very poorly developed/preserved, typically down-ice dipping S1  
206 fabric (purple on Figs. 2 to 7). This fabric, where present, is cross-cut by a highly heterogeneous,  
207 subhorizontal to very gently inclined S2 fabric (pale green on Figs. 2 to 7). In the upper part of the till  
208 (C1M to C3M) S2 occurs within weakly to moderately well-defined, lenticular bands (Figs. 2 and 3)  
209 and is interpreted as having formed coplanar to the bed of the overriding ice. A weakly developed  
210 asymmetrical to S-shaped fabric geometry (S-C-type fabric) within the bands of S2 records a sinistral  
211 (in this plane of section) SE-directed (down-ice) sense of shear (Figs. 3, 4, 5 and 7). Lower within the  
212 till sequence (C4M to C6M), however, the banded appearance of S2 is less apparent as this fabric has  
213 been variably overprinted by a later foliation (see below).

214 Poorly to rarely developed, arcuate grain alignments and turbate structures (van der Meer,  
215 1983; Menzies, 2000) occur within the microlithons between S1 and S2, and are locally truncated  
216 against these foliations. Turbate structures are interpreted to have formed where larger clasts rotate  
217 through angles up to, and greater than 360° entraining the adjacent finer grained matrix (van der  
218 Meer, 1993, 1997; Menzies, 2000; Hiemstra and Rijdsdijk, 2003; Lea and Palmer, 2014). Their variable  
219 preservation within the S1 and S2 microlithons suggests that this rotational deformation occurred  
220 prior to the imposition of the clast microfabrics.

221 The dominant fabric is an up-ice dipping S3 microfabric (dark green on Figs. 2 to 7). This fabric  
222 cross-cuts S2, with the earlier S1 being preserved within the microlithons separating the S3 domains.  
223 In detail S3 is composed of two components: (i) a moderately to steeply (40° to 50°) up-ice dipping  
224 foliation; and (ii) a more gently inclined (20° to 40°) foliation (Figs. 2, 3 and 4). The later, down-ice  
225 dipping (20° to 40°) S4, microfabric is heterogeneous, potentially reflecting the partitioning of  
226 deformation into increasingly narrower zones of shear during the later stages of bed deformation. S2  
227 and S3 are deformed by S4 resulting in a distinctive S-shape to sigmoidal fabric geometry

228 comparable to an extensional crenulation cleavage (ECC fabric) associated with extensional shears  
229 formed in brittle-ductile shear zones (Passchier and Trouw, 1996). This fabric geometry once again  
230 records a sinistral, SE-directed sense of shear (Figs. 2, 3, 5 and 6) consistent with the ice movement  
231 direction in the study area (see Fig. 1c).

232 The same microfabric relationships (S2 to S4) were recorded in all thin sections (see Figs. 2 to 7)  
233 indicating that not only were these fabrics developed in response to the same overall stress regime,  
234 but that this regime (dominated by SE-directed shear) was maintained throughout the deposition of  
235 the entire till sequence. This conclusion is supported by the rose diagrams shown on Figs. 2 to 7  
236 which indicate that the orientation of S2, S3 and S4 remains essentially constant throughout the till  
237 with very little modification due to compaction. The geometry of these fabrics is consistent with  
238 their development in response to the formation of Y (S2), R (S3) and P-type (S4) Reidel shears (c.f.  
239 Spagnolo *et al.*, 2016) within a subhorizontal subglacial shear zone formed beneath the overriding  
240 OPIS (Fig. 10).

241 S1 to S4 shear related fabrics are cross-cut by a subvertical, anastomosing S5 fabric which locally  
242 wraps around the larger granule to pebble sized clasts (Figs. 3 to 7). The overall intensity of this  
243 fabric increases down-wards through the till (C2M to C6M) where it locally overprints the earlier  
244 developed foliations. The patches of micritic carbonate within the matrix of the till occur within, or  
245 close to the areas possessing a well-developed S5 fabric (Figs. 4 to 7), indicating that development of  
246 this fabric may have accompanied the passage of CaCO<sub>3</sub>-bearing fluids through the sediment (see  
247 section 5).

### 248 **4.3. Effect of grain size on microfabric development**

249 To assess the effects of grain size on clast microfabric development the long axis data were divided  
250 into three sets: (i) < 0.25mm (fine-sand and below); (ii) 0.25 to 0.5 mm (medium-sand); and (iii) > 0.5  
251 mm in size (coarse-sand and above). The orientation data derived for these sets are shown on Figs.  
252 11 and 12. Although the same clast microfabrics are present within all three clast sizes (Fig. 12), they  
253 are most pronounced within the finer grained components with the data highlighting a change in the  
254 orientation of the dominant fabric downwards through the till (Fig. 11). In the “upper” part of the till  
255 (C1M, C2M, C3M), the up-ice dipping S3 is dominant and its relative intensity appears to increase  
256 downwards (Figs. 11 and 12). This variation in the intensity of fabric development may record a  
257 progressive change in the relative intensity of deformation/magnitude cumulative strain imposed at  
258 the ice/bed interface during till accretion (Boulton, 1996; Boulton and Hindmarsh, 1987; Evans *et al.*,  
259 2006 and references therein). In the “lower” part of the till (C4M, C5M, C6M) the relative intensity of  
260 the shear fabrics decreases with S3 being replaced by the down-ice dipping S4 as the dominant

261 foliation (Figs. 11 and 12). However, the fabrics (S2, S3, S4) in the “lower” part of the till have been  
262 strongly modified by the imposition of S5 (see Figs. 5 to 7). The boundary between these “upper”  
263 and “lower” till units appears to be relatively sharp and located between samples C3M and C4M.  
264 However, no obvious boundary was observed at this level within the trench (see Fig. 1c).

## 265 **5. Variation in microfabric intensity within the OPIS till**

266 The results of the micromorphological study indicate that there is a significant variation in the  
267 relative intensity of microfabric development within the till.

### 268 **5.1. Automated clast microfabric analysis**

269 The results of the automated approach to quantify the variation in clast microfabric strength are  
270 shown in Fig. 8. The thin sections used (X1M, Y1M, Z1M) represent a traverse down the flank of an  
271 MSGL (Fig. 1c) designed to investigate any potential lateral changes in the style and relative intensity  
272 of fabric development across this subglacial landform. Importantly this method has revealed a  
273 similar pattern of microfabric development within each of the thin sections as those analysed using  
274 manual methodology of Phillips *et al.* (2011) (compare Figs. 2, 3 and 8). It is clear from Fig. 8 that all  
275 of the microfibrils (S2, S3, S4, S5) are heterogeneously developed, even within a single thin section,  
276 reflecting the partitioning of deformation on a microscale within the bed of the OPIS. S3 is the  
277 dominant fabric and is most intensely developed within the samples located close to the crest of the  
278 MSGL (X1M) and within the adjacent trough (Z1M) (Fig. 8). In contrast, both S2 and S4 are more  
279 weakly developed on top of the landform (X1M) in comparison to its flanks (Y1M, Z1M) the strength  
280 of S5 increases down the flank of the MSGL towards the adjacent trough.

### 281 **5.2. Statistical clast microfabric analysis**

282 The variation in E1 eigenvalues calculated for the samples from site C (C1M to C6M) are illustrated  
283 on Fig. 13. The red colours represent areas of the thin sections with higher E1 values (0.65-0.68),  
284 corresponding to relatively stronger fabric development, and purple lower values (0.53-0.51),  
285 highlighting areas where the clast microfibrils are less well-developed. This approach reveals that  
286 fabric strength not only varies within an individual thin section, but also vertically through the till  
287 (Figs. 13 and 14a), supporting the results of the grain size fabric analysis (Figs. 11 and 12). The  
288 eigenvalues are typically higher within the upper part of the till (C1M, C2M, C3M; Table 2), recording  
289 an overall relative increase in fabric strength upwards through the till (Fig. 13); although sample C3M  
290 has the highest E1 values compared to the other two samples. In contrast, the lower three samples  
291 all possess low eigenvalues (Figs. 13 and 14a; Table 2) corresponding to much weaker fabric  
292 strengths. Although it is tempting to suggest that the lower part of the till is more weakly deformed,

293 this may simply reflect the overprinting of the earlier shear related fabrics (S2 to S4) by the later S5  
294 microfabric (see Figs. 5 to 7).

295 The eigenvalues ( $< 0.6$ ; Fig. 13 and Table 2) can be used to suggest that the amount of shear  
296 being transmitted into the bed of the OPIS was relatively low. In the absence of any obvious strain  
297 markers (e.g. deformed clasts known to have been originally circular in shape) estimating the  
298 magnitude of the shear strains imparted by the overriding ice remains problematic. Several workers  
299 have used the relative abundance of selected microstructures (e.g. microshears, grain stacks) as a  
300 proxy for estimating strain in subglacial traction tills (Larsen *et al.*, 2006, 2007; Narloch *et al.*, 2012).  
301 However, the development of such features can be strongly lithologically controlled; e.g.  
302 microshears defined by a unistrial plasmic fabrics will only form in clay-rich sediments. Furthermore,  
303 their identification is qualitative and potentially subjective (Leighton *et al.*, 2012; Neudorf *et al.*,  
304 2013). Phillips *et al.* (2013) suggested that shear strain curves established from experimental  
305 deformation studies (e.g. Thomason and Iverson, 2006) can be used to provide a minimum estimate  
306 of the shear strains experienced by subglacial traction tills. In the absence of strain curves for a  
307 range of naturally occurring tills, the potential closest “match” to the sand-rich OPIS till is the  
308 Douglas till strain curve of Thomason and Iverson (2006) (Fig. 14b). When projected onto this curve  
309 the range of average E1 values for the OPIS till suggests that microfabric development occurred in  
310 response to shear strains of  $< 15$  (Fig. 14b). If this approach is valid then it supports the suggestion  
311 that the amount of shear being transmitted into the bed of the OPIS by the overriding ice was  
312 relatively low (cf. Larsen *et al.*, 2007; Narloch *et al.*, 2012; Phillips *et al.*, 2013).

## 313 **6. Implications for bed deformation beneath the OPIS**

### 314 ***6.1. Foliation development in response to progressive simple shear***

315 Results of this detailed microscale study reveal that deformation within the bed of the OPIS was  
316 dominated by foliation development which lacked any concomitant folding and/or faulting (c.f.  
317 Spagnolo *et al.*, 2016). The microfabrics define a set of up-ice and down-ice dipping Riedel shears (S3  
318 – P-type and S4 – R-type shears; Fig. 10), as well as a subhorizontal shear foliation (S2 – Y-type  
319 shears; Fig. 10) with S2 having formed parallel to the ice-bed interface (Figs. 10 and 15). The  
320 consistency of the geometry and orientation (Figs. 2 to 7, and 15) of these microfabrics indicate that  
321 not only did subglacial deformation occur in response to the same overall stress regime, but also  
322 that they have undergone very little modification due to compaction/loading subsequent to  
323 formation which would have led to the “flattening” (decrease in dip) of the fabrics at depth within  
324 the till (see rose diagrams on Fig. 15). The consistency of the data also indicates that ductile shearing

325 beneath the OPIS was spatially uniform and occurred within an essentially subhorizontal subglacial  
326 shear zone (see Fig. 10). Furthermore kinematic indicators (S-C and ECC-type microfabrics) record a  
327 consistent SE-directed (sinistral) sense of shear, coincident with the long axes of the MSGL and  
328 regional pattern of ice flow across the Wielkopolska Lowland (Przybylski, 2008; Spagnolo *et al.*,  
329 2016).

330 The cross-cutting relationships displayed by S<sub>2</sub>, S<sub>3</sub> and S<sub>4</sub> can be interpreted as reflecting  
331 temporal changes in the style of deformation being accommodated within the till. However, the  
332 consistent SE-directed sense of shear recorded by these fabrics clearly indicates that they formed in  
333 response to the same overall stress regime imposed during a single progressive shear event rather  
334 than completely separate phases of deformation. S<sub>2</sub> defines a series of subhorizontal Y-type shear  
335 planes indicating that the earlier stages of ductile deformation were dominated by shear occurring  
336 coplanar to the base of the overriding ice. S<sub>2</sub> is cross-cut by the up-ice dipping S<sub>3</sub> foliation which  
337 defines a number of P-type shears indicating that initial layer-parallel shear was superseded by  
338 compressional deformation. S<sub>3</sub> is then cross-cut by the later down-ice dipping S<sub>4</sub> which records the  
339 nucleation and growth of apparently late-stage extensional R-type shears within the bed of the OPIS.  
340 The same relationships were observed in all the thin sections and are interpreted as recording  
341 temporal changes in the style of deformation imposed on the till during its evolution.

## 342 **6.2. Clast microfabric development in tills**

343 The individual clast microfabrics within the OPIS till formed as a result of the passive rotation of  
344 coarse-silt to sand-grade particles into the plane of the developing foliation(s) (Fig. 16) reflecting the  
345 stress field imposed by the overriding ice (c.f. Hiemstra and Rijdsdijk, 2003; Phillips *et al.*, 2011). Once  
346 aligned, further rotation will cease and the clasts will maintain their preferred alignment unless  
347 there is a change in the orientation of this stress field within the evolving Reidel shears. As  
348 deformation continues, the microfabric domains defining the shear fabrics will propagate laterally as  
349 more grains become aligned. Further deformation within the microshears will be accommodated by  
350 either sliding of the grains past one another (Fig. 16a) and/or the partitioning of shear into the  
351 intervening finer grained matrix. If the matrix contains a significant modal proportion of clay  
352 minerals then this may lead to the formation of a unistrial plasmic fabric (van der Meer, 1993;  
353 Menzies, 2000; 2012; Hiemstra and Rijdsdijk, 2003) coplanar to the evolving clast microfabric.

354 The grain size of the sediment also influenced microfabric development with preferred clast  
355 alignments being most apparent within the finer grained (< 0.25 µm; 0.25 to 0.5 µm) components  
356 (matrix) of the till (Figs. 11 and 12). During deformation, it is suggested that the larger grains  
357 (cobbles, pebbles) were the first to stop rotating, becoming “locked” into position with subsequent

358 increments of deformation being partitioned into the still “active” matrix (c.f. Evans *et al.*, 2016;  
359 Evans, 2018). Consequently the matrix of the till continued to respond to shear long after the larger  
360 clasts have become immobile and therefore provide the most complete record of subglacial  
361 deformation. Results from several micromorphological studies (Phillips *et al.*, 2011, 2013, 2018)  
362 suggest that larger sand, granule to pebble sized clasts (where present) control the spacing of the  
363 developing microfabric domains, influencing the pattern of deformation partitioning within the till  
364 (Fig. 16). By the way of analogy, in metamorphic rocks the mica domains, defining the schistosity in  
365 amphibolite facies pelitic rocks (metamorphosed mudstones), are thought to nucleate upon the  
366 margins of ridged porphyroblasts (e.g. garnet), propagating laterally away from these relatively  
367 higher strain areas as deformation/fabric development continues (Bell, 1985; Bell and Rubenach,  
368 1983; Bell *et al.*, 1986; Vernon, 1989; Johnson, 1990). It is possible that a similar process also occurs  
369 in tills with the microfibrils nucleating upon the larger clasts due to the concentration of strain  
370 along the margins of these rigid grains. The evolving foliation then propagates away from this  
371 nucleation point into the adjacent matrix. The presence of larger rigid clasts will affect/distort (on a  
372 microscale) the stress and strain field imposed upon the till matrix (Fig. 16b) leading to the  
373 development of anastomosing microfabric, wrapping around these coarse sand to pebble-sized  
374 grains (Fig. 16c).

375         Evans *et al.* (2006) suggested that deformation (fabric development, folding, faulting) within  
376 the bed will only occur when the intergranular pore water pressure falls and a coherent “till-matrix  
377 framework” develops (also see Evans, 2018). Consequently, the nucleation and subsequent  
378 evolution of the clast microfibrils is likely to be controlled by the water content and packing of the  
379 till. However, increasing the packing of the constituent grains within the sediment will lead directly  
380 to an increase in its peak frictional strength and its resistance to deformation, which may be  
381 overcome by an increase in the magnitude of the imposed shear stress. Consequently, there is likely  
382 to be a critical range in intergranular porewater content/pressure and sediment packing for foliation  
383 development to occur within tills. For example, foliation development within a “dry” till comprising  
384 dense, closely packed grains will be limited due to the high percentage of inter granular contacts  
385 restricting grain rotation. In contrast, a shear stress applied to a water-saturated till is likely to  
386 induce dilation or even localised liquefaction which will not only inhibit fabric development, but also  
387 lead to the overprinting of earlier developed microstructures (Evans *et al.*, 2006; Phillips *et al.*, 2011;  
388 2013; 2018). The inherent spatial variation in the water content and packing of the till will lead to  
389 the small-scale partitioning of deformation and heterogeneous fabric development within the bed of  
390 the OPIS (Figs. 2 to 8).

### 391 **6.3. Evidence of intergranular fluid flow and dewatering of the bed**

392 The matrix of the till forming the bed of the OPIS contains clay-lined or filled voids, intergranular  
393 pore spaces and fractures (Fig. 9). The clay is petrographically similar to clay cutan within soils,  
394 suggesting that it was deposited by water flowing through the till so that the laminated nature of  
395 these fines records several phases of fluid flow. Although clay infiltration can occur during pedogenic  
396 processes, these features occur well below the base of the modern soil (see Fig. 1d) indicating that it  
397 was unrelated to recent pedogenic processes. Observations from contemporary (Alley *et al.*, 1986;  
398 Tulaczyk *et al.*, 1998) and palaeo (Ó Cofaigh *et al.*, 2007) ice stream beds indicate that the tills are  
399 typically highly porous and weak, with a water content close to the liquid limit. The clay within the  
400 OPIS till forms between 5 and 15 % of the matrix and is observed variably infilling intergranular pore  
401 spaces, consistent with the nature of tills described from modern ice stream beds (Alley *et al.*, 1986;  
402 Tulaczyk *et al.*, 1998; Ó Cofaigh *et al.*, 2007). Significant compaction of the OPIS till would have led to  
403 an increase in its packing, reducing in its porosity and permeability. Consequently the preferred  
404 interpretation is that fluid flow and clay infiltration probably occurred shortly after till deposition; a  
405 conclusion supported by the dark coloration of the clay resulting from its replacement by Fe/Mn  
406 during 'maturation' (van der Meer, 2007 pers. comm.; Phillips and Auton, 2008).

407 In the lower part of the OPIS till the earlier shear fabrics (S<sub>2</sub> to S<sub>4</sub>) are variably overprinted  
408 by S<sub>5</sub> (Fig. 15). The intergranular clays show very little, if any, evidence of disruption/fragmentation  
409 suggesting that they post-date any significant deformation and/or reorganisation of the structure  
410 and packing of the till during the imposition of S<sub>5</sub>. Consequently, it is suggested that S<sub>5</sub> pre-dated  
411 the intergranular fluid flow and clay infiltration, with the imposition of this late stage foliation  
412 probably accompanying the dewatering of the till driven by the ice overburden pressure; a  
413 conclusion supported by the relative increase in the intensity of this fabric downward through the till  
414 (Fig. 15). The micritic carbonate within the matrix of the till (red areas on Fig. 15) exhibits a close  
415 spatial relationship to S<sub>5</sub>, indicating that diagenesis may have accompanied the imposition of this  
416 fabric and recording the passage of a CaCO<sub>3</sub>-bearing fluid phase through the sediment. If the  
417 carbonate was pedogenic then it should increase upwards towards the base of the modern soil,  
418 possibly forming a calcified crust at the base of the soil profile. Several studies have examined the  
419 dissolution of carbonate in glacial environments (e.g. Fairchild *et al.*, 1993; 1994; Menzies and Brand,  
420 2007) with McGillen and Fairchild (2005) suggesting that this process may be facilitated by the  
421 crushing and comminution of carbonate grains within subglacial traction tills. Menzies and Brand  
422 (2007) argued that carbonate cementation of ice-contact sands and gravels exposed within core of a  
423 large drumlin in New York State occurred in response to a reduction in hydrostatic pressure and  
424 release of CO<sub>2</sub> from the meltwater escaping from beneath the Laurentide Ice Sheet. Consequently, it

425 is possible that carbonate diagenesis within the OPIS till may have occurred penecontemporaneous  
426 with subglacial deformation. As noted above, S5 clearly overprints the shear related fabrics within  
427 the till and therefore post-dated subglacial shearing. However, due to the potentially diachronous  
428 nature of deformation within the OPIS till, dewatering and imposition of S5 within the structural  
429 lower parts of the sequence is thought to have been initiated whilst subglacial shearing was  
430 continuing at a higher level within the bed (Fig. 15) (see below).

#### 431 **6.4. The nature of bed deformation beneath the OPIS: deformation** 432 **partitioning or incremental progressive simple shear during till accretion**

433 Spagnolo *et al.* (2016) argued that the till beneath the OPIS was being continuously accreted at the  
434 top of a shallow-deforming bed (cf. Tulaczyk, 1999; Iverson *et al.*, 1998; Fuller and Murray, 2000;  
435 Piotrowski *et al.*, 2001, 2004; Evans *et al.*, 2006; Cuffey and Paterson, 2010; Iverson, 2011; Evans,  
436 2018). However, the possibility that pervasive bed deformation may have occurred to greater depths  
437 beneath this ice stream needs to be considered. Quantitative data presented here provides evidence  
438 that fabric (S2, S3, S4, S5) strength varies (on a microscale) both laterally and vertically within the  
439 bed of the OPIS (Figs. 13 and 15). If this variation in fabric strength can be used as a proxy for the  
440 relative intensity of shear imposed on the till it may reflect either: (i) the partitioning of deformation  
441 within the subglacial shear zone formed beneath this ice stream, within the deforming bed and  
442 potentially encompassing the entire thickness of the till; or (ii) the variation in the magnitude of the  
443 shear being transmitted into the deforming bed during the progressive accretion of the OPIS till.

444 A number of studies have suggested that subglacial shear zones migrate through the bed  
445 due to spatial and temporal fluctuations in water content/pressure (Tulaczyk, 1999; Truffer *et al.*,  
446 2000; Evans *et al.*, 2006; Kjær *et al.*, 2006; Lee and Phillips, 2008) and/or the ability of these  
447 sediments to drain intergranular porewater (Piotrowski *et al.*, 2004). Deformation within the shear  
448 zone can either be 'pervasive' (homogeneous) and transmitted throughout the entire bed (van der  
449 Meer *et al.*, 2003; Menzies *et al.*, 2006), or heterogeneous where the bed comprises a 'mosaic' of  
450 actively deforming and stable (non-deforming) zones (Piotrowski and Kraus, 1997; Piotrowski *et al.*,  
451 2004; Lee and Phillips, 2008). Tills within the beds of ice streams are thought to be water-saturated  
452 and weak (Alley *et al.*, 1986; Tulaczyk *et al.*, 1998; Ó Cofaigh *et al.*, 2007), and therefore able to  
453 accommodate a significant proportion of the forward motion of the overriding ice. The consistency  
454 of the orientation and geometry of the shear fabrics (S2 to S4; Fig. 15) throughout the OPIS till,  
455 coupled with the very low preliminary estimates of shear strain (Fig. 14) may be used to support the  
456 presence of weak, water-saturated sediments beneath this palaeo-ice stream, facilitating the  
457 transmission of shear throughout its entire bed (Hart and Boulton, 1991; van der Wateren *et al.*,  
458 2000; van der Meer *et al.*, 2003; Menzies *et al.*, 2006). In detail, the relative intensity of the shear

459 fabrics vary both laterally and vertically (Figs. 8 and 13). However, there is no obvious macroscale  
460 evidence for the presence of a significant décollement surface (thrust) within the bed of the OPIS till  
461 (see Fig. 1d). So as a result, it cannot be argued that forward motion of the OPIS was accommodated  
462 by deformation at a deeper level within the sediment pile (Fig. 17a). Consequently, variations in the  
463 relative intensity of foliation development within the OPIS till is more likely to record the small-scale  
464 partitioning of deformation during subglacial shear.

465 The deforming bed model for glacier motion predicts an increase in cumulative strain  
466 upwards toward the ice-bed interface (Boulton, 1986; Boulton and Hindmarsh, 1987; Evans *et al.*,  
467 2006) with deformation being accommodated by a weak, water-saturated layer located immediately  
468 adjacent to, or at the ice-bed interface (Fig. 17b). The continuous deposition of a soft (weak),  
469 compositionally homogenous (well-mixed; see section 4.1) till layer at the top of the bed of the OPIS  
470 (Spagnolo *et al.*, 2016) means that over time deformation will have progressively shifted upwards as  
471 till accreted (Fig. 17b). The trenches in the study area reveal that the till sequence beneath the OPIS  
472 is at least 1.2 to 1.4 m thick, with Ground Penetrating Radar data indicating that the diamicton may  
473 be in the order of 2 to 3 m thick (Spagnolo *et al.*, 2016). The soft deforming layer responsible for till  
474 accretion at the top of this sequence is likely to have only been a few tens of centimetres thick  
475 (Menzies 1982; Alley *et al.*, 1986, 1987; Boulton, 1996; Hindmarsh, 1998; Larsen *et al.*, 2004, 2007;  
476 Evans *et al.*, 2006; Stokes *et al.*, 2013a) and localised in nature reflecting the spatial and temporal  
477 changes in bed conditions. Phillips *et al.* (2018) suggested that the term ‘transient mobile zone’ for  
478 this actively deforming layer in order to emphasize the spatial and temporal variations in subglacial  
479 deforming bed processes proposed by a number of researchers (e.g. Piotrowski and Kraus, 1997;  
480 Boyce and Eyles, 2000; van der Meer *et al.*, 2003; Larsen *et al.*, 2004, 2007; Piotrowski *et al.*, 2004,  
481 2006; Evans *et al.*, 2006; Meriano and Eyles, 2009; Evans, 2018). The focusing of deformation into  
482 this water-saturated mobile layer would have effectively switched off deformation at a deeper level  
483 within the bed. As a consequence of this progressive till accretion-deformation, the relative age of  
484 subglacial shearing beneath the OPIS would be diachronous, becoming progressively younger  
485 toward the top of the bed. The observed vertical variation in the relative intensity of the shear  
486 fabrics (S2 to S4; Figs. 13 and 15) may therefore be interpreted as recording changes in the  
487 magnitude of the cumulative strain being recorded by the till during this accretion-deformation  
488 process (Fig. 17c) (cf. Larsen *et al.*, 2004). The variation in fabric strength may reflect the degree of  
489 coupling between the ice and the underlying bed; the greater the fabric intensity the higher the  
490 degree of ice-bed coupling and transmission of shear into the bed.

491 Preliminary estimates of the shear strains involved are low ( $< 15$ ; Fig. 14b) indicating that the  
492 amount of shear being transmitted into the bed of the OPIS was relatively small (cf. Larsen *et al.*,  
493 2006; Narloch *et al.*, 2012; Phillips *et al.*, 2013). Consequently, fast flow of this ice stream would  
494 have been largely accommodated by either basal sliding and/or flow deformation within a weak,  
495 water-saturated layer located at the top of the accreting till sequence (Evans *et al.*, 2006; Phillips *et al.*,  
496 2013; Spagnolo *et al.*, 2016). Evidence of the latter is potentially provided by the rotational  
497 turbate structures (van der Meer, 1993, 1997; Menzies, 2000; Hiemstra and Rijdsdijk, 2003). These  
498 structures are truncated by the shear fabrics, indicating that rotational deformation occurred prior  
499 to foliation development within the OPIS till. Turbate structures form where larger clasts are able to  
500 rotate through angles of up to, or  $> 360^\circ$ , entraining the adjacent finer grained matrix (van der Meer,  
501 1993; Menzies, 2000; Lachniet *et al.*, 2001; Hiemstra and Rijdsdijk, 2003; Phillips, 2006; Lea and  
502 Palmer, 2014). This requires either very high shear strains or the lowering of the strength of the  
503 sediment enabling clast rotation at the much lower strains (Evans *et al.*, 2006), the latter being more  
504 likely due to the very low shear strain estimates obtained for the OPIS till. The presence of an active  
505 layer at the top of the bed would have markedly reduced or even prevented transmission of the  
506 shear into the underlying till. Furthermore, this layer is likely to have been highly mobile, facilitating  
507 the advection of well-mixed, far-travelled sediment down-ice and the continuous accretion of till at  
508 the top of a shallow deforming bed (Spagnolo *et al.*, 2016). Spatial and temporal fluctuations in the  
509 water content within this active layer will have affected the degree of ice-bed coupling, leading to  
510 the observed complex pattern of cumulative strain (Figs. 13, 14 and 15).

511 Microtextural evidence (sections 5 and 6) indicates dewatering, consolidation and  
512 cementation of the OPIS till. These processes could have led to an increase in the shear strength of  
513 the till potentially resulting in the increased “stabilisation” of the sediment within the cores of the  
514 MSGL as they grew beneath the OPIS. Till consolidation or the presence of a relatively hard/stiff core  
515 has been invoked in the initiation of some subglacial landforms (e.g. Menzies and Brand 2007;  
516 Menzies *et al.*, 2016), although this concept is challenged by the regular spatial distribution of these  
517 landforms (e.g. Spagnolo *et al.*, 2016). In specific case presented here, the results suggest that till  
518 consolidation may have been initiated at a lower level within the MSGL whilst till deformation and  
519 accretion continued above (see sections 6.3. and 6.4), thus providing no support for the idea that a  
520 stiffened core is required for MSGL initiation.

521 Although it is acknowledged that the detailed micromorphological/microstructural study  
522 presented here has largely focused upon a single site (site C), the results are applicable to the wider  
523 footprint of the OPIS as well as other contemporary and palaeo-ice streams. Deformation beneath

524 glaciers and ice sheets is widely viewed as being dominated by simple shear within a subglacial shear  
525 zone (e.g. van der Wateren *et al.*, 2000; Hart, 2007; Lee and Phillips, 2008; Benn and Evans 2010).  
526 The style of deformation identified within the bed of the OPIS is consistent with this assumption,  
527 with comparable microscale shear fabrics being recognised in subglacial traction tills (*sensu* Evans *et*  
528 *al.*, 2006) from other glaciated terrains (e.g. Germany - van der Wateren *et al.*, 2000; England  
529 (Norfolk) - Vaughan-Hirsh *et al.*, 2013; British Columbia, Canada - Neudorf *et al.*, 2013; Central  
530 Poland - Narloch *et al.*, 2012, 2013; Baltic Coast, northern Germany - Brumme, 2015; Gehrmann *et*  
531 *al.*, 2017; Scotland - Phillips *et al.*, 2011, 2018; Switzerland - Phillips *et al.*, 2013). In structural  
532 geology, Pumpelly's rule states that small deformation structures are a key to understanding the  
533 structural evolution of an area as they mimic the styles and orientations of a larger-scale structures  
534 of the same generation. Consequently, the geometry and the relationships displayed between the  
535 range of microscale (and macro-) structures found within subglacially deformed sediments can not  
536 only be used to establish the overall stress regime responsible for deformation (e.g. van der Wateren  
537 *et al.*, 2000; Vaughan-Hirsh *et al.*, 2013; Gehrmann *et al.*, 2017), but also aid in the reconstruction of  
538 the regional pattern of ice movement (e.g. Brumme, 2015). The bed of the OPIS across the  
539 Wielkopolska Lowland region is very gently undulating (Fig. 1c) (Przybylski, 2008; Spagnolo *et al.*,  
540 2016) and can, in general, be considered to be represented by an essentially subhorizontal ductile-  
541 brittle shear zone. The thick (c. 30 m) sequence of Quaternary sediments which blanket the area  
542 result in an absence of any major bedrock highs which would have imposed significant changes on  
543 the stress regime active within the bed of this palaeo-ice stream. Furthermore, gravel to cobble-  
544 sized clasts, which would have locally influenced (modified) microscale fabric development, are rare  
545 with the OPIS till. Consequently, the proposed model of progressive ductile shearing during till  
546 accretion is considered to be applicable across the bed of the Odra palaeo-ice stream with changes  
547 in microfabric geometry reflecting local changes in the physical properties of the sediment (e.g. grain  
548 size, porewater content) during fast ice flow.

## 549 **7. Conclusions**

550 The detailed microstructural study of the thick subglacial traction till formed within the bed of the  
551 Weichselian Odra palaeo-ice stream contributes to our understanding of the deformation processes  
552 occurring within its bed and it how it evolved over time.

- 553 • The massive, compositionally homogenous nature of the till indicates that the sediment  
554 being accreted to the bed was well-mixed and included far travelled material derived from  
555 Palaeozoic rocks of the Baltic Basin and crystalline basement of Scandinavian, consistent  
556 with the till being laid down by ice advancing from the NE.

- 557 • Deformation within the bed of the ice stream was dominated by foliation development (S1  
558 to S5) recording progressive ductile shearing within a subhorizontal subglacial shear zone.  
559 Cross-cutting relationships displayed by these shear fabrics (S2, S3 and S4) reflect temporal  
560 changes in the style of deformation being accommodated within the till during a single  
561 progressive shear event as the till accreted vertically. Kinematic indicators (S-C and ECC-type  
562 microfabrics) within the till record a consistent SE-directed (sinistral) sense of shear  
563 coincident with the regional pattern of ice flow across the Wielkopolska Lowland.
- 564 • The clast microfabrics reflect the stress field imposed by the overriding ice and form as a  
565 result of the passive rotation of detrital grains into the plane of the developing foliation(s).  
566 Larger clasts (pebbles, cobbles) become “locked” into position at an earlier stage within the  
567 deformation history and with subsequent shear being partitioned into the matrix of the till.  
568 These larger clasts then control the partitioning of deformation within the till and the  
569 spacing of the evolving microfabric domains.
- 570 • Evidence of intergranular pore water flowing through the bed of the OPIS is provided by the  
571 presence of clay-filled pore spaces within the till matrix having potentially occurred during or  
572 shortly after deposition. Further evidence of fluid flow through the bed is provided by the  
573 anastomosing, subvertical S5 fabric which formed in response to the dewatering and  
574 consolidation of the till in response to deposition from the deforming layer and final  
575 shutdown of the ice stream. Dewatering of the bed was accompanied by the growth of  
576 diagenetic micritic carbonate indicating that the escaping fluid contained dissolved CaCO<sub>3</sub>  
577 derived from the dissolution of detrital limestone.
- 578 • Bed deformation beneath the OPIS occurred in response to incremental progressive simple  
579 shear during till accretion. The relative age of deformation was diachronous (younger  
580 towards the top of the bed) as a thin deforming layer migrated progressively upwards in  
581 response to till accretion at the top of the ice stream bed. Focusing of deformation into this  
582 “active” layer or “transient mobile zone” effectively switched off deformation deeper within  
583 the bed. Variations in the relatively intensity of the microfabrics may record changes in the  
584 magnitude of the cumulative strain being imposed on the till during this accretion-  
585 deformation process and the degree of coupling between the ice and the underlying bed.  
586 Preliminary estimates of the shear strains involved are low, indicating that the amount of  
587 shear being transmitted into the bed of the OPIS was relatively small. This has implications  
588 for the mechanism responsible for the forward motion of this palaeo-ice stream.

## 589 **8. Acknowledgements**

590 The authors would like to thank various colleagues for numerous discussions on the nature of glacier  
591 bed deformation over the years, in particular Dave Evans, Jaap van der Meer, Jonathan Lee, Clive  
592 Auton, Jon Merritt, Chris Clark, Chris Stokes, Wojciech Wysota, Iza Szuman and Adriano Ribolini.  
593 Adrian Palmer (Royal Holloway University of London) is thanked for his expertise in making the thin  
594 sections. David J.A. Evans and one anonymous reviewer are thanked for their constructive reviews of  
595 this paper. ERP publishes with the permission of the Executive Director of the British Geological  
596 Survey, Natural Environmental Research Council.

## 597 **9. References**

- 598 Alley, R.B., 1989. Water pressure coupling of sliding and bed deformation: 1. Water system. *Journal*  
599 *of Glaciology* 35, 108-118.
- 600 Alley, R.B., Blankenship, D.D., Bentley, C.R., Rooney, S.T., 1986. Deformation of till beneath ice  
601 stream B, West Antarctica. *Nature* 322, 57-59.
- 602 Alley, R.B., Blankenship, D.D., Bentley, C.R., Rooney, S.T., 1987a. Till beneath ice stream B: 3. Till  
603 deformation: evidence and implications. *Journal of Geophysical Research* 92, 8921-8929.
- 604 Alley, R.B., Blankenship, D.D., Rooney, S.T., Bentley, C.R., 1987b. Till beneath ice stream B: 4. A  
605 coupled ice-till flow model. *Journal of Geophysical Research* 92, 8931-8940.
- 606 Bamber, J.L., Vaughan, D.G., Joughin, I., 2000. Widespread complex flow in the interior of the  
607 Antarctic Ice Sheet. *Science* 287, 1248–1250.
- 608 Baroni, C., Fasano, F., 2006. Micromorphological evidence of warm-based glacier deposition from  
609 the Ricker Hills Tillite (Victoria Land, Antarctica). *Quaternary Science Reviews*. 25, 976-992.
- 610 Bell, T.H., 1985. Deformation partitioning and porphyroblast rotation in metamorphic rocks: a radical  
611 reinterpretation. *Journal of Metamorphic Geology* 3, 109-118.
- 612 Bell, T.H., Rubenach, M.J., 1983. Sequential porphyroblast growth and crenulation cleavage  
613 development during progressive deformation. *Tectonophysics* 92, 171-194.
- 614 Bell, T.H., Rubenach, M.J., Fleming, P.D., 1986. Porphyroblast nucleation, growth and dissolution in  
615 regional metamorphic rocks as a function of deformation partitioning during foliation development.  
616 *Journal of Metamorphic Geology* 4, 37-67.

617 Bennett, M.R., 2003. Ice streams as the arteries of an ice sheet: their mechanics, stability and  
618 significance. *Earth Science Reviews* 61, 309–339.

619 Bingham, R.G., Vaughan, D.G., King, E.C., Davies, D., Cornford, S.L., Smith, A.M., Arthern, R.J.,  
620 Brisbourne, A.M., Rydt, J., Graham, A.G., Spagnolo, M., 2017. Diverse landscapes beneath Pine Island  
621 Glacier influence ice flow. *Nature communications* 8, 1618.

622 Boulton, G.S., 1996. Theory of glacial erosion, transport and deposition as a consequence of  
623 subglacial deformation. *Journal of Glaciology* 42, 43-62.

624 Boulton, G.S., Hindmarsh, R.C.A., 1987. Sediment deformation beneath glaciers: rheology and  
625 geological consequences. *Journal of Geophysical Research* 92, 9059-9082.

626 Boulton, G.S., Dobbie, K.E., Zatsepin, S., 2001. Sediment deformation beneath glaciers and its  
627 coupling to the subglacial hydraulic system. *Quaternary International* 86, 3-28.

628 Boyce, J.I., Eyles, N., 2000. Architectural element analysis applied to glacial deposits: internal  
629 geometry of a late Pleistocene till sheet, Ontario, Canada. *Bulletin of the Geological Society of*  
630 *America* 112, 98-118.

631 Brumme, J., 2015. Three-dimensional microfabric analyses of Pleistocene tills from the cliff section  
632 Dwasieden on Rügen (Baltic Sea coast): micromorphological evidence for subglacial polyphase  
633 deformation (PhD thesis). Ernst-Moritz-Arndt-Universität, Greifswald, 210pp.

634 Clarke, G.K.C., 1987. Subglacial till: a physical framework for its properties and processes. *Journal of*  
635 *Geophysical Research* 92, 9023-9036.

636 Cuffey, K.M., Paterson, W.S.B., 2010. *The Physics of Glaciers Fourth Edition*. Academic Press, Oxford.

637 Denis, M., Guiraud, M., Konaté, M., Buoncristiani, J.-F., 2010. Subglacial deformation and water-  
638 pressure cycles as a key for understanding ice stream dynamics: evidence from the Late Ordovician  
639 succession of the Djado Basin (Niger). *International Journal of Earth Science (Geol Rundsch)* 99,  
640 1399-1425.

641 Dyke, A.S., Morris, T.F., 1988. Drumlin fields, dispersal trains, and ice streams in Arctic Canada.  
642 *Canadian Geographer* 32, 86–90.

643 Evans, D.J., Phillips, E.R., Hiemstra, J.F., Auton, C.A., 2006. Subglacial till: formation, sedimentary  
644 characteristics and classification. *Earth Science Reviews* 78, 115-176.

645 Evans, D.J.A., Roberts, D.H., Evans, S.C., 2016. Multiple subglacial till deposition: A modern exemplar  
646 for Quaternary palaeoglaciology. *Quaternary Science Reviews* 145, 183-203.

647 Evans, D.J.A., 2018. *Till: A Glacial Process Sedimentology*. John Wiley and Sons Ltd, UK.

648 Fairchild, I.J., Bradby, L., Spiro, B., 1993. Carbonate diagenesis in ice. *Geology* 21, 901-904.

649 Fairchild, I.J., Bradby, L., Spiro, B., 1994. Reactive carbonate in glacial systems: a preliminary  
650 synthesis of its creation, dissolution and reincarnation. In Deymoux, M., Miller, J.M.G., Domack,  
651 E.W., Young, G.M. (Eds) *International Geological Correlation Project 260: Earth's Glacial record*.  
652 Cambridge University Press. 176-192.

653 Fuller, S., Murray, T., 2002. Sedimentological investigations in the forefield of an Icelandic surge-type  
654 glacier: implications for the surge mechanism. *Quaternary Science Reviews* 21, 1503-1520.

655 Gehrman, A., Hüneke, H., Meschede, M., Phillips, E. 2017. 3D microstructural architecture of  
656 deformed glacial sediments associated with large-scale glaciectonism, Jasmund Peninsula (NE  
657 Rügen), Germany. *Journal of Quaternary Science* 32, 213-230. DOI: 10.1002/jqs.2843.

658 Hart, J.K., Boulton, G.S., 1991. The interrelation of glaciectonic and glaciodepositional processes  
659 within the glacial environment. *Quaternary Science Reviews* 10, 335-350.

660 Hiemstra, J.F., van der Meer, J.J.M., 1997. Pore-water controlled grain fracturing as an indicator for  
661 subglacial shearing in tills. *Journal of Glaciology* 43, 446-454.

662 Hiemstra, J.F., Rijdsdijk, K.F. 2003. Observing artificially induced strain: implications for subglacial  
663 deformation. *Journal of Quaternary Science* 18, 373-383.

664 Hiemstra, J.F., Rijdsdijk, K.F., Evans, D.J.A., van der Meer, J.J.M., 2005. Integrated micro- and macro-  
665 scale analyses of Last Glacial maximum Irish Sea diamicts from Abermaw and Treath y Mwnt, Wales,  
666 UK. *Boreas* 34, 61-74.

667 Hindmarsh, R.C.A., 1998b. The stability of a viscous till sheet coupled with ice flow, considered at  
668 wavelengths less than the ice thickness. *Journal of Glaciology* 44, 285-292.

669 Hindmarsh, R.C., 2009. Consistent generation of ice-streams via thermo-viscous instabilities  
670 modulated by membrane stresses. *Geophysical Research Letters* 36(6).

671 Hodgson, D.A., 1994. Episodic ice streams and ice shelves during retreat of the northwestern most  
672 sector of the Late Wisconsinan Laurentide Ice Sheet over the central Canadian Arctic archipelago.  
673 *Boreas* 23, 14-28.

674 Humphrey, N., Kamb, B., Fahnestock, M., Engelhardt, H., 1993. Characteristics of the bed of the  
675 lower Columbia Glacier, Alaska. *Journal of Geophysical Research* 98, 837-846.

676 Iverson, N.R., 2011. Shear resistance and continuity of subglacial till: hydrology rules. *Journal of*  
677 *Glaciology* 56, 1104-1114.

678 Iverson, N.R., Hooyer, T.S., Baker, R.W. 1998. Ring-shear studies of till deformation: Coulomb-plastic  
679 behaviour and distributed strain in glacier beds. *Journal of Glaciology* 44, 634-642.

680 Johnson, S.E. 1990. Lack of porphyroblast rotation in the Otago schists. New Zealand: implications  
681 for crenulation cleavage development, folding and deformation partitioning. *Journal of*  
682 *Metamorphic Geology* 8, 13-30.

683 King, E.C., Hindmarsh, R.C., Stokes, C.R., 2009. Formation of mega-scale glacial lineations observed  
684 beneath a West Antarctic. *Nature Geoscience* 2, 585–588.

685 Kjaer, K.H., Houmark-Nielsen, M., Richardt, N., 2003. Ice-flow patterns and dispersal of erratics at  
686 the southwestern margin of the last Scandinavian Ice Sheet: signature of palaeo-ice streams. *Boreas*  
687 32, 130-148.

688 Kjær, K.H., Larsen, E., van der Meer, J.J.M., Ingólfsson, Ó., Krüger, J., Benediktsson, I.Ó., Knudsen,  
689 C.G., Schomacker, A., 2006. Subglacial decoupling at the sediment/bedrock interface: a new  
690 mechanism for rapid flowing ice. *Quaternary Science Reviews* 25, 2704-2712.

691 Khatwa, A., Tulaczyk, S., 2001. Microstructural interpretations of modern and Pleistocene  
692 subglacially deformed sediments: the relative role of parent material and subglacial processes.  
693 *Journal of Quaternary Science* 16, 507-517.

694 Kozarski, S., 1988. Time and dynamics of the last Scandinavian ice-sheet retreat from northwestern  
695 Poland. *Geographica Polonica* 55, 91-101.

696 Kyrke-Smith, T.M., Katz, R.F., Fowler, A.C., 2015. Subglacial hydrology as a control on emergence,  
697 scale, and spacing of ice streams. *Journal of Geophysical Research: Earth Surface* 120, 1501-1514.

698 Lachniet, M.S., Larson, G.J., Lawson, D.E., Evenson, E.B., Alley, R.B., 2001. Microstructures of  
699 sediment flow deposits and subglacial sediments: a comparison. *Boreas* 30, 254-262.

700 Larsen, N. K., Piotrowski, J. A. and Kronborg, C. 2004. A multiproxy study of a basal till: a time-  
701 transgressive accretion and deformation hypothesis. *Journal of Quaternary Science* 19, 9–21

702 Larsen, N.K., Piotrowski, J.A., Christiansen, F., 2006. Microstructures and micro-shears as proxy for  
703 strain in subglacial diamicts: implications for basal till formation. *Geology*. 34, 889-892.

704 Larsen, N.K., Piotrowski, J.A., Menzies, J., 2007. Microstructural evidence of low-strain, time  
705 transgressive subglacial deformation. *Journal of Quaternary Science*. 22, 593-608.

706 Lea, J.M., Palmer, A., 2014. Quantification of turbate microstructures through a subglacial till:  
707 dimensions and characteristics. *Boreas*. 10.1111/bor.12073. ISSN 0300-9483.

708 Lee, J.R., Phillips, E.R., 2008. Progressive soft sediment deformation within a subglacial shear zone –  
709 a hybrid mosaic-pervasive deformation model for Middle Pleistocene glaciotectionised sediments  
710 from eastern England. *Quaternary Science Reviews* 27, 1350-1362.

711 Leighton, I.D., Hiemstra, J.F., Weidman, C.T., 2012. Recognition of micro-scale deformation  
712 structures in glacial sediments – pattern perception, observer bias and the influence of experience.  
713 *Boreas*, 10.1111/j.1502-3885.2011.00246.x. ISSN 0300-9483.

714 McGillen, M.R., Fairchild, I.J., 2005. An experimental study of incongruent dissolution of CaCO<sub>3</sub> under  
715 analogue glacial conditions. *Journal of Glaciology* 51, 383-390.

716 Marks, L., 2012. Timing of the Late Vistulian (Weichselian) glacial phases in Poland. *Quaternary  
717 Science Reviews* 44, 81-88.

718 Margold, M., Stokes, C.R., Clark, C.D. 2015. Ice Streams in the Laurentide Ice Sheet: Identification,  
719 characteristics and comparison to modern ice sheets. *Earth-Science Reviews* 143, 117-146.

720 Menzies, J., 1982. Till hummock (proto-drumlin) at the ice glacier bed interface. In: Davidson-Arnott,  
721 R., Nickling, W., Fahey, B.D. (Eds.), *Research in Glacial, Glacio-Fluvial and Glacio-Lacustrine Systems*.  
722 *Proceedings of the 6th Guelph Symposium on Geomorphology*, 33-47.

723 Menzies, J., 2000. Micromorphological analyses of microfabrics and microstructures indicative of  
724 deformation processes in glacial sediments. In: A.J. Maltman, B. Hubbard, M.J. Hambrey (eds.).  
725 *Deformation of glacial materials*. Geological Society of London, Special Publication 176, 245-257.

726 Menzies, J., 2012. Strain pathways, till internal architecture and microstructures – perspectives on a  
727 general kinematic model – a ‘blueprint’ for till development. *Quaternary Science Reviews* 50, 105-  
728 124.

729 Menzies, J., Maltman, A.J., 1992. Microstructures in diamictons - evidence of subglacial bed  
730 conditions. *Geomorphology* 6, 27-40.

731 Menzies, J., Brand, U., 2007. The internal sediment architecture of a drumlin, Port Byron, New York  
732 State, USA. *Quaternary Science Reviews* 26, 322-335.

733 Menzies, J., Zaniewski, K., Dreger, D., 1997. Evidence from microstructures of deformable bed  
734 conditions within drumlins, Chimney Bluffs, New York State. *Sedimentary Geology* 111, 161-175.

735 Menzies, J., van der Meer, J.J.M., Rose, J., 2006. Till – a glacial "tectomict", a microscopic  
736 examination of a till's internal architecture. *Geomorphology* 75, 172-200.

737 Menzies, J., Hess, D.P., Rice, J.M., Wagner, K.G., Ravier, E., 2016. A case study in the New York  
738 Drumlin Field, an investigation using microsedimentology, resulting in the refinement of a theory of  
739 drumlin formation. *Sedimentary Geology* 338, 84-96.

740 Meriano, M., Eyles, N., 2009. Quantitative assessment of the hydraulic role of subglaciofluvial  
741 interbeds in promoting deposition of deformation till (Northern Till, Ontario). *Quaternary Science*  
742 *Reviews* 28, 608-620.

743 Narloch, W., Piotrowski, J.A., Wysota, W., Larsen, N.K., Menzies, J., 2012. The signature of strain  
744 magnitude in tills associated with the Vistula Ice Stream of the Scandinavian Ice Sheet, central  
745 Poland. *Quaternary Science Reviews* 57, 105-120.

746 Narloch, W., Wysota, W., Piotrowski, J.A., 2013. Sedimentological record of subglacial conditions and  
747 ice sheet dynamics of the Vistula Ice Stream (north-central Poland) during the Last Glaciation.  
748 *Sedimentary Geology* 293, 30-44.

749 Neudorf, C. M., Brennand, T. A., Lian, O. B., 2013. Till-forming processes beneath parts of the  
750 Cordilleran Ice Sheet, British Columbia, Canada: macroscale and microscale evidence and a new  
751 statistical technique for analysing microstructure data. *Boreas*, 10.1111/bor.12009. ISSN 0300-9483.

752 O' Cofaigh, C., Evans, J., Dowdeswell, J.A., Larter, R. 2007. Till characteristics, genesis and transport  
753 beneath Antarctic paleo-ice streams. *Journal of Geophysical Research* 112, F03006.

754 Palmer, A.P. 2005. The micromorphological description, interpretation and palaeoenvironmental  
755 significance of lacustrine clastic laminated sediments. Unpublished PhD thesis, University of London.

756 Passchier, C.W., Trouw, R.A.J., 1996. *Microtectonics*. Springer.

757 Patterson, C.J., 1997. Southern Laurentide ice lobes were created by ice streams: Des Moines Lobe in  
758 Minnesota, USA. *Sedimentary Geology* 111, 249-261.

759 Patterson, C.J., 1998. Laurentide glacial landscapes: the role of ice streams. *Geology* 26, 643-646.

760 Phillips, E.R., 2006. Micromorphology of a debris flow deposit: evidence of basal shearing,  
761 hydrofracturing, liquefaction and rotational deformation during emplacement. *Quaternary Science*  
762 *Reviews* 25, 720-738.

763 Phillips, E., Merritt, J.W., Auton, C.A., Gollidge, N.R., 2007. Microstructures developed in subglacially  
764 and proglacially deformed sediments: faults, folds and fabrics, and the influence of water on the  
765 style of deformation. *Quaternary Science Reviews* 26, 1499-1528.

766 Phillips, E.R., Auton, C.A., 2000. Micromorphological evidence for polyphase deformation of  
767 glaciolacustrine sediments from Strathspey, Scotland. In: Maltman, A.J., Hubbard, B., Hambrey, M.J.  
768 (Eds). *Deformation of glacial materials*. The Geological Society of London, Special Publication 176,  
769 279-291.

770 Phillips, E.R., Auton, C.A., 2008. Microtextural analysis of a glacially 'deformed' bedrock: implications  
771 for inheritance of preferred clast orientations in diamictons. *Journal of Quaternary Science* 23, 229-  
772 240.

773 Phillips, E., Merritt, J., 2008. Evidence for multiphase water-escape during rafting of shelly marine  
774 sediments at Clava, Inverness-shire, NE Scotland. *Quaternary Science Reviews* 27, 988–1011.

775 Phillips, E.R., van der Meer, J.J.M., Ferguson, A., 2011. A new 'microstructural mapping'  
776 methodology for the identification and analysis of microfabrics within glacial sediments. *Quaternary*  
777 *Science Reviews* 30, 2570-2596.

778 Phillips, E., Everest, J., Reeves, H., 2013. Micromorphological evidence for subglacial multiphase  
779 sedimentation and deformation during overpressurized fluid flow associated with hydrofracturing.  
780 *Boreas* 42, 395–427.

781 Phillips, E.R., Lipka, E., van der Meer, J.J.M. 2013. Micromorphological evidence of liquefaction and  
782 sediment deposition during basal sliding of glaciers. *Quaternary Science Reviews* 81, 114-137.

783 Phillips, E., Evans, D.J.A., van der Meer, J.J.M., Lee, J.R. 2018. Microscale evidence of liquefaction and  
784 its potential triggers during soft-bed deformation within subglacial traction tills. *Quaternary Science*  
785 *Reviews* 181, 123-143.

786 Piotrowski, J.A., 1994a. In Warren, W.P., Croot, D. (eds) *Formation and Deformation of Glacial*  
787 *Deposits*, Balkema, Rotterdam. 3-8.

788 Piotrowski, J.A., 1994b. Tunnel-valley formation in northwest Germany—geology, mechanisms of  
789 formation and subglacial bed conditions for the Bornhöved Tunnel Valley. *Sedimentary Geology* 89,  
790 107-141.

791 Piotrowski, J.A., Kraus, A.M., 1997. Response of sediment to ice sheet loading in northwestern  
792 Germany: effective stresses and glacier bed stability. *Journal of Glaciology* 43, 495-502.

793 Piotrowski, J.A., Tulaczyk, S., 1999. Subglacial conditions under the last ice sheet in northwest  
794 Germany: ice-bed separation and enhanced basal sliding? *Quaternary Science Reviews* 18, 737-751.

795 Piotrowski, J.A., Mickelson, D.M., Tulaczyk, S., Krzyszowski, D., Junge, F. 2001. Were subglacial  
796 deforming beds beneath past ice sheets really widespread? *Quaternary International* 86, 139-150.

797 Piotrowski, J.A., Larsen, N.K., Junge, F., 2004. Soft subglacial beds: a mosaic of deforming and stable  
798 spots. *Quaternary Science Reviews* 23, 993-1000.

799 Piotrowski, J.A., Larsen, N.K., Menzies, J., Wysota, W., 2006. Formation of subglacial till under  
800 transient bed conditions: deposition, deformation, and basal decoupling under a Weichselian ice  
801 sheet lobe, central Poland. *Sedimentology* 53, 83-106.

802 Przybylski, B., 2008. Geomorphic traces of a Weichselian ice stream in the Wielkopolska Lowland,  
803 western Poland. *Boreas* 37, 286-296.

804 Spagnolo, M., Clark, C.D., Ely, J.C., Stokes, C.R., Anderson, J.B., Andreassen, K., Graham, A.G.C., King,  
805 E.C., 2014. Size, shape and spatial arrangement of mega-scale glacial lineations. *Earth Surface*  
806 *Processes and Landforms* 39, 1432-1448.

807 Spagnolo, M., Phillips, E., Piotrowski, J.A., Rea, B.R., Clark, C.D., Stokes, C.R., Carr, S.J., Ely, J.C.,  
808 Adriano Ribolini, A., Wysota, W., Izabela Szuman, I., 2016. Ice stream motion facilitated by a shallow-  
809 deforming and accreting bed. *Nature Communications* DOI: 10.1038/ncomms10723.

810 Spagnolo, M., Bartholomaeus, T.C., Clark, C.D., Stokes, C.R., Atkinson, N., Dowdeswell, J.A., Ely, J.C.,  
811 Graham, A., Hogan, K.A., King, Larter, R.D., E., Livingstone, S.J., Pritchard, H.D., 2017. The periodic  
812 topography of ice stream beds: insights from the Fourier spectra of mega-scale glacial lineations.  
813 *Journal of Geophysical Research: Earth Surface* 122, 1355-1373.

814 Stokes, C.R., Clark, C.D., 1999. Geomorphological criteria for identifying Pleistocene ice streams.  
815 *Annals of Glaciology* 28, 67-75.

816 Stokes, C.R., Clark, C.D., 2001. Palaeo-ice streams. *Quaternary Science Reviews* 20, 1437-1457.

817 Stokes, C.R., Clark, C.D., 2002. Are long subglacial bedforms indicative of fast ice flow? *Boreas* 31,  
818 239-249.

819 Stokes, C.R., Clark, C.D., 2003. The Dubawnt Lake palaeo-ice stream: evidence for dynamic ice sheet  
820 behaviour on the Canadian Shield and insights regarding the controls on ice stream location and  
821 vigour. *Boreas* 32, 263-279.

822 Stokes, C.R., Fowler, A.C., Clark, C.D., Hindmarsh, R.C.A., Spagnolo, M., 2013a. The instability theory  
823 of drumlin formation and its explanation of their varied composition and internal structure.  
824 *Quaternary Science Reviews* 62, 77–96.

825 Stokes, C.R., Spagnolo, M., Clark, C.D., Ó Cofaigh, C., Lian, O.B., Dunstone, R.B., 2013b. Formation of  
826 mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 1. Size, shape and spacing from a  
827 large remote sensing dataset. *Quaternary Science Reviews* 77, 190–209.

828 Thomason, J.F., Iverson, N.R., 2006 Microfabric and micro-shear evolution in deformed till.  
829 *Quaternary Science Reviews* 25, 1027-1038.

830 Truffer, M., Harrison, W.D., Echelmeyer, K.A., 2000. Glacier motion dominated by processes deep in  
831 underlying till. *Journal of Glaciology* 46, 213e221. <https://doi.org/10.3189/172756500781832909>.

832 Tulaczyk, S., 1999 Ice sliding over weak, fine-grained tills: dependence of ice-till interactions on till  
833 granulometry. *Special Papers Geological Society of America* 337, 159-177.

834 Tulaczyk, S., Kamb, B., Scherer, R.P., Engelhardt, H.F., 1998. Sedimentary processes at the base of a  
835 West Antarctic ice stream: constraints from textural and compositional properties of subglacial  
836 debris. *Journal of Sedimentary Research* 68, 487-496.

837 van der Meer, J.J.M., 1987. Micromorphology of glacial sediments as a tool in distinguishing genetic  
838 varieties of till. *Geological Survey of Finland Special Paper* 3, 77-89.

839 van der Meer, J.J.M., 1993. Microscopic evidence of subglacial deformation. *Quaternary Science*  
840 *Reviews* 12, 553-587.

841 van der Meer, J.J.M., 1997. Particle and aggregate mobility in till: microscopic evidence of subglacial  
842 processes. *Quaternary Science Reviews* 16, 827-831.

843 van der Meer, J.J.M., Menzies, J., Rose, J., 2003. Subglacial till, the deformable glacier bed.  
844 *Quaternary Science Reviews* 22, 1659-1685.

845 van der Meer, J.J.M., Kjær, K.H., Krüger, J., Rabassa, J., Kilfeather, A.A., 2009. Under pressure: clastic  
846 dykes in glacial settings. *Quaternary Science Reviews* 28, 708-720.

847 van der Wateren, F.M., Kluiving, S.J., Bartek, L.R., 2000. Kinematic indicators of subglacial shearing.  
848 In: A.J. Maltman, B. Hubbard, M.J. Hambrey (eds.) *Deformation of glacial materials*. Geological  
849 Society of London, Special Publication. 176, 259-278.

850 Vaughan-Hirsch, D.P., Phillips, E., Lee, J.R., Hart, J.K., 2013. Micromorphological analysis of poly-  
851 phase deformation associated with the transport and emplacement of glaciotectionic rafts at West  
852 Runton, north Norfolk, UK. *Boreas* 42, 376–394.

853 Vernon, R.H., 1989. Porphyroblast-matrix microstructural relationships: recent approaches and  
854 problems. In Daly et al. (Eds), *Evolution of metamorphic belts*. Geological Society of London, Special  
855 Publication 43, 83-102.

856

## 857 **10. Figures**

858 **Fig. 1.** (a) and (b) Maps showing the location of the study area in western Poland; (c) Digital  
859 Elevation Model (DEM) of the Środa Wielkopolska area showing the well-developed NW-SE-trending  
860 megascale glacial lineations and locations of the trenches excavated into these subglacial landforms.  
861 Also shown is the SE-directed regional ice flow across the area; and (d) An example of a trench dug  
862 into the Quaternary sediments forming the landforms showing the position of the samples collected  
863 for thin sectioning. Note that the samples were collected from below the base of the soil layer.

864 **Fig. 2.** Microstructural map and high resolution scan of thin section C1M. The orientation of the long  
865 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose  
866 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
867 calculated for each area (see text for details).

868 **Fig. 3.** Microstructural map and high resolution scan of thin section C2M. The orientation of the long  
869 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose  
870 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
871 calculated for each area (see text for details).

872 **Fig. 4.** Microstructural map and high resolution scan of thin section C3M. The orientation of the long  
873 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose

874 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
875 calculated for each area (see text for details).

876 **Fig. 5.** Microstructural map and high resolution scan of thin section C4M. The orientation of the long  
877 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose  
878 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
879 calculated for each area (see text for details).

880 **Fig. 6.** Microstructural map and high resolution scan of thin section C5M. The orientation of the long  
881 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose  
882 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
883 calculated for each area (see text for details).

884 **Fig. 7.** Microstructural map and high resolution scan of thin section C6M. The orientation of the long  
885 axes of sand to granule sized clasts included within the diamicton are shown on a series of rose  
886 diagrams. The thin section has been subdivided into 16 subareas and the E1 and E2 eigenvalues  
887 calculated for each area (see text for details).

888 **Fig. 8.** Microstructural maps and automated clast density maps of the main clast microfabrics (S2, S3,  
889 S4, S5) identified within samples X1M, Y1M and Z1M.

890 **Fig. 9.** Photomicrographs showing the fine-grained, dusty looking carbonate which locally replaces  
891 the matrix to the diamicton (a to c) and clay lined and filled pore spaces (d to f).

892 **Fig. 10. (a)** Diagram showing the relationships between the different sets of Riedel shears developed  
893 within the diamicton in response to deformation imposed by the overriding ice stream; and **(b)**  
894 Example of a detailed microstructural map of sample C1M. The coloured polygons represent the  
895 different generations of clast microfabrics, which define the Riedel shears, subhorizontal shear fabric  
896 and up-ice dipping foliation.

897 **Fig. 11.** Graphs showing the effects of grain size on clast microfabric development within the  
898 diamicton at site C. The long axis data for each thin section (C1M to C6M) are divided into three sets:  
899 (i) grains < 0.25 mm in length; (ii) grains between 0.25 to 0.5 mm in size; and (iii) grains over > 0.5  
900 mm in length. The number of grains in each of the three classes is plotted against the orientation of  
901 their long axis (0° represents horizontal). The large “spike” in the data set for the finest grains and  
902 total clasts at 0° results from the unavoidable “snapping” to the horizontal of short long axes dipping  
903 at very low angles (-2° to +2°) during digitisation using CorelDraw. On Figure 12 the data are plotted  
904 on a series of rose diagrams showing the dip of the long axes within the 2D plane of the thin section.

905 **Fig. 12.** Rose diagrams showing the variation in dip of the long axes of coarse silt to sand sized clasts  
906 within the 2D plane of the thin sections C1M to C6M. The data are divided into three sets: (i) grains <  
907 0.25mm in length; (ii) grains between 0.25 to 0.5 mm in size; and (iii) grains over > 0.5 mm in length.

908 **Fig. 13.** Variation in E1 eigenvalues calculated for the thin sections (C1M to C6M) showing the  
909 variation in relative fabric strength both with an individual thin section and vertically through the  
910 diamicton at site C. Red colours represent areas of the thin sections with higher E1 values (0.65-  
911 0.68) and purple low E1 values (0.53-0.51).

912 **Fig. 14. (a)** Plot showing the variation in average E1 eigenvalue calculated for each thin section (C1M  
913 to C6M) with respect to depth within the diamicton sequence; and **(b)** Plot of E1 eigenvalue against  
914 shear strain. The shear strain curves for the Batestown (gravelly) and Douglas (sand-rich) tills are  
915 taken from Thomason and Iverson (2006) and are used to obtain an estimate of the range of shear  
916 strains encountered by the diamicton exposed at site C.

917 **Fig. 15.** Diagram showing the variation in clast microfabric development at site C.

918 **Fig. 16. (a)** Cartoon showing the passive rotation of elongate clasts into the plane of the developing  
919 microfabric. Further deformation of clasts aligned within this fabric is thought to occur in response  
920 to grain sliding; **(b)** Strain field diagram modified from Bell *et al.* (1986) used here to show the  
921 proposed geometry of the strain field formed in response to the presence of large immobile clasts  
922 within a weaker deforming matrix; and **(c)** Schematic diagram showing the development of  
923 anastomosing clast microfabrics within a till defined by the preferred shape alignment of finer  
924 grained clasts. The spacing of the microfabric domains is controlled by the grain size and spacing of  
925 the larger sand to pebble sized clasts in response to deformation partitioning within the till.

926 **Fig. 17.** Schematic profiles through the bed of a glacier and the resulting idealised cumulative strain  
927 curves: **(a)** Pervasive deformation throughout the subglacial shear zone with the amount of strain  
928 increasing upwards towards the ice-bed interface; **(b)** Deformation partitioning within the subglacial  
929 shear zone with localised detachments forming at deeper levels within the bed; and **(c)** Deformation  
930 confined to the “active” layer located at the top of the bed with shearing migrating upwards keeping  
931 pace with till accretion (cf. Larsen *et al.*, 2004).

## 11. Tables

**Table 1.** Detrital clast assemblage identified within the silty sand subglacial traction till exposed at Ste C.




Sample Number	Detrital Components	
	<i>major</i>	<i>minor to accessory</i>
C1M	monocrystalline quartz, feldspar (plagioclase, K-feldspar)	polycrystalline quartz, opaque minerals, altered volcanic rocks, clinopyroxene, microcline, mudstone, sericitised granitic rock, amphibole, glauconitic material, micrographic intergrowth, ?alkali granite, garnet
C2M	monocrystalline quartz, feldspar (plagioclase, K-feldspar)	cryptocrystalline quartz (chert), opaque minerals, pyroxene, siltstone, mudstone, microcline, altered igneous rock, glauconitic material, micrographic intergrowth, amphibole, feldspar-chlorite rock, biotite-schistose metamorphic rock, fine sandstone (hematitic cement)
C3M	monocrystalline quartz	polycrystalline quartz, biotite, metamorphic rock, indurated quartz-arenite, echinoderm fragments, plagioclase, carbonate mineral(s), micritic limestone, brachiopod fragments, K-feldspar, glauconitic material, amphibole, opaque minerals, cryptocrystalline quartz (chert), very fine-grained micaceous material, epidote, siltstone/very fine-grained sandstone, foraminifera, garnet, ?zircon, muscovite
C4M	monocrystalline quartz, feldspar (plagioclase, K-feldspar)	micrographic intergrowth, carbonate minerals, mudstone, microcline, glauconitic material, amphibole, indurated siltstone, bioclastic limestone, glauconite-bearing micritic limestone, biotite, ?alkali granitic rock, micritic limestone, hematized siltstone, opaque minerals, altered granitic rock, ?garnet, brachiopod fragments, echinoderm fragments, quartz-epidote-rock, amphibolite
C5M	monocrystalline quartz	polycrystalline quartz, siltstone, amphibole, opaque minerals, carbonate minerals, micritic limestone, glauconitic material, microcline, plagioclase, altered granitic rock, devitrified igneous rock/felsite, zircon, muscovite, garnet, bioclastic limestone, glauconitic sandstone, biotite metamorphic rock
C6M	monocrystalline quartz	plagioclase, K-feldspar, carbonate minerals, bioclastic limestone, glauconitic material, micritic bioclastic limestone, muscovite-granite, amphibole, opaque minerals, biotite-granite, laminated siltstone, amphibolite, altered igneous rock, microcline, garnet, calcareous siltstone, very fine-grained sandstone/siltstone, zircon, epidote, tourmaline



**Table 2.** Average E1 and E2 eigenvalues calculated for clast microfabrics developed within subglacial traction till samples C1M to C6M.

Depth below surface (cm)	Sample	Eigenvalues								Median	Average
		E1	E1	E1	E1	E1	E1	E1	E1		
21	C1M a-h	0.555	0.582	0.578	0.580	0.582	0.573	0.570	0.549	0.575	0.571
24	C1M i-p	0.604	0.575	0.591	0.601	0.561	0.605	0.576	0.571	0.584	0.586
46	C2M a-h	0.583	0.588	0.589	0.631	0.600	0.565	0.606	0.562	0.588	0.590
49	C2M i-p	0.579	0.594	0.588	0.562	0.582	0.584	0.597	0.580	0.583	0.583
71	C3M a-h	0.649	0.642	0.618	0.635	0.654	0.658	0.651	0.629	0.646	0.642
74	C3M i-p	0.638	0.640	0.645	0.626	0.596	0.617	0.600	0.627	0.627	0.624
96	C4M a-h	0.559	0.540	0.523	0.550	0.564	0.561	0.553	0.586	0.556	0.555
99	C4M i-p	0.545	0.552	0.530	0.517	0.547	0.531	0.542	0.560	0.544	0.541
121	C5M a-h	0.521	0.566	0.541	0.528	0.557	0.554	0.557	0.571	0.555	0.549
124	C5M i-p	0.526	0.552	0.562	0.571	0.559	0.556	0.573	0.613	0.561	0.564
146	C6M a-h	0.573	0.556	0.534	0.571	0.568	0.551	0.568	0.582	0.568	0.563
154	C6M i-p	0.556	0.566	0.561	0.569	0.542	0.535	0.530	0.570	0.559	0.554
		E2	E2	E2	E2	E2	E2	E2	E2	Median	Average
21	C1M a-h	0.444	0.417	0.421	0.419	0.417	0.426	0.429	0.450	0.424	0.428
24	C1M i-p	0.395	0.424	0.408	0.398	0.438	0.394	0.423	0.428	0.415	0.413
46	C2M a-h	0.416	0.411	0.410	0.368	0.399	0.434	0.393	0.437	0.411	0.409
49	C2M i-p	0.420	0.405	0.411	0.437	0.417	0.415	0.402	0.419	0.416	0.416



71	C3M a-h	0.350	0.357	0.381	0.364	0.345	0.341	0.348	0.370	0.353	0.357
74	C3M i-p	0.361	0.359	0.354	0.373	0.403	0.382	0.399	0.372	0.372	0.375
96	C4M a-h	0.440	0.459	0.476	0.449	0.435	0.438	0.446	0.413	0.443	0.444
99	C4M i-p	0.454	0.447	0.469	0.482	0.452	0.468	0.457	0.439	0.455	0.458
121	C5M a-h	0.478	0.43	0.458	0.471	0.442	0.445	0.442	0.428	0.444	0.450
124	C5M i-p	0.473	0.447	0.437	0.428	0.440	0.443	0.426	0.386	0.438	0.435
146	C6M a-h	0.426	0.443	0.465	0.428	0.431	0.448	0.431	0.417	0.431	0.436
154	C6M a-p	0.443	0.433	0.438	0.430	0.457	0.464	0.469	0.429	0.440	0.445






- S1 clast microfabric
- S2 clast microfabric
- S3 clast microfabric
- S4 clast microfabric
- subhorizontal Y-type Riedel shear
- moderately down-ice dipping R-type shears
- steeply down-ice dipping R'-type shears


- sense of shear
- orientation of clast long axis
- alignment of clast long axes
- clay lined fractures, voids and pore spaces




Area B: N = 1179




Area C: N = 1191




Area D: N = 1133




Area E: N = 1154




Area F: N = 1083



Area G: N = 1242




Area A: N = 1192




**Sample C1M**


orientation 122 degrees




Area H: N = 967




Area P: N = 745




Area J: N = 1099




Area K: N = 1244




Area L: N = 1064




Area M: N = 1123



Area N: N = 1003





















Area O: N = 1021



- S1 clast microfabric
- S2 clast microfabric
- S3 clast microfabric
- S4 clast microfabric
- S5 clast microfabric
- subhorizontal Y-type Riedel shear
- moderately down-ice dipping R-type shears
- steeply down-ice dipping R'-type shears
- subvertical anastomosing dewatering foliation

- $\leftarrow$  sense of shear
- $\leftarrow$  orientation of clast long axis
- $\leftarrow$  alignment of clast long axes
- $\leftarrow$  clay lined fractures, voids and pore spaces





- S1 clast microfabric
- S2 clast microfabric
- S3 clast microfabric
- S4 clast microfabric
- S5 clast microfabric
- subhorizontal Y-type Riedel shear
- moderately down-ice dipping R-type shears
- steeply down-ice dipping R'-type shears
- subvertical anastomosing dewatering foliation
- micritic carbonate cement



















**Sample C3M** orientation 122 degrees


subhorizontal Y-type Riedel shears

moderately inclined, up-ice dipping foliation (S3) forming dominant fabric

subhorizontal foliation (S2) defining Y-type Riedel shears

- $\leftarrow$  sense of shear
- $\leftarrow$  orientation of clast long axis
- $\leftarrow$  alignment of clast long axes
- $\leftarrow$  clay lined fractures, voids and pore spaces






- S1 clast microfabric
- S2 clast microfabric
- S3 clast microfabric
- S4 clast microfabric
- S5 clast microfabric
- subhorizontal Y-type Riedel shear
- moderately down-ice dipping R-type shears
- steeply down-ice dipping R'-type shears
- subvertical anastomosing dewatering foliation
- micritic carbonate cement

**Sample C4M**  
orientation 122 degrees

subhorizontal foliation (S2) defining Y-type Riedel shears

moderately inclined, up-ice dipping foliation (S3) forming dominant fabric



sinistral (down-ice) shear sense defined by asymmetrical microfabric

early foliation (S1) deformed by small-scale crenulations







subvertical anastomosing foliation (S5)

**Sample C4M**

anastomosing zones of subvertical clast microfabric (S5)

moderately inclined down-ice dipping foliation (S4) defining extensional Riedel shears

down-ice dipping R-type Riedel shears



Area B: N = 1335


Area C: N = 1218

Area D: N = 1165



Area E: N = 1427

Area F: N = 1204


Area G: N = 1409




Area A: N = 1417




Area H: N = 1004




Area P: N = 1150




Area I: N = 1669




Area K: N = 1348




Area L: N = 1295




Area M: N = 1500




Area N: N = 1373



Area O: N = 1373




Area J: N = 1508



**Sample C5M**  
orientation 122 degrees

subhorizontal foliation (S2) defining Y-type Riedel shears  
moderately inclined, up-ice dipping foliation (S3) forming dominant fabric



















**Sample C5M**  
anastomosing zones of subvertical clast microfabric (S5)  
moderately inclined down-ice dipping extensional Riedel shears  
down-ice dipping R-type Riedel shears

- S1 clast microfabric
  - S2 clast microfabric
  - S3 clast microfabric
  - S4 clast microfabric
  - S5 clast microfabric
  - subhorizontal Y-type Riedel shear
  - moderately down-ce dipping R-type shears
  - steeply down-ce dipping R'-type shears
  - subvertical anastomosing dewatering foliation
  - micritic carbonate cement
- sense of shear  
 orientation of clast long axis  
 alignment of clast long axes  
 clay lined fractures, voids and pore spaces


sinistral (down-ice) shear sense defined by asymmetrical microfabric

way up  
↑







way up  
↑




way up  
↑



**Sample C4M**  
orientation 122 degrees




Area B: N = 1643    Area C: N = 1591    Area D: N = 1479    Area E: N = 1579    Area F: N = 1529    Area G: N = 1703  
Area A: N = 1066    Area H: N = 1404    Area I: N = 1482    Area J: N = 1852    Area K: N = 1936    Area L: N = 1899    Area M: N = 1836    Area N: N = 1494    Area O: N = 1836



- S1 clast microfabric
- S2 clast microfabric
- S3 clast microfabric
- S4 clast microfabric
- S5 clast microfabric
- subhorizontal Y-type Riedel shear
- moderately down-ice dipping R-type shears
- steeply down-ice dipping R'-type shears
- subvertical anastomosing dewatering foliation
- micritic carbonate cement

**Sample C6M**  
orientation 122 degrees

subhorizontal foliation (S2) defining Y-type Riedel shears  
moderately inclined, up-ice dipping foliation (S3) forming dominant fabric




























**Sample C6M**  
orientation 122 degrees

anastomosing zones of subvertical clast microfabric (S5)  
moderately inclined down-ice dipping foliation (S4) defining extensional Riedel shears  
down-ice dipping R-type Riedel shears

- $\leftarrow$  sense of shear
- $\leftarrow$  orientation of clast long axis
- $\leftarrow$  alignment of clast long axes
- $\leftarrow$  clay lined fractures, voids and pore spaces

sinistral (down-ice) shear sense defined by asymmetrical microfabric

early foliation (S1) deformed by small-scale crenulations  
subvertical anastomosing foliation (S5)



**Sample C6M**  
orientation 122 degrees

orientation 122 degrees

way up

Area B: N = 1934

Area C: N = 1501

Area D: N = 1736

Area E: N = 1740

Area F: N = 1562

Area G: N = 1632

Area A: N = 1590

Area H: N = 1297

Area I: N = 1046

Area P: N = 1078

Area J: N = 1299

















Area K: N = 1725


Area L: N = 1706

Area M: N = 1526


Area N: N = 1429

Area O: N = 1616







Sample: C5M




Sample: C6M



Sample: C6M




Sample: C1M




Sample: C2M


all photomicrographs taken in plane polarised light



Sample: C2M


scale bar = 1 mm























— grain size < 0.25 mm  
 — grain size 0.25-0.5 mm  
 — grain size > 0.5 mm  
 — total clasts

— moving average (7 per.) < 0.25 mm  
 — moving average (7 per.) 0.25-0.5 mm  
 — moving average (7 per.) > 0.5 mm  
 — moving average (7 per.) total clasts









- |                      |                           |  |                            |                             |
|----------------------|---------------------------|--|----------------------------|-----------------------------|
| S1 clast microfabric | S4 clast microfabric      | subhorizontal Y-type Riedel shears (S2)        | sense of shear             | up-ice dipping S4 fabric    |
| S2 clast microfabric | S5 clast microfabric      | moderately down-ice dipping R-type shears (S3) | subhorizontal S2 fabric    | orientation of main fabrics |
| S3 clast microfabric | micritic carbonate cement | areas of well-developed S5                     | down-ice dipping S3 fabric | fluid pathways              |

**a**

sense of sliding/displacement of grains past each other

grain sliding within the plane of evolving clast microfabric

plane of evolving clast microfabric

grain rotation into plane of evolving clast microfabric

progressive rotation of grain

**b**

larger grains influencing the partitioning of deformation into anastomosing higher strain zones

anastomosing higher strain zones which form the focus for clast microfabric development

low strain areas

larger clastic grains set in a finer grained matrix

partitioning of deformation within the matrix due to the presence of large clasts

**c**

down-ice dipping clast microfabric

areas with poorly aligned clasts

up-ice dipping clast microfabric

wrapping of clast microfabric around larger grains

microfabric domains

larger grains

partitioning of deformation within the matrix due to the presence of large clasts

