



UNIVERSITY OF LEEDS

This is a repository copy of *Associations between risk factors for schizophrenia and concordance in four monozygotic twin samples*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/133440/>

Version: Accepted Version

Article:

Pepper, EJ, Pathmanathan, S, McIlrae, S et al. (2 more authors) (2018) Associations between risk factors for schizophrenia and concordance in four monozygotic twin samples. *American Journal of Medical Genetics, Part B, Neuropsychiatric Genetics*, 177 (5). pp. 503-510. ISSN 1552-4841

<https://doi.org/10.1002/ajmg.b.32640>

(c) 2018 Wiley periodicals, Inc. This is the peer reviewed version of the following article: Pepper, EJ, Pathmanathan, S, McIlrae, S et al. (2 more authors) (2018) Associations between risk factors for schizophrenia and concordance in four monozygotic twin samples. *American Journal of Medical Genetics, Part B, Neuropsychiatric Genetics*, 177 (5). pp. 503-510. ISSN 1552-4841 , which has been published in final form at <https://doi.org/10.1002/ajmg.b.32640>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

TABLE 1 Characteristics of the monozygotic twin samples

Sample	Setting	Ascertainment	Main-lifetime diagnosis of schizophrenia for current study	Mean age of proband at first contact with psychiatric services (S.D.)	Gender (% male)	Mean age of co-twin at last follow-up (S.D.)	Concordance pairwise	Concordance probandwise [†]
Maudsley register (Cardno et al., 1999; includes sample of Gottesman & Shields, 1972)	London, UK	Twin register based on admissions to psychiatric clinical services (inpatient and outpatient) (twins born 1887-1968)	DSM-III-R	25.0y (9.4)	59.6%	44.7y (13.5)	13/40 (32.5%)	20/47 (42.6%)
Slater & Shields (1953)	London, UK	Psychiatric clinical services – standing population of hospitals and inpatient and outpatient admissions (twins born 1861-1916)	DSM-IV	29.8y (8.8)	34.8%	43.2y (10.7)	10/20 (50.0%)	13/23 (56.5%)
Fischer (1973)	Denmark	Linked national twin register with central psychiatric register (twins born 1870-1920)	DSM-IV	33.4y (9.0)	41.7%	60.0y (17.2)	4/21 (19.0%)	7/24 (29.2%)
Kringlen (1967a,b)	Norway	Linked national twin register with central register of psychosis (twins born 1901-1930)	DSM-IV	26.6y (8.3)	67.5%	46.9y (10.7)	8/32 (25.0%)	16/40 (40.0%)

Samples combined	27.9y (9.3)	54.5%	47.8y (14.2)	35/113 (31.0%)	56/134 (41.8%)
---------------------	-------------	-------	--------------	-------------------	-------------------

No overlap between twins in Maudsley register and Slater & Shields samples with DSM schizophrenia.

†The number of probandwise twin pairs in each sample is given by the denominator of the probandwise concordance. There was a total of 134 pairs in the samples combined.