

This is a repository copy of *Alexithymia has no clinically relevant association with outcome of multimodal treatment tailored to needs of patients suffering from somatic symptom and related disorders:A Clinical Prospective Study*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/133503/>

Version: Accepted Version

Article:

de Vroege, Lars, Emons, Wilco H M, Sijtsma, Klaas et al. (1 more author) (2018)
Alexithymia has no clinically relevant association with outcome of multimodal treatment tailored to needs of patients suffering from somatic symptom and related disorders:A Clinical Prospective Study. *Frontiers in psychiatry*. 292. ISSN 1664-0640

<https://doi.org/10.3389/fpsy.2018.00292>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Author's Proof

Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)). Do not forget to reply to the queries.

We do not accept corrections in the form of edited manuscripts.

In order to ensure the timely publication of your article, please submit the corrections within 48 hours.

If you have any questions, please contact science.production.office@frontiersin.org.

Author Queries Form

Q1	The citation and surnames of all of the authors have been highlighted. Please check all of the names carefully and indicate if any are incorrect. Please note that this may affect the indexing of your article in repositories such as PubMed.	
Q2	Confirm that the email address in your correspondence section is accurate.	
Q3	If you decide to use previously published, copyrighted figures in your article, please keep in mind that it is your responsibility, as the author, to obtain the appropriate permissions and licenses and to follow any citation instructions requested by third-party rights holders. If obtaining the reproduction rights involves the payment of a fee, these charges are to be paid by the authors.	
Q4	Ensure that all the figures, tables and captions are correct.	
Q5	Verify that all the equations and special characters are displayed correctly.	
Q6	Ensure, if it applies to your study, the ethics statement is included in the article.	
Q7	Ensure to add all grant numbers and funding information, as after publication this is no longer possible.	
Q8	Could you please confirm if all author affiliations are fine as listed?	
Q9	Please provide doi for the following references. “(5); (26); (28).”	
Q10	Please provide the name of the editors, book title, city name, and page range for “(43).”	
Q11	Please provide the publisher name for “(52).”	
Q12	Please provide the city name for “(53).”	

Q13	Please provide the name of the editors and city name for "(54)."	
Q14	Please provide meaning of the bold values provided in "Tables 3, 4, and 5."	
Q15	Please provide the meaning of "*", lower case letter b" in "Table 2."	
Q16	Please cite "(67)." inside the text.	



Alexithymia Has No Clinically Relevant Association With Outcome of Multimodal Treatment Tailored to Needs of Patients Suffering From Somatic Symptom and Related Disorders. A Clinical Prospective Study

OPEN ACCESS

Edited by:

Michael Noll-Hussong,
Universitätsklinikum des Saarlandes,
Germany

Reviewed by:

Domenico De Berardis,
Azienda Usl Teramo, Italy
Federica Galli,
Università degli Studi di Milano, Italy
Mattie Tops,
VU University Amsterdam,
Netherlands
Rory Allen,
Goldsmiths, University of London,
United Kingdom
Gianfranco Spalletta,
Fondazione Santa Lucia (IRCCS), Italy

*Correspondence:

Lars de Vroege
l.devroege@ggzbreburg.nl

Specialty section:

This article was submitted to
Psychosomatic Medicine,
a section of the journal
Frontiers in Psychiatry

Received: 31 December 2017

Accepted: 12 June 2018

Published: xx June 2018

Citation:

de Vroege L, Emons WHM, Sijtsma K
and van der Feltz-Cornelis CM (2018)
Alexithymia Has No Clinically Relevant
Association With Outcome of
Multimodal Treatment Tailored to
Needs of Patients Suffering From
Somatic Symptom and Related
Disorders. A Clinical Prospective
Study. *Front. Psychiatry* 9:292.
doi: 10.3389/fpsy.2018.00292

Lars de Vroege^{1,2*}, Wilco H. M. Emons³, Klaas Sijtsma³ and
Christina M. van der Feltz-Cornelis^{1,2,4}

¹ Department Tranzo, Tilburg School of Social and Behavioral Sciences, Tilburg University, Tilburg, Netherlands, ² Clinical Centre of Excellence for Body, Mind and Health, GGz Breburg, Tilburg, Netherlands, ³ Department of Methodology and Statistics, Tilburg School of Social and Behavioral Sciences, Tilburg University, Tilburg, Netherlands, ⁴ Department of Health Sciences, Hull York Medical School, University of York, York, United Kingdom

Introduction: Alexithymia may moderate the effectiveness of treatment and may predict impaired general functioning of patients suffering from somatic symptom and related disorders (SSRD).

Aim: We compared alexithymia levels in a clinical prospective study with 234 consecutive patients suffering from SSRD from the Centre of Excellence for Body, Mind, and Health, Tilburg using the Bermond-Vorst Alexithymia Questionnaire, with general population norm scores. Second, we explored treatment outcomes of a multimodal treatment tailored to patient needs by Shared Decision Making (SDM) and Patient Related Outcome Monitoring (PROM) in patients with SSRD. Third, we explored whether alexithymia is associated with treatment outcome. Fourth, we explored if the presence of a chronic medical condition (e.g., diabetes mellitus, cardiovascular diseases) affects the association of alexithymia with treatment outcomes.

Results: Compared to norm scores, SSRD patients showed elevated scores on the subscales identifying, verbalizing, and fantasizing, and on the cognitive dimension. All patients benefited from treatment in terms of anxiety, depression, and physical symptoms. The association of alexithymia with treatment outcome was significant, but the effect size was negligible (range odds ratios 1.02–1.25). The association between alexithymia and treatment outcome was stronger in patients suffering from chronic medical conditions compared to patients without chronic medical conditions. However, the effect size of this association was negligible (range odds ratio 0.94–1.12).

Discussion: Alexithymia scores are elevated in patients with SSRD compared to general population scores, but the level of alexithymia has no clinically relevant association with treatment outcome both in SSRD patients with and without comorbid chronic medical conditions.

Keywords: alexithymia, treatment outcome, somatic symptom and related disorders, depression, anxiety, physical symptoms, general functioning

INTRODUCTION

Nemiah and Sifneos (1) introduced the concept of alexithymia to describe an emotional deficiency in patients with classic psychosomatic disorders, such as asthma, and hypertension. Patients were unaware of their feelings or were incapable to verbalize them, and they were unable to fantasize about their inner thoughts, feelings, and attitudes. Although the concept originated from psychoanalytical research, in time it also incorporated other perspectives, such as those originating from cognitive behavioral research and from stress research. In the 1990s, alexithymia was described as a combination of the following features: (a) difficulty identifying and describing feelings, (b) difficulty distinguishing feelings and bodily sensations caused by emotional arousal, (c) constricted imaginal processes, and (d) a cognitive style characterized by a preoccupation with the details of external events (2). These characteristics are related to stress and adaptation, and have repercussions for psychotherapeutic treatment possibilities.

Although alexithymic patients were prepared to participate in therapy consisting of a psychodynamic oriented multimodal therapy (3), alexithymia has also been described as interfering with psychotherapy (4), such as group psychotherapy, individual psychodynamic psychotherapy, and supportive therapy. Recent studies found that specifically focusing on alexithymia during treatment improved treatment outcomes in terms of symptom reduction and general functioning (5–9), but the results of a 2013 systematic review (10) were inconclusive, and was suggested that development of evidence-based treatments are necessary (10). Since most psychotherapeutic approaches rely on the patients' access to their emotions, patients unable to address these emotions provide a challenge for therapists. We do not know of studies exploring the association between alexithymia and treatment outcome at symptom level in patients suffering from somatic symptom and related disorder (SSRD).

Because alexithymia was found to be related to impoverished general functioning in somatoform disorders (2, 11–21), following the classification of the Diagnostic and Statistical Manual of Mental Disorders (DSM)-IV-TR (22) and other precursors of SSRD, as described in DSM-5 (23), general functioning also may be a relevant outcome of treatment. As far as we know, studies exploring this aspect of treatment outcome in patients suffering from SSRD have not been done. This study explores whether alexithymia has a moderating effect in treating depressive, anxiety, physical symptoms, and general functioning in SSRD patients.

OBJECTIVES AND HYPOTHESES

The first objective of this study was to estimate the level of alexithymia of patients suffering from SSRD and compare this level to known norm scores for the general population. We expected the alexithymia scores of SSRD patients to be higher.

The second objective was to explore outcomes of a multimodal treatment tailored to patient needs by Shared Decision Making (SDM) and Patient Related Outcome Monitoring (PROM) (24) with respect to depression, anxiety, physical symptoms, and general functioning in patients with SSRD, independent of alexithymia scores at baseline. We expected the treatment outcome to be improved.

The third objective was to examine the association of alexithymia with treatment outcomes with respect to depression, anxiety, physical symptoms, and general functioning. We predicted the patients with high levels of alexithymia show less favorable treatment outcomes than patients with low levels of alexithymia.

The fourth objective was to examine the influence of chronic medical conditions (e.g., diabetes mellitus, cardiovascular diseases) on the association between alexithymia and treatment outcome. We hypothesized a stronger association between alexithymia and treatment outcomes in patients suffering from chronic medical condition and alexithymia than in patients without chronic medical condition.

METHODS

Study Design

The study uses data from a longitudinal observational design in a clinical setting. The sample consisted of patients suffering from SSRD who were treated at the Clinical Centre of Excellence for Body, Mind and Health (Dutch abbreviation: CLGG), a department of GGz Breburg, Tilburg, the Netherlands. We assessed alexithymia at intake, and we assessed outcome measures including depression, anxiety, physical symptoms, and general functioning at intake before treatment and at discharge. All patients who were referred to CLGG between August 2013 and April 2016 were included in the study.

The standard intake procedure at the CLGG consists of questionnaire assessment during intake (referred to as baseline measurement), case history assessment, physical assessment, psychiatric evaluation, and psycho-diagnostic assessment. The Bermond-Vorst Alexithymia Questionnaire (BVAQ) was self-administered during the psycho-diagnostic assessment at intake. Level of education was determined using Verhage coding (25),

which includes seven levels ranging from low (levels 1 through 4), medium (level 5) to high (levels 6–7). Throughout treatment, patient's progress was evaluated using computerized Routine Outcome Monitoring (ROM) (26). We used ROM data with regard to depression, anxiety, physical symptoms and general functioning scores. Patients were informed at intake about the scientific research conducted at CLGG. Patients who did not give their consent to use their data were excluded from the dataset. Data were coded in order to create an anonymous dataset. The Commission of Scientific Research of GGz Breburg approved of this study (file number: CWO 2014-09).

Setting and Participants

Inclusion/exclusion criteria were evaluated for all patients that were referred to CLGG. Inclusion criteria were (1) completion of the intake, and (2) age at least equal to 18 years. Patients were excluded if they (1) were engaged in personal or professional injury procedures, (2) were not able to come to CLGG, (3) did not complete questionnaires from the ROM during intake and during treatment, and (4) had an IQ below 80 [assessed during intake using the Dutch Adult Reading Test (27)]. In addition, they were excluded (5) if the primary treatment focus was not related to physical symptoms. Other exclusion criteria were (6) presence of psychosis or psychotic features that hampered treatment, (7) an active suicide risk (threatening), and (8) substance dependency. Comorbid conditions and DSM-5 classifications were assessed in a clinical interview during intake.

Treatment at CLGG was of multimodal, in accordance with the multidisciplinary guideline for medically unexplained symptoms and somatic disorders (28, 29), and tailored to the needs and treatment expectations of the patient. Treatment consisted of cognitive behavioral therapy (CBT), acceptance and commitment therapy (ACT), or problem solving treatment (PST) provided by trained psychologists, in combination with pharmacotherapy provided by a physician or psychiatrist. The psychotherapeutic treatments were provided sequentially and were tailored to the needs and treatment expectations of the patients. During treatment, every 3 months both psychotherapeutic and pharmacotherapeutic treatment were adjusted based on progress in terms of PROM (24) and in a SDM (25) by multidisciplinary team consultation. Patients were treated for 1 year on average, using this multimodal approach.

Figure 1 shows a flow chart of the study. Two hundred and thirty-five patients filled out the BVAQ at intake. One patient, who gave no consent, was excluded from the study. Of the remaining 234 patients, 145 (62.0%) completed treatment. Of the patients who completed treatment, 142 patients (97.9%) filled out the Physical Symptom Checklist (PSC), 142 (97.9%) filled out the Generalized Anxiety Disorder questionnaire (GAD-7), 144 (99.3%) filled out the Patient Health Questionnaire for assessing depression (PHQ-9), and 126 (86.9%) filled out the 36-item Short Form Health Survey (SF-36), both at intake and at discharge.

Instruments

We assessed depression, anxiety, physical symptoms, and general functioning before, during and after treatment by means of

questionnaires with PROM. Alexithymia was only assessed at intake.

Bermond-Vorst Alexithymia Questionnaire (BVAQ)

Alexithymia was assessed using the 40-item BVAQ (30). The BVAQ provides valid and reliable measures of cognitive and affective dimensions of alexithymia (30–34). The Toronto Alexithymia Scale (35, 36), another instrument widely used in alexithymia research, covers cognitive factors of alexithymia but not affective dimensions. We preferred the BVAQ because it has a broader scope.

Responses to the items were scored on a five-point Likert scale. Higher scores indicate higher levels of alexithymia. The BVAQ consists of five subscales containing eight items each. The subscales are identifying, verbalizing, analyzing, fantasizing, and emotionalizing, each in accordance with the five-factor model of alexithymia (30). The five subscales constitute a cognitive dimension and an affective dimension. Scores on the cognitive dimension were obtained by adding the scores of the subscales identifying, analyzing, and verbalizing (scores ranged from 24 through 120). Scores on the affective dimension were obtained by adding the scores of the subscales emotionalizing and fantasizing (scores ranged from 16 through 80). In our SSRD study sample, coefficient alpha (37) for the cognitive dimension equaled 0.90 and for the affective dimension it equaled 0.68.

The Patient Health Questionnaire (PHQ-9)

Depression was assessed using the PHQ-9 (38). The PHQ-9 is a reliable 9-item self-report questionnaire, with higher scores indicating higher levels of depressive symptoms (38). Item scores ranged from 0 (not at all) to 3 (nearly every day), and total scores ranged from 0 to 27 (38). Cutoff points of 5, 10, 15, and 20 represent mild, moderate, moderately severe and severe levels of depression (39).

Generalized Anxiety Disorder Questionnaire (GAD-7)

Anxiety was assessed using the GAD-7. The GAD-7 is a reliable 7-item self-report questionnaire that measures symptoms of anxiety during the last 2 weeks (40). GAD-7 scores range from 0 to 21, and cutoff scores of 5, 10, and 15 represent mild, moderate and severe levels of anxiety (39).

Physical Symptom Checklist (PSC)

Physical symptoms were measured using the PSC (41), which is a 51-item questionnaire. The total score on the PSC ranges from 0 to 51 and represents the number of physical symptoms that were regularly or often present in the last week (41). De Waal and Van Hemert (42) provided normative data.

36-Item Short Form Health Survey (SF-36)

We used the SF-36 (43) to assess general functioning. Studies confirmed the SF-36's validity and reliability (44–46). The SF-36 is a self-report questionnaire that contains 36 items, which are distributed across eight scales. Using the developers' scoring algorithm (47), the eight subscales were converted into two summary measures, a physical component summary measure (PCS) and a mental component summary measure (MCS). Scores range from 0 to 100, where higher scores on the PCS and MCS




FIGURE 1 | Flowchart of Patients included in the Study. Sample Sizes are given for Patients who completed Treatment and Questionnaire Assessment. BVAQ, Bermond-Vorst Alexithymia Questionnaire.

indicate better general functioning. Normative data are available in Maglante, Hays (48).

Treatment Outcome Variables

Raw Change

For each outcome, a change score variable was created by subtracting the score after treatment from the scores at intake. This way, change scores represented treatment outcomes with respect to depression, anxiety, physical symptoms, and general functioning (PCS and MCS).

Reliable Change

To examine in more detail to what extent alexithymia predicts reliable changes at the individual level, we adopted Jacobson and Truax' (49) framework. Using their reliable change index (RCI), we determined which patients showed reliable change on the PHQ-9, the GAD-7, the PSC, the PCS of the SF-36, or the MCS of the SF-36. A dichotomous change variable was created, reflecting change at a 90% confidence level; that is, scores equal to 0 reflected no reliable change (i.e., $-1.645 < RCI < 1.645$) and scores equal to 1 reflected reliable change (i.e., $RCI < -1.645$ or $RCI > 1.645$).

Clinical Change

To further study the association between alexithymia and clinical change at the individual level, we defined a categorical variable called clinical remission. A patient shows clinical remission if change is reliable and his/her score at intake exceeds a clinical cutoff but not any more after the treatment. The following clinical cutoffs were used to define remission. For both the PHQ-9 and GAD-7, we used a score of 5, which identifies at least mild levels of depression or anxiety. For the PSC, we also used 5 as the cutoff. This cutoff coincides with the 75th percentile of PSC scores in normative data from general practitioner's offices (42). This means that remission is observed if after the treatment the patient's PSC score is no longer among the highest 25% in the

general population. To define remission on the PCS and MCS of the SF-36, the mean scores in the general population were used (48). In particular, the cutoffs for remission were 50 for the PCS and 54 for the MCS after treatment. Furthermore, to speak of clinical remission, patients must also have shown reliable change. This results in a clinical change variable having three levels: 0 = no reliable change (i.e., $|RCI| < 1.645$), 1 = reliable change but no remission, and 2 = remission.

Statistical Methods

Objective 1: Level of alexithymia was described by means of normed scores. These normed scores were obtained using normative data from the general population (50). In particular, normed scores have a mean zero and a standard deviation equal to 1. Positive norm scores suggest above average levels compared to the general population. We used one-sample *t*-tests to test whether mean differences between patients suffering from SSRD and the normal group with respect to the normed scores were significant.

Objective 2: We studied mean differences between the raw scores at intake and at discharge for the PHQ-9, GAD-7, PSC, and PCS and MCS of the SF-36, using the paired-sample *t*-tests, for the complete group. For each outcome measure, effect-size Cohen's *d* was obtained following Rosner (51). Effect sizes equal to $d = 0.2$ are considered small, $d = 0.5$ medium, and $d \geq 0.8$ large (52).

Objective 3: We used linear regression analysis to explore the association of alexithymia with depression, anxiety, physical symptoms, and general functioning. The raw change-score variables for the PHQ-9, the GAD-7, the PSC, the PCS of the SF-36, and the MCS of the SF-36 were used as dependent variables.

We used multinomial logistic regression to explore the association of alexithymia with clinical change for depression, anxiety, physical symptoms, and general functioning. Logistic regression and multinomial regression analysis provide insight into the predictive value of alexithymia in the clinical context.

Objective 4: Chronic medical condition and the two interaction terms of the dimension of alexithymia with chronic medical condition were used as independent variables in regression models.

We used logistic regression to investigate if the cognitive and affective dimensions, chronic medical condition, and the interactions between the dimensions of alexithymia and chronic medical condition predict reliable change at the individual level. For each outcome variable, the regression analyses were done as follows. First, we estimated the full model that included as predictors the background variables (age, gender, and education level), the first-order effects of the cognitive and affective alexithymia dimensions, and chronic medical conditions, and the interaction effects between the alexithymia dimensions and medical conditions. To study the interaction effects, we centered the independent variables to avoid potential problems with respect to multicollinearity (53). Second, in case some of the interaction effects were non-significant, we re-estimated the model without the non-significant interaction effects.

For logistic regression analysis, we used Nagelkerke's R -square to gauge effect size. Formally, the pseudo R -square does not represent proportions of explained variance, but we interpreted the pseudo R -square as if it did provide the proportion of the variation the model explained (54). We used the guidelines of Cohen (55) to interpret Nagelkerke's pseudo R -square (i.e., R -square = 0.02 was considered small, R -square = 0.13 was considered medium, R -square \geq 0.26 was considered large). All analyses were done by means of the Statistical Package for the Social Sciences version 22 (56).

RESULTS

Sample Characteristics

Two hundred and two patients (86.3%) were diagnosed with a somatic symptom disorder, 10 patients (4.3%) were diagnosed with an illness anxiety disorder, and 22 patients (9.4%) were diagnosed with a conversion disorder.

Table 1 (upper panel) describes the socio-demographic characteristics of the SSRD patient sample. The SSRD sample consisted of 234 patients (59.0% females). The sample had a mean (M) age of 42.8 [standard deviation (SD) = 12.56; range: 19 to 79]. Seven patients had missing values on the BVAQ items. One of these patients had six missing item scores and was excluded from further analyses. The only missing item score for the remaining six patients was imputed using two-way imputation (57, 58).

The level of alexithymia of patients suffering from SSRD compared to known norm scores for the general population.

Table 1 (lower panel) shows the means for the raw scores (column 2) and normed scores (column 5) on the BVAQ. Using a significance level equal to 0.007 (alpha of 0.05 divided by 7, equal to the number of used tests), significant mean differences with respect to the norm scores were found for the subscales verbalizing [$t(233) = 4.239, p < 0.001$], fantasizing [$t(233) = 3.770, p < 0.001$], identifying [$t(233) = 7.759, p < 0.001$], and emotionalizing [$t(233) = -8.106, p < 0.001$]. A significant mean difference was

found for the cognitive dimension [$t(233) = 4.944, p < 0.001$]. For the subscales of the cognitive dimension, we found elevated levels of identifying ($M = 0.69$; range -1.97 to 4.26) and verbalizing ($M = 0.39$; range -2.87 to 3.08) compared to the general population. No significant mean differences were found between SSRD patients and the general population for analyzing [$t(233) = -0.500, p = 0.618$] and the affective dimension [$t(233) = -1.632, p = 0.104$]. For the subscales of the affective dimension in the BVAQ, we found lowered levels of emotionalizing ($M = -0.55$; range -2.96 to 2.48), but higher mean values for fantasizing ($M = 0.27$; range -2.51 to 2.42).

Treatment outcomes of a multimodal treatment tailored to patients needs by SDM and PROM in terms of depression, anxiety, physical symptoms, and general functioning.

Table 2 shows the mean scores before and after treatment for the PSC, GAD-7, PHQ-9, and the MCS and PCS of the SF-36. Results suggested substantive mean improvements of the treatment outcomes. PSC means before and after treatment improved significantly [$t(141) = 4.207, p < 0.001, d = 1.82$], the mean scores on the PHQ-9 also differed significantly before and after treatment [$t(143) = 4.837, p < 0.001, d = 1.43$], and the mean scores on the GAD-7 differed significantly before and after treatment [$t(141) = 5.090, p < 0.001, d = 1.21$]. Mean MCS and PCS scores for the SF-36 did not differ significantly before and after treatment [respectively, $t(125) = 0.757, p = 0.450, d = 0.92$ and $t(126) = 1.494, p = 0.138, d = 1.06$].

The association of alexithymia with treatment outcomes in terms of depression, anxiety, physical symptoms, and general functioning.

Alexithymia had significant association with treatment outcome regarding anxiety (**Table 3**). The cognitive and affective dimensions did not significantly predict change with respect to depression, physical symptoms scores, and general health functioning.

Table 4 shows the results of the logistic regression analyses. The cognitive dimension had a significant negative main association on treatment outcome with respect to anxiety (*Odds Ratio* (OR) = 1.02, 95% CI = [1.00, 1.05]). The squared semi-partial correlation for this dimension was 0.05, which means that 5% of the total variability of the treatment outcome for anxiety is uniquely associated with the cognitive dimension. The affective dimension was associated with a significant positive association on treatment outcome with respect to general mental health functioning ($OR = 1.25, 95\%CI = [1.09, 1.44]$). Removing the affective dimension decreased R -square to .15, which renders the association substantial.

Table 5 shows the results for predicting reliable and clinical change. With regard to general functioning (MCS of the SF-36), the affective dimension was significantly associated with clinical change in the group of patients with no remission vs. the group of patients with no clinical change and no remission ($OR = 1.24, 95\%CI = [1.08, 1.42]$). Results for clinical change and remission vs. no clinical change and no remission could not be computed, because none of the patients showed remission on the MCS. No significant associations were found between anxiety and for the PCS of the SF-36.

TABLE 1 | Socio-demographic characteristics and descriptive statistics for the BVAQ in the SSRD sample at intake ($N = 234$).

Characteristic	<i>M (SD)</i>	Min/Max	<i>n (%)</i>	Normed scores ^a (min/max)	<i>p</i>
BACKGROUND VARIABLES					
Gender					
Men			96 (41.0)		
Women			138 (59.0)		
Age	42.78 (12.56)	19/79			
Educational level*					
Low (1–4)			56 (24.9)		
Medium (5)			103 (45.8)		
High (6–7)			66 (29.2)		
(Missing value)			(9)		
Marital status					
Married/living together			150 (71.4)		
Divorced			11 (5.2)		
Widow(er)			1 (0.5)		
Single			48 (22.9)		
(Missing value)			(24)		
PSC	16.57 (8.08)	0/38			
GAD-7	11.51 (5.47)	0/21			
PHQ-9	14.21 (6.07)	0/27			
SF-36 ($n = 225$) ^b					
PCS	40.48 (5.44)	27.49/57.43			
MCS	44.01 (5.16)	21.30/55.55			
Comorbidity at intake					
Comorbid anxiety			2 (0.9)		
Comorbid depression			25 (11.4)		
Comorbid depression and anxiety			193 (87.7)		
BVAQ SCORES					
Cognitive dim.	67.12 (17.64)	32/106		0.43 (–2.38/3.25)	< 0.001
Identifying	22.10 (7.02)	8/40		0.69 (–1.97/4.26)	< 0.001
Analyzing	19.61 (6.02)	8/34		–0.04 (–2.52/2.97)	0.618
Verbalizing	25.41 (8.29)	8/40		0.39 (–2.87/3.08)	< 0.001
Affective dim.	45.48 (8.88)	19/66		–0.10 (–2.79/2.45)	0.104
Fantasizing	26.74 (6.92)	8/40		0.27 (–2.51/2.42)	< 0.001
Emotionalizing	18.74 (5.16)	8/32		–0.55 (–2.96/2.48)	< 0.001

SSRD, Somatic Symptom and Related Disorders; PSC, Physical Symptom Checklist; GAD-7, Generalized Anxiety Disorder questionnaire; PHQ-9, Patient Health Questionnaire; SF-36, 36-item Short Form Health Survey; MCS, Mental Component Summary; PCS, Physical Component Summary; BVAQ, Bermond-Vorst Alexithymia Questionnaire.

^aNormed scores were based on normative data (50).

Influence of chronic medical conditions (e.g., diabetes mellitus, cardiovascular diseases, and others) on the association between alexithymia and treatment outcome.

The cognitive dimension and medical condition showed a significant interaction association (see **Table 3**). Simple effects analysis suggested a negative association for patients without a chronic medical condition ($B = -0.08$, $p = 0.022$), and a non-significant association for patients with a chronic medical condition ($B = 0.04$, $p = 0.329$).

The interaction between the cognitive dimension and chronic medical condition had a significant association with reliable change with regard to depression ($OR = 0.95$, $95\%CI = [0.91, 0.99]$) (**Table 4**). The squared semi-partial correlation for this interaction was 0.06, which means that 6% of the total

variability of treatment outcome for depression is uniquely associated with the interaction between cognitive dimension of alexithymia and chronic medical condition. The OR equal to 0.95 suggests that the association between alexithymia (cognitive dimension) and treatment outcome in terms of depression is negative for patients with chronic medical condition compared to patients without a chronic medical condition. Nevertheless, these correlations and ORs are very small and we render them clinically irrelevant.

Regarding physical symptoms, a significant interaction effect between cognitive dimension and chronic medical condition was found for the PSC ($OR = 0.95$, $95\%CI = [0.91, 1.00]$). The OR equal to 0.95 suggests that the association between alexithymia (cognitive dimension) and treatment outcome in

TABLE 2 | Mean scores on the PHQ-9, GAD-7, PSC, and SF-36 of the SSRD Sample at Intake and After Treatment.

Outcome measure	N	Measurement occasion				p
		At intake		After treatment		
		M	SD	M	SD	
PSC	142	16.26	7.68	13.54	9.22	<0.001
GAD-7	142	11.18	5.41	9.09	6.17	<0.001
PHQ-9	144	13.94	6.12	11.50	7.36	<0.001
SF-36	126					
PCS		41.10	5.44	40.42	5.48	0.138
MCS		43.47	5.73	43.03	5.82	0.450

M, Mean; SD, Standard Deviation; PHQ-9, Patient Health Questionnaire-9; GAD-7, General Anxiety Disorder questionnaire; PSC, Physical Symptom Checklist; SF-36, 36-item Short Form Health Survey; MCS, Mental Component Summary; PCS, Physical Component Summary.

n = the number of patients who completed the treatment and who filled out the questionnaire both at intake and after treatment.

TABLE 3 | Linear regression of raw change scores for the GAD-7 on the BVAQ dimensions and covariates.

Predictors	Change scores for the GAD-7		
	B	95% CI	P
Cognitive dimension	-0.08	[-0.15, -0.01]	0.021
Affective dimension	-0.01	[-0.13, 0.12]	0.911
Chron med cond	0.76	[-0.97, 2.49]	0.386
Int cogn_med	0.12	[0.02, 0.22]	0.022
Int aff_med	-0.15	[-0.35, 0.05]	0.140

GAD-7, Generalized Anxiety Disorder questionnaire; 95% CI, 95% confidence interval for B; Chron med cond, chronic medical condition; Int cogn_med, interaction between the cognitive dimension of alexithymia and chronic medical condition; Int aff_med, interaction between the affective dimension and chronic medical condition.

terms of physical symptoms is negative for patients with chronic medical condition compared to patients without a chronic medical condition. The squared semi-partial correlation for this interaction was 0.05, which means that 5% of the total variability of treatment outcome for physical symptoms is uniquely associated with this interaction.

Table 5 shows the results for predicting reliable and clinical change. Regarding depression, the interaction between the cognitive dimension and chronic medical condition is significant in patients with clinical change and no remission vs. patients with no clinical change and no remission ($OR = 0.94$, $95\%CI = [0.89, 0.99]$). The interaction between the affective dimension and chronic medical condition is significant in patients with clinical change and no remission vs. patients with no clinical change and no remission ($OR = 1.12$, $95\%CI = [1.00, 1.24]$).

To conclude, our results suggest some associations of alexithymia with clinical change and the influence of chronic medical condition on the association between alexithymia and

treatment outcome with respect to depression, anxiety, physical symptoms, and general functioning. However, the estimated ORs of ~ 1.00 suggests that these associations are very small and negligible.

DISCUSSION

Main Findings

Alexithymia in patients suffering from SSRD was compared to normative data for the general population. The scores of SSRD patients on emotionalizing were lower compared to the norm scores, while elevated scores were found for other BVAQ subscales. The results suggest that SSRD patients show reduced abilities to identify, verbalize, and fantasize, and tend to be aroused by emotional events. This confirms the first hypothesis.

The results also suggest that patients suffering from SSRD improve after multimodal and tailored treatment with regard to anxiety, depression, and physical symptoms (Cohen's d ranged 1.21–1.82). This confirms the second hypothesis.

Even though our results suggest some associations of alexithymia with treatment outcome, the odds ratios were close to 1.0 (range 1.02–1.25). Therefore, we render the association of alexithymia not clinically relevant with regard to treatment outcome in terms of depression, anxiety, physical symptoms, and general functioning. This is not what we expected based on the literature.

The cognitive dimension of alexithymia affects treatment outcomes for patients suffering from chronic medical condition but not for patients free of chronic medical conditions with regard to depression and physical symptoms, but odds ratios were equal to 0.95 so we render the association also not clinically relevant. This is not what we expected.

SSRD patients in this study received multimodal treatment tailored to the patients' needs in a SDM model based on repeated PROM, and improved significantly after treatment. However, the level of alexithymia at baseline was not associated with a clinically relevant difference in treatment outcome, although our results suggested that SSRD patients have difficulties with identification and verbalization of emotions. Two possible explanations are the following. First, as treatment outcome was positive independent of suffering from alexithymia, it might be that alexithymia is not a clinically relevant factor needing specific attention when treating patients with SSRD. This might be a possible explanation, if in a randomized clinical trial design the positive association of this multimodal treatment model would be confirmed and again the association with alexithymia would be clinically irrelevant. Second, it could be that the treatment, although yielding positive outcomes, could have better outcomes for patients with high alexithymia scores if the treatment would address them. In that case, treatment should focus on improving identification and verbalization of emotions (6).

Treatment options for SSRD include affective mentalizing as prominent factor because affective mentalization is involved in the onset and prolongation of physical symptomatology and the interpersonal problems that co-occur with these physical symptoms (59, 60). The link between emotional states and bodily distress and how to restore this link could be improved

TABLE 4 | Logistic regression analyses predicting reliable change regarding depression, physical symptoms, and general functioning.

Predictors	Outcome variable											
	Depression (PHQ-9)			Anxiety (GAD-7)			Physical symptoms (PSC)			General functioning (MSC of the SF-36)		
	OR	95% CI	R ^{2a}	OR	95% CI	R ^{2a}	OR	95% CI	R ^{2a}	OR	95% CI	R ^{2a}
			0.10			0.08			0.14			0.43
Cogn dim	1.03	[1.00, 1.06]		1.02	[1.00, 1.05]		1.01	[0.98, 1.04]		1.00	[0.96, 1.04]	
Aff dim	1.03	[0.98, 1.09]		0.98	[0.94, 1.03]		0.99	[0.94, 1.05]		1.25	[1.09, 1.44]	
Chron med cond	1.06	[0.50, 2.26]		0.93	[0.45, 1.92]		1.38	[0.63, 2.98]		0.42	[0.08, 2.28]	
Cogn × medical condition	0.95	[0.91, 0.99]		–	–		0.95	[0.91, 1.00]		–	–	
Aff × medical condition	1.01	[0.92, 1.10]		–	–		1.10	[1.00, 1.21]		–	–	

PHQ-9, Patient Health Questionnaire; GAD-7, Generalized Anxiety Disorder questionnaire; PSC, Physical Symptom Checklist; MCS, Mental Component Summary; SF-36, 36-item Short Form Health Survey; OR, Odds Ratio; 95% CI, 95% confidence interval; Chron med cond, chronic medical condition; Int cognmed, interaction term of the cognitive dimension of alexithymia and chronic medical condition; Int aff, interaction term of the affective dimension and chronic medical condition. Results for chronic medical condition, and for the interaction terms of the alexithymia dimension and chronic medical condition (Model 3 and 4; respectively) yielded no significant results. ^aNagelkerke's pseudo R-square. All coefficients in bold are significant at the 5% significance level.

TABLE 5 | Multinomial logistic regression analyses predicting clinical change regarding depression, physical symptoms, and general functioning.

Predictors	Depression (PHQ-9)			Physical symptoms (PSC)			General functioning (MCS of the SF-36)		
	OR	95% CI	p	OR	95% CI	p	OR	95% CI	p
CLINICAL CHANGE/NO REMISSION vs. NO CLINICAL CHANGE/NO REMISSION									
Affective dimension	1.02	[0.95, 1.09]	0.561	1.00	[0.94, 1.06]	0.880	1.24	[1.08, 1.42]	0.003
Cognitive dimension	1.03	[1.00, 1.07]	0.091	1.01	[0.98, 1.04]	0.572	0.90	[0.95, 1.04]	0.630
Chron med cond	0.86	[0.33, 2.22]	0.751	1.25	[0.53, 2.94]	0.618	0.39	[0.07, 2.12]	0.276
Int aff_med	1.00	[0.89, 1.11]	0.944	1.12	[1.00, 1.24]	0.048	–	–	–
Int cogn_med	0.94	[0.89, 0.99]	0.021	0.95	[0.91, 1.00]	0.054	–	–	–
CLINICAL CHANGE/REMISSION vs. NO CLINICAL CHANGE/NO REMISSION									
Affective dimension	1.03	[0.95, 1.12]	0.424	1.01	[0.90, 1.15]	0.822	–	–	–
Cognitive dimension	1.02	[0.98, 1.07]	0.301	1.00	[0.94, 1.07]	0.977	–	–	–
Chron med cond	1.46	[0.52, 4.12]	0.613	1.19	[0.22, 6.40]	0.836	–	–	–
Int aff_med	0.97	[0.91, 1.02]	0.230	1.03	[0.86, 1.25]	0.732	–	–	–
Int cogn_med	1.03	[0.91, 1.16]	0.643	0.95	[0.87, 1.05]	0.319	–	–	–

PHQ-9, Patient Health Questionnaire; PSC, Physical Symptom Checklist; MCS, Mental Component Summary; SF-36, 36-Item Short Form Health Survey; OR, Odds Ratio; 95% CI, 95% confidence interval; Chron med cond, chronic medical condition; Int cogn_med, interaction term of the cognitive dimension of alexithymia and chronic medical condition; Int aff_med, interaction term of the affective dimension and chronic medical condition. Results not reported because interaction effect was not significant, or effect could not be estimated because no patients showed remission on the MCS.

by enhancing ones capacity of emotional awareness. A recent study suggested that women with fibromyalgia might benefit from an emotional disclosure or expression intervention (61). Our results suggested that SSRD patients have difficulties with identification and verbalization of emotions. Therefore, treatment of SSRD patients should focus on improving identification and verbalization of emotions which was also suggested by a previous study (6).

Nevertheless, based on our results we conclude that the influence of alexithymia is clinically irrelevant. Previous studies also found a relationship between alexithymia and interpersonal

dysfunction, aggression, and personality disorders (62–64). This association is not yet explored amongst patients suffering from SSRD. Personality characteristics such as interpersonal dysfunction, aggressive behavior or coping strategies may also increase insights in the personal characteristics of patients suffering from SSRD and might offer treatment options. Studies focusing on these kinds of personality characteristics are warranted in order to establish such new therapies. Future studies should also include other patient groups (e.g., depressed patients), to explore differences in emotion regulation between patients having SSRD and other patients. This way,

researchers are able to explore whether or not impoverished emotional regulation is a specific feature of SSRD or a common feature of patients suffering from other mental disorders.

Strengths and Limitations of the Study

The use of norm scores of a large sample from the general population as a reference group accounts as a major strength, because gender and age-specific norms could be used from a large and representable group. Furthermore, the sample consisted of consecutive patients visiting a Clinical Centre of Excellence for patients with Somatic Symptom Disorder, who were referred by their general practitioner, their medical specialist or their psychiatrist or psychotherapist after an average treatment duration of 7 years without solace. The patients had high complexity levels at biological and psychological symptom levels (65), suffered from comorbidity, complex treatment histories, and high levels of social vulnerability. Hence, this study provides us with findings relevant for such a patient group with SSRD, but results may not necessarily be generalizable to the general population. However, although the composition of patient populations may differ across regions, it is unlikely that the underlying mechanisms of treatment

outcome differ across specialty mental health institutions. Unfortunately, our sample was too small to explore the relationship between alexithymia with treatment outcome for different SSRD categories (e.g., somatic symptom disorder and illness anxiety disorder). Our sample was heterogeneous with respect to SSRD diagnoses. Furthermore, we did not include neuropsychological aspects in this study whom may have negatively influence treatment outcome since patients suffering from SSRD experience significant cognitive problems (66).

Implications for Research

Future studies on the relationship between alexithymia and treatment outcome should differentiate between SSRD categories. Future studies should also include large samples and evaluate effectiveness of multimodal tailored treatment supported by SDM and PROM in a randomized design.

AUTHOR CONTRIBUTIONS

LdV drafted the manuscript. LdV and WE were responsible for design and analysis of the data. WE, KS, and CvdF-C revised the draft. All authors approved of the final manuscript.

REFERENCES

- Nemiah JC, Sifneos PE. Psychosomatic illness: a problem in communication. *Psychother Psychosomat.* (1970) 18:154–60. doi: 10.1159/000286074
- Cox BJ, Kuch K, Parker JDA, Schulman ID, Evans RJ. Alexithymia in somatoform disorder patients with chronic pain. *J Psychosomat Res.* (1994) 38:523–7. doi: 10.1016/0022-3999(94)90049-3
- Leweke F, Bausch S, Leichsenring F, Walter B, Stingl M. Alexithymia as a predictor of outcome of psychodynamically oriented inpatient treatment. *Psychother Res.* (2009) 19:323–31. doi: 10.1080/10503300902870554
- Ogrodniczuk JS, Piper WE, Joyce AS. Effect of alexithymia on the process and outcome of psychotherapy: a programmatic review. *Psychiatry Res.* (2011) 190:43–8. doi: 10.1016/j.psychres.2010.04.026
- Beck T, Breuss M, Kunnig M, Schüßler G. The first step is the hardest-emotion recognition in patients with somatoform disorders. *Zeitschrift für Psychosomatische Medizin und Psychotherapie* (2013) 59: 385–90.
- Cameron K, Ogrodniczuk J, Hadjipavlou G. Changes in alexithymia following psychological intervention: a review. *Harv Rev Psychiatry* (2014) 22:162–78. doi: 10.1097/HRP.0000000000000036
- Gay MC, Hanin D, Luminet O. Effectiveness of an hypnotic imagery intervention on reducing alexithymia. *Contemp Hypnos.* (2008) 25:1–13. doi: 10.1002/ch.344
- Ogrodniczuk JS, Sochting I, Piper WE, Joyce AS. A naturalistic study of alexithymia among psychiatric outpatients treated in an integrated group therapy program. *Psychol Psychother.* (2012) 85:278–91. doi: 10.1111/j.2044-8341.2011.02032.x
- Tulipani C, Morelli F, Spedicato MR, Maiello E, Todarello O, Porcelli P. Alexithymia and cancer pain: the effect of psychological intervention. *Psychother Psychosomat.* (2010) 79:156–63. doi: 10.1159/000286960
- Samur D, Tops M, Schlinkert C, Quirin M, Cuijpers P, Koole SL. Four decades of research on alexithymia: moving toward clinical applications. *Front Psychol.* (2013) 4:861. doi: 10.3389/fpsyg.2013.00861

- Bach M, Bach D. Predictive value of alexithymia: a prospective study in somatizing patients. *Psychother Psychosomat.* (1995) 64:43–8. doi: 10.1159/000288989
- Burba B, Oswald R, Grigaliunien V, Neverauskiene S, Jankuviene O, Chue P. A controlled study of alexithymia in adolescent patients with persistent somatoform pain disorder. *Can J Psychiatry* (2006) 51:468–71. doi: 10.1177/070674370605100709
- Cohen K, Auld F, Brooker H. Is alexithymia related to psychosomatic disorder and somatizing? *J Psychosomat Res.* (1994) 38:119–27.
- Duddu V, Isaac MK, Chaturvedi SK. Alexithymia in somatoform and depressive disorders. *J Psychosomat Res.* (2003) 54:435–8. doi: 10.1016/S0022-3999(02)00440-3
- Kooijman CG. The status of alexithymia as a risk factor in medically unexplained physical symptoms. *Comprehens Psychiatry* (1998) 39:152–9. doi: 10.1016/S0010-440X(98)90075-X
- Kosturek A, Gregory RJ, Sousou AJ, Trief P. Alexithymia and somatic amplification in chronic pain. *Psychosomatics* (1998) 39:399–404. doi: 10.1016/S0033-3182(98)71298-8
- Moreno-Jiménez B, Blanco BL, Rodríguez-Muñoz A, Hernández EG. The influence of personality factors on health-related quality of life of patients with inflammatory bowel disease. *J Psychosomat Res.* (2007) 62:39–46. doi: 10.1016/j.jpsychores.2006.07.026
- Taylor GJ, Parker JDA, Bagby M, Acklin MW. Alexithymia and somatic complaints in psychiatric out-patients. *J Psychosomat Res.* (1992) 36:417–24. doi: 10.1016/0022-3999(92)90002-J
- Verissimo R, Mota-Cardoso R, Taylor G. Relationships between alexithymia, emotional control, and quality of life in patients with inflammatory bowel disease. *Psychother Psychosomat.* (1998) 67:75–80. doi: 10.1159/000012263
- Von Rimscha S, Moergeli H, Weidt S, Straumann D, Hegemann S, Ruffer M. Alexithymia and health-related quality of life in patients with dizziness. *Psychopathology* (2013) 46:377–83. doi: 10.1159/000345357
- Wolf LD, Hentz JG, Ziemba KS, Kirilin KA, Noe KH, Hoerth MT, et al. Quality of life in psychogenic nonepileptic seizures and epilepsy: the role of somatization and alexithymia. *Epilepsy Behav.* (2015) 43:81–8. doi: 10.1016/j.yebeh.2014.12.010

- 1027 22. American Psychiatric Association. *DSM-IV-TR: Diagnostic and Statistical*
1028 *Manual of Mental Disorders, Text Revision*. Washington, DC: American
1029 Psychiatric Association (2000).
1030 23. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental*
1031 *Disorders (DSM-5)*. Washington, DC: American Psychiatric Association
1032 (2013).
1033 24. Black N. Patient reported outcome measures could help transform healthcare.
1034 *Br Med J*. (2013) 346:f167. doi: 10.1136/bmj.f167
1035 25. Verhage F. *Intelligence and Age: Study With Dutch People Aged 12-77*. Assen:
1036 *Van Gorcum* (1964).
1037 26. Van der Feltz-Cornelis CM, Andrea H, Kessels E, Duivenvoorden HJ, Biemans
1038 H, Metz M. Shared decision making in combinatie met ROM bij patiënten met
1039 gecombineerde lichamelijke en psychische klachten; een klinisch-empirische
1040 verkenning [article in Dutch] English: does routine outcome monitoring
1041 have a promising future? An investigation into the use of shared decision-
1042 making combined with ROM for patients with a combination of physical and
1043 psychiatric symptoms. *Tijdschrift voor Psychiatrie* (2014) 56:375–84.
1044 27. Schmand B, Lindeboom J, Van Harskamp F. *Dutch Adult Reading Test*.
1045 Netherlands: Swets & Zeitlinger, Lisse (1992).
1046 28. Van der Feltz-Cornelis C, Swinkels J, Blankenstein A, Hoedeman R, Keuter
1047 E. The Dutch multidisciplinary guideline entitled 'Medically unexplained
1048 physical symptoms and somatoform disorder'. *Nederlands tijdschrift voor*
1049 *geneeskunde* (2010) 155:A1244–A.
1050 29. Van der Feltz-Cornelis CM, Hoedeman R, Keuter EJ, Swinkels JA.
1051 Presentation of the Multidisciplinary guideline medically unexplained
1052 physical symptoms (MUPS) and Somatoform Disorder in the Netherlands:
1053 disease management according to risk profiles. *J Psychosom Res*. (2012)
1054 72:168–9. doi: 10.1016/j.jpsychores.2011.11.007
1055 30. Vorst HC, Bermond B. Validity and reliability of the Bermond-
1056 Vorst Alexithymia Questionnaire. *Pers Individ Diff*. (2001) 30:413–34.
1057 doi: 10.1016/S0191-8869(00)00033-7
1058 31. Bermond B, Clayton K, Liberova A, Luminet O, Maruszewski T, Ricci
1059 Bitti PE, et al. A cognitive and an affective dimension of alexithymia
1060 in six languages and seven populations. *Cogn Emot*. (2007) 21:1125–36.
1061 doi: 10.1080/02699930601056989
1062 32. Deborde A-S, Berthoz S, Wallier J, Fermanian J, Falissard B, Jeammet P,
1063 et al. The bermond-vorst alexithymia questionnaire cutoff scores: a study
1064 in eating-disordered and control subjects. *Psychopathology* (2007) 41:43–9.
1065 doi: 10.1159/000109955
1066 33. Müller J, Bühner M, Ellgring H. The assessment of alexithymia: psychometric
1067 properties and validity of the Bermond-Vorst alexithymia questionnaire. *Pers*
1068 *Individ Diff*. (2004) 37:373–91. doi: 10.1016/j.paid.2003.09.010
1069 34. Zech E, Luminet O, Rimé B, Wagner H. Alexithymia and its measurement:
1070 confirmatory factor analyses of the 20-item Toronto Alexithymia Scale and
1071 the Bermond-Vorst Alexithymia Questionnaire. *Eur J Pers*. (1999) 13:511–32.
1072 35. Bagby RM, Parker JD, Taylor GJ. The twenty-item Toronto Alexithymia
1073 Scale–I. Item selection and cross-validation of the factor structure. *J*
1074 *Psychosom Res*. (1994) 38:23–32.
1075 36. Bagby RM, Taylor GJ, Parker JD. The Twenty-item Toronto Alexithymia
1076 Scale–II. Convergent, discriminant, and concurrent validity. *J Psychosom Res*.
1077 (1994) 38:33–40.
1078 37. Cronbach LJ. Coefficient alpha and the internal structure of tests.
1079 *Psychometrika* (1951) 16:297–334.
1080 38. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief
1081 depression severity measure. *J Gen Inter Med*. (2001) 16:606–13.
1082 doi: 10.1046/j.1525-1497.2001.016009606.x
1083 39. Kroenke K, Spitzer RL, Williams JB, Löwe B. The patient health questionnaire
1084 somatic, anxiety, and depressive symptom scales: a systematic review. *Gen*
1085 *Hosp Psychiatry* (2010) 32:345–59. doi: 10.1016/j.genhosppsych.2010.03.006
1086 40. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing
1087 generalized anxiety disorder: the GAD-7. *Arch Int Med*. (2006) 166:1092–7.
1088 doi: 10.1001/archinte.166.10.1092
1089 41. Van Hemert A. *Lichamelijke Klachten Vragenlijst [in Dutch]*. Leiden: Leids
1090 Universitair Medisch Centrum (2003).
1091 42. De Waal MW, Van Hemert A. *Spreadsheet Normscores Dutch Respondents*
1092 *in the General Population and in a General Practitioner's Population*. Available
1093 online at: [http://www.psychiatrieweb.mywebhome.nl/pw.somatisatie/files/](http://www.psychiatrieweb.mywebhome.nl/pw.somatisatie/files/docs/lkv31norm.pdf)
1094 [docs/lkv31norm.pdf](http://www.psychiatrieweb.mywebhome.nl/pw.somatisatie/files/docs/lkv31norm.pdf) (2013).
1095 43. Ware JE, Keller SD, Kosinski M. SF-36: Physical and mental health summary
1096 scales: a user's manual. Health Assessment Lab (1994).
1097 44. Aaronson NK, Muller M, Cohen PD, Essink-Bot M-L, Fekkes M, Sanderman
1098 R, et al. Translation, validation, and norming of the Dutch language version
1099 of the SF-36 Health Survey in community and chronic disease populations. *J*
1100 *Clin Epidemiol*. (1998) 51:1055–68.
1101 45. Garratt AM, Ruta DA, Abdalla MI, Russell IT. SF 36 health survey
1102 questionnaire: II. Responsiveness to changes in health status in four common
1103 clinical conditions. *Q Health Care* (1994) 3:186–92. doi: 10.1136/qshc.3.4.186
1104 46. McHorney CA, Ware Jr. JE, Raczek AE. The MOS 36-item short-form
1105 health survey (SF-36): II. Psychometric and clinical tests of validity in
1106 measuring physical and mental health constructs. *Med Care* (1993) 13:247–63.
1107 doi: 10.1097/00005650-199303000-00006
1108 47. Ware J, Kosinski M, Keller S. *SF-36 Physical and Mental Health Summary*
1109 *Scales: A Manual for Users of Version 1. 2nd Edn*. Lincoln, RI: QualityMetric.
1110 Inc (2001).
1111 48. Maglinte GA, Hays RD, Kaplan RM. US general population norms for
1112 telephone administration of the SF-36v2. *J Clin Epidemiol*. (2012) 65:497–502.
1113 doi: 10.1016/j.jclinepi.2011.09.008
1114 49. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining
1115 meaningful change in psychotherapy research. *J Consul Clin Psychol*. (1991)
1116 59:12. doi: 10.1037/0022-006X.59.1.12
1117 50. De Vroegte L, Emons WH, Sijtsma K, van der Feltz-Cornelis CM.
1118 Psychometric properties of the bermond-vorst alexithymia questionnaire
1119 (BVAQ) in the general population and a clinical population. *Front Psychiatry*
1120 (2018) 9:111. doi: 10.3389/fpsy.2018.00111
1121 51. Rosner B. *Fundamentals of Biostatistics*. Nelson Education (2015).
1122 52. Cohen J. Statistical power analysis for the behavioral sciences Lawrence
1123 Earlbaum Associates. Hillsdale, NJ (1988).
1124 53. Cohen J, Cohen P, West SG, Aiken LS. *Applied Multiple Regression/Correlation*
1125 *Analysis for the Behavioral Sciences*. Routledge (2013).
1126 54. Nagelkerke NJ. Maximum likelihood estimation of functional relationships.
1127 In: *Maximum Likelihood Estimation of Functional Relationships*. Springer
1128 (1992). p. 11–61.
1129 55. Cohen J. A power primer. *Psychol Bull*. (1992) 112:155–9.
1130 doi: 10.1037/0033-2909.112.1.155
1131 56. IBM Corp. *IBM SPSS Statistics for Windows, Version 22.0*. IBM Corp Armonk,
1132 NY (2011).
1133 57. Bernaards CA, Sijtsma K. Influence of imputation and EM methods on
1134 factor analysis when item nonresponse in questionnaire data is nonignorable.
1135 *Multivar Behav Res*. (2000) 35:321–64. doi: 10.1207/S15327906MBR3503_03
1136 58. Van Ginkel JR, Van der Ark LA, Sijtsma K, Vermunt JK. Two-way imputation:
1137 a Bayesian method for estimating missing scores in tests and questionnaires,
1138 and an accurate approximation. *Comput Stat Data Anal*. (2007) 51:4013–27.
1139 doi: 10.1016/j.csda.2006.12.022
1140 59. Luyten P, Van Houdenhove B, Lemma A, Target M, Fonagy P. A
1141 mentalization-based approach to the understanding and treatment of
1142 functional somatic disorders. *Psychoanalyt Psychother*. (2012) 26:121–40.
1143 doi: 10.1080/02668734.2012.678061
1144 60. Tominaga T, Choi H, Nagoshi Y, Wada Y, Fukui K. Relationship between
1145 alexithymia and coping strategies in patients with somatoform disorder.
1146 *Neuropsychiatr Dis Treat* (2014) 10:55–62. doi: 10.2147/NDT.S55956
1147 61. Geenen R, van Ooijen-van der Linden L, Lumley MA, Bijlsma JW, van
1148 Middendorp H. The match–mismatch model of emotion processing styles
1149 and emotion regulation strategies in fibromyalgia. *J Psychosom Res*. (2012)
1150 72:45–50. doi: 10.1016/j.jpsychores.2011.09.004
1151 62. Nicolò G, Semerari A, Lysaker PH, Dimaggio G, Conti L, D'Angerio S,
1152 et al. Alexithymia in personality disorders: correlations with symptoms
1153 and interpersonal functioning. *Psychiatry Res*. (2011) 190:37–42.
1154 doi: 10.1016/j.psychres.2010.07.046
1155 63. Fossati A, Acquarini E, Feeney JA, Borroni S, Grazioli F, Giarolli LE,
1156 et al. Alexithymia and attachment insecurities in impulsive aggression.
1157 *Attach Hum Dev*. (2009) 11:165–82. doi: 10.1080/146167308026
1158 25235
1159 64. Singleton RA, Bruce C. Straits, and margaret miller straits. *Approach Soc. Res*.
1160 (1993):260–77.
1161 65. Van Eck van der Sluijs JF, de Vroegte L, van Manen AS, Rijnders CAT, van
1162 der Feltz-Cornelis CM. Complexity assessed by the INTERMED in patients
1163 1164

- 1141 with somatic symptom disorder visiting a specialized outpatient mental
1142 health care setting: a cross-sectional study. *Psychosomatics* (2017) 58:427–36.
1143 doi: 10.1016/j.psym.2017.02.008
1144 66. de Vroege L, Timmermans A, Kop WJ, and van der Feltz-Cornelis
1145 CM. Neurocognitive dysfunctioning and the impact of comorbid
1146 depression and anxiety in patients with somatic symptom and related
1147 disorders: a cross-sectional clinical study. *Psychol Med.* (2017) 4:1–11.
1148 doi: 10.1017/S0033291717003300
1149 67. Bailey PE, Henry JD. Alexithymia, somatization and negative
1150 affect in a community sample. *Psychiatry Res.* (2007) 150:13–20.
1151 doi: 10.1016/j.psychres.2006.05.024

Conflict of Interest Statement: The authors declare that the research was
1198 conducted in the absence of any commercial or financial relationships that could
1199 be construed as a potential conflict of interest.
1200

*Copyright © 2018 de Vroege, Emons, Sijtsma and van der Feltz-Cornelis. This is an
1201 open-access article distributed under the terms of the Creative Commons Attribution
1202 License (CC BY). The use, distribution or reproduction in other forums is permitted,
1203 provided the original author(s) and the copyright owner are credited and that the
1204 original publication in this journal is cited, in accordance with accepted academic
1205 practice. No use, distribution or reproduction is permitted which does not comply
1206 with these terms.*
1207

Q16

1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197

1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254