



This is a repository copy of *Study of the 3D Euler equations using Clebsch potentials: dual mechanisms for geometric depletion (vol 31, pg R25, 2018)*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/134288/>

Version: Accepted Version

---

**Article:**

Ohkitani, K. (2018) Study of the 3D Euler equations using Clebsch potentials: dual mechanisms for geometric depletion (vol 31, pg R25, 2018). *Nonlinearity*, 31 (8). p. 3973. ISSN 0951-7715

<https://doi.org/10.1088/1361-6544/aac6aa>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Addendum: “Study of the 3D Euler equations using Clebsch potentials: dual mechanisms for geometric depletion”

**Koji Ohkitani**

School of Mathematics and Statistics  
The University of Sheffield  
Hicks Building, Hounsfield Road Sheffield S3 7RH, U.K.  
E-mail: [K.Ohkitani@sheffield.ac.uk](mailto:K.Ohkitani@sheffield.ac.uk)

**Abstract.** After the publication of [1], it has come to the author’s attention that a class of Clebsch potentials for the Kida-Pelz flow, similar to what was derived in Appendix **B** of [1], has been studied in detail in [2]. We also note that there are typos in the formulas for one such example in [3], and these are corrected in [1].

## References

- [1] OHKITANI, K. 2018 Study of the 3D Euler equations using Clebsch potentials: dual mechanisms for geometric depletion *Nonlinearity* **31**, R25–R51.
- [2] HE, P. AND YUE Y. 2016 Construction of initial vortex-surface fields and Clebsch potentials for flows with high-symmetry using first integrals *Phys. Fluids* **28**, 037101-1–13.
- [3] OHKITANI, K. 2008 A geometrical study of 3D incompressible Euler flows with Clebsch potentials a long-lived Euler flow and its power-law energy spectrum *Physica D* **237**, 2020–2027.