

This is a repository copy of *Clostridium difficile: Investigating Transmission Patterns Between Infected and Colonized Patients Using Whole Genome Sequencing.*

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/134934/

Version: Supplemental Material

Article:

Kong, LY, Eyre, DW, Corbeil, J et al. (18 more authors) (2019) Clostridium difficile: Investigating Transmission Patterns Between Infected and Colonized Patients Using Whole Genome Sequencing. Clinical Infectious Diseases, 68 (2). pp. 204-209. ISSN 1058-4838

https://doi.org/10.1093/cid/ciy457

© 2018, The Author(s). Published by Oxford University Press for the Infectious Diseases Society of America. This is an author produced version of a paper published in Clinical Infectious Diseases. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Supplementary	7 Table. Detailed	participation rates	among hospitals in	initial cohort.
Suppremental	Tuore. Detuneu	purificipation rates	unions noopituito m	minulai comort.

	Hospital 1 RVH	Hospital 2 HSL	Hospital 3 CHUS	Hospital 4 HMR	Hospital 5 HDQ	Hospital 6 OGH	Total
Admissions							12304
Eligible patients	1823	<mark>x</mark>	1326	<mark>x</mark>	2538	1326	9502
Participants (% eligible)	1078 (59)	<mark>932</mark>	861 (65)	<mark>1159</mark>	850 (33)	542 (41)	5422 (57)
Patients testing positive in original cohort (colonized, infected)	118 (63, 55)	82 (54, 28)	64 (42, 22)	171 (112, 59)	80 (64, 16)	53 (37, 16)	568 (372, 196)
Isolates successfully sequenced from original cohort (colonized, infected)	111 (62, 49)	62 (50, 12)	53 (34, 19)	155 (110, 45)	75 (62, 13)	46 (35, 11)	502 (353, 149)
Number of CDI cases not enrolled in study	77	N/A	0	N/A	N/A	N/A	N/A
Infection control CDI isolates sequenced	52	0	0	0	0	0	52
Total isolates successfully sequenced (colonized, infected)	163 (62, 101)	62 (50, 12)	53 (34, 19)	155 (110, 45)	75 (62, 13)	46 (35, 11)	554 (353, 201)
Proportion of all CDI cases included in analyses	77% (101/132)	N/A	100% (19/19)	N/A	N/A	N/A	N/A