



UNIVERSITY OF LEEDS

This is a repository copy of *From ideation to action: Differentiating between those who think about suicide and those who attempt suicide in a national study of young adults*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/135184/>

Version: Accepted Version

Article:

Wetherall, K, Cleare, S, Eschle, S et al. (4 more authors) (2018) From ideation to action: Differentiating between those who think about suicide and those who attempt suicide in a national study of young adults. *Journal of Affective Disorders*, 241. pp. 475-483. ISSN 0165-0327

<https://doi.org/10.1016/j.jad.2018.07.074>

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

From ideation to action: differentiating between those who think about suicide and those who attempt suicide in a national study of young adults

Karen Wetherall, Seonaid Cleare, Sarah Eschle, Eamonn Ferguson, Daryl B. O'Connor, Ronan E O'Carroll, Rory C O'Connor

Corresponding Author: Karen Wetherall, Suicidal Behaviour Research Laboratory, Institute of Health & Wellbeing, University of Glasgow, Glasgow, G12 0XH, T: 0141 232 2181, E: Karen.wetherall@glasgow.ac.uk

Abstract

Background: Although many suicide risk factors have been identified, there is still relatively little known about the factors that differentiate those who think about suicide from those who make a suicide attempt.

Aims: Using the integrated motivational-volitional model (IMV) of suicidal behaviour as a framework, this study hypothesised that i) motivational and volitional phase factors would differentiate non-suicidal controls from those who had a history of suicidal ideation or suicide attempts, and ii) only volitional phase factors would differentiate between those who had a history of suicidal ideation and those who had attempted suicide.

Method: The Scottish Wellbeing Study (n=3508) is a nationally representative study of young people (18-34 years) recruited throughout Scotland. Using multinomial regression analysis, three groups (non-suicidal control (n=2534), lifetime suicide ideation (n=498) and lifetime suicide attempt (n=403) groups) were compared on motivational and volitional phase variables.

Results: Consistent with the IMV model, motivational and volitional phase variables differentiated the control group from both the ideation and attempt groups. Only volitional phase variables differentiated between the suicide attempt group and the suicidal ideation group; with those reporting a suicide attempt being higher on acquired capability, mental imagery about death, impulsivity, and being more likely to know a friend who had made a suicide attempt. Having a family member or friend die by suicide or a family member attempt suicide did not differentiate between the groups.

Limitations: The findings were based on cross-sectional data derived from self-report measures.

Conclusions: These findings provide further support for the IMV model, and highlight potential targets for intervention and suicide risk assessment.

Keywords: suicide, integrated motivational-volitional model (IMV), ideation-to-action framework, theory

1. Introduction

Suicide is a global health problem, and although suicide affects people across the lifespan, it is the second leading cause of death of 16-29 year olds worldwide (World Health Organisation, 2014), as well as being the leading cause of death among people under 50 in the UK (Samaritans, 2017). Recent research has identified a wide range of social, psychological and biological factors that act to increase suicide risk (O'Connor and Nock, 2014), although these factors often do not distinguish between those who will think about suicide and those who will go on to act on suicidal thoughts (Klonsky and May, 2014). With around 60% of transitions from suicidal ideation to a first attempt occurring within a year of ideation onset (Nock et al., 2008), it is crucial that we identify factors that distinguish those whose suicidal thoughts may transition into suicidal behaviours (Kessler et al., 2005).

In light of this, recent models of suicidal behaviour have adopted an *ideation-to-action* framework, where the development of suicidal ideation and the transition to a suicide attempt are viewed as distinct processes (Klonsky et al., 2017). The first theoretical model to emphasise this distinction was the interpersonal-psychological theory of suicide (IPT; Joiner, 2005), proposing that suicidal desire (comprised of perceived burdensomeness and thwarted belongingness) alone was insufficient to lead to a serious suicide attempt/death by suicide. A suicidal individual must also have the capability to act upon that desire characterised by a lowered physical pain sensitivity and high fearlessness about death that overrides the instinct towards self-preservation (Joiner, 2005). Although there has been considerable evidence for the key premises underpinning the IPTS (Chu et al., 2017), a recent systematic review of IPTS studies found limited evidence for an interaction between perceived burdensomeness, thwarted belongingness and acquired capability in association

with suicide attempts, with the authors concluding that the relationships between the variables may be less straightforward than originally presented (Ma et al., 2016). Therefore, models of suicidal behaviour may need to account for a more complex relationship between suicidal ideation and the transition to a suicide attempt.

[Figure 1]

In this vein, the integrated motivational-volitional model of suicidal behaviour (IMV; O'Connor, 2011) was proposed in 2011 and refined in 2018 (O'Connor and Kirtley, 2018). The IMV model is a tri-partite framework (Figure 1) that builds upon previous theories to map the context in which suicide may occur (the pre-motivational phase), the development of suicidal ideation (the motivational phase) and the transition of suicidal thoughts into suicidal behaviours (the volitional phase). Building upon the *cry of pain hypothesis* (Williams, 1997), the motivational phase focuses on feelings of defeat and entrapment as the key drivers of suicidal ideation. Importantly for the present study, within the final phase of the model (volitional phase), it is argued that a group of factors, labelled volitional moderators, governs the transition from thinking about suicide to attempting suicide. In addition to Joiner's concept of acquired capability, these factors include impulsivity, planning, exposure to the suicidal acts of others, access to means, past suicidal behaviour and mental imagery about death (O'Connor and Kirtley, 2018).

There has been support for the main facets of the IMV model (e.g., Dhingra et al., 2016; O'Connor et al., 2013; Wetherall et al., 2018), including a growing body of evidence demonstrating that volitional moderators do indeed differentiate between those who think about suicide and those who engage in suicidal behaviour (O'Connor et al., 2016; O'Connor and Kirtley, 2018). For example, in one study of adolescents, only volitional phase variables

(self-harm by friends and family, thinking about peers' self-harm, impulsivity) and stress differentiated between those with thoughts of self-harm and those who engaged in self-harm (O'Connor et al., 2012). Similarly, in a test of the IMV facets with students, within a multivariable model, only the volitional phase factors (exposure to suicide, impulsivity and fearlessness about death) distinguished between those who had experienced suicidal ideation and those who had attempted suicide (Dhingra et al., 2015). Additionally, in a recent cohort study, exposure to the self-harm of others (alongside psychiatric disorder) was key to differentiating between adolescents who had made a suicide attempt compared to those who had thought about but not attempted suicide (Mars et al., 2018).

A final model utilising the *ideation-to-action* framework is the more recent three-step theory (3ST; Klonsky and May, 2015). The initial steps tap the development and escalation of suicidal ideation with a combination of pain, hopelessness and a lack of connectedness, and in the final step ideation progresses to an attempt when the capability for suicide is present. The concept of acquired capability has been a consistent component across all three models explored, with recent evidence suggesting that when those high on capability become agitated, suicidal intensity increases, thereby facilitating suicidal behaviour by providing sufficient energy and arousal (Ribeiro et al., 2015). Therefore, this concept, along with the additional volitional factors of impulsivity, exposure to suicide and mental imagery about death, are key variables to be explored more fully as factors that can differentiate those who think about suicide from those who will make a suicide attempt.

Current study

This study aimed to investigate a key premise of the IMV model; namely that volitional phase variables govern the transition from suicidal ideation to suicide attempts when

motivational phase variables are controlled for (O'Connor & Kirtley, 2018). Although a small number of studies have investigated the psychological factors associated with behavioural enactment (e.g., Dhingra et al., 2015), to our knowledge this is the most detailed study of its kind and the first study to do so in a nationally representative sample. To this end, the Scottish Wellbeing Study (O'Connor et al., 2018), a nationally representative interview-based survey of young adults aged 18 to 34 years across Scotland, was conducted. In short, we hypothesised that i) motivational and volitional phase factors would differentiate non-suicidal controls from those who had a history of suicidal ideation or suicide attempts, and ii) only volitional phase factors would differentiate between those who had a history of suicidal ideation and those who had attempted suicide in a multivariable model.

2. Method

2.1 Sample and procedure

The data are from the Scottish Wellbeing Study (O'Connor et al., 2018) which is a nationally representative sample of young people aged 18 to 34 years (n=3508) from across Scotland. Recruitment was conducted by Ipsos MORI, a social research organisation, between 25th March 2013 and 12th December 2013. A quota sampling methodology was utilised; quotas were based on age (three quota groups), sex and working status (for more details, see O'Connor et al., 2018). Following written consent, participants completed an hour-long interview, carried out face-to-face in their homes, using Computer Assisted Personal Interviewing (CAPI), with confidential completion of sensitive questions (including suicidal history) on a personal computer. Participants were compensated £25 for their time. Ethical approval was obtained from the University of Stirling (Psychology Department) ethics

committee as well as from the US Department of Defense Human Research Protections Office.

2.2 Measures

2.2.1 Outcome measure: Lifetime history of suicidal ideation and attempts. This was assessed with two items drawn from the Adult Psychiatric Morbidity Survey (APMS; McManus et al., 2007): “Have you ever seriously thought of taking your life, but not actually attempted to do so?” and “Have you ever made an attempt to take your life, by taking an overdose of tablets or in some other way?”. Responses to these questions were “no”, “yes” or “would rather not say”. These items were used to create a 3 category variable indicating if participants had (i) no history of suicidal ideation/ attempt (control group), (ii) had experienced suicidal ideation but had never attempted suicide (suicidal ideation group), or (iii) had reported having attempted suicide in the past (suicidal attempt group).

2.2.2 Motivational phase risk factors

2.2.2.1 Defeat. The Defeat Scale (Gilbert & Allan, 1998) is a 16-item self-report measure of perceived failed struggle and loss of rank (e.g., “I feel that I have not made it in life”). This scale has good psychometric properties and is significantly correlated with depressive symptoms (Griffiths et al., 2014). In the present study the measure had high internal reliability (Cronbach’s $\alpha = 0.96$).

2.2.2.2 Entrapment. The 16-item Entrapment Scale (Gilbert & Allan, 1998) is a measure of the sense of being unable to escape feelings of defeat and rejection (e.g., I am in a situation I feel trapped in). This measure consists of 10 items reflecting external entrapment (entrapment by external situations), and 6 items tapping internal entrapment (entrapment

by one's own thoughts and feelings). The scale has good psychometric properties (Griffiths et al., 2014) and demonstrated high internal consistency in the present study (Cronbach's $\alpha = 0.96$).

2.2.2.3 Perceived burdensomeness and thwarted belongingness. These were assessed using the 12-item Interpersonal Needs Questionnaire (INQ; Van Orden et al., 2012). The INQ includes 7-items to tap burdensomeness (e.g., "I feel like a burden on the people in my life") and 5-items to assess belongingness (e.g., "I feel disconnected from other people"). The scales have been shown to have good internal consistency and construct validity (Van Orden et al., 2012), including in this study (perceived burdensomeness Cronbach's $\alpha = 0.87$, thwarted belongingness Cronbach's $\alpha = 0.84$).

2.2.2.4 Goal Disengagement and Goal Reengagement. The 10-item goal adjustment scale (GAS; Wrosch et al., 2003) consists of a 4-item goal disengagement (e.g., "If I have to stop pursuing an important goal in my life its easy for me to stop thinking about the goal and let it go") subscale and a 6-item goal reengagement (e.g., "If I have to stop pursuing an important goal in my life I start working on other new goals") subscale. Both subscales have shown good validity (Wrosch et al., 2003), and in the present study they had adequate to good internal consistency (goal disengagement Cronbach's $\alpha = 0.70$, goal reengagement Cronbach's $\alpha = 0.87$).

2.2.2.5 Social support. The 7-item ENRICHD Social Support Instrument (ESSI; Mitchell et al., 2003), taps four defining attributes of social support: emotional, instrumental, informational, and appraisal (e.g., "Is there someone available to give you good advice about a problem?"). It has been found to be a valid and reliable measure of social support

(Vaglio et al., 2004), and displayed good internal reliability in the present study (Cronbach's $\alpha = 0.87$).

2.2.2.6 Resilience. Resilience was measured using the 10-item Brief Resilience Scale (BRS; Campbell-Sills and Stein, 2007), adapted from the 25-item Connor-Davidson Resilience Scale (CD-RISC; Connor and Davidson, 2003). This 10-item version (e.g., "Coping with stress can strengthen me") has good psychometric properties and is highly correlated with the original 25-item version (Campbell-Sills and Stein, 2007), and in the present study it displayed excellent internal consistency (Cronbach's $\alpha = 0.90$).

2.2.3 Volitional phase risk factors

2.2.3.1 Acquired capability. The Acquired Capability for Suicide Scale (ACSS; Van Orden et al., 2008) is a 5-item measure designed to assess one's fearlessness about death and physical pain sensitivity (e.g., "The pain involved in dying frightens me"). The scale has demonstrated convergent and discriminant validity (Van Orden et al., 2008), and in this study the ACSS had a relatively low internal consistency of 0.63 (Cronbach's α).

2.2.3.2 Impulsivity. This was assessed using the 30-item Barratt Impulsiveness Scale Version 11 (BIS-11; Patton et al., 1995); a self-report questionnaire that accounts for the multi-faceted nature of the construct (i.e., attentional, motor and non-planning impulsiveness) that provides a general impulsiveness score (e.g., "I act on the spur of the moment"). The BIS is a commonly used scale that has been shown to correlate with behavioural measures of impulsivity (Martins et al., 2004), and it displayed good internal validity in the present study (Cronbach's $\alpha = 0.83$).

2.2.3.3 Mental imagery. Eight questions were asked to establish the frequency with which participants imagine death related imagery when they feel down or distressed, including engaging in self-harm or suicidal behaviour (e.g., "...images of yourself planning/preparing to harm yourself or make a suicide attempt"). Greater presence of suicide-related imagery has been linked to higher levels of suicidal ideation (Holmes et al., 2007). The scale displayed good internal reliability (Cronbach's $\alpha = 0.84$).

2.2.3.4 Exposure to suicide. Participants were asked three items to establish whether they had friends or family who attempted or died by suicide (e.g., "Has anyone among your family attempted suicide?"). These items have been used in previous research (O'Connor et al., 2012) and have been shown to differentiate between those who think about suicide and those who attempt suicide (Dhingra et al., 2015).

2.2.4 Covariates: Demographic characteristics and mood

2.2.4.1 Demographic characteristics. We recorded the following demographic information: age, gender, marital status (married vs. not married), ethnicity (white vs. non-white) and economic activity (employed, inactive and unemployed).

2.2.4.2 Depressive symptoms. The Beck Depression Inventory-II (BDI-II; Beck et al., 1996) is a well-established measure tapping a range of depressive symptoms (e.g., self-dislike, loss of energy) containing 21 items. It has been shown to yield reliable, internally consistent, and valid scores in many different populations (e.g., Dozois et al., 1998), and in this study, it displayed high internal reliability (Cronbach's $\alpha = 0.95$).

2.3 Statistical analysis

Data analysis was conducted using SPSS version 22. The missing data included items missed by participants and participants selecting 'would rather not say'. We used every participant's data as long as they had completed 75% or more of a psychological scale, this resulted in minimal missing data, <1% on any variable (range 0.31% to 0.86%; including those who had refused). These small amounts of missing data were checked against demographic characteristics and as there were no significant associations, expectation maximisation (EM) was applied to replace missing items for each scale. The multinomial regression model included only those who completed >75% of every measure (n=3330; 95% of total sample), with a small proportion of the data EM replaced. More information on the EM replacement method is included in the supplementary materials.

Additionally, the data were weighted to ensure that the attained sample based on the quota variables was in line with the population in the sample frame using rim weighting. Overall, as the quotas were almost always met (30-34 year olds, full-time students and full-time workers were slightly under-represented) the effect of the weights was small. All analyses and reporting of data were conducted with the weights on. More information on the rim weighting is included in the supplementary materials.

To investigate the respective influence of the motivational and volitional phase variables, initial univariate multinomial regression analyses were conducted (See Table S1 in the supplementary materials). To control for the number of comparisons the Holm-Bonferroni correction method (Holm, 1979) was applied. In order to identify which variables independently distinguished between the groups, a multivariable multinomial logistic regression was performed. Specifically, demographic and mood variables were entered as covariates (age, gender, marital status, ethnicity, economic activity and depressive

symptoms), followed by the motivational phase variables (defeat, entrapment, perceived burdensomeness, thwarted belongingness, goal disengagement, goal reengagement, social support and resilience) and then the volitional phase variables (acquired capability, impulsivity, mental images, exposure to suicide death (family & friend), exposure to suicide attempt by friend, exposure to suicide attempt by family) were entered. Odds ratios (OR) indicating the likelihood of each variable's association with the higher risk group were reported (i.e., the ideation and attempt groups relative to the controls, and the attempt group relative to the ideation group), with those greater than one indicating increased risk and less than one decreased risk.

To better understand how well the volitional phase measures distinguish between those who have thought of suicide only and those who have made a suicide attempt at an individual level, the sensitivity (i.e., proportion of the sample high on a volitional phase variable that were correctly identified as having made a suicide attempt) and specificity (i.e., the proportion of the sample that were low on a volitional phase variables and had not made a suicide attempt) of each of the volitional phase variables is reported, along with their positive predictive value (i.e., the probability that the individual high on a volitional phase variable had attempted suicide) and negative predictive value (i.e., the probability that the individual low on a volitional phase variable had not attempted suicide). A cut-off score (mean +1SD) was created for the continuous variables to indicate those 'high' and 'low' on a particular volitional phase variable.

3. Results

3.1 Sample characteristics

In the primary analysis (n=3330), the majority of the sample had no suicidal history (n=2470; 74.6%), 14.3% (n=481) had experienced suicidal ideation in their lifetime but had never made a suicide attempt, and 11% (n=379) had attempted suicide in their lifetime. The descriptive statistics by group membership (i.e., ideation vs attempt vs control) and univariate differences for those who responded to the suicidal history questions (n=3435) are provided in Table 1. With demographics, the univariate multinomial regression analyses indicated that those with suicidal ideation were more likely to be male, not married and unemployed compared to controls, and those who had reported a suicide attempt were more likely to be female, older and unemployed than both the controls and those in the suicidal ideation group.

Members of the control group scored significantly lower on all of the psychological risk factors compared to those in the suicide ideation and suicide attempt groups; this included depressive symptoms, defeat, entrapment, acquired capability and impulsivity. Those in the suicide attempt group reported more frequent exposure to the suicidal behaviour of others, with almost 50% having been exposed to a friend making a suicide attempt, compared to just 16% for the control group. The control group reported higher levels of protective factors such as resilience and social support. A similar pattern emerged between the two suicidal history groups; those in the suicide attempt group more strongly endorsed the motivational and volitional phase risk factors compared to those in the suicide ideation group.

[Insert Table 1 about here]

3.2 Multivariable multinomial regression analyses

The results of the multinomial regression analyses are presented in Table 2. The model was statistically significant ($\chi^2(42) = 1528.60, p < 0.001$; pseudo R-square (Cox and Snell) = 0.37;).

Those in the control group were significantly lower than both suicidal history groups on a combination of motivational (defeat and burdensomeness) and volitional phase factors (acquired capability, mental images, exposure to suicide attempt by family or friend).

Additionally, those in the suicide attempt group were more likely to be female, older, and higher on impulsivity than controls. Depressive symptoms did not distinguish between any of the groups when all motivational and volitional factors were accounted for.

Similarly, those who reported a suicide attempt were older (OR=1.07 [95% CI= 1.03-1.10]) and more likely to be female (OR=0.49 [95% CI= 0.36-0.67]) than those in the ideation group. However, consistent with the IMV model, the only psychological factors that distinguished those in the suicide attempt group from those in the suicidal ideation group were volitional phase variables; none of the mood or motivational phase variables significantly differentiated between these groups. In comparison to those in the suicidal ideation group, those who reported a suicide attempt scored significantly higher on levels of acquired capability (OR=1.10 [95% CI= 1.06-1.14]), impulsivity (OR=1.02 [95% CI= 1.01-1.04]), mental images about death (OR=1.07 [95% CI= 1.03-1.10]) and they were significantly more likely to have been exposed to a suicide attempt of a friend (OR=1.49 [95% CI= 1.09-2.06]).

[Insert Table 2 about here]

3.3 Sensitivity and specificity of the volitional phase variables in differentiating between suicide ideation and suicide attempts groups

The findings of the sensitivity and specificity analyses are displayed in Table 3. High acquired capability and impulsivity scores and exposure to a suicide death identified those who had made a suicide attempt just over one third of the time, and exposure to a family member or a friend making a suicide attempt identified two fifths of those who had made a suicide attempt. Mental imagery was the most sensitive factor, identifying over half of those who had made an attempt. The specificity of the variables was higher (range 61.9 – 77.4%) indicating that generally being low on any of the volitional phase variables helped identify those who had not made a suicide attempt. All the volitional variables, when taken together, identified around 45% of those who had made a suicide attempt, and nearly 80% of those who had not. The positive predictive values (PPV) were relatively similar across the individual volitional variables (range 39.6% - 50.0%), as were the negative predictive values (NPV; range 43.1%-49.4%); indicating that the probability of individuals high (or low) on any of the volitional phase variables having attempted suicide (or not) was similar across the variables. The PPV and NPV both increased when all volitional variables were taken into account. Additionally, the variance explained by the volitional variables taken together was around 11%.

[Insert Table 3 about here]

4. Discussion

We tested a key premise of the integrated motivational-volitional model (IMV, O'Connor, 2011; O'Connor & Kirtley, 2018), namely that volitional phase factors are key to governing the transition from suicidal ideation to a suicide attempt. We hypothesised that i) motivational and volitional phase factors would differentiate non-suicidal controls from those who had a history of suicidal ideation or suicide attempts, and ii) only volitional phase

factors would differentiate between those who had a history of suicidal ideation and those who had attempted suicide in a multivariable analysis. Findings yielded clear evidence in support of both hypotheses. Specifically, a combination of motivational and volitional phase variables distinguished the control group from both the suicide ideation group and the suicide attempt group. Whereas, apart from some demographic differences (those in the attempt group being older and female), only volitional phase variables differentiated between those with a history of suicidal ideation and those who had reported a suicide attempt; with the latter group reporting higher levels of acquired capability, impulsivity, mental imagery about death and they were more likely to have been exposed to the suicide attempt of a friend.

This study adds to the growing literature highlighting the importance of the volitional phase factors within the IMV model (e.g., O'Connor et al., 2012; Dhingra et al., 2015) and the *ideation-to-action* framework more generally (Klonsky et al., 2017). It is also unique as it is the first study of its kind to investigate the role of volitional phase factors in a large, nationally representative sample. Although motivational phase variables, including key components of IPT (e.g., perceived burdensomeness) and the IMV model (e.g., defeat), are useful to identify who may think of suicide, they are not key drivers of behavioural enactment. In light of the recent concerns that most risk factors do not distinguish between those suicidal individuals who are/are not at increased risk of making a suicide attempt (Klonsky and May, 2014), the present volitional phase findings are important as they address this dearth in the research literature. In addition, they highlight potential targets for interventions and therapies, as well as for risk assessment, consistent with a recent call to action to identify more specific markers of suicide risk (Holmes et al., 2018). Although,

consistent with other research in this area (Quinlivan et al., 2017; Steeg et al., 2018), the sensitivity of the individual volitional phase variables is relatively low, therefore limiting their utility in assessing risk at an individual level. Nonetheless, the volitional phase variables should be used to inform treatment targets within a clinical formulation.

Consistent with previous findings (e.g., Dhingra et al., 2015; Mars et al., 2018), exposure to suicide in others, in particular to the suicide attempt of a friend, was most strongly associated with belonging to the suicide attempt group. Contrary to our predictions, the other exposure variables of suicide attempt by family member or death by suicide of either a family member or a friend, did not significantly differentiate between those in the suicidal ideation and the suicide attempt groups. It would be useful to explore why these other types of exposure did not differentiate between the groups. Interestingly, Mars et al. (2018) found a dose response effect with adolescents, whereby exposure to self-harm in both family and friends was 5 times higher in their suicide attempt group compared to those reporting suicide ideation only. A number of mechanisms have been suggested to explain this relationship; including that exposure to suicidal peers increases risk due to suicide modelling via social learning (Insel & Gould, 2008) and cognitive accessibility (Biddle et al., 2012). Contagion may also be more likely due to assortative relating processes whereby similar individuals are more likely to associate (Joiner, 2003), and there may even be evidence for a genetic basis to imitation (Brent and Melhem, 2008). Although further research is needed to better understand the mechanisms behind this phenomenon, ultimately the present study highlights the importance of exposure to suicide as a key risk factor for a suicide attempt.

Additionally, recent research suggests that exposure to suicidal or self-harming behaviours may act as painful and provocative life experiences which feed into acquired capability (Klonsky et al., 2017). Although measures of acquired capability were only weakly associated with suicide attempt history in a recent meta-analysis (Chu et al., 2017), the concept of having to override an innate instinct for survival appears important in understanding the transition to a suicide attempt (Klonsky and May, 2015). Specifically, having fearlessness about death and reduced pain sensitivity appear to be important mechanisms in increasing the ability to act upon one's thoughts of suicide (Smith et al., 2010). Indeed, Kirtley and colleagues (2016) in a systematic review found a pervasive relationship between lower pain sensitivity and self-harm more generally but highlighted the dearth of research in this area (Kirtley et al., 2016). A better understanding of how capability for suicide develops requires urgent attention, in particular whether its effects can be buffered by protective interventions such as safety planning (Stanley and Brown, 2012).

Impulsivity could also increase acquired capability through more exposure to painful events (Anestis et al., 2014). Although impulsivity is an established risk factor, traditionally thought to facilitate suicidal behaviours by increasing the likelihood of enacting suicidal thoughts (Mann et al., 1999), more recent findings have questioned the nature of this relationship. As in this study, a meta-analysis found the relationship between trait impulsivity and suicidal behaviour was relatively small (Anestis et al., 2014). Arguably, the research fails to differentiate between state and trait impulsivity; as an individual high in trait impulsivity may plan a suicide attempt (and vice versa) (Gvion and Apter, 2011). Therefore, impulsivity remains a problematic concept that may be difficult to target in interventions; trait

impulsivity may not accurately reflect the individual's suicidal intentions, but from a clinician's perspective it may be useful to be aware of this.

The finding that mental imagery related to death distinguishes between those who have made a suicide attempt and those who have not is important and novel, and as it was also the most sensitive factor in identifying who had made a suicide attempt therefore may potentially be the most useful for assessing suicide risk. Our finding is consistent with Holmes et al. (2007) who found that 'flash forwards', defined as imagined future acts of suicide or self-harm are associated with suicide risk. They may be important targets for intervention, with evidence showing that a reduction in suicidal imagery is associated with less suicidal thoughts over time (Ng et al., 2016). However, to be effective, the key mechanisms need to be explored further as there is competing evidence. For example, it has been suggested that imagery increases the cognitive availability of powerful images (Florentine and Crane, 2010), potentially leading to more distress (Holmes and Mathews, 2005); however, for some the images may also function as a deterrent for suicidal behaviour (Crane et al., 2012). In contrast, it is also suggested that habituation may occur, whereby the fear of the (suicidal) act is reduced thereby facilitating behavioural enactment (Crane et al., 2012). In short, we need to advance our understanding of how experiencing suicide 'flash forwards' increases suicide risk, and then how best to intervene to reduce suicide risk.

Limitations

Although this study had many strengths, a number of potential limitations should be noted. Firstly, the data were cross-sectional; therefore causality or directionality cannot be inferred. Secondly, as with much psychological research, the measures here are reliant on self-report, therefore they are subject to memory and reporting biases. Indeed, suicidal

ideation in particular may be subject to mis-reporting (Mars et al., 2016), and as the former was assessed using a single item, we were not able to tap the intensity or severity of thoughts. Thirdly, although the sample was representative of young people across Scotland, it may not be generalisable to other populations, in particular to clinical groups who are at increased risk of suicidal behaviour. Finally, as the effect sizes of the volitional phase variables were relatively small (OR range = 1.02 – 1.49) their potential to predict suicide risk in clinical practice may be limited; this is supported by the relatively low sensitivity of the volitional variables.

Despite these limitations, the current research is unique and represents the most robust test to date of the volitional phase of the integrated motivational-volitional model of suicidal behaviour (O'Connor et al., 2016). In the multivariable analyses, only volitional phase factors (acquired capability, exposure to a friend's suicide attempt, mental imagery and impulsivity) differentiated between those who reported suicide ideation and those who reported a lifetime suicide attempt. It extends our understanding of the factors which aid the transition from suicidal thoughts to attempts and it provides strong support for the ideation-to-action framework (Klonsky et al., 2017). As highlighted, future research would benefit from more prospective studies utilising high-risk populations, as well as further exploration of how these particular volitional factors emerge, how best to incorporate them into risk assessment protocols and how to optimally target them in interventions.

References

- Anestis, M.D., Soberay, K.A., Gutierrez, P.M., Hernandez, T.D., Joiner, T.E., 2014. Reconsidering the link between impulsivity and suicidal behavior. *Pers Soc Psychol Rev* 18, 366-386.

- Beck, A.T., Steer, R.A., Brown, G.K., 1996. BDI-II, Beck depression inventory : manual. Psychological Corp. ; Harcourt Brace, San Antonio, Tex.; Boston.
- Biddle, L., Gunnell, D., Owen-Smith, A., Potokar, J., Longson, D., Hawton, K., Kapur, N., Donovan, J., 2012. Information sources used by the suicidal to inform choice of method. *J Affect Disord* 136, 702-709.
- Brent, D.A., Melhem, N., 2008. Familial transmission of suicidal behavior. *Psychiatr Clin North Am* 31, 157-177.
- Campbell-Sills, L., Stein, M.B., 2007. Psychometric analysis and refinement of the Connor-Davidson Resilience Scale (CD-RISC): Validation of a 10-item measure of resilience. *Journal of Traumatic Stress* 20, 1019-1028.
- Chu, C., Buchman-Schmitt, J.M., Stanley, I.H., Hom, M.A., Tucker, R.P., Hagan, C.R., Rogers, M.L., Podlogar, M.C., Chiurliza, B., Ringer, F.B., Michaels, M.S., Patros, C.H.G., Joiner, T.E., 2017. The interpersonal theory of suicide: A systematic review and meta-analysis of a decade of cross-national research. *Psychol Bull* 143, 1313-1345.
- Connor, K.M., Davidson, J.R.T., 2003. Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depression and Anxiety* 18, 76-82.
- Dhingra, K., Boduszek, D., O'Connor, R.C., 2016. A structural test of the Integrated Motivational-Volitional model of suicidal behaviour. *Psychiatry Research* 239, 169-178.
- Dhingra, K., Boduszek, D., O'Connor, R.C., 2015. Differentiating suicide attempters from suicide ideators using the Integrated Motivational–Volitional model of suicidal behaviour. *Journal of Affective Disorders* 186, 211-218.
- Dozois, D.J.A., Dobson, K.S., Ahnberg, J.L., 1998. A psychometric evaluation of the Beck Depression Inventory–II. American Psychological Association, US, pp. 83-89.

Florentine, J.B., Crane, C., 2010. Suicide prevention by limiting access to methods: a review of theory and practice. *Soc Sci Med* 70, 1626-1632.

Griffiths, A.W., Wood, A.M., Maltby, J., Taylor, P.J., Tai, S., 2014. The prospective role of defeat and entrapment in depression and anxiety: A 12-month longitudinal study. *Psychiatry Research* 216, 52-59.

Gvion, Y., Apter, A., 2011. Aggression, impulsivity, and suicide behavior: a review of the literature. *Arch Suicide Res* 15, 93-112.

Holm, S., 1979. A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal of Statistics* 6, 65-70.

Holmes, E.A., Crane, C., Fennell, M.J., Williams, J.M., 2007. Imagery about suicide in depression--"Flash-forwards"? *J Behav Ther Exp Psychiatry* 38, 423-434.

Holmes, E.A., Ghaderi, A., Harmer, C.J., Ramchandani, P.G., Cuijpers, P., Morrison, A.P., Roiser, J.P., Bockting, C.L.H., O'Connor, R.C., Shafran, R., Moulds, M.L., Craske, M.G., The *Lancet Psychiatry* Commission on psychological treatments research in tomorrow's science. *The Lancet Psychiatry* 5, 237-286.

Holmes, E.A., Mathews, A., 2005. Mental imagery and emotion: a special relationship? *Emotion* 5, 489-497.

Insel, B.J., Gould, M.S., 2008. Impact of modeling on adolescent suicidal behavior. *Psychiatr Clin North Am* 31, 293-316.

Joiner, T., 2005. *Why people die by suicide*. Harvard University Press.

Joiner, T.E., 2003. Contagion of suicidal symptoms as a function of assortative relating and shared relationship stress in college roommates. *Journal of Adolescence* 26, 495-504.

Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., Walters, E.E., 2005. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62, 593-602.

Kirtley, O.J., O'Carroll, R.E., O'Connor, R.C., 2016. Pain and self-harm: A systematic review. *J Affect Disord* 203, 347-363.

Klonsky, E.D., May, A.M., 2014. Differentiating suicide attempters from suicide ideators: a critical frontier for suicidology research. *Suicide Life Threat Behav* 44, 1-5.

Klonsky, E.D., May, A.M., 2015. The Three-Step Theory (3ST): A New Theory of Suicide Rooted in the "Ideation-to-Action" Framework. *International Journal of Cognitive Therapy* 8, 114-129.

Klonsky, E.D., Saffer, B.Y., Bryan, C.J., 2017. Ideation-to-Action Theories of Suicide: A Conceptual and Empirical Update. *Current Opinion in Psychology*.

Ma, J., Batterham, P.J., Calear, A.L., Han, J., 2016. A systematic review of the predictions of the Interpersonal-Psychological Theory of Suicidal Behavior. *Clin Psychol Rev* 46, 34-45.

Mann, J.J., Waternaux, C., Haas, G.L., Malone, K.M., 1999. Toward a clinical model of suicidal behavior in psychiatric patients. *Am J Psychiatry* 156, 181-189.

Mars, B., Cornish, R., Heron, J., Boyd, A., Crane, C., Hawton, K., Lewis, G., Tilling, K., Macleod, J., Gunnell, D., 2016. Using Data Linkage to Investigate Inconsistent Reporting of Self-Harm and Questionnaire Non-Response. *Arch Suicide Res* 20, 113-141.

Mars, B., Heron, J., Klonsky, E.D., Moran, P., O'Connor, R.C., Tilling, K., Wilkinson, P., Gunnell, D., 2018. What distinguishes adolescents with suicidal thoughts from those who have attempted suicide? A population-based birth cohort study. *Journal of Child Psychology and Psychiatry*.

Martins, S.S., Tavares, H., da Silva Lobo, D.S., Galetti, A.M., Gentil, V., 2004.

Pathological gambling, gender, and risk-taking behaviors. *Addict Behav* 29, 1231-1235.

McManus, S., Meltzer, H., Brugha, T., Bebbington, P., Jenkins, R., 2007. Adult psychiatric morbidity in England, 2007. Results of a household survey, Leeds: NHS Information Centre for Health and Social Care.

Mitchell, P.H., Powell, L., Blumenthal, J., Norten, J., Ironson, G., Pitula, C.R., Froelicher, E.S., Czajkowski, S., Youngblood, M., Huber, M., Berkman, L.F., 2003. A short social support measure for patients recovering from myocardial infarction: the ENRICH Social Support Inventory. *J Cardiopulm Rehabil* 23, 398-403.

Ng, R.M.K., Di Simplicio, M., McManus, F., Kennerley, H., Holmes, E.A., 2016. 'Flash-forwards' and suicidal ideation: A prospective investigation of mental imagery, entrapment and defeat in a cohort from the Hong Kong Mental Morbidity Survey. *Psychiatry Res* 246, 453-460.

Nock, M.K., Borges, G., Bromet, E.J., Alonso, J., Angermeyer, M., Beautrais, A., Bruffaerts, R., Chiu, W.T., de Girolamo, G., Gluzman, S., de Graaf, R., Gureje, O., Haro, J.M., Huang, Y., Karam, E., Kessler, R.C., Lepine, J.P., Levinson, D., Medina-Mora, M.E., Ono, Y., Posada-Villa, J., Williams, D., 2008. Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *The British Journal of Psychiatry* 192, 98.

O'Connor, R.C., Cleare, S., Eschle, S., Wetherall, K., Kirtley, O.J., 2016. The Integrated Motivational-Volitional Model of Suicidal Behaviour, *The International Handbook of Suicide Prevention*, pp. 220 - 240.

O'Connor, R.C., Rasmussen, S., Hawton, K., 2012. Distinguishing adolescents who think about self-harm from those who engage in self-harm. *British Journal of Psychiatry* 200, 330-335.

O'Connor, R.C., Smyth, R., Ferguson, E., Ryan, C., Williams, J.M.G., 2013. Psychological Processes and Repeat Suicidal Behavior: A Four-Year Prospective Study. *Journal of Consulting and Clinical Psychology* 81, 1137-1143.

O'Connor, R.C., Kirtley, O.J., 2018. The Integrated Motivational-Volitional Model of Suicidal Behaviour. . *Philosophical Transactions of the Royal Society B*.

O'Connor, R.C., Nock, M.K., 2014. The psychology of suicidal behaviour. *The Lancet Psychiatry* 1, 73-85.

O'Connor, R.C., Wetherall, K., Cleare, S., Eschle, S., Drummond, J., Ferguson, E., O'Connor, D.B., O'Carroll, R.E., in press. Suicide attempts and non-suicidal self-harm: a national prevalence study of young adults. *BJPsych Open*.

Patton, J.H., Stanford, M.S., Barratt, E.S., 1995. Factor structure of the Barratt impulsiveness scale. *J Clin Psychol* 51, 768-774.

Quinlivan, L., Cooper, J., Meehan, D., Longson, D., Potokar, J., Hulme, T., Marsden, J., Brand, F., Lange, K., Riseborough, E., Page, L., Metcalfe, C., Davies, L., Connor, R., Hawton, K., Gunnell, D., Kapur, N., 2017. Predictive accuracy of risk scales following self-harm: multicentre, prospective cohort study. *The British Journal of Psychiatry* 210, 429.

Ribeiro, J.D., Bender, T.W., Buchman, J.M., Nock, M.K., Rudd, M.D., Bryan, C.J., Lim, I.C., Baker, M.T., Knight, C., Gutierrez, P.M., Joiner, T.E., Jr., 2015. An investigation of the interactive effects of the capability for suicide and acute agitation on suicidality in a military sample. *Depress Anxiety* 32, 25-31.

Smith, P.N., Cukrowicz, K.C., Poindexter, E.K., Hobson, V., Cohen, L.M., 2010. The acquired capability for suicide: a comparison of suicide attempters, suicide ideators, and non-suicidal controls. *Depress Anxiety* 27, 871-877.

Snowcroft, E., 2017. Samaritans: suicide statistics report 2017 (Including data for 2013-2015). Samaritans.

Stanley, B., Brown, G.K., 2012. Safety Planning Intervention: A Brief Intervention to Mitigate Suicide Risk. *Cognitive and Behavioral Practice* 19, 256-264.

Steeg, S., Quinlivan, L., Nowland, R., Carroll, R., Casey, D., Clements, C., Cooper, J., Davies, L., Knipe, D., Ness, J., O'Connor, R.C., Hawton, K., Gunnell, D., Kapur, N., 2018. Accuracy of risk scales for predicting repeat self-harm and suicide: a multicentre, population-level cohort study using routine clinical data. *BMC Psychiatry* 18, 113.

Vaglio, J., Conard, M., Poston, W.S., O'Keefe, J., Haddock, C.K., House, J., Spertus, J.A., 2004. Testing the performance of the ENRICH Social Support Instrument in cardiac patients. *Health and Quality of Life Outcomes* 2, 24-24.

Van Orden, K.A., Cukrowicz, K.C., Witte, T.K., Joiner, T.E., 2012. Thwarted Belongingness and Perceived Burdensomeness: Construct Validity and Psychometric Properties of the Interpersonal Needs Questionnaire. *Psychological Assessment* 24, 197-215.

Van Orden, K.A., Witte, T.K., Gordon, K.H., Bender, T.W., Joiner Jr, T.E., 2008. Suicidal desire and the capability for suicide: Tests of the interpersonal-psychological theory of suicidal behavior among adults. *Journal of Consulting and Clinical Psychology* 76, 72-83.

Wetherall, K., Robb, K.A., O'Connor, R.C., 2018. An Examination of Social Comparison and Suicide Ideation Through the Lens of the Integrated Motivational-Volitional Model of Suicidal Behavior. *Suicide Life Threat Behav.*

Williams, J.M.G., 1997. *Cry of pain: Understanding suicide and self harm*. Penguin, Harmondsworth, England.

World Health Organisation, 2014. Preventing Suicide: A global imperative, http://www.who.int/mental_health/suicide-prevention/world_report_2014/en/.

Wrosch, C., Scheier, M.F., Miller, G.E., Schulz, R., Carver, C.S., 2003. Adaptive self-regulation of unattainable goals: goal disengagement, goal reengagement, and subjective well-being. *Pers Soc Psychol Bull* 29, 1494-1508.

Figure 1: The integrated motivational-volitional model of suicidal behaviour (O'Connor et al., 2011; O'Connor et al., 2016; O'Connor & Kirtley, 2018)




Table 1: Descriptive statistics and univariate multinomial regression analyses for non-suicidal control, suicide ideation and suicide attempts groups (n=3435)

	Total (n=3435)	Control (n=2534)	Ideations (n=498)	Attempts (n=403)	Chi-square (χ^2)	Significant differences (OR)^a
<i>Categorical Variables</i>	%	%	%	%		
Gender ¹ (female)	49.4	48.5	45.1	60.5	24.39***	C < I (1.16); C > A (0.62); I > A (0.53)
Ethnicity ² (white)	93.8	93.6	93.5	95.1	1.61	
Marital status ³ (not married ^b)	83.1	81.3	88.7	86.2	19.20***	C > I (0.56); I < A (1.30)
Economic activity ^{4 c}					57.58***	
Employed	61.8	64.7	56.2	50.8		C > I (0.56); C > A (0.35); I > A (0.63)
Economically Inactive	27.9	27.1	30.8	30.5		C > A (0.54)
Unemployed	10.2	8.2	13.0	18.7		
Exposure to suicide death (family & friend)	19.9	15.9	28.0	34.5	88.98***	C < I (2.05); C < A (2.77); I < A (1.35)
Exposure to suicide attempt (family)	20.9	15.7	31.3	40.6	149.15***	C < I (2.45); C < A (3.68); I < A (1.50)
Exposure to suicide attempt (friend)	22.0	16.0	32.9	46.0	196.05***	C < I (2.57); C < A (4.48); I < A (1.74)
<i>Continuous Variables</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>		
Age	25.70 (4.86)	25.64 (4.90)	25.33 (4.70)	26.54 (4.70)	13.49	C < A (1.04); I < A (1.05)
Depressive symptoms	10.83 (11.23)	7.27 (7.58)	18.69 (12.09)	22.93 (15.20)	945.45***	C < I (1.11); C < A (1.14); I < A (1.02)
Defeat	16.24 (13.24)	11.95 (9.64)	26.39 (13.63)	29.98 (16.09)	982.74***	C < I (1.10); C < A (1.12); I < A (1.02)
Entrapment	10.27 (13.79)	5.94 (9.06)	20.64 (15.90)	24.32 (18.88)	889.45***	C < I (1.09); C < A (1.10); I < A (1.01)
Burdensomeness	13.81 (7.58)	11.48 (4.90)	19.24 (9.16)	21.31 (10.30)	862.23***	C < I (1.17); C < A (1.19); I < A (1.02)
Belongingness	11.56 (6.55)	10.06 (5.54)	15.26 (6.99)	16.17 (7.70)	473.47***	C < I (1.13); C < A (1.15); I < A (1.02)
Goal Disengagement	12.56 (3.20)	12.48 (3.15)	12.93 (3.26)	12.55 (3.37)	7.75*	C < I (1.04)

Goal Reengagement	14.10 (4.28)	13.73 (4.07)	14.81 (4.39)	15.56 (5.04)	78.77***	C < I (1.06); C < A (1.10); I < A (1.04)
Social Support	25.78 (4.59)	26.56 (3.91)	23.93 (5.39)	23.18 (5.72)	262.20***	C > I (0.89); C > A (0.87); I > A (0.98)
Resilience	28.22 (7.44)	29.60 (6.59)	25.40 (7.52)	23.43 (9.27)	320.31***	C > I (0.93); C > A (0.90); I > A (0.97)
Acquired capability	14.01 (4.26)	13.62 (4.08)	14.43 (4.24)	15.96 (4.75)	109.53***	C < I (1.05); C < A (1.14); I < A (1.09)
Mental Images	13.47 (5.18)	11.69 (3.51)	17.30 (5.08)	19.71 (6.56)	1255.37***	C < I (1.31); C < A (1.41); I < A (1.07)
Impulsivity	61.84 (10.86)	59.97 (10.14)	65.20 (10.93)	69.12 (11.27)	305.90***	C < I (1.05); C < A (1.08); I < A (1.03)

*p<0.05, **p<0.01, ***p<0.001

† C = control, I = Ideations, A = Attempts

¹ Female as reference, ² Not married as reference, ³ Non-white as reference, ⁴Unemployed as reference

^aOnly statistically significant (p<0.05) associations reported, after Holm-Bonferroni sequential correction applied to each set of comparisons.

^b not married includes single, separated, divorced and widowed

^c Economic activity reduced to 3 categories based upon the APMS dataset; '*Employed*' are economically active people, '*Economically Inactive*' includes students, those looking after home, long term sick or disabled, or retired, '*Unemployed*' are those out of work but are available to start work.

Table 2: Multinomial logistic regression of variables associated with suicidal history group membership (n=3330)

Model variables	Ideation vs. Control ^a		Attempts vs. control ^a		Attempts vs. ideation ^b	
	OR	95% CI	OR	95% CI	OR	95% CI
Age	1.00	0.98 - 1.03	1.07***	1.04 - 1.10	1.07***	1.03 - 1.10
Gender ¹	1.07	0.84 - 1.37	0.52***	0.39 - 0.70	0.49***	0.36 - 0.67
Ethnicity ²	0.93	0.57 - 1.50	1.23	0.65 - 2.31	1.33	0.69 - 2.54
Marital status ³	0.78	0.54 - 1.13	0.91	0.60 - 1.44	1.16	0.74 - 1.83
Economic Activity ⁴						
Employed ^c	1.03	0.69 - 1.52	0.69	0.45 - 1.08	0.68	0.44 - 1.04
Inactive	0.87	0.57 - 1.33	0.57	0.36 - 0.93	0.66	0.41 - 1.06
Depressive symptoms	1.00	0.98 - 1.02	1.01	0.99 - 1.03	1.01	0.98 - 1.03
Defeat	1.04***	1.02 - 1.06	1.03**	1.01 - 1.06	0.99	0.97 - 1.02
Entrapment	1.00	0.98 - 1.01	0.98	0.97 - 1.00	0.99	0.97 - 1.00
Burdensomeness	1.06***	1.04 - 1.09	1.07***	1.04 - 1.10	1.01	0.99 - 1.03
Belongingness	1.01	0.98 - 1.04	1.00	0.97 - 1.04	1.00	0.96 - 1.03
Goal Disengagement	1.02	0.98 - 1.06	0.98	0.93 - 1.02	0.96	0.92 - 1.00
Goal Reengagement	0.99	0.97 - 1.02	1.00	0.97 - 1.04	1.01	0.97 - 1.04
Social Support	1.01	0.98 - 1.04	1.02	0.98 - 1.06	1.01	0.97 - 1.05
Resilience	1.01	0.99 - 1.03	0.99	0.97 - 1.02	0.98	0.96 - 1.01
Acquired capability	1.03*	1.00 - 1.06	1.13***	1.10 - 1.18	1.10***	1.06 - 1.14
Impulsivity	1.01	1.00 - 1.02	1.03***	1.02 - 1.05	1.02*	1.01 - 1.04
Mental Images	1.18***	1.15 - 1.22	1.26***	1.22 - 1.31	1.07***	1.03 - 1.10
Exposure to suicide death (family & friend)	1.12	0.83 - 1.51	0.92	0.65 - 1.30	0.82	0.58 - 1.17
Exposure to family attempt	1.54***	1.16 - 2.03	1.93***	1.40 - 2.66	1.26	0.91 - 1.73
Exposure to friend	1.37*	1.04 - 1.80	2.04***	1.49 - 2.80	1.49***	1.09 - 2.06

attempt						
---------	--	--	--	--	--	--

OR = Odds ratio

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (statistically significant ($p < 0.05$) associations after Holm-Bonferroni sequential correction applied to each set of comparisons)

^acontrol as reference, ^bideation as reference

^c Economic activity reduced to 3 categories based upon the APMS dataset; '*Employed*' are economically active people, '*Economically Inactive*' includes students, those looking after home, long term sick or disabled, or retired, '*Unemployed*' are those out of work but are available to start work.

¹ Female as reference, ² Not married (including single, separated, divorced and widowed) as reference, ³ Non-white as reference, ⁴Unemployed as reference.

Table 3: The accuracy of the volitional phase variables in identifying who had or had not made a suicide attempt.

<i>Volitional Factors</i>	<i>Sensitivity (%)</i>	<i>Specificity (%)</i>	<i>PPV (%)</i>	<i>NPV (%)</i>
Acquired capability	37.1	77.4	39.6	43.1
Impulsivity	37.8	75.4	39.9	44.7
Mental images	54.4	61.9	37.4	46.2
Exposure to suicide death (family & friend)	34.6	72.2	50.0	42.1
Exposure to family attempt	40.8	68.2	41.	49.4
Exposure to friend attempt	39.4	67.1	39.4	46.9
All volitional factors	45.7	79.1	63.4	64.8

PPV = Positive Predictive Value, NPV = Negative Predictive Value