



This is a repository copy of *Coherent quantum transport in hybrid Superconductor-2DEG-Superconductor planar Josephson junctions*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/137726/>

Version: Accepted Version

Proceedings Paper:

Delfanazari, K., Puddy, R.K., Ma, P. et al. (8 more authors) (2017) Coherent quantum transport in hybrid Superconductor-2DEG-Superconductor planar Josephson junctions. In: 2017 16th International Superconductive Electronics Conference (ISEC). 16th International Superconductive Electronics Conference (ISEC), 12-16 Jun 2017, Naples, Italy. IEEE . ISBN 978-1-5090-5868-6

<https://doi.org/10.1109/ISEC.2017.8314214>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Coherent quantum transport in hybrid Superconductor-2DEG-Superconductor planar Josephson junctions

K. Delfanazari^{1,2,*}, R. K. Puddy², P. Ma², T. Yi², M. Cao², Y. Gul³, I. Farrer^{2,4}, D. A. Ritchie², H. J. Joyce¹, M. J. Kelly^{1,2} and C. G. Smith²

¹Electrical Engineering Division, University of Cambridge, Cambridge CB3 0FA, UK

²Cavendish Laboratory, University of Cambridge, Cambridge CB3 0HE, UK

³Department of Electronic and Electrical Engineering, University College London, London WC1E 7JE, UK

⁴Department of Electronic and Electrical Engineering, University of Sheffield, Mappin Street, Sheffield, S1 3JD, UK

*kd398@cam.ac.uk

The study of hybrid superconductor-semiconductor (S-Sm) structures have received renewed interest because of the recently reported detection of Majorana fermions at the S-Sm interface in nanowire and topological insulator based devices.

Here, we report the experimental results on proximity induced superconductivity in a high-mobility two-dimensional electron gas (2DEG) in InGaAs heterostructures. Eight symmetric S-Sm-S Josephson junctions were fabricated on a single InGaAs chip.

The superconducting electrodes were made of Niobium (Nb) and the quantum transport of junctions were measured using a lock-in technique at temperature 40 mK. The subharmonic energy gap structures (SGS) were observed at $V=2\Delta_{\text{Nb}}/ne$ on dV/dI (V) due to Andreev reflections at the S-Sm interfaces. The SGS features suppressed significantly with increasing temperature and magnetic field, leading to a shift of the SGSs toward zero bias.

The differential conductance dI/dV versus magnetic field B curves show a maxima at zero B and decreases by increasing B . When decreasing the magnetic field to below +0.4 T, there is hysteresis in B accompanied by periodic oscillations of conductance with a periodicity of ~ 15 mT between +0.05 and 0 T. The same effect could also be seen for negative fields. Both the hysteresis and periodic oscillation effects are strongly temperature dependent.

Our observation paves the way for development of coherent circuits applicable in novel quantum multiplexers and processors.