



This is a repository copy of *OC-0537: Prophylactic irradiation of tracts (PIT) in patients with pleural mesothelioma: a phase III trial.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/140717/>

Version: Accepted Version

---

**Proceedings Paper:**

Bayman, N., Appel, W., Ashcroft, L. et al. (15 more authors) (2018) OC-0537: Prophylactic irradiation of tracts (PIT) in patients with pleural mesothelioma: a phase III trial. In: Radiotherapy and Oncology. ESTRO 37, 20-24 Apr 2018, Barcelona, Spain. Elsevier , S286-S287.

[https://doi.org/10.1016/S0167-8140\(18\)30847-8](https://doi.org/10.1016/S0167-8140(18)30847-8)

---

Article available under the terms of the CC-BY-NC-ND licence  
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Prophylactic irradiation of tracts (PIT) in patients with pleural mesothelioma: a phase III trial**

### **Same authors as world lung**

*Title: abstract titles should be brief and reflect the content of the abstract. The title (maximum 100 characters) is important since it focuses attention (it is the “showcase” for the presentation). Do not use capital letters in the title except for words that are always capitalised and do not use non-standard abbreviations.*

*Body of the abstract: abstracts should be structured in such a way as to include (1) Purpose/Objective; (2) Material/methods; (3) Results; (4) Conclusion.*

*The on-line abstract submission procedure: the on-line abstract submission system will not accept abstracts that exceed 3,000 characters (body of the abstract, including spaces).*

### **Purpose/Objective**

It has been widespread practice across Europe to irradiate diagnostic or therapeutic chest wall (CW) intervention sites in patients with malignant pleural mesothelioma (MPM) post-procedure - a practice known as prophylactic irradiation of tracts (PIT). This study aims to determine the efficacy of PIT in reducing the incidence of CW metastases following a chest wall procedure in MPM.

### **Material/Methods**

In this multicentre phase III randomised controlled trial, MPM patients following a chest wall procedure were randomised 1: 1 to receive PIT (within 42-days of procedure) or no PIT. Large thoracotomies, needle biopsy sites and indwelling pleural catheters were excluded. PIT was delivered at a dose of 21Gy in 3 fractions over 3 consecutive weekdays using a single electron field adapted to maximise coverage of the tract from skin surface to pleura. The primary outcome was the incidence of CW metastases within 6 months from randomisation, assessed in the intention-to-treat population. Stratification factors included epithelioid histology and intention to give chemotherapy. Trial registration number NCT01604005.

### **Results**

375 patients (186 PIT and 189 no PIT) were randomised between 06/2012-12/2015 from 54 UK centres. Comparing PIT vs no PIT, %male patients was 89.8/88.4%, median age 72.8/74.6 years, %ECOG PS (0,1,2) 32.2/56.5, 11.3/23.8, 56.1/20.1%, %confirmed epithelioid histology 79.6/74.1%, and %with intention to give chemotherapy 71.5/71.4%. The chest wall procedures were VATS (58.1/51.3%), open surgical biopsy (2.7/5.3%), local-anaesthetic-thoracoscopy (26.9/27.0%), chest drain (5.9/8.5%) and others (6.5/7.9%) for the PIT vs no PIT arm respectively. Radiotherapy was received as intended by 181/186 patients in the PIT arm. The proportion of CW metastases by 6 months was 6/186 (3.2%) vs 10/189 (5.3%) for the PIT vs no PIT arm respectively (odds ratio 0.60 [95% CI 0.17-1.86]; p=0.44) and by 12 months 15/186 (8.1%) versus 19/189 (10.1%) respectively (OR=0.79 [95% CI 0.36-1.69]; p=0.59). Cumulative incidence of CW metastases at 6months/12 months/24 months was 3.3/8.5/10.0% in the PIT arm vs 5.6/10.9/18.7% in the no PIT arm. Evaluable patients who developed CW metastases reported a mean increase in visual analogue scale pain score of 13.3 (p<0.01) compared to baseline. Skin toxicity was the most common radiotherapy-

related adverse event in the PIT arm with 96(51.6%) grade 1, 19(10.2%) grade 2, and 1(0.5%) grade 3 radiation dermatitis (CTCAE V4.0). There were no other grade 3 or higher radiotherapy-related adverse events.

### **Conclusion**

There was no significant difference in incidence CWM between the 2 groups and the increase in VAS pain score in patients with CWM was below the 20% increase which we considered clinically significant. There therefore is no role for the routine use of PIT following diagnostic or therapeutic CW procedures in patients with MPM.

2890 characters