



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/140852/>

Version: Accepted Version

---

**Article:**

Joyce, W, Egginton, S, Farrell, AP et al. (2019) Adrenergic and adenosinergic regulation of the cardiovascular system in an Antarctic icefish: Insight into central and peripheral determinants of cardiac output. *Comparative biochemistry and physiology. Part A, Molecular & integrative physiology*, 230. pp. 28-38. ISSN: 1095-6433

<https://doi.org/10.1016/j.cbpa.2018.12.012>

---

© 2018 Elsevier Inc. All rights reserved. Licensed under the Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# **Adrenergic and Adenosinergic Regulation of the Cardiovascular System in an Antarctic Icefish: Insight into Central and Peripheral Determinants of Cardiac Output**

William Joyce<sup>1,\*</sup>, Stuart Egginton<sup>2</sup>, Anthony P. Farrell<sup>3</sup>, and Michael Axelsson<sup>4</sup>

<sup>1</sup> *Department of Zoophysiology, Aarhus University, 8000 Aarhus C, Denmark*

<sup>2</sup> *School of Biomedical Sciences, University of Leeds, Leeds, UK*

<sup>3</sup> *Department of Zoology, and Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC 45*

<sup>4</sup> *Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg, Sweden*

\* author for correspondence: [william.joyce@bios.au.dk](mailto:william.joyce@bios.au.dk)

## Abstract

Icefishes characteristically lack the oxygen-binding protein haemoglobin and therefore are especially reliant on cardiovascular regulation to augment oxygen transport when oxygen demand increases, such as during activity and warming. Using both *in vivo* and *in vitro* experiments, we evaluated the roles for adrenaline and adenosine, two well-established cardio- and vasoactive molecules, in regulating the cardiovascular system of the blackfin icefish, *Chaenocephalus aceratus*. Despite increasing cardiac contractility (increasing twitch force and contraction kinetics in isometric myocardial strip preparations) and accelerating heart rate ( $f_H$ ), adrenaline (5 nmol kg<sup>-1</sup> bolus intra-arterial injection) did not significantly increase cardiac output ( $\dot{Q}$ ) *in vivo* because it elicited a large decrease in vascular conductance ( $G_{sys}$ ). In contrast, and despite preliminary data suggesting a direct negative inotropic effect of adenosine on isolated atria and little effect on isolated ventricle strips, adenosine (500 nmol kg<sup>-1</sup>) generated a large increase in  $\dot{Q}$  by increasing  $G_{sys}$ , a change reminiscent of that previously reported during both acute warming and invoked activity. Our data thus illustrate how  $\dot{Q}$  in *C. aceratus* may be much more dependent on peripheral control of vasomotor tone than direct regulation of the heart.

Keywords: Channichthyidae, adrenaline, adenosine, heart rate, conductance

## Introduction

Icefishes (Family: Channichthyidae) distinctively lack the expression of haemoglobin (Ruud, 1954; Sidell and O'Brien, 2006) and the high values reported for cardiac output ( $\dot{Q}$ ) are thought of as a compensatory mechanism for the associated  $\sim 10$ -fold reduction in arterial carrying capacity for oxygen. However, we have recently demonstrated, with direct measurements of ventral aortic blood flow in the blackfin icefish *Chaenocephalus aceratus*, that routine  $\dot{Q}$  ( $26.6 \text{ ml min}^{-1} \text{ kg}^{-1}$ ; Joyce et al., 2018a) is much lower than the majority of previous estimates in this species ( $60 - 150 \text{ ml min}^{-1} \text{ kg}^{-1}$ ; Hemmingsen et al., 1972; Høleton, 1970), which means that icefishes have a considerable scope in which to increase  $\dot{Q}$ . Indeed, during acute warming ( $0$  to  $8^\circ\text{C}$ ) and moderate activity, the blackfin icefish substantially increased  $\dot{Q}$  up to a maximum of  $86.3 \text{ ml min}^{-1} \text{ kg}^{-1}$  (Joyce et al., 2018a). This increase was also associated with an up to five-fold increase in systemic vascular conductance ( $G_{\text{sys}}$ ) from  $15$  to  $75 \text{ ml kPa}^{-1} \text{ min}^{-1} \text{ kg}^{-1}$  (i.e., systemic vascular resistance,  $R_{\text{sys}}$ , decreased). In the absence of haemoglobin, these large cardiovascular changes are crucial to augment oxygen transport when oxygen consumption increases (Joyce et al., 2018a) because it is not possible to greatly change the arterio-venous oxygen difference, nor is it possible to increase haematocrit, as occurs prominently in other Antarctic fishes (Axelsson, 2005; Franklin et al., 1993; Joyce et al., accepted). The aim of the present study was to address the mechanisms responsible for cardiovascular regulation in icefish. We chiefly focussed on adrenaline and adenosine, two well-established cardio- and vasoactive molecules, as potential mediators of the cardiovascular response to stressors such as warming and activity.

The importance of adrenergic modulation of cardiovascular performance in Antarctic fishes remains ambiguous. Plasma catecholamines (i.e., adrenaline and noradrenaline) are not elevated during moderate exercise in *C. aceratus*, or the red-blooded species *Notothenia coriiceps* and *Trematomus bernacchii* (Davison et al., 1995; Egginton, 1997). However, in two red-blooded Nototheniids, *Pagothenia borchgrevinki* and *Trematomus bernacchii*, plasma noradrenaline and adrenaline increased in response to severe hyperthermia ( $10^\circ\text{C}$ ; Forster et al., 1998). Likewise, Whiteley and Egginton (1999) observed high catecholamine levels in both red- and white-blooded Antarctic fishes immediately following capture by trawling, suggesting adrenergic control may be

recruited *in extremis* (Whiteley and Egginton, 1999). This study nonetheless revealed a low capacity for catecholamines synthesis in Antarctic fishes (Whiteley and Egginton, 1999).

Catecholamines characteristically increase heart rate ( $f_H$ ) and cardiac contractility in teleost fishes *via* direct stimulation of cardiac  $\beta$ -adrenoceptors, either via sympathetic autonomic innervation of the heart or humoral release into the circulation (Farrell and Smith, 2017). Nevertheless, in *in situ* perfused hearts, maximum adrenergic stimulation had little or no effect on maximum  $\dot{Q}$ ,  $V_S$  or  $f_H$  in icefish (*C. aceratus* and *Chionodraco rastrospinosus*) (Egginton et al., submitted). Skov et al. (2009) revealed a positive inotropic effect of adrenaline in *C. aceratus*, in isometric ventricular strip preparations, while atrial preparations were insensitive to adrenaline (Skov et al., 2009). Catecholamines are also potent modulators of vasomotor tone and decrease  $G_{sys}$ , but typically increase branchial vascular conductance ( $G_{branch}$ ) in teleost fishes (Pettersson and Nilsson, 1980; Sandblom and Gräns, 2017; Wood, 1974), offering the potential for dynamic cardiorespiratory coupling.

The capability of the cardiovascular system to respond to adrenergic stimulation has largely been elucidated by pharmacological interventions. Indeed, adrenaline injection in *P. borchgrevinki* elevated  $\dot{Q}$ , exclusively by increasing stroke volume ( $V_S$ ) without changing  $f_H$  (Axelsson et al., 1994; Sandblom et al., 2012). Because ventral ( $P_{va}$ ) and dorsal ( $P_{da}$ ) aortic pressure increased in proportion to the increase in  $\dot{Q}$ , it was revealed that  $G_{sys}$  did not change and branchial conductance ( $G_{branch}$ ) increased only modestly (Axelsson et al., 1994; Sandblom et al., 2012). In *T. bernacchii*, adrenaline injection increased both  $V_S$  and  $f_H$ , as well as total vascular resistance ( $R_{tot}$ , the sum of  $R_{sys}$  and  $R_{branch}$ ) (Axelsson et al., 1992). In *C. aceratus*, noradrenaline injection increased  $P_{da}$  without affecting  $f_H$  (Egginton, 1997), but as this response was not measured simultaneously with blood flow the effects of adrenergic stimulation on  $\dot{Q}$  and vascular conductance *in vivo* remain unknown.

Adenosine is also capable of exerting diverse cardiovascular effects when it is released from cells as oxygen demand increases or oxygen supply decrease (Mubagwa et al., 1996), such as may occur during exercise and/or warming. In this way,  $\dot{Q}$  is controlled

peripherally rather than centrally by the autonomic nervous system. In fishes, including the red-blooded Antarctic nototheniid *P. borchgrevinki*, adenosine injection slowed the heart, increased  $G_{\text{sys}}$  and decreased  $G_{\text{branch}}$  (Sundin et al., 1999; Sundin and Nilsson, 1996). In both *N. coriiceps* and *C. aceratus*, exposure to high temperature (critical thermal maximum;  $CT_{\text{max}}$ ) resulted in decreased cardiac ATP and increased cardiac ADP levels (O'Brien et al., 2018), which is a signature for adenosine release (Mugabwa et al., 1996), making adenosine a potential driver of the increased  $G_{\text{sys}}$  observed during warming in *C. aceratus* (Joyce et al., 2018a). In the present study, we used a pharmacological approach to provide insight into the adrenergic and adenosinergic control of the cardiovascular system in *C. aceratus*, using a combination of *in vivo* and *in vitro* experiments.

## Materials and Methods

### *Experimental animals*

Adult *Chaenocephalus aceratus* of both sexes were captured at 100-200 m depth using an otter trawl deployed from the *ARSV Laurence M. Gould* in Dallmann Bay (64°10'S, 62°35'W) and off the south-western shore of Low Island (63°24'S, 62°10'W). The fish were held on the ship in aerated sea water at ambient temperature for up to 2 days during transportation to Palmer Station, Antarctica. Full details of animal husbandry are described elsewhere (Joyce et al., 2018a). Briefly, the fish were maintained in 700 or 1700 L tanks in aerated seawater at  $0 \pm 1^\circ\text{C}$ . The fish did not feed in captivity. The fish were allowed to recover from transportation for at least 72 hours prior to experiments. All experiments were approved and permitted by the University of Alaska, Fairbanks Institutional Animal Use and Care Committee (570217-9).

The *in vivo* data reported here were acquired from 12 fish (body mass:  $0.87 \pm 0.18$  kg, mean  $\pm$  SD) instrumented for our study on the cardiovascular effects of activity and warming (Joyce et al., 2018a). All of the data in the present manuscript were acquired approximately 12 h (overnight) after a heating protocol, during which environmental temperature had been raised to  $8^\circ\text{C}$  (i.e., substantially lower than  $CT_{\text{max}}$ : 12 – 14 °C (Beers and Sidell 2011; Joyce et al. 2018a; O'Brien et al. 2018)). A rapid cooling was used to

restore ambient water temperature to 0-1°C within 1-2 h. All cardiovascular variables stabilized at or close to baseline values overnight, suggesting the fish were fully recovered before the present study commenced.

### *In vivo* pharmacological study

#### *Surgery and Instrumentation for the in vivo measurements*

The surgery was described previously in detail (Joyce et al., 2018a). Briefly, fishes were anaesthetised in cold ( $0\pm 1^\circ\text{C}$ ) seawater containing MS-222 ( $140\text{ mg l}^{-1}$ ) and maintained in an anaesthetized state (gills irrigated with  $70\text{ mg l}^{-1}$  MS-222) on a surgery table. Surgery was performed in a cold room at  $\sim 2\text{-}3^\circ\text{C}$ . A Transonic flow probe (4PSB or 2.5PSL) was placed around the ventral aorta to measure total cardiac output ( $\dot{Q}$ ). The efferent branchial artery was occlusively cannulated using PE-50 with a 2F polyurethane tip to measure dorsal aortic pressure ( $P_{\text{da}}$ ) and, in some of the fish, the afferent branchial artery in the same gill arch was cannulated using a PE-50 cannula for the measurement ventral aortic pressure ( $P_{\text{va}}$ ). The cannulae and leads from the flow probe were sutured to the skin and the ventral incision was closed with 3-0 surgical silk. Post-surgery, the gills were irrigated with fresh seawater until voluntary ventilation resumed and each fish was then placed individually in a triangular 12.4 l respirometer.

The flow probe was connected to a Transonic flow meter (T402; Transonic Systems, USA). The cannulae were attached to pressure transducers (Medizintechnik, Kirchseeon, Germany), which were calibrated against a static water column before every experiment, and output signals pre-amplified by a Senselab 4CHAMP amplifier, (Somedic sales, Hörby, Sweden). The flow meter and 4CHAMP amplifier were connected to a PowerLab data acquisition system (ADInstruments, Castel Hill, Australia), interfaced to a computer running LabChart Pro (version 7; ADInstruments, Bella Vista, Australia).

#### *Pharmacological Protocol*

All experiments were conducted on resting fish at ambient temperature ( $0\text{-}1^\circ\text{C}$ ). Adrenaline ( $5\text{ nmol kg}^{-1}$ ; equivalent to that previously used in bald rockcod (*P.*

*borchgrevinki*) by Sandblom et al. (2012) and less than the 10 nmol kg<sup>-1</sup> used by Axelsson et al. (1994) also in *P. borchgrevinki*) was injected into the dorsal aorta (n=8). Once all parameters had returned to baseline levels, adenosine (500 nmol kg<sup>-1</sup>; intermediate between levels previously used in epaulette sharks (*Hemiscyllium ocellatum*; Stensl kken et al., 2004) and rainbow trout (*Oncorhynchus mykiss*; Sundin and Nilsson, 1996), both 1  mol kg<sup>-1</sup>; and bald rockcod (*P. borchgrevinki*; 10 nmol kg<sup>-1</sup>, Sundin et al., 1999) was likewise injected into the dorsal aorta (n=8). Once all cardiovascular parameters returned to baseline levels, subsequently (or in some cases independently in another cohort of resting fish) the icefish were treated with the muscarinic cholinergic antagonist atropine (n=12). When all parameters had stabilised ~25 min post-injection, the atropinised animals were subsequently treated with the  -adrenergic antagonists sotalol (N=8) or propranolol (n=4). We opted to use both  -adrenergic antagonists, in different animals, to mitigate for their possible respective side effects (e.g., Altimiras et al., 1997). All antagonists were administered as 2 mg kg<sup>-1</sup> boluses injected into the dorsal aorta (Altimiras et al., 1997; Axelsson et al., 1992; Campbell et al., 2009). The adenosine but not the adrenaline injection was repeated after double-autonomic blockade (n=7).

### Calculations

Stroke volume ( $V_S$ ) was calculated as:

$$V_S = \dot{Q} / f_H$$

Given that central venous pressure is negligible in *C. aceratus* (Joyce et al., 2018a), we assumed it to be zero in the calculation of systemic conductance ( $G_{sys}$ ):

$$G_{sys} = \dot{Q} / P_{da}$$

Branchial conductance ( $G_{branch}$ ) was calculated as:

$$G_{branch} = \dot{Q} / P_{va} - P_{da}$$

Autonomic tones on  $f_H$  were calculated using the equations provided by Altimiras et al. (1997), using R-R interval (i.e., 60/ $f_H$ ):

$$\text{Cholinergic tone (\%)} = ((R-R)_{cont} - (R-R)_{atr}) / (R-R)_{db} * 100$$

$$\text{Adrenergic tone (\%)} = ((\text{R-R})_{\text{db}} - (\text{R-R})_{\text{atr}}) / (\text{R-R})_{\text{db}} * 100$$

Where:

(R-R)<sub>cont</sub> = control R-R interval

(R-R)<sub>atr</sub> = R-R interval after cholinergic blockade with atropine

(R-R)<sub>db</sub> = R-R interval after double autonomic blockade (atropine and propranolol or sotalol)

### *In vitro* myocardial preparations

The *in vitro* heart strip data used 5 additional *C. aceratus* (body mass:  $1.53 \pm 1.54$  kg, mean  $\pm$  SD) that were killed by a sharp blow to the head followed by pithing before the heart was dissected. Excised hearts were placed in ice-cold physiological saline (composed of (mM): NaCl (250), KCl (2.5), MgCl<sub>2</sub> (0.9), CaCl<sub>2</sub> (2.5), TES acid (3.1), TES sodium salt (6.1), glucose (5.6), pH 7.95-8) (Egginton et al. submitted) and carefully dissected. The saline solution was based on that previously published for marine fishes (Farrell et al., 2013) and modified to allow for the higher Na<sup>+</sup> concentration that characterises Antarctic teleosts (Egginton, 1994). Each fish provided one atrial strip (length:  $9.7 \pm 2.6$  mm; mass  $32.6 \pm 22.4$  mg, mean  $\pm$  SD); also included are preliminary data from one ventricular strip preparation (length: 13.5 mm; mass 29.3 mg). Each strip was vertically suspended between a hook and metal rod attached to a force transducer (model P/N 730; Somedic SenseLab AB, Hörby, Sweden) using 4/0 surgical silk. The signal from the force transducer was preamplified using a 4CHAMP amplifier, (Somedic sales, Hörby, Sweden) and processed by a PowerLab data acquisition system and LabChart Pro v.7 (ADInstruments, Bella Vista, Australia). The preparations were immersed in 10 ml physiological saline in water-jacketed organ baths maintained at 1°C for 10 min before stimulation started, then stabilised for 60 min at low tension before being stretched, over a second period of 60 min, to produce maximum force. The preparations were initially stimulated at 0.2 Hz with 15 ms pulses at 150% the threshold voltage required to elicit contraction (50 – 100 V) using a Grass S9 stimulator (Quincy, MA, USA).

During the experimental protocol, we performed a force-frequency trial, increasing stimulation frequency from 0.2 to 0.5 Hz, thus covering most of the *in vivo*  $f_H$  range for this species. The preparations were then exposed to a saturating concentration of adrenaline bitartrate (5  $\mu\text{M}$ ), which was given 10 min to take full effect, before another force frequency trial. This adrenaline dose exceeded the plasma concentration invoked in our *in vivo* investigation but was chosen to stimulate a maximum response *in vitro*. A preliminary experiment using one atrial and one ventricular strip preparation investigated the effects of adenosine (100  $\mu\text{M}$ ) (Aho and Vornanen, 2002) at the end of the adrenaline protocol (*i.e.*, still in the presence of adrenaline) to provide illustrative data until a dedicated study can be performed.

### *Statistical Analysis*

For the *in vivo* study, all cardiovascular parameters were analysed in 20-s blocks immediately before and following the injections of adrenaline and adenosine. The effects of atropine, sotalol and propranolol ( $\beta$ -adrenergic blockade) were given 20 min to take full effect. A one-way ANOVA was used to investigate the cardiovascular effects of adrenaline and adenosine (both before and after double blockade) over time. Dunnett's multiple comparison post-hoc test was used to reveal significant changes from the pre-injection period (60 s prior to recording). Because adrenaline elicited complex interactions between  $f_H$  and  $V_s$ , linear regressions were used to investigate the relationship between how  $f_H$  and  $V_s$  changed (relative to pre-injection values) at 20-s intervals for the period 100 s after an adrenaline infusion. A repeated-measured one-way ANOVA and Tukey's post-hoc test was also used to compare cardiovascular parameters in control conditions, following muscarinic receptor blockade and following double autonomic blockade. An unpaired t-test was used to investigate differences in  $\beta$ -adrenergic tone achieved with propranolol and with sotalol.

For the *in vitro* experiments, contractile performance was analysed using twitch force, time to peak, maximum rate of contraction and the maximum rate of relaxation with the LabChart Pro Peak Analysis plug-in. Forces were normalised to cross-sectional area (assuming a density of 1.0  $\text{mg mm}^{-3}$  and uniform thickness of the strips). A two-way

ANOVA, followed by a Sidak's post-hoc test, investigated the effects of adrenaline and frequency on contractile parameters.

Statistical analysis was performed with GraphPad Prism (v. 7.0d). Statistical significance was assigned when  $P \leq 0.05$  and data are presented as means  $\pm$  s.e.m, unless stated otherwise.

## Results

### *In vivo effects of adrenaline*

The mean cardiovascular changes in response to adrenaline are presented in Figure 1.  $\dot{Q}$  initially decreased from 34.4 to 26.2 ml min<sup>-1</sup> kg<sup>-1</sup> but quickly recovered to above 42 ml min<sup>-1</sup> kg<sup>-1</sup>, although this delayed increase was not significantly higher than pre-treatment (Fig. 1a). Neither  $f_H$  (Fig. 1b) or  $V_S$  (Fig. 1c) was solely responsible for the initial decrease in  $\dot{Q}$ . Because both  $P_{da}$  and  $P_{va}$  increased (Fig. 1d) with the decrease in  $\dot{Q}$ , the decrease in  $f_H$  was most likely a hypertensive bradycardia baroreflex. The most typical response to adrenaline was transient bradycardia (Fig. 2a), although in some cases this consisted of only occasional prolonged beats (Fig. 2b), in others it lasted for several minutes (Fig. 2c). Consequently, mean  $f_H$  did not significantly change for the initial few minutes. The overall cardiovascular response was also influenced by  $V_S$ , which tended to decrease with increasing cardiac afterload (i.e., an increase in  $P_{va}$ ), with  $f_H$  and  $V_S$  changing reciprocally. Therefore, to quantify this interaction, we present the relationship between  $\Delta f_H$  and  $\Delta V_S$  over the first 100 s following an adrenaline injection (Fig. 3). In fish displaying a strong bradycardia,  $V_S$  did not significantly decrease, i.e.,  $V_S$  was presumably compensated by an increase in cardiac filling time. Subsequently, a tachycardia developed (likely a direct action of adrenaline) and  $f_H$  maximally reached 12.1 beats min<sup>-1</sup>, greater than the control value of 8.8 beats min<sup>-1</sup> ( $P < 0.05$ ). Adrenaline significantly decreased  $G_{sys}$  from 16.9 to  $< 10.0$  ml kPa<sup>-1</sup> min<sup>-1</sup> kg<sup>-1</sup>, without affecting  $G_{branch}$ , thereby indicating systemic vasoconstriction (Fig. 1).

### *In vivo effects of adenosine before and after autonomic blockade*

Adenosine injection significantly decreased  $P_{da}$  and  $P_{va}$ , and significantly increased  $\dot{Q}$  and  $G_{sys}$  (Figs. 4 and 5), while  $G_{branch}$  was unchanged.  $\dot{Q}$  peaked at  $65.2 \text{ ml min}^{-1} \text{ kg}^{-1}$  while  $G_{sys}$  reached  $64.6 \text{ ml kPa}^{-1} \text{ min}^{-1} \text{ kg}^{-1}$ .  $f_H$  exhibited a delayed, but large increase from 8.3 to 14.7 beats  $\text{min}^{-1}$  ( $P < 0.05$ ), a response that was abolished by double autonomic blockade, which suggested that this tachycardia was likely a hypotensive baroreflex exerted through withdrawal of vagal tone and/or sympathetic drive. Indeed, injection of adenosine after double blockade elicited a small but significant bradycardia (Fig. 4d).

Muscarinic blockade (atropine injection) increased  $\dot{Q}$  through an increase in  $f_H$  despite a modest decrease in  $V_S$  (Table 1).  $P_{da}$  and  $P_{va}$  increased in proportion to  $\dot{Q}$ , demonstrating that  $G_{sys}$  was unaffected by muscarinic blockade. In contrast,  $G_{branch}$  increased substantially in 4 out of 5 fish following muscarinic blockade (in two fish it doubled, but in one animal there was a small decrease, thus it was statistically unchanged overall). After atropinisation,  $\beta$ -adrenergic blockade significantly decreased  $f_H$  and increased  $V_S$  but had no other effects (Table 1), supporting the earlier suggestion that an adrenaline injection could have a direct (but slowed) chronotropic cardiac effect.

Cholinergic tone was calculated to be  $103.0 \pm 10.3 \%$  and adrenergic tone  $19.5 \pm 1.0 \%$ . There was no significant difference between adrenergic tonus calculated using either sotalol or propranolol ( $P = 0.70$ ).

#### *In vitro effects stimulation frequency, adrenaline and adenosine in atrial preparations*

Increasing stimulation frequency in atrial preparations revealed a negative force-frequency effect (Fig. 6A), shorter time-to-peak force (Fig. 6B) and a slowing of maximum contractile kinetics (Fig. 6C and D) ( $P < 0.05$ ). Adrenaline significantly ( $P < 0.05$ ) increased maximum isometric force by approximately 30% (Fig. 6A), a positive inotropic effect that reached statistical significance only at 0.2, 0.3 and 0.4 Hz. Adrenaline decreased ( $P < 0.05$ ) time-to-peak only at 0.3 Hz (Fig. 6B) but increased the maximum rate of contraction at all frequencies (Fig. 6C) ( $P < 0.05$ ). Adrenaline also invoked a significant ( $P < 0.05$ ) positive lusitropic effect at 0.2 and 0.3 Hz (Fig. 6D).

The preliminary data for adenosine on one atrial strip revealed a clear 50% decrease in maximum isometric force at all frequencies (Fig. 7A) but the ventricular strip did not respond to adenosine (Fig. 7B).

## Discussion

### *The effects of adrenaline*

The role of catecholamines in the regulation of the cardiovascular system in notothenioids, including icefishes, has proven controversial (Egginton, 1997; Skov et al., 2009; Whiteley and Egginton, 1999), in part because plasma catecholamine levels are normally unchanged during the stress associated with exercise (Egginton, 1997), unlike the more typical teleostean response (Reid et al., 1998). Nevertheless, plasma catecholamines can increase under extreme conditions, such as following capture by trawling (Whiteley and Egginton, 1999). Therefore, to provide further insight into the likely functional relevance of circulating catecholamines, our pharmacological study investigated the effects of adrenaline on the cardiovascular system of *C. aceratus*.

An  $\alpha$ -adrenoceptor-mediated vasoconstriction of the systemic circulation is a well-established mechanism for regulating  $G_{\text{sys}}$  and central arterial blood pressure in fishes (Farrell, 1981; Wahlqvist, 1980; Wood, 1976; Wood and Shelton, 1975), including in Antarctic notothenioids (Sandblom et al., 2010, 2009). Thus, the immediate increase in both  $P_{\text{da}}$  and  $P_{\text{va}}$  as a result of a fall in  $G_{\text{sys}}$  (with no change in  $G_{\text{branch}}$ ) when adrenaline was injected into the dorsal aorta of *C. aceratus* is consistent with such a mechanism. Indeed, the hypertensive response to adrenaline is also in accordance with previous brief reports for *C. aceratus* (Egginton, 1997; Hemmingsen et al., 1972). Nevertheless, the hypertension in *C. aceratus* was considerably more pronounced than that reported in *P. borchgrevinki* (Axelsson et al., 1994; Sandblom et al., 2012), but more akin to that in *T. bernacchii* (Axelsson et al., 1992), despite similar doses of adrenaline being used in all three studies. This increase in arterial blood pressure with adrenaline injections and the well-established barostatic reflex in fishes (Jones and Milsom, 1982; Randall and Stevens, 1967; Stevens et al., 1972) helps with our interpretation of the variable cardiac responses of *C. aceratus* to an injection of adrenaline.

A hypertensive bradycardia following adrenaline injection is typical of temperate fish (Helgason and Nilsson, 1973; Stevens et al., 1972; Wood and Shelton, 1980), although it was absent in the Antarctic notothenioids *P. borchgrevinki* and *T. bernacchii*, despite the fact that the former does develop bradycardia in response to the hypertension elicited by angiotensin (Axelsson et al., 1992; 1994). In *C. aceratus*, we observed a biphasic response to adrenaline, consisting of an initial decrease in  $\dot{Q}$  followed by a recovery that matched or in some cases exceeded baseline conditions. The initial decline in  $\dot{Q}$  was due to a combination of hypertensive bradycardia and a direct decrease in  $V_s$  (due to increased afterload). However, in animals with a pronounced bradycardia, the increased cardiac filling time compensated for the increase in afterload, meaning that  $V_s$  was maintained (Fig. 3). In Figure 3 the lines of best fit always intersect the x-axis at negative  $V_s$ , confirming that in the theoretical absence of chronotropic changes, the primary response to the increase in afterload is a decrease in  $V_s$ . This is consistent with the results of perfused heart studies that have demonstrated that icefish hearts are exceptionally sensitive to afterload (Egginton et al., submitted; Acierno et al., 1997; Tota et al., 1991).

The delayed recovery in  $\dot{Q}$  we observed was probably attributable to the slowly developing positive inotropic (and chronotropic) action of adrenaline, which took 5-10 minutes to reach maximal effect in the *in vitro* cardiac strip preparations. The vascular response (increase in  $P_{da}$  and  $P_{va}$ ) may also have been quicker to develop because the adrenaline bolus was injected into the dorsal aorta and hence perfused the systemic circulation before reaching the heart. The positive chronotropic action we observed contrasted with the finding that maximum adrenergic stimulation did not increase  $f_H$  in perfused *C. aceratus* hearts (Egginton et al., submitted). However, this earlier work required hearts to be perfused with a relatively high tonic adrenaline concentration during control conditions (50 nM), which may have saturated the sino-atrial  $\beta$ -adrenoceptors before the effects of additional adrenaline were investigated.

Skov et al. (2009) reported no effect of stimulation frequency or adrenaline treatment on contractile performance in *C. aceratus* atria, although this was not supported by the present data. This is remarkable because most previous investigations on fish atrial preparations have revealed clear positive inotropic effects of adrenergic stimulation

(Fløysand and Helle, 1994; Gesser, 1996; Keen et al., 1992; Lennard and Huddart, 1992; Meghji and Burnstock, 1984). Given the important role that atria have in filling the fish ventricle (Farrell, 1991), we re-evaluated the effect of adrenaline on *C. aceratus* atrial tissue. In contrast to Skov and colleagues, we observed a clear negative force-frequency effect and positive inotropic effect of adrenaline. The discrepancy may be ascribed to the higher concentration used in the present study (5 as opposed to 1  $\mu\text{M}$ ), which unequivocally demonstrated that adrenaline has the potential to modulate atrial contractility in this species. Plasma catecholamines after trawling have not been measured in *C. aceratus*, but in another icefish, *Champscephalus gunnari*, total plasma catecholamine concentration (noradrenaline and adrenaline) slightly exceeded 1  $\mu\text{mol l}^{-1}$  (adrenaline: 826  $\text{nmol l}^{-1}$ , noradrenaline 238  $\text{nmol l}^{-1}$ ), and in another notothenioid (*Dissostichus mawsoni*), reached even higher levels (adrenaline: 2601  $\text{nmol l}^{-1}$ , noradrenaline 841  $\text{nmol l}^{-1}$ ) (Whiteley and Egginton, 1999), suggesting that 1  $\mu\text{M}$  may not have been high enough to represent the physiological extreme.

### *The effects of adenosine*

In fish (Sundin et al., 1999; Sundin and Nilsson, 1996), as well as mammals (Mubagwa et al., 1996) and reptiles (Joyce and Wang, 2014), adenosine typically dilates the systemic vasculature. In *C. aceratus*, we confirmed that adenosine elicited a large peripheral vasodilatation, as evidenced by the several-fold increase in  $G_{\text{sys}}$  following injection. Quantitatively, the effect of adenosine on the systemic vasculature was similar to that in *P. borchgrevinki*, although we did not observe an effect on  $G_{\text{branch}}$ , in contrast to the decreased  $G_{\text{branch}}$  seen in *P. borchgrevinki* (Sundin et al., 1999).

Most striking was the a large increase in  $\dot{Q}$  with adenosine infusion that peaked at 65.2  $\text{ml min}^{-1} \text{kg}^{-1}$ , thereby exceeding peak  $\dot{Q}$  evoked during moderate burst activity demonstrated previously at the same temperature (53  $\text{ml min}^{-1} \text{kg}^{-1}$ ; Joyce et al., 2018a). Before autonomic blockade, a delayed but prominent tachycardia was observed after adenosine infusion. Because this was abolished by atropine and  $\beta$ -adrenergic antagonists, we deduce that it was a hypotensive baroreflex (mediated by the autonomic nervous system) and not a direct effect of adenosine or other non-adrenergic non-cholinergic mechanisms. This response developed relatively slowly ( $> 2$  min) in

comparison to that observed temperate teleosts, in which a baroreflex appears almost immediately following hypotension (Sandblom and Axelsson, 2005). This difference may be attributable to the low temperature slowing the responsiveness in the icefish, and is consistent with the slow positive chronotropic effect we observed in response to adrenaline infusion. Following double-blockade, adenosine elicited a slight bradycardia (1.3 beats  $\text{min}^{-1}$ / 8.7% fall), which was smaller than that previously reported in *P. borchgrevinki* (~5 beats  $\text{min}^{-1}$ / 21.7% fall) (Sundin et al., 1999).

In rainbow trout (*Oncorhynchus mykiss*), adenosine exerts a clear negative inotropic effect on atrial tissue, and a smaller negative inotropic effect on the ventricle (Aho and Vornanen, 2002; Meghji and Burnstock, 1984). This is similar to the response in other animals, including most mammals (Hollander and Webb, 1957) and reptiles (Joyce et al., 2014). However, in carp, a positive inotropic effect of adenosine has been reported in ventricular tissue (Vornanen and Tuomennoro, 1999). It was therefore instructive to perform a preliminary experiment to determine the effect of adenosine on the icefish heart, while urging caution given the low sample size. As adenosine exerted a clear negative inotropic on atrial tissue, and apparently little effect on ventricular tissue, this indicates that the increase in  $\dot{Q}$  observed *in vivo* was unlikely to be a result of a positive inotropic effect on the heart. Therefore, the increase in  $\dot{Q}$ , in both blocked and unblocked fish, is likely attributable to a peripheral vascular effect. It has been similarly argued, by comparing the effects of exercise and vasodilators, that  $\dot{Q}$  is largely determined by peripheral vascular conductance in humans (Bada et al., 2012; González-Alonso et al., 2008). The finding that large changes in  $\dot{Q}$  were attained after double-blockade, in the absence of any change in  $f_H$ , is consistent with the view developed by pacing studies that the control of  $f_H$  *per se* is of little importance in determining  $\dot{Q}$  (Joyce et al., 2018b; Munch et al., 2014).

### *The effects of dual autonomic blockade*

The teleost fish heart operates under dual (i.e., cholinergic and adrenergic) autonomic control (Burnstock, 1969; Farrell and Smith, 2017; Vornanen, 2017; Wang, 2012). Cholinergic tone is especially pronounced in red-blooded Antarctic fishes, which, in conjunction with the near-freezing temperatures, results in low resting  $f_H$  (< 15 beats

min<sup>-1</sup> (Axelsson et al., 1992; Campbell et al., 2009; Lowe et al., 2005; Sandblom and Axelsson, 2011)). In *C. aceratus*, Hemmingsen et al. (1972) observed that atropine elicited only a small increase in  $f_H$  from 16 to 18 beats min<sup>-1</sup>, implying that cholinergic tone could be relatively weak in this haemoglobinless species. However, in ECG-instrumented fish (a minimally invasive surgery) we recently reported cholinergic tone of 55% and adrenergic tone of 12% (Joyce et al. 2018a), which are similar to those reported in red-blooded relatives (Sandblom and Axelsson 2011; Egginton and Campbell 2016). In the present study, the cholinergic tone was even higher (103.0%), whilst adrenergic tone was more similar (19.5%). Given that stress is characteristically associated with reduced or loss of cholinergic tone in fish (Sandblom and Axelsson, 2011; Wahlqvist and Nilsson, 1980), this supports our contention that by the time this protocol was conducted the fish had sufficiently recovered from the warming protocol on the previous day. The large increase in  $f_H$  following atropinisation was associated with an increase in  $\dot{Q}$ , although  $V_s$  decreased, likely due to the tachycardia decreasing cardiac filling time (Altimiras and Axelsson, 2004).

### *Conclusions*

Adrenaline induces a range of cardiovascular effects including decreased  $G_{sys}$ , increased  $f_H$ , and increased atrial contractility. However, effects on the vasculature appears inconsistent with cardiovascular changes (i.e., increased  $G_{sys}$ ) observed during activity and warming *in vivo* (Joyce et al., 2018a). This suggests that circulating adrenaline is of little importance in regulating cardiovascular physiology *in vivo*, and helps explain why plasma catecholamines do not increase during exercise in this species (Egginton, 1997). Sympathetic innervation of the heart may provide an alternative route for adrenergic control of cardiac performance. Although this was not detected in the red-blooded nototheniid *P. borchgrevinki* (Sandblom et al., 2010), it remains to be investigated in icefishes.

Adenosine elicited a several-fold increase in  $G_{sys}$  that approached the maximum previously observed at high temperature (8°C) during activity (65 vs. 75 ml kPa<sup>-1</sup> min<sup>-1</sup> kg<sup>-1</sup>) (Joyce et al., 2018a). Proof that adenosine is involved in the reported responses would require evidence that adenosinergic-receptor antagonists attenuate the activity- and

warming-induced vascular response. However, such an experiment may prove futile because other vasodilators (such as nitric oxide, bradykinin, and prostacyclin) likely act in synchrony (Hellsten et al., 2012; Joyner, 2013; Joyner and Dempsey, 2018; Lamb et al., 2018). Nevertheless, our data clearly illustrate that  $\dot{Q}$  may be largely determined by peripheral factors, distinct from direct autonomic cardiac control (Guyton, 1968). It may nevertheless be fruitful for future studies to further investigate how adrenergic stimulation affects blood flow in combination with potentially sympatholytic vasodilators, as opposed to their independent effects.

## Acknowledgements

This study would not have been possible without the contributions of Drs. Kristin M. O'Brien (University of Alaska Anchorage) and Elizabeth L. Crockett (Ohio University), who led our project on the effects of temperature on Antarctic fish physiology. We are furthermore indebted to Amanda Biederman, Anna Rix and Elizabeth Evans for assistance with fishing and animal care. We are also grateful to the Masters and crew of the ARSV Laurence M. Gould, and the winter support staff at Palmer Station. Funding was provided by grants from the National Science Foundation (ANT 1341602 to E.L.C. and ANT 1341663 to K.M.O.) and the Swedish Research council (Vetenskapsrådet; grant 2015-05286). APF is supported by the Canadian NSERC and a Canada Research Chair.

## References

- Acierno, R., Agnisola, C., Tota, B., Sidell, B.D., 1997. Myoglobin enhances cardiac performance in antarctic icefish species that express the protein. *Am. J. Physiol.* 273, R100-106. <https://doi.org/10.1152/ajpregu.1997.273.1.R100>
- Aho, E., Vornanen, M., 2002. Effects of adenosine on the contractility of normoxic rainbow trout heart. *J. Comp. Physiol. B, Biochem. Syst. Environ. Physiol.* 172, 217–225. <https://doi.org/10.1007/s00360-001-0245-9>
- Altimiras, J., Aissaoui, A., Tort, L., Axelsson, M., 1997. Cholinergic and Adrenergic Tones in the Control of Heart Rate in Teleosts. How Should They be Calculated? *Comparative Biochemistry and Physiology Part A: Physiology* 118, 131–139. [https://doi.org/10.1016/S0300-9629\(96\)00402-1](https://doi.org/10.1016/S0300-9629(96)00402-1)
- Altimiras, J., Axelsson, M., 2004. Intrinsic autoregulation of cardiac output in rainbow trout (*Oncorhynchus mykiss*) at different heart rates. *Journal of Experimental Biology* 207, 195–201. <https://doi.org/10.1242/jeb.00742>

- Axelsson, M., 2005. The Circulatory System and Its Control, in: Fish Physiology, Physiology of Polar Fishes. Academic Press, pp. 239–280.  
[https://doi.org/10.1016/S1546-5098\(04\)22006-4](https://doi.org/10.1016/S1546-5098(04)22006-4)
- Axelsson, M., Davison, B., Forster, M., Nilsson, S., 1994. Blood Pressure Control in the Antarctic Fish *Pagothenia Borchgrevinki*. *Journal of Experimental Biology* 190, 265–279.
- Axelsson, M., Davison, W., Forster, M.E., Farrell, A.P., 1992. Cardiovascular responses of the red-blooded antarctic fishes *Pagothenia bernacchii* and *P. borchgrevinki*. *J. Exp. Biol.* 167, 179–201.
- Bada, A.A., Svendsen, J.H., Secher, N.H., Saltin, B., Mortensen, S.P., 2012. Peripheral vasodilatation determines cardiac output in exercising humans: insight from atrial pacing. *J Physiol* 590, 2051–2060.  
<https://doi.org/10.1113/jphysiol.2011.225334>
- Beers, J.M., Sidell, B.D., 2011. Thermal tolerance of Antarctic notothenioid fishes correlates with level of circulating hemoglobin. *Physiol. Biochem. Zool.* 84, 353–362. <https://doi.org/10.1086/660191>
- Burnstock, G., 1969. Evolution of the Autonomic Innervation of Visceral and Cardiovascular Systems in Vertebrates. *Pharmacol Rev* 21, 247–324.
- Campbell, H., Davison, W., Fraser, K.P.P., Peck, L.S., Egginton, S., 2009. Heart rate and ventilation in Antarctic fishes are largely determined by ecotype. *J. Fish Biol.* 74, 535–552. <https://doi.org/10.1111/j.1095-8649.2008.02141.x>
- Davison, W., Axelsson, M., Forster, M., Nilsson, S., 1995. Cardiovascular responses to acute handling stress in the Antarctic fish *Trematomus bernacchii* are not mediated by circulatory catecholamines. *Fish Physiol Biochem* 14, 253–257.  
<https://doi.org/10.1007/BF00004316>
- Egginton, S., 1997. A comparison of the response to induced exercise in red- and white-blooded Antarctic fishes. *J Comp Physiol B* 167, 129–134.  
<https://doi.org/10.1007/s003600050056>
- Egginton, S., 1994. Stress response in two Antarctic teleosts (*Notothenia coriiceps* Richardson and *Chaenocephalus aceratus* Lönnberg) following capture and surgery. *J Comp Physiol B* 164, 482–491. <https://doi.org/10.1007/BF00714586>
- Egginton, S., Campbell, H.A., 2016. Cardiorespiratory responses in an Antarctic fish suggest limited capacity for thermal acclimation. *J. Exp. Biol.* 219, 1283–1286.  
<https://doi.org/10.1242/jeb.130963>
- Farrell, A.P., 1991. From Hagfish to Tuna: A Perspective on Cardiac Function in Fish. *Physiological Zoology* 64, 1137–1164.  
<https://doi.org/10.1086/physzool.64.5.30156237>
- Farrell, A.P., 1981. Cardiovascular Changes in the Lingcod (*Ophiodon Elongatus*) Following Adrenergic and Cholinergic Drug Infusions. *Journal of Experimental Biology* 91, 293–305.
- Farrell, A.P., Altimiras, J., Franklin, C.E., Axelsson, M., 2013. Niche expansion of the shorthorn sculpin (*Myoxocephalus scorpius*) to Arctic waters is supported by a thermal independence of cardiac performance at low temperature. *Can. J. Zool.* 91, 573–580. <https://doi.org/10.1139/cjz-2013-0038>
- Farrell, A.P., Smith, F., 2017. 4 - Cardiac Form, Function and Physiology, in: Gamperl, A.K., Gillis, T.E., Farrell, A.P., Brauner, C.J. (Eds.), *Fish Physiology, The Cardiovascular System*. Academic Press, pp. 155–264.  
<https://doi.org/10.1016/bs.fp.2017.07.001>

- Fløysand, R., Helle, K.B., 1994. Effects of temperature and adrenaline on the atrial myocardium of the cultured Atlantic salmon (*Salmo salar*). *Fish Physiol Biochem* 13, 23–30. <https://doi.org/10.1007/BF00004116>
- Forster, M.E., Davison, W., Axelsson, M., Sundin, L., Franklin, C.E., Gieseg, S., 1998. Catecholamine release in heat-stressed Antarctic fish causes proton extrusion by the red cells. *J Comp Physiol B* 168, 345–352. <https://doi.org/10.1007/s003600050153>
- Franklin, C.E., Davison, W., McKenzie, J.C., 1993. The Role of the Spleen During Exercise in the Antarctic Teleost, *Pagothenia Borchgrevinki*. *Journal of Experimental Biology* 174, 381–386.
- Gesser, H., 1996. Cardiac force-interval relationship, adrenaline and sarcoplasmic reticulum in rainbow trout. *J Comp Physiol B* 166, 278–285. <https://doi.org/10.1007/BF00262872>
- González-Alonso, J., Mortensen, S.P., Jeppesen, T.D., Ali, L., Barker, H., Damsgaard, R., Secher, N.H., Dawson, E.A., Dufour, S.P., 2008. Haemodynamic responses to exercise, ATP infusion and thigh compression in humans: insight into the role of muscle mechanisms on cardiovascular function. *J. Physiol. (Lond.)* 586, 2405–2417. <https://doi.org/10.1113/jphysiol.2008.152058>
- Guyton, A.C., 1968. Regulation of cardiac output. *Anesthesiology* 29, 314–326.
- Helgason, S.S., Nilsson, S., 1973. Drug effects on pre- and post-branchial blood pressure and heart rate in a free-swimming marine teleost, *Gadus morhua*. *Acta Physiol. Scand.* 88, 533–540. <https://doi.org/10.1111/j.1748-1716.1973.tb05482.x>
- Hellsten, Y., Nyberg, M., Jensen, L.G., Mortensen, S.P., 2012. Vasodilator interactions in skeletal muscle blood flow regulation. *J Physiol* 590, 6297–6305. <https://doi.org/10.1113/jphysiol.2012.240762>
- Hemmingsen, E.A., Douglas, E.L., Johansen, K., Millard, R.W., 1972. Aortic blood flow and cardiac output in the hemoglobin-free fish *Chaenocephalus aceratus*. *Comparative Biochemistry and Physiology Part A: Physiology* 43, 1045–1051. [https://doi.org/10.1016/0300-9629\(72\)90176-4](https://doi.org/10.1016/0300-9629(72)90176-4)
- Holeton, G.F., 1970. Oxygen uptake and circulation by a hemoglobinless antarctic fish (*Chaenocephalus aceratus* Lonnberg) compared with three red-blooded antarctic fish. *Comparative Biochemistry and Physiology* 34, 457–471. [https://doi.org/10.1016/0010-406X\(70\)90185-4](https://doi.org/10.1016/0010-406X(70)90185-4)
- Hollander, P.B., Webb, J.L., 1957. Effects of adenine nucleotides on the contractility and membrane potentials of rat atrium. *Circ. Res.* 5, 349–353.
- Jones, D.R., Milsom, W.K., 1982. Peripheral Receptors Affecting Breathing and Cardiovascular Function in Non-Mammalian Vertebrates. *Journal of Experimental Biology* 100, 59–91.
- Joyce, W., Axelsson, M., Egginton, S., Farrell, A.P., Crockett, E.L., O'Brien, K.M., Accepted. The effects of thermal acclimation on cardio-respiratory performance in an Antarctic fish (*Notothenia coriiceps*). *Conservation Physiology*.
- Joyce, W., Egginton, S., Farrell, A.P., Crockett, E.L., O'Brien, K.M., Axelsson, M., 2018a. Exploring nature's natural knockouts: In vivo cardiorespiratory performance of Antarctic fishes during acute warming. *Journal of Experimental Biology* jeb.183160. <https://doi.org/10.1242/jeb.183160>
- Joyce, W., Else, R.M., Wang, T., Crossley, D.A., 2018b. Maximum heart rate does not limit cardiac output at rest or during exercise in the American alligator (*Alligator mississippiensis*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 315, R296–R302. <https://doi.org/10.1152/ajpregu.00027.2018>

- Joyce, W., Gesser, H., Wang, T., 2014. Purinoceptors exert negative inotropic effects on the heart in all major groups of reptiles. *Comp. Biochem. Physiol., Part A Mol. Integr. Physiol.* 171, 16–22. <https://doi.org/10.1016/j.cbpa.2014.02.005>
- Joyce, W., Wang, T., 2014. Adenosinergic regulation of the cardiovascular system in the red-eared slider *Trachemys scripta*. *Comp. Biochem. Physiol., Part A Mol. Integr. Physiol.* 174, 18–22. <https://doi.org/10.1016/j.cbpa.2014.04.002>
- Joyner, M.J., 2013. Physiology and Redundancy. *Physiology* 28, 136–137. <https://doi.org/10.1152/physiol.00015.2013>
- Joyner, M.J., Dempsey, J.A., 2018. Physiological Redundancy and the Integrative Responses to Exercise. *Cold Spring Harb Perspect Med* 8. <https://doi.org/10.1101/cshperspect.a029660>
- Keen, J.E., Farrell, A.P., Tibbits, G.F., Brill, R.W., 1992. Cardiac physiology in tunas. II. Effect of ryanodine, calcium, and adrenaline on force–frequency relationships in atrial strips from skipjack tuna, *Katsuwonus pelamis*. *Can. J. Zool.* 70, 1211–1217. <https://doi.org/10.1139/z92-168>
- Lamb, I.R., Novielli, N.M., Murrant, C.L., 2018. Capillary response to skeletal muscle contraction: evidence that redundancy between vasodilators is physiologically relevant during active hyperaemia. *J. Physiol. (Lond.)* 596, 1357–1372. <https://doi.org/10.1113/JP275467>
- Lennard, R., Huddart, H., 1992. The effect of thermal stress and adrenaline on the dynamics of the flounder (*Platichthys flesus*) heart. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology* 101, 307–310. [https://doi.org/10.1016/0742-8413\(92\)90278-F](https://doi.org/10.1016/0742-8413(92)90278-F)
- Lowe, C.J., Seebacher, F., Davison, W., 2005. Thermal sensitivity of heart rate and insensitivity of blood pressure in the Antarctic nototheniid fish *Pagothenia borchgrevinkii*. *J Comp Physiol B* 175, 97–105. <https://doi.org/10.1007/s00360-004-0466-9>
- Meghji, P., Burnstock, G., 1984. Actions of some autonomic agents on the heart of the trout (*Salmo gairdneri*) with emphasis on the effects of adenylyl compounds. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology* 78, 69–75. [https://doi.org/10.1016/0742-8413\(84\)90049-5](https://doi.org/10.1016/0742-8413(84)90049-5)
- Mubagwa, K., Mullane, K., Flameng, W., 1996. Role of adenosine in the heart and circulation. *Cardiovasc. Res.* 32, 797–813.
- Munch, G.D.W., Svendsen, J.H., Damsgaard, R., Secher, N.H., González-Alonso, J., Mortensen, S.P., 2014. Maximal heart rate does not limit cardiovascular capacity in healthy humans: insight from right atrial pacing during maximal exercise. *J Physiol* 592, 377–390. <https://doi.org/10.1113/jphysiol.2013.262246>
- O'Brien, K.M., Rix, A.S., Egginton, S., Farrell, A.P., Crockett, E.L., Schlauch, K., Woolsey, R., Hoffman, M., Merriman, S., 2018. Cardiac mitochondrial metabolism may contribute to differences in thermal tolerance of red- and white-blooded Antarctic nototheniid fishes. *Journal of Experimental Biology* 221, jeb177816. <https://doi.org/10.1242/jeb.177816>
- Pettersson, K., Nilsson, S., 1980. Drug induced changes in cardio-vascular parameters in the Atlantic cod, *Gadus morhua*. *J Comp Physiol B* 137, 131–138. <https://doi.org/10.1007/BF00689211>
- Randall, D.J., Stevens, E.D., 1967. The role of adrenergic receptors in cardiovascular changes associated with exercise in salmon. *Comparative Biochemistry and Physiology* 21, 415–424. [https://doi.org/10.1016/0010-406X\(67\)90803-1](https://doi.org/10.1016/0010-406X(67)90803-1)

- Reid, S.G., Bernier, N.J., Perry, S.F., 1998. The adrenergic stress response in fish: control of catecholamine storage and release. *Comp. Biochem. Physiol. C, Pharmacol. Toxicol. Endocrinol.* 120, 1–27.
- Ruud, J.T., 1954. Vertebrates without erythrocytes and blood pigment. *Nature* 173, 848–850.
- Sandblom, E., Axelsson, M., 2011. Autonomic control of circulation in fish: A comparative view. *Autonomic Neuroscience* 165, 127–139. <https://doi.org/10.1016/j.autneu.2011.08.006>
- Sandblom, E., Axelsson, M., 2005. Baroreflex mediated control of heart rate and vascular capacitance in trout. *Journal of Experimental Biology* 208, 821–829. <https://doi.org/10.1242/jeb.01470>
- Sandblom, E., Axelsson, M., Davison, W., 2009. Circulatory function at sub-zero temperature: venous responses to catecholamines and angiotensin II in the Antarctic fish *Pagothenia borchgrevinki*. *J. Comp. Physiol. B, Biochem. Syst. Environ. Physiol.* 179, 165–173. <https://doi.org/10.1007/s00360-008-0299-z>
- Sandblom, E., Davison, W., Axelsson, M., 2012. Cold Physiology: Postprandial Blood Flow Dynamics and Metabolism in the Antarctic Fish *Pagothenia borchgrevinki*. *PLOS ONE* 7, e33487. <https://doi.org/10.1371/journal.pone.0033487>
- Sandblom, E., Gräns, A., 2017. 7 - Form, Function and Control of the Vasculature, in: Gamperl, A.K., Gillis, T.E., Farrell, A.P., Brauner, C.J. (Eds.), *Fish Physiology, The Cardiovascular System*. Academic Press, pp. 369–433. <https://doi.org/10.1016/bs.fp.2017.06.001>
- Sandblom, E., Olsson, C., Davison, W., Axelsson, M., 2010. Nervous and humoral catecholaminergic control of blood pressure and cardiac performance in the Antarctic fish *Pagothenia borchgrevinki*. *Comp. Biochem. Physiol., Part A Mol. Integr. Physiol.* 156, 232–236. <https://doi.org/10.1016/j.cbpa.2010.02.004>
- Sidell, B.D., O'Brien, K.M., 2006. When bad things happen to good fish: the loss of hemoglobin and myoglobin expression in Antarctic icefishes. *J. Exp. Biol.* 209, 1791–1802. <https://doi.org/10.1242/jeb.02091>
- Skov, P.V., Bushnell, P.G., Tirsgaard, B., Steffensen, J.F., 2009. The role of adrenaline as a modulator of cardiac performance in two Antarctic fishes. *Polar Biol* 32, 215. <https://doi.org/10.1007/s00300-008-0522-0>
- Stensløkken, K.-O., Sundin, L., Renshaw, G.M.C., Nilsson, G.E., 2004. Adenosinergic and cholinergic control mechanisms during hypoxia in the epaulette shark (*Hemiscyllium ocellatum*), with emphasis on branchial circulation. *J. Exp. Biol.* 207, 4451–4461. <https://doi.org/10.1242/jeb.01291>
- Stevens, E.D., Bennion, G.R., Randall, D.J., Shelton, G., 1972. Factors affecting arterial pressures and blood flow from the heart in intact, unrestrained lingcod, *Ophiodon elongatus*. *Comparative Biochemistry and Physiology Part A: Physiology* 43, 681–695. [https://doi.org/10.1016/0300-9629\(72\)90255-1](https://doi.org/10.1016/0300-9629(72)90255-1)
- Sundin, L., Axelsson, M., Davison, W., Forster, M.E., 1999. Cardiovascular responses to adenosine in the antarctic fish *pagothenia borchgrevinki*. *Journal of Experimental Biology* 202, 2259–2267.
- Sundin, L., Nilsson, G.E., 1996. Branchial and systemic roles of adenosine receptors in rainbow trout: an in vivo microscopy study. *Am. J. Physiol.* 271, R661–669. <https://doi.org/10.1152/ajpregu.1996.271.3.R661>
- Tota, B., Acierno, R., Agnisola, C., 1991. Mechanical performance of the isolated and perfused heart of the haemoglobinless Antarctic icefish *Chionodraco hamatus*

- (Lönnberg): effects of loading conditions and temperature. *Phil. Trans. R. Soc. Lond. B* 332, 191–198. <https://doi.org/10.1098/rstb.1991.0049>
- Vornanen, M., 2017. 3 - Electrical Excitability of the Fish Heart and Its Autonomic Regulation, in: Gamperl, A.K., Gillis, T.E., Farrell, A.P., Brauner, C.J. (Eds.), *Fish Physiology, The Cardiovascular System*. Academic Press, pp. 99–153. <https://doi.org/10.1016/bs.fp.2017.04.002>
- Vornanen, M., Tuomennoro, J., 1999. Effects of acute anoxia on heart function in crucian carp: importance of cholinergic and purinergic control. *Am. J. Physiol.* 277, R465–475.
- Wahlqvist, I., 1980. Effects of catecholamines on isolated systemic and branchial vascular beds of the cod, *Gadus morhua*. *J Comp Physiol B* 137, 139–143. <https://doi.org/10.1007/BF00689212>
- Wahlqvist, I., Nilsson, S., 1980. Adrenergic control of the cardio-vascular system of the Atlantic cod, *Gadus morhua*, during “stress.” *J Comp Physiol B* 137, 145–150. <https://doi.org/10.1007/BF00689213>
- Wang, T., 2012. Chapter 141 - Evolution of the Cardiovascular Autonomic Nervous System in Vertebrates, in: Robertson, D., Biaggioni, I., Burnstock, G., Low, P.A., Paton, J.F.R. (Eds.), *Primer on the Autonomic Nervous System (Third Edition)*. Academic Press, San Diego, pp. 669–673. <https://doi.org/10.1016/B978-0-12-386525-0.00141-4>
- Whiteley, N.M., Egginton, S., 1999. Antarctic fishes have a limited capacity for catecholamine synthesis. *Journal of Experimental Biology* 202, 3623–3629.
- Wood, C.M., 1976. Pharmacological properties of the adrenergic receptors regulating systemic vascular resistance in the rainbow trout. *J Comp Physiol B* 107, 211–228. <https://doi.org/10.1007/BF00691227>
- Wood, C.M., 1974. A Critical Examination of the Physical and Adrenergic Factors Affecting Blood Flow Through the Gills of the Rainbow Trout. *Journal of Experimental Biology* 60, 241–265.
- Wood, C.M., Shelton, G., 1980. Cardiovascular dynamics and adrenergic responses of the rainbow trout in vivo. *Journal of Experimental Biology* 87, 247–270.
- Wood, C.M., Shelton, G., 1975. Physical and adrenergic factors affecting systemic vascular resistance in the rainbow trout: a comparison with branchial vascular resistance. *J. Exp. Biol.* 63, 505–523.

## Figure Legends

Figure 1: The effects of adrenaline (5 nmol kg<sup>-1</sup>) on cardiovascular parameters in *C. aceratus*. Intra-arterial adrenaline infusion is marked with an arrow (ADR). (a)  $\dot{Q}$ , cardiac output (n=8); (b)  $f_H$ , heart rate (n=8); (c)  $V_S$ , stroke volume (n=8); (d)  $P_{va}$ , ventral aortic pressure (n=6);  $P_{da}$ , dorsal aortic pressure (n=8); (e)  $G_{branch}$ , branchial conductance (n=6); (f)  $G_{sys}$ , systemic conductance (n=8). All values are means  $\pm$  s.e.m. Asterisks denote significant ( $P<0.05$ ) changes from the first (control) measurement within a given variable (repeated measures one-way ANOVA followed by Dunnett's multiple comparison test).

Figure 2. Original traces from 3 icefish depicting the variable cardiovascular effects of adrenaline. Arterial infusion is marked with an arrow (ADR).  $\dot{Q}$ , cardiac output;  $P_{va}$ , ventral aortic pressure;  $P_{da}$ , dorsal aortic pressure;  $f_H$ , heart rate. Fish 1 (a,d,g) represents a typical biphasic response consisting of an initial hypertensive bradycardia and subsequent tachycardia; Fish 2 (b,e,h) exhibits a near absence of bradycardia; Fish 3 (c,f,i) depicts a persistent bradycardia.

Figure 3: Mutual dependency of heart rate and stroke volume 20 – 100 seconds following adrenaline infusion. The change in heart rate ( $\Delta f_H$ ) and change in stroke volume ( $\Delta V_S$ ) are calculated relative to pre-injection values for each individual fish. In animals in which heart rate increased, stroke volume fell and *vice versa*. This resulted in a universal decrease in  $\dot{Q}$ , despite the changes in  $f_H$  and  $V_S$  alone not being statistical significant.

Figure 4: The effects of adenosine (500 nmol kg<sup>-1</sup>) on cardiovascular parameters in *C. aceratus*. Arterial infusion is marked with an arrow (ADO). (a)  $\dot{Q}$ , cardiac output (n=8 before double blockade, n=7 after double blockade); (b)  $f_H$ , heart rate (n=8, n=7 after double blockade); (c)  $V_S$ , stroke volume (n=8, n=7 after double blockade); (d)  $P_{va}$ , ventral aortic pressure (n=6, n=5 after double blockade);  $P_{da}$ , dorsal aortic pressure (n=8, n=7 after double blockade); (e)  $G_{branch}$ , branchial conductance (n=6, n=5 after double blockade); (f)  $G_{sys}$ , systemic conductance (n=8, n=7 after double blockade). All values are means  $\pm$  s.e.m. Asterisks denote significant ( $P<0.05$ ) changes from the first (control) measurement within a given variable (repeated measures one-way ANOVA followed by Dunnett's multiple comparison test).

Figure 5: The effect of adenosine on cardiovascular dynamics in a representative icefish before and after double-autonomic blockade. Intra-arterial adenosine infusion is marked with an arrow (ADO).  $\dot{Q}$ , cardiac output;  $P_{va}$ , ventral aortic pressure;  $P_{da}$ , dorsal aortic pressure;  $f_H$ , heart rate.

Figure 6. The effect of adrenaline (5  $\mu\text{M}$ ) and stimulation frequency on *in vitro* contractile performance of atrial strips from *C. aceratus*. Asterisks show significant differences after adrenaline treatment at a given frequency. Dissimilar letters represent stimulation frequency-dependent changes within a given treatment (lower case: control; capital letters: adrenaline treated; two-way ANOVAs followed by Sidak post-hoc test.). Values are mean + s.e.m. (n=5).

Figure 7. Preliminary data showing the effects of adenosine (100  $\mu\text{M}$ ) on *in vitro* contractile performance (force-stimulation frequency response) in one atrial and one ventricular preparation from *C. aceratus*.




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.

	$\dot{Q}$ (ml min <sup>-1</sup> kg <sup>-1</sup> )	$f_H$ (beats min <sup>-1</sup> )	$V_S$ (ml kg <sup>-1</sup> )	$P_{da}$ (kPa)	$P_{va}$ (kPa)	$P_{cv}$ (kPa)	$G_{sys}$ (ml kPa <sup>-1</sup> min <sup>-1</sup> kg <sup>-1</sup> )	$G_{branch}$ (ml kPa <sup>-1</sup> min <sup>-1</sup> kg <sup>-1</sup> )
control	35.1 (±3.0) <sup>a</sup>	8.5 (±0.5) <sup>a</sup>	4.16 (± 0.35) <sup>a</sup>	2.15 (± 0.35) <sup>a</sup>	2.49 (± 0.06) <sup>a</sup>	0.27 (± 0.2) <sup>a</sup>	17.1 (± 1.7) <sup>a</sup>	86.0 (±19.7) <sup>a</sup>
muscarinic blockade	43.9 (±4.3) <sup>b</sup>	18.9 (±0.6) <sup>b</sup>	2.32 (±0.24) <sup>b</sup>	2.84 (± 0.24) <sup>b</sup>	3.16 (± 0.06) <sup>b</sup>	0.00 (±0.01) <sup>a</sup>	15.5 (± 1.5) <sup>a</sup>	114.6 (±22.4) <sup>a</sup>
double- blockade	42.3 (±4.3) <sup>b</sup>	15.2 (±0.5) <sup>c</sup>	2.78 (±0.29) <sup>c</sup>	2.71 (± 0.29) <sup>b</sup>	3.16 (± 0.09) <sup>b</sup>	0.06 (±0.05) <sup>a</sup>	16.2 (± 1.9) <sup>a</sup>	87.0 (±16.7) <sup>a</sup>

Table 1. The effects of muscarinic blockade (atropine; 2 mg kg<sup>-1</sup>) and double autonomic blockade (atropine and propranolol or sotalol; 2 mg kg<sup>-1</sup>) on cardiac output ( $\dot{Q}$ , n=10), heart rate ( $f_H$ , n=12), stroke volume ( $V_S$ , n=10), dorsal aortic pressure ( $P_{da}$ , n=12), ventral aortic pressure ( $P_{va}$ , n=7), central venous pressure ( $P_{cv}$ , n=3), systemic conductance ( $G_{sys}$ , n=10) and branchial conductance ( $G_{branch}$ , n=5) in icefish at ambient temperature. Dissimilar letters denote significant differences ( $P<0.05$ ; repeated measures one-way ANOVA and Tukey's post-hoc test)