White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Leeds.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/140903/

Version: Accepted Version

Article:
Agler, J, Lykova, Z and Young, NJ (2019) A geometric characterization of the symmetrized bidisc. Journal of Mathematical Analysis and Applications, 473 (2). pp. 1377-1413. ISSN: 0022-247X

https://doi.org/10.1016/j.jmaa.2019.01.027

© 2019 Elsevier Inc. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/.

Reuse
This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

A GEOMETRIC CHARACTERIZATION OF THE SYMMETRIZED BIDISC

JIM AGLER, ZINAIDA LYKOVA, AND N. J. YOUNG

Abstract. The symmetrized bidisc

$$G \stackrel{\text{def}}{=} \{(z+w, zw) : |z| < 1,\ |w| < 1\}$$

has interesting geometric properties. While it has a plentiful supply of complex geodesics and of automorphisms, there is nevertheless a unique complex geodesic $\mathcal{R}$ in G that is invariant under all automorphisms of G. Moreover, G is foliated by those complex geodesics that meet $\mathcal{R}$ in one point and have nontrivial stabilizer. We prove that these properties, together with two further geometric hypotheses on the action of the automorphism group of G, characterize the symmetrized bidisc in the class of complex manifolds.

Contents

Introduction 2
1. The action of automorphisms on G 6
1.1. The action of $\operatorname{Aut}\mathbb{D}$ on G 6
1.2. The action of $\operatorname{Aut} G$ on the royal variety 7
1.3. Orbits in G as manifolds 7
1.4. The sharp direction in G 13
1.5. Flat geodesics and the action of $\operatorname{Aut} G$ 14
1.6. Synchrony in G 16
2. Royal manifolds 18
2.1. Regularity properties of royal manifolds 21
2.2. Flat fibrations over royal discs 25
2.3. Synchrony in Ω 26
2.4. The sharp direction in Ω 28
2.5. Sharpness of the action of $\operatorname{Aut}\Omega$ 30

Date: 9th January 2019.

2010 *Mathematics Subject Classification.* Primary: 32A07, 53C22, 54C15, 47A57, 32F45; Secondary: 47A25, 30E05.

Partially supported by National Science Foundation Grants DMS 1361720 and 1665260, a Newcastle URC Visiting Professorship and the Engineering and Physical Sciences Research Council grant EP/N03242X/1 .

2.6. A characterization of G 34
3. A characterization of G via flat co-ordinates 39
4. Asymmetry of domains 41
References 42
Index 44

Introduction

By a *domain* we mean a connected open set in $\mathbb{C}^n$ for some integer $n \geq 1$. A domain is *homogeneous* if the automorphisms of the domain act transitively. It is *symmetric* if every point of the domain is an isolated fixed point of an involutive automorphism of the domain.

The nature of a bounded symmetric homogeneous domain in $\mathbb{C}^n$ is captured by the great classification theorem of Élie Cartan [8], an early triumph of the theory of several complex variables [14, 16]. It states that any such domain is isomorphic to a product of domains, each of which is isomorphic to a domain of one of six concrete types. The theorem is fundamental to the complex geometry and function theory of bounded symmetric homogeneous domains.

In this paper we are interested in irreducible domains Ω which narrowly miss being homogeneous, in the sense that the action of the automorphisms of Ω splits the domain into a one-parameter family of orbits. Such domains are said to have *cohomogeneity* 1, and have an extensive theory [15, 11] in both the mathematical and physics literatures.

One familiar domain that has cohomegeneity 1 is the annulus

$$\mathbb{A}_q \stackrel{\text{def}}{=} \{z \in \mathbb{C} : q < |z| < q^{-1}\} \tag{0.1}$$

where $0 < q < 1$. The orbits here are the sets

$$\{z : |z| = t\} \cup \{z : |z| = t^{-1}\} \tag{0.2}$$

where $q < t \leq 1$.

For a higher-dimensional example, consider the domain

$$G \stackrel{\text{def}}{=} \{(z + w, zw) : |z| < 1,\ |w| < 1\} \tag{0.3}$$

in $\mathbb{C}^2$, known as the *symmetrized bidisc*. The automorphisms of G are the maps of the form

$$(z + w, zw) \mapsto (m(z) + m(w), m(z)m(w)) \tag{0.4}$$

for some automorphism m of the unit disc $\mathbb{D}$. The orbits in G are therefore generically 3-dimensional real manifolds, and there is a one-parameter family of them.

Another domain, now in $\mathbb{C}^3$, having a one-parameter family of orbits is the *tetrablock*, which comprises the points $(x^1, x^2, x^3) \in \mathbb{C}^3$ such that

$$1 - x^1 z - x^2 w + x^3 zw \neq 0 \tag{0.5}$$

for all $z, w \in \mathbb{C}$ such that $|z| \leq 1$ and $|w| \leq 1$.

An ambitious project would be to classify bounded domains in $\mathbb{C}^n$, and more generally, complex manifolds, for which the orbits under the automorphisms of the manifold comprise a one-parameter family. By way of a start we shall here characterize in geometric terms our archetypal example G defined in equation (0.3). This domain has been studied by numerous authors over the past 20 years, and has proved to be a domain with a very rich complex geometry and function theory: see, besides many other papers, [4, 10, 13, 17, 22, 19, 25, 2]. G is significant for the theory of invariant distances [18], because it has Lempert's property, that the Carathéodory and Kobayashi metrics coincide [21], despite the fact that G is not convex (nor even biholomorphic to a convex domain [10]). It plays a role in operator theory [7, 23] and even has applications to a problem in the theory of robust control (for example, [27]); indeed the control application was the original motivation for the study of G. In an earlier paper [3] we characterized G in terms of the Carathéodory extremal functions that it admits. Here we give another characterization, this time in terms of its complex geodesics and automorphisms.

An *automorphism* of a complex manifold is a bijective holomorphic self-map of the manifold; such a map automatically has a holomorphic inverse. For any complex manifold Ω we denote by $\operatorname{Aut}\Omega$ the automorphism group of Ω with the compact-open topology. A *complex geodesic* of G can be defined as the range of an analytic map $f : \mathbb{D} \to G$ that has an analytic left inverse, where $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$.

We draw attention to two striking geometric properties of G.

(1) There exists a unique complex geodesic $\mathcal{R}$ in G that is invariant under all automorphisms of G. Moreover, every automorphism of $\mathcal{R}$ extends to a unique automorphism of G;
(2) for every $s \in \mathcal{R}$ there exists a unique geodesic F_s in G having a nontrivial stabilizer in $\operatorname{Aut} G$ and such that

$$F_s \cap \mathcal{R} = \{s\}.$$

Moreover, the geodesics $\{F_s : s \in \mathcal{R}\}$ foliate G.[1]

We call $\mathcal{R}$ the *royal variety* and the sets F_s the *flat geodesics* of G.

Could it be that properties (1) and (2) suffice to characterize G? In the present paper we show that the answer is yes under some further geometric hypotheses, which we now describe.

We say that a properly embedded analytic disc[2] D in a complex manifold Ω is a *royal disc* if it has properties analogous to those of $\mathcal{R}$ in (1), that is, D is invariant under every automorphism of Ω, and every automorphism of D extends to a unique automorphism of Ω. A *royal manifold* is a pair (Ω, D) where Ω is a complex manifold and D is a royal disc in Ω.

If (Ω, D) is a royal manifold then a collection $\mathcal{E} = \{E_\lambda : \lambda \in D\}$ of properly embedded analytic discs[3] in Ω is a *flat fibration over* D if it has properties similar to those of $\{F_s : s \in \mathcal{R}\}$ in (2), that is, $E_\lambda \cap D = \{\lambda\}$ for every $\lambda \in D$, $\mathcal{E}$ is a partition of Ω and, for every automorphism θ of Ω and every $\lambda \in D$, $\theta(E_\lambda) = E_{\theta(\lambda)}$. The triple $(\Omega, D, \mathcal{E})$ is then called a *flatly fibered royal manifold*.

The orbits in $(\Omega, D, \mathcal{E})$ have a natural parametrization by $[0, \infty)$. For any $\mu \in \Omega$ there is a unique $\lambda \in D$ such that $\mu \in E_\lambda$; we define the *Poincaré parameter* $P(\mu)$ to be the Poincaré distance from μ to λ in the disc E_λ (see Definition 2.19). Two points μ_1, μ_2 in Ω lie in the same orbit if and only if $P(\mu_1) = P(\mu_2)$.

Flatly fibered royal manifolds can enjoy two geometric properties: synchrony and sharpness. *Synchrony* is a condition which relates the actions of $\operatorname{Aut}\Omega$ on D and on the discs in $\mathcal{E}$. To be precise, if θ is an automorphism of Ω which fixes a point $\lambda \in D$, then it follows easily from the definition of a flat fibration over a royal manifold that the eigenspaces of the operator $\theta'(\lambda)$ on the tangent space $T_\lambda\Omega$ to Ω at λ are the tangent spaces $T_\lambda D$ and $T_\lambda E_\lambda$. We say that $(\Omega, D, \mathcal{E})$ is *synchronous* if, for every $\lambda \in D$, the eigenvalue of $\theta'(\lambda)$ corresponding to $T_\lambda E_\lambda$ is the square of the eigenvalue of $\theta'(\lambda)$ corresponding to $T_\lambda D$.

Sharpness is a condition on the action of $\operatorname{Aut}\mathbb{D}$ on Ω in a flatly fibered royal manifold $(\Omega, D, \mathcal{E})$. The definition of $(\Omega, D, \mathcal{E})$ implies that every $m \in \operatorname{Aut}\mathbb{D}$ induces an automorphism $\Theta(m)$ of Ω. For $\alpha \in \mathbb{D}$, let B_α denote the automorphism

$$B_\alpha(z) = \frac{z - \alpha}{1 - \bar{\alpha} z} \tag{0.6}$$

[1] That is, every point of G lies in some F_s and no point of G lies in two distinct F_s

[2] not assumed to be a geodesic

[3] again, not assumed to be geodesics

of $\mathbb{D}$. We say that $\operatorname{Aut}\Omega$ *acts sharply* at a point $\mu \in \Omega \setminus D$ if, in local co-ordinates,

$$\mathrm{e}^{2P(\mu)}(\Theta(B_{it})(\mu) - \mu) = i\,(\Theta(B_t)(\mu) - \mu) + o(t) \tag{0.7}$$

as $t \to 0$ in $\mathbb{R}$.

The geometric content of the sharpness condition at μ relates to the derivative at zero of the map $\alpha \mapsto \Theta(B_\alpha)(\mu)$ from $\mathbb{D}$ to Ω. This map is *a priori* a real-linear map from $T_0\mathbb{D}$ to $T_\mu\Omega$; now $T_0\mathbb{D}(= \mathbb{C})$ and $T_\mu\Omega$ are both complex vector spaces, and the sharpness condition is equivalent to the statement that the derivative at zero is also a *complex* linear map.

If we denote by $\mu^\sharp$ the range of this complex-linear derivative, then it is easy to see that $\mu^\sharp$ is the unique nonzero complex linear subspace of $T_\mu\Omega$ that is contained in the 3-dimensional real tangent space at μ to the orbit of μ in Ω.

The sharp direction $\mu^\sharp$ is a covariant line bundle over Ω which has interesting geometric properties. For example, in G, the sharp direction $\mu^\sharp$ is characterized by the fact that the complex geodesic $\mathcal{C}$ through μ with direction $\mu^\sharp$ has the *closest point property*, meaning that, for any point $\lambda \in \mathcal{C}$, if F_s, $s \in \mathcal{R}$, is the flat geodesic containing μ, then the closest point to λ in F_s is μ.

Our main result, Theorem 2.30 in the body of the paper, gives a precise version of the following statement, which holds under suitable regularity conditions.

Theorem A. *Let Ω be a complex manifold. Ω is isomorphic to G if and only if there exist a royal disc D in Ω and a flat fibration $\mathcal{E}$ of Ω over D such that $(\Omega, D, \mathcal{E})$ is a synchronous flatly fibered royal manifold and $\operatorname{Aut}\Omega$ acts sharply on $E_\lambda \setminus \{\lambda\}$ for some $\lambda \in D$.*

Formal definitions of synchrony and sharp action are given in Subsections 2.3 and 2.5. The appropriate notion of regularity is described in Subsection 2.1.

Remarkably, Theorem A implies that if $(\Omega, D, \mathcal{E})$ is a synchronous flatly fibered royal domain with suitable regularity, and $\operatorname{Aut}\Omega$ acts sharply, then both D and the leaves in $\mathcal{E}$ are complex geodesics of Ω. It suggests that G might be characterized also in terms of the properties of its complex geodesics, and in a future paper we shall show that it is so.

In Section 3 we give in Theorem 3.2 a characterization of G in terms of the existence of global co-ordinates ranging over the bidisc and satisfying certain partial differential equations. These co-ordinates are related to the flat geodesics in G.

In a short final section we discuss the relevance of the notion of symmetric space to the question of classification and show that the annulus, the symmetrized bidisc and the tetrablock, besides being inhomogeneous, also fail to be symmetric in É. Cartan's sense.

If U and Ω are complex manifolds, we denote by $\Omega(U)$ the set of holomorphic mappings from U into Ω.

We have used the expression *properly embedded analytic disc* in a complex manifold Ω. By this phrase we mean a proper injective analytic map $k : \mathbb{D} \to \Omega$ such that $k'(z) \neq 0$ for all $z \in \mathbb{D}$. The range of such a map k will also be called a properly embedded analytic disc.

1. The action of automorphisms on G

In this section we study the orbit structure of G under the action of $\operatorname{Aut} G$.

1.1. **The action of $\operatorname{Aut}\mathbb{D}$ on G.** As we stated in the introduction (see equation (0.4)), every automorphism m of $\mathbb{D}$ induces a map $\gamma_m : G \to G$ via the formula

$$\gamma_m(z + w, zw) = (m(z) + m(w), m(z)m(w)) \tag{1.1}$$

for $z, w \in \mathbb{D}$. It is easy to check that this formula defines a map $\gamma_m \in G(G)$ and that $\gamma_m \in \operatorname{Aut} G$.

Proposition 1.1. *The map $\gamma : \operatorname{Aut}\mathbb{D} \to \operatorname{Aut} G$ given by*

$$\gamma(m) = \gamma_m \tag{1.2}$$

for $m \in \operatorname{Aut}\mathbb{D}$ is a continuous isomorphism of topological groups.

The fact that γ is an isomorphism of groups is proved in [6, Theorem 5.1] or [18]. It is routine to show that γ is continuous with respect to the compact-open topologies on $\operatorname{Aut}\mathbb{D}$ and $\operatorname{Aut} G$.

The following statements are elementary.

Proposition 1.2. (1) $\operatorname{Aut}\mathbb{D}$ *and* $\operatorname{Aut} G$ *are Lie groups.*
(2) *For any $s \in G$ the map*

$$e_s : \operatorname{Aut}\mathbb{D} \to G \text{ given by } e_s(m) = \gamma_m(s) \tag{1.3}$$

is real-analytic.

The map e_s, where $s \in G$, will be called the *evaluation map* at s on $\operatorname{Aut}\mathbb{D}$.

1.2. **The action of** $\operatorname{Aut} G$ **on the royal variety.** The *royal variety* in G is defined to be the set

$$\begin{aligned}\mathcal{R} &= \{s = (s^1, s^2) \in G : (s^1)^2 = 4s^2\} \\ &= \{(2z, z^2) : z \in \mathbb{D}\}\end{aligned}$$

(we use superscripts to denote the components of a point in $\mathbb{C}^d$). Thus $\mathcal{R} = R(\mathbb{D})$ where

$$R(z) = (2z, z^2) \qquad \text{for } z \in \mathbb{D}. \tag{1.4}$$

Clearly

$$\gamma_m(R(z)) = R(m(z)) \qquad \text{for } z \in \mathbb{D} \text{ and } m \in \operatorname{Aut}\mathbb{D}. \tag{1.5}$$

The observations (1.2) and (1.5) have three consequences, summarized in the following proposition.

Proposition 1.3. (1) *Every automorphism of G leaves $\mathcal{R}$ invariant.*
(2) *Every automorphism of G is uniquely determined by its values on $\mathcal{R}$.*
(3) *Every automorphism of $\mathcal{R}$ has a unique extension to an automorphism of G.*

In statement (3), automorphisms of $\mathcal{R}$ are with respect to the structure of $\mathcal{R}$ as a complex manifold.

We can summarize these three statements by saying that the restriction map $\gamma \mapsto \gamma|\mathcal{R}$ is an isomorphism from $\operatorname{Aut} G$ to $\operatorname{Aut}\mathcal{R}$. The following commutative diagram describes the situation, where $\iota_{\mathcal{R}}$ denotes the injection of $\mathcal{R}$ into G and $m \in \operatorname{Aut}\mathbb{D}$.

$$\begin{array}{ccccc} \mathbb{D} & \xrightarrow{R} & \mathcal{R} & \xrightarrow{\iota_{\mathcal{R}}} & G \\ {\scriptstyle m}\downarrow & & {\scriptstyle \gamma_m|\mathcal{R}}\downarrow & & {\scriptstyle \gamma_m}\downarrow \\ \mathbb{D} & \xrightarrow{R} & \mathcal{R} & \xrightarrow{\iota_{\mathcal{R}}} & G \end{array} \tag{1.6}$$

1.3. **Orbits in** G **as manifolds.** For any complex manifold U and any $\lambda \in U$, we denote by $\operatorname{Orb}_U(\lambda)$ the orbit of λ under the action of the group of automorphisms of U:

$$\operatorname{Orb}_U(\lambda) = \{\gamma(\lambda) : \gamma \in \operatorname{Aut} U\}.$$

Consider the case that $U = G$ and $\lambda = s \in G$.

In view of Proposition 1.1, for any $s \in G$,

$$\operatorname{Orb}_G(s) = \{\gamma_m(s) : m \in \operatorname{Aut}\mathbb{D}\}, \tag{1.7}$$

so that $\operatorname{Orb}_G(s)$ is the range of the evaluation map e_s of equation (1.3).

$\operatorname{Aut}\mathbb{D}$ is a 3-dimensional real-analytic manifold, for which we shall need local co-ordinates.

Lemma 1.4. *For $(r,\alpha) \in \mathbb{R}\times\mathbb{D}$ let $m_{r,\alpha} \in \operatorname{Aut}\mathbb{D}$ be given by the formula*

$$m_{r,\alpha}(z) = \mathrm{e}^{ir}\frac{z-\alpha}{1-\bar\alpha z}, \qquad z\in\mathbb{D}. \tag{1.8}$$

Let

$$\begin{aligned} U_1 &= \{m_{r,\alpha} : -\pi < r < \pi,\ \alpha\in\mathbb{D}\}, \\ U_2 &= \{m_{r,\alpha} : 0 < r < 2\pi,\ \alpha\in\mathbb{D}\} \end{aligned} \tag{1.9}$$

and define

$$\varphi_1 : U_1 \to (-\pi,\pi)\times\mathbb{D} \quad \text{ by } \quad \varphi_1(m_{r,\alpha}) = (r,\alpha),$$

and similarly for $\varphi_2 : U_2 \to (0,2\pi)\times\mathbb{D}$. Then (U_1,φ_1) and (U_2,φ_2) are charts in $\operatorname{Aut}\mathbb{D}$ which together comprise a real-analytic atlas $\mathcal{A}$ for the group manifold. The identity automorphism $\mathrm{id}_{\mathbb{D}} = m_{0,0}$ belongs to U_1.

Proof. The automorphisms of $\mathbb{D}$ consist of the maps $m_{r,\alpha}$ for $r\in[-2\pi,2\pi]$ and $\alpha\in\mathbb{D}$, and therefore $\operatorname{Aut}\mathbb{D} = U_1\cup U_2$. If $-\pi < r < 0$ then

$$\varphi_2\circ\varphi_1^{-1}(r,\alpha) = (r+2\pi,\alpha)$$

and similarly when $0<r<\pi$. The transition map is therefore real-analytic from $\varphi_1(U_1\cap U_2)$ to $\varphi_2(U_1\cap U_2)$. □

Proposition 1.5. *If $s\notin\mathcal{R}$ then the evaluation map $e_s : \operatorname{Aut}\mathbb{D}\to \operatorname{Orb}_G(s)$ is a local homeomorphism and a two-to-one covering map, given explicitly by*

$$e_s(m_{r,\alpha}) = \frac{(\mathrm{e}^{ir}(-2\alpha+(1+|\alpha|^2)s^1-2\bar\alpha s^2), \mathrm{e}^{2ir}(\alpha\alpha-\alpha s^1+s^2))}{1-\bar\alpha s^1+\bar\alpha\bar\alpha s^2}.$$

Proof. Consider a point $s=(z+w,zw)\in G$ where $z,w\in\mathbb{D}$ and $z\neq w$. Let υ be the unique automorphism of $\mathbb{D}$ that maps z to w and w to z. Note that υ is not the identity automorphism $\mathrm{id}_{\mathbb{D}}$ since $z\neq w$. For $m_1,m_2\in\operatorname{Aut}\mathbb{D}$,

$$\begin{aligned} \gamma_{m_1}(s) = \gamma_{m_2}(s) &\Leftrightarrow \pi(m_1(z),m_1(w)) = \pi(m_2(z),m_2(w)) \\ &\Leftrightarrow \begin{cases} \text{either } m_1(z)=m_2(z) \text{ and } m_1(w)=m_2(w) \\ \text{or } m_1(z)=m_2(w) \text{ and } m_1(w)=m_2(z) \end{cases} \\ &\Leftrightarrow \begin{cases} \text{either } m_1 = m_2 \\ \text{or } m_2^{-1}\circ m_1 = \upsilon. \end{cases} \end{aligned} \tag{1.10}$$

Thus $e_s : m\mapsto\gamma_m(s)$ is two-to-one from $\operatorname{Aut}\mathbb{D}$ to $\operatorname{Orb}_G(s)$.

To prove that e_s is a local homeomorphism, choose any point $e_s(\beta)$ of $\operatorname{Orb}_G(s)$, where $\beta\in\operatorname{Aut}\mathbb{D}$. Choose a neighborhood U of $\mathrm{id}_{\mathbb{D}}$ such that

$$m_2^{-1}\circ m_1 \neq \upsilon \quad \text{ for all } m_1,m_2\in U. \tag{1.11}$$

We claim that $\beta \circ U$ is a neighborhood of β on which e_s is injective. Certainly it is a neighborhood of β, and if $e_s|\beta \circ U$ is not injective then there exist distinct points $m_1, m_2 \in U$ such that $e_s(\beta \circ m_1) = e_s(\beta \circ m_2)$. That is, $\gamma_\beta \circ \gamma_{m_1}(s) = \gamma_\beta \circ \gamma_{m_2}(s)$, and therefore $\gamma_{m_1}(s) = \gamma_{m_2}(s)$. Hence, by the equivalence (1.10), $m_2^{-1} \circ m_1 = v$. This equation contradicts the statement (1.11). Thus e_s is locally injective on $\operatorname{Aut}\mathbb{D}$.

Choose a compact neighborhood V of $\mathrm{id}_\mathbb{D}$ contained in U. Since a continuous bijective map from a compact space to a Hausdorff space is a homeomorphism, $e_s|V$ is a homeomorphism onto its range. It follows by homogeneity that e_s is a local homeomorphism on $\operatorname{Aut}\mathbb{D}$. Indeed, consider any $m \in \operatorname{Aut}\mathbb{D}$ and its neighborhood $m \circ V$. Define $L_m : \operatorname{Aut}\mathbb{D} \to \operatorname{Aut}\mathbb{D}$ by $L_m(\theta) = m^{-1} \circ \theta$ for $\theta \in \operatorname{Aut}\mathbb{D}$. In the commutative diagram

$$\begin{array}{rcl} m \circ V & \stackrel{e_s|m\circ V}{\longrightarrow} & e_s(m \circ V) \\ {\scriptstyle L_m|m\circ V}\big\downarrow & & \big\uparrow{\scriptstyle \gamma(m)|e_s(V)} \\ V & \stackrel{e_s|V}{\longrightarrow} & e_s(V) \end{array} \tag{1.12}$$

the map $e_s|m \circ V$ is expressed as the composition of three homeomorphisms, and so is itself a homeomorphism. Thus e_s is a local homeomorphism.

The formula for $e_s(m_{r,\alpha})$ is a simple calculation. □

For any $s \in G$, the map $e_s'(\mathrm{id}_\mathbb{D})$ is a real-linear map from the tangent space $T_{\mathrm{id}_\mathbb{D}} \operatorname{Aut}\mathbb{D}$ to $T_s \operatorname{Orb}_G(s)$. The space $T_{\mathrm{id}_\mathbb{D}} \operatorname{Aut}\mathbb{D}$ is the Lie algebra of $\operatorname{Aut}\mathbb{D}$, so we shall denote it by $\operatorname{Lie}(\operatorname{Aut}\mathbb{D})$ (though we shall not use its Lie structure, only its real-linear structure).

For every $s \in G$ we define a real-linear subspace $\mathcal{V}(s)$ of $\mathbb{C}^2$ by

$$\mathcal{V}(s) \stackrel{\text{def}}{=} \operatorname{span}_\mathbb{R} \left\{ i \begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix}, \begin{pmatrix} 2 - (s^1)^2 + 2s^2 \\ s^1 - s^1 s^2 \end{pmatrix}, i \begin{pmatrix} 2 + (s^1)^2 - 2s^2 \\ s^1 + s^1 s^2 \end{pmatrix} \right\}. \tag{1.13}$$

Theorem 1.6. (1) *If $s \in \mathcal{R}$, then $\operatorname{Orb}_G(s)$ is a one-dimensional complex manifold properly embedded in G.*

(2) *If $s \in G \setminus \mathcal{R}$, then $\operatorname{Orb}_G(s)$ is a three-dimensional real-analytic manifold properly embedded in G.*

Moreover, in either case, the tangent space to $\operatorname{Orb}_G(s)$ at s is $\mathcal{V}(s)$ and

$$\mathcal{V}(s) = \operatorname{ran} e_s'(\mathrm{id}_\mathbb{D}). \tag{1.14}$$

In the sequel the notation $T_s \operatorname{Orb}_G(s)$ denotes the complex tangent space if $s \in \mathcal{R}$ and the real tangent space if $s \notin \mathcal{R}$. Thus, for all $s \in G$,

$$T_s \operatorname{Orb}_G(s) = \operatorname{ran} e_s'(\mathrm{id}_\mathbb{D}). \tag{1.15}$$

Proof. Consider $s \in G$. We shall calculate the rank of the real linear operator $e_s'(m)$ for $m \in \operatorname{Aut}\mathbb{D}$. Let $\iota : \operatorname{Orb}_G(s) \to \mathbb{C}_r^2$ denote the inclusion map.

Lemma 1.7. *For any tangent vector (r, α) at $(0, 0)$ to $(-\pi, \pi) \times \mathbb{D}$, let $v_{r,\alpha}(s)$ denote the tangent vector $(e_s \circ \varphi_1^{-1})'(0, 0)(r, \alpha)$ in $T_sG \subset \mathbb{C}^2$. Then*

$$v_{r,\alpha}(s) = ir \begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix} - \alpha \begin{pmatrix} 2 \\ s^1 \end{pmatrix} + \bar{\alpha} \begin{pmatrix} (s^1)^2 - 2s^2 \\ s^1 s^2 \end{pmatrix}. \tag{1.16}$$

Proof. We have $(r, \alpha) \in \mathbb{R} \times \mathbb{C}$. Define a path $\kappa(t) = (tr, t\alpha)$ in $(-\pi, \pi) \times \mathbb{D}$ for $|t| < \varepsilon$, where ε is small enough.

Then let

$$f_s = \iota \circ e_s \circ \varphi_1^{-1} : (-1, 1) \times \mathbb{D} \to \mathbb{C}_r^2$$

and define $v_{r,\alpha}(s) \in \mathbb{C}^2$ by the formula

$$\begin{aligned} v_{r,\alpha}(s) &= \frac{\mathrm{d}}{\mathrm{d}t} f_s \circ \kappa(t)\big|_{t=0} && (1.17) \\ &= \frac{\mathrm{d}}{\mathrm{d}t} \iota \circ e_s \circ \varphi_1^{-1} \circ \kappa(t)\big|_{t=0} \\ &= \frac{\mathrm{d}}{\mathrm{d}t} \iota \circ e_s \circ m_{\kappa(t)}\big|_{t=0} \\ &= \frac{\mathrm{d}}{\mathrm{d}t} \iota \circ e_s \circ m_{tr,t\alpha}\big|_{t=0} \\ &= \frac{\mathrm{d}}{\mathrm{d}t} \iota \circ \gamma_{m_{tr,t\alpha}}(s)\big|_{t=0}. && (1.18) \end{aligned}$$

From equation (1.8), for any $z \in \mathbb{D}$,

$$\frac{\mathrm{d}}{\mathrm{d}t} m_{tr,t\alpha}(z)\big|_{t=0} = irz - \alpha + \bar{\alpha} z^2.$$

Hence, by equations (1.1) and (1.18), if $s = (z_1 + z_2, z_1 z_2)$,

$$\begin{aligned} v_{r,\alpha}(s) &= \frac{\mathrm{d}}{\mathrm{d}t} \begin{pmatrix} m_{tr,t\alpha}(z_1) + m_{tr,t\alpha}(z_2) \\ m_{tr,t\alpha}(z_1) m_{tr,t\alpha}(z_2) \end{pmatrix}\Bigg|_{t=0} \\ &= \begin{pmatrix} irz_1 - \alpha + \bar{\alpha}(z_1)^2 + irz_2 - \alpha + \bar{\alpha}(z_2)^2 \\ (irz_1 - \alpha + \bar{\alpha}(z_1)^2) z_2 + (irz_2 - \alpha + \bar{\alpha}(z_2)^2) z_1 \end{pmatrix} \\ &= ir \begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix} - \alpha \begin{pmatrix} 2 \\ s^1 \end{pmatrix} + \bar{\alpha} \begin{pmatrix} (s^1)^2 - 2s^2 \\ s^1 s^2 \end{pmatrix}. && (1.19) \end{aligned}$$

□

Continuing the proof of Theorem 1.6, by the Chain Rule we have, from equation (1.17),

$$\begin{aligned} v_{r,\alpha}(s) &= f'_s(\kappa(0))\kappa'(0) \\ &= f'_s(0,0)\begin{pmatrix} r \\ \alpha \end{pmatrix}. \end{aligned} \tag{1.20}$$

Thus the range of the real linear map $f'_s(0,0) : \mathbb{R} \times \mathbb{C} \to \mathbb{C}^2$ is the set

$$\begin{aligned} \operatorname{ran} f'_s(0,0) &= \{v_{r,\alpha} : r \in \mathbb{R}, \alpha \in \mathbb{C}\} \\ &= \left\{ ir\begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix} - \alpha\begin{pmatrix} 2 \\ s^1 \end{pmatrix} + \bar{\alpha}\begin{pmatrix} (s^1)^2 - 2s^2 \\ s^1 s^2 \end{pmatrix} : r \in \mathbb{R},\ \alpha \in \mathbb{C} \right\}. \end{aligned}$$

On taking (r, α) to be successively $(1, 0)$, $(0, -1)$ and $(0, -i)$ we find that, for any $s \in G$,

$$\operatorname{ran} f'_s(0,0) = \mathcal{V}(s),$$

the real vector space introduced in equation (1.13). Thus

$$\iota'(s) \operatorname{ran} e'_s(\mathrm{id}_{\mathbb{D}}) = \operatorname{ran} f'_s(0,0) = \mathcal{V}(s) \tag{1.21}$$

for all $s \in G$. In the sequel we shall suppress the inclusion map $\iota'(s)$ and regard $\operatorname{ran} e'_s(\mathrm{id}_{\mathbb{D}})$ as a subspace of $\mathbb{C}^2_r$.

Now consider $s \in G \setminus \mathcal{R}$. By Lemma 1.8 below, $\dim_{\mathbb{R}} \mathcal{V}(s) = 3$, and so, by equation (1.21), $e'_s(\mathrm{id}_{\mathbb{D}})$ has rank 3. We claim that $e'_s(m)$ has rank 3 for all $m \in \operatorname{Aut} \mathbb{D}$. Indeed, on differentiating the relation

$$e_s(m) = \gamma_m(s) = \gamma_m \circ \gamma_{\mathrm{id}_{\mathbb{D}}}(s) = \gamma_m \circ e_s(\mathrm{id}_{\mathbb{D}}),$$

we find (since $e_s(\mathrm{id}_{\mathbb{D}}) = s$) that

$$e'_s(m) = \gamma'_m(s) e'_s(\mathrm{id}_{\mathbb{D}}).$$

Since γ_m is an automorphism of G, $\gamma'_m(s)$ is a nonsingular real linear transformation of $\mathbb{C}^2$. Thus

$$\operatorname{rank}_{\mathbb{R}} e'_s(m) = \operatorname{rank}_{\mathbb{R}} e'_s(\mathrm{id}_{\mathbb{D}}) = 3$$

for every $m \in \operatorname{Aut} \mathbb{D}$.

We wish to deduce that $\operatorname{Orb}_G(s)$ is a real 3-dimensional C^∞-manifold which (modulo the identification map $\iota'(s)$) lies in $\mathbb{C}^2$. The following statement is [24, Theorem 5.2].

A subset M of $\mathbb{R}^n$ is a k-dimensional manifold if and only if, for every point $s \in M$ there exist an open neighborhood V of s in $\mathbb{R}^n$, an open set W in $\mathbb{R}^k$ and an injective differentiable function $f : W \to \mathbb{R}^n$ such that

(1) *$f(W) = M \cap V$,*
(2) *$f'(y)$ has rank k for every $y \in W$.*

We shall apply this criterion in the case $n = 4,\ k = 3,\ M = \mathrm{Orb}_G(s)$. Consider any point $e_s(m) \in \mathrm{Orb}_G(s)$, where $m \in \operatorname{Aut}\mathbb{D}$, say $m \in U_j$, $j = 1$ or 2. By Proposition 1.5, e_s is a local homeomorphism, and so we may choose an open neighborhood N of m in U_j such that $e_s|N$ is a homeomorphism from N to an open subset of $\mathrm{Orb}_G(s)$. Since $\mathrm{Orb}_G(s)$ has the relative topology induced by G, there is an open set V in $\mathbb{R}^4$ such that $e_s(N) = V \cap \mathrm{Orb}_G(s)$.

Let $W = \varphi_j(N)$. Then the map $f = e_s \circ \varphi_j^{-1}$ satisfies conditions (1) and (2). It follows that $\mathrm{Orb}_G(s)$ is a real 3-dimensional C^∞ manifold in $\mathbb{C}^2$.

The linear map $(e_s \circ \varphi_1^{-1})'(0,0)$ maps the tangent space $T_{(0,0)}(-\pi,\pi)\times\mathbb{D}$ into the tangent space $T_s\,\mathrm{Orb}_G(s) \subset \mathbb{C}^2$. We have seen that the range of $(e_s \circ \varphi_1^{-1})'(0,0)$ is $\mathcal{V}(s)$. Hence

$$\mathcal{V}(s) \subseteq T_s\,\mathrm{Orb}_G(s).$$

Since both spaces have real dimension 3, the inclusion holds with equality.

In the case that $s \in \mathcal{R}$, say $s = (2\zeta, \zeta^2)$ for some $\zeta \in \mathbb{D}$,

$$\mathrm{Orb}_G(s) = \mathcal{R} = \{(2z, z^2) : z \in \mathbb{D}\},$$

which is a one-dimensional complex manifold properly embedded in G by the map $R : \mathbb{D} \to G$. The complex tangent space to $\mathcal{R}$ at s is $\mathbb{C}(1,\zeta)$, and, by equation (1.13),

$$\begin{aligned}\mathcal{V}(s) &= \operatorname{span}_{\mathbb{R}} \left\{ i\zeta \begin{pmatrix}1\\ \zeta\end{pmatrix}, (1-\zeta^2)\begin{pmatrix}1\\ \zeta\end{pmatrix}, i(1+\zeta^2)\begin{pmatrix}1\\ \zeta\end{pmatrix} \right\} \\ &= \mathbb{C}\begin{pmatrix}1\\ \zeta\end{pmatrix}.\end{aligned}$$

Thus $T_s\,\mathrm{Orb}_G(s) = \mathcal{V}(s)$ in the sense of complex manifolds. □

Lemma 1.8. *For any $s \in G$, the real vector space $\mathcal{V}(s)$ defined by equation* (1.13) *satisfies*

$$\dim_{\mathbb{R}} \mathcal{V}(s) = \begin{cases} 3 & \text{if } (s^1)^2 \neq 4s^2 \\ 2 & \text{if } (s^1)^2 = 4s^2. \end{cases} \tag{1.22}$$

Proof. It is clear from the definition (1.13) that $\mathcal{V}(s)$ is a real vector subspace of $\mathbb{C}^2$ of real dimension at most 3.

Suppose that scalars $\lambda, \mu, \nu \in \mathbb{R}$ satisfy

$$\lambda i \begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix} + \mu \begin{pmatrix} 2 - (s^1)^2 + 2s^2 \\ s^1 - s^1 s^2 \end{pmatrix} + \nu i \begin{pmatrix} 2 + (s^1)^2 - 2s^2 \\ s^1 + s^1 s^2 \end{pmatrix} = 0. \tag{1.23}$$

Multiply on the left by the row matrix $\begin{pmatrix} 2s^2 & -s^1 \end{pmatrix}$ to obtain

$$-((s^1)^2 - 4s^2)\left((1+s^2)\mu + (1-s^2)\nu i\right) = 0. \tag{1.24}$$

Consider the first case in equation (1.22), namely, that $(s^1)^2 \neq 4s^2$ (equivalently, $s \notin \mathcal{R}$). By equation (1.24)

$$(1 + s^2)\mu + (1 - s^2)\nu i = 0,$$

whence

$$\mu + i\nu = -s^2(\mu - i\nu).$$

Since $s \in G$, we have $|s^2| < 1$, and so necessarily $\mu = \nu = 0$. Since $(s^1)^2 \neq 4s^2$, at least one of s^1, s^2 is nonzero, and so, by equation (1.23), $\lambda = 0$. Hence the three spanning vectors for $\mathcal{V}(s)$ in equation (1.13) are linearly independent. We have shown that $\dim_{\mathbb{R}} \mathcal{V}(s) = 3$ when $s \notin \mathcal{R}$.

Next consider a point $s \in \mathcal{R}$. On substituting $s^2 = \frac{1}{4}(s^1)^2$ in equation (1.13) we obtain

$$\mathcal{V}(s) = \operatorname{span}_{\mathbb{R}} \left\{ is^1 \begin{pmatrix} 2 \\ s^1 \end{pmatrix}, (4 - (s^1)^2) \begin{pmatrix} 2 \\ s^1 \end{pmatrix}, i(4 + (s^1)^2) \begin{pmatrix} 2 \\ s^1 \end{pmatrix} \right\}.$$

Since each of these vectors is a complex scalar multiple of the vector $\begin{pmatrix} 2 & s^1 \end{pmatrix}^T$, it follows that $\dim_{\mathbb{R}} \mathcal{V}(s) \leq 2$.

In fact $\dim_{\mathbb{R}} \mathcal{V}(s) = 2$. For otherwise the second and third spanning vectors for $\mathcal{V}(s)$ are linearly dependent over $\mathbb{R}$, and so there exist $\mu, \nu \in \mathbb{R}$, not both zero, such that

$$\mu(4 - (s^1)^2) + \nu i(4 + (s^1)^2) = 0$$

and consequently

$$4(\mu + \nu i) = (s^1)^2(\mu - \nu i).$$

Thus $|s^1| = 2$, contrary to choice of $s \in G$. Therefore

$$\dim_{\mathbb{R}} \mathcal{V}(s) = 2 \quad \text{when } s \in \mathcal{R}.$$

□

1.4. **The sharp direction in G.** By Theorem 1.6, for any $s \in G \setminus \mathcal{R}$, the tangent space $\mathcal{V}(s)$ at s to the orbit $\operatorname{Orb}_G(s)$ is a real 3-dimensional subspace of $\mathbb{C}^2$. Accordingly $\mathcal{V}(s)$ contains a unique 2-real-dimensional subspace that is also a one-dimensional complex subspace of $\mathbb{C}^2$, equal to $\mathcal{V}(s) \cap i\mathcal{V}(s)$. On the other hand, for $s \in \mathcal{R}$, the tangent space $\mathcal{V}(s)$ is already a complex subspace of $\mathbb{C}^2$.

Definition 1.9. *For any $s \in G$, the* sharp direction at s *is the unique nonzero complex subspace of $\mathcal{V}(s)$ in $\mathbb{C}^2$ and is denoted by $s^\sharp$. Thus*

$$s^\sharp = \mathcal{V}(s) \cap i\mathcal{V}(s).$$

The sharp direction is covariant with automorphisms of G, in the following sense.

Proposition 1.10. *If $\gamma \in \operatorname{Aut} G$ and $s \in G$, then*

$$\gamma(s)^\sharp = \gamma'(s)s^\sharp.$$

Proof. Since γ is a differentiable self-map of $\operatorname{Orb}_G(s)$, its derivative $\gamma'(s)$ is a real-linear map between the tangent spaces $\mathcal{V}(s)$ and $\mathcal{V}(\gamma(s))$. Since furthermore $\gamma'(s)$ is a nonsingular complex linear map from $T_sG \sim \mathbb{C}^2$ to $T_{\gamma(s)} \sim \mathbb{C}^2$, it maps the complex subspace $s^\sharp$ of $\mathbb{C}^2$ to a nonzero complex subspace of $\mathbb{C}^2$. Hence $\gamma'(s)s^\sharp$ is a nonzero complex subspace of $\mathcal{V}(\gamma(s))$. Hence $\gamma'(s)s^\sharp = \gamma(s)^\sharp$. □

1.5. **Flat geodesics and the action of** $\operatorname{Aut} G$**.** In the introduction we defined the flat geodesics of G to be the geodesics that meet the royal geodesic $\mathcal{R}$ exacly once and are stabilized by a nontrivial automorphism of G. This definition has the merit that it is geometrical in character, but in practice (for example, to show that the flat geodesics foliate G) it is often simpler to use the fact that the flat geodesics in G are the sets of the form

$$F^\beta \stackrel{\text{def}}{=} \{(\beta + \bar{\beta}z, z) : z \in \mathbb{D}\} \tag{1.25}$$

for some $\beta \in \mathbb{D}$. One can check that the point $s \in G$ lies on the unique F^β with

$$\beta = \frac{s^1 - \overline{s^1}s^2}{1 - |s^2|^2} \in \mathbb{D}. \tag{1.26}$$

More details can be found in [18, 10, 5] and [2, Appendix A].

Let us at least sketch a proof that the set F^β is indeed a flat geodesic according to the definition in the introduction. Firstly, a straightforward calculation shows that any automorphism of G maps F^β to a set of the form $F^{\beta'}$ for some $\beta' \in \mathbb{D}$. Clearly F^β is a complex geodesic in G: for any $\beta \in \mathbb{D}$ the co-ordinate function s^2 is a holomorphic left inverse of the properly embedded analytic disc $z \mapsto (\beta + \bar{\beta}z, z)$ in G. It is simple to check that F^β meets $\mathcal{R}$ exactly once, say at the point $s(\beta) \in \mathcal{R}$. Choose a nontrivial automorphism θ of the analytic disc $\mathcal{R}$ that fixes $s(\beta)$, and let γ be the unique extension of θ to a (necessarily nontrivial) automorphism of G. Then $\gamma(F^\beta) = F^{\beta'}$, and since γ fixes $s(\beta)$, it follows that $F^{\beta'}$ meets $\mathcal{R}$ at $s(\beta)$. Distinct sets F^β are disjoint, and therefore $\beta = \beta'$. That is, F^β is stabilized by a nontrivial automorphism of G.

The converse statement, that every flat geodesic is an F^β, follows from the classification into five types of the complex geodesics in G given in [2, Chapter 7].

We summarize the main geometric properties of flat geodesics.

Proposition 1.11. (1) *Through each point s in G there passes a unique flat geodesic F_s.*

(2) *Every flat geodesic intersects the royal geodesic* $\mathcal{R}$ *in exactly one point.*
(3) *Automorphisms of* G *carry flat geodesics to flat geodesics.*

The following lemma is a reformulation of the first two of these facts.

Lemma 1.12. *The family*

$$\mathcal{F} \stackrel{\text{def}}{=} \{F_s : s \in \mathcal{R}\}$$

is a partition of G.

Definition 1.13. *For any* $s \in G$, *the complex tangent space at* s *to the unique flat geodesic through* s *will be called the* flat direction *at* s, *and will be denoted by* $s^\flat$.

Thus, if $s^1 = \beta + \bar{\beta}s^2$, then

$$s^\flat = \mathbb{C}(\bar{\beta}, 1), \tag{1.27}$$

which is a one-dimensional complex subspace of $\mathbb{C}^2$. The map $s \mapsto s^\flat$ is a covariant line bundle which is a sub-bundle of TG.

Facts (1)-(3) in Proposition 1.11 imply the following description of the action of $\operatorname{Aut} G$ on $\mathcal{F}$.

Lemma 1.14. *If* $\gamma \in \operatorname{Aut} G$ *and* $s \in \mathcal{R}$, *then* $\gamma(F_s) = F_{\gamma(s)}$.

Proof. Fix $\gamma \in \operatorname{Aut} G$ and $s \in \mathcal{R}$. By Fact 3, there exists $t \in \mathcal{R}$ such that $\gamma(F_s) = F_t$, and Condition (i) in Proposition 1.3 implies that $\gamma(s) \in \mathcal{R}$. Therefore $\gamma(s) \in \mathcal{R} \cap F_t$. Hence by Fact 2, $t = \gamma(s)$. □

We shall call $\{F_s : s \in \mathcal{R}\}$ the *flat fibration of* G.

Proposition 1.15. *For all* $s \in G$ *the spaces* $s^\sharp$ *and* $s^\flat$ *are unequal.*

This statement will follow from explicit formulae for the sharp and flat directions. We already know that, for $s \in F^\beta$, $s^\flat$ is given by equation (1.27).

Proposition 1.16. *For any* $\beta \in \mathbb{D}$ *and any* $s \in F^\beta$,

$$s^\sharp = \mathbb{C}\left(1, \frac{\beta - \frac{1}{2}s^1}{1 - \frac{1}{2}\bar{\beta}s^1}\right). \tag{1.28}$$

Proof. Consider $s \in G \setminus \mathcal{R}$. Let

$$v_1 = i\begin{pmatrix} s^1 \\ 2s^2 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 2 - (s^1)^2 + 2s^2 \\ s^1 - s^1 s^2 \end{pmatrix}, \quad v_3 = i\begin{pmatrix} 2 + (s^1)^2 - 2s^2 \\ s^1 + s^1 s^2 \end{pmatrix}.$$

By Theorem 1.6, $\{v_1, v_2, v_3\}$ constitutes a basis for $T_s \operatorname{Orb}_G(s)$. Let

$$c_1 = -2s^1, \quad c_2 = -i(1 - s^2), \quad c_3 = 1 + s^2$$

and note that c_2, c_3 are nonzero. One finds that $c_1v_1 + c_2v_2 + c_3v_3 = 0$, and therefore

$$(\operatorname{Re} c_1)v_1 + (\operatorname{Re} c_2)v_2 + (\operatorname{Re} c_3)v_3 = -i(\operatorname{Im} c_1)v_1 - i(\operatorname{Im} c_2)v_2 - i(\operatorname{Im} c_3)v_3.$$

Hence

$$v \stackrel{\text{def}}{=} (\operatorname{Re} c_1)v_1 + (\operatorname{Re} c_2)v_2 + (\operatorname{Re} c_3)v_3 \neq 0$$

and

$$v \in T_s \operatorname{Orb}_G(s) \cap iT_s \operatorname{Orb}_G(s) = s^\sharp.$$

Further calculation yields the formula

$$v = 2i(1 - |s^2|^2) \begin{pmatrix} 1 - \bar{\beta}\frac{1}{2}s^1 \\ \beta - \frac{1}{2}s^1 \end{pmatrix},$$

in agreement with equation (1.28). This proves the proposition in the case that $s \in G \setminus \mathcal{R}$. For $s = (2z, z^2) \in \mathcal{R}$ equation (1.28) is easily checked. □

The fact that $s^\sharp \neq s^\flat$ can now be verified by a simple comparison of equations (1.28) and (1.27).

Corollary 1.17. *The tangent bundle of G is the direct sum of the sharp bundle and the flat bundle:*

$$T_sG = s^\sharp \oplus s^\flat \qquad \text{for all } s \in G.$$

1.6. **Synchrony in G.** There is a subtle relationship between the action of an automorphism of G on the royal variety and its action on any flat geodesic.

For any complex manifold U and λ in U, denote by $\operatorname{Aut}_\lambda U$ the stabilizer of λ in $\operatorname{Aut} U$ (also known as the isotropic subgroup of $\operatorname{Aut} U$ at λ). For any $s_0 \in \mathcal{R}$, the sets $\mathcal{R}$ and F_{s_0} are embedded analytic discs in G that intersect transversally at the point s_0. Every θ in $\operatorname{Aut}_{s_0} G$ determines an automorphism of the analytic variety $\mathcal{R} \cup F_{s_0}$. For an automorphism of a general variety there need be no connection between the action on two leaves beyond what is implied by the condition that the restrictions of the automorphism to the two leaves must agree at any common point. However, in the context of the domain G, in the light of Condition (ii) in Proposition 1.3, the action of θ on F_{s_0} is uniquely determined by the action of θ on $\mathcal{R}$. The following propositions describe this dependence explicitly.

We denote the unit circle $\{z : |z| = 1\}$ in the complex plane by $\mathbb{T}$. For $\eta \in \mathbb{T}$ let ρ_η denote the element of $\operatorname{Aut}_0 \mathbb{D}$ defined by $\rho_\eta(z) = \eta z$. Clearly $\operatorname{Aut}_0 \mathbb{D} = \{\rho_\eta : \eta \in \mathbb{T}\}$.

Proposition 1.18. *If $s_0 \in \mathcal{R}$ and $\theta \in \operatorname{Aut}_{s_0} G$, then $\theta'(s_0)$ has eigenspaces $T_{s_0}\mathcal{R}$ and $T_{s_0}F_{s_0}$ with corresponding eigenvalues η and η^2 for some $\eta \in \mathbb{T}$.*

Proof. Since θ leaves invariant both $\mathcal{R}$ and F_{s_0}, it follows that $\theta'(s_0)$ leaves invariant the tangent spaces $T_{s_0}\mathcal{R}$ and $T_{s_0}F_{s_0}$. These two one-dimensional tangent spaces are thus eigenspaces of $\theta'(s_0)$.

Observe that γ_{ρ_η} is the restriction to G of the linear operator on $\mathbb{C}^2$ with matrix $\operatorname{diag}\{\eta, \eta^2\}$, and hence

$$\gamma'_{\rho_\eta}(s_0) \sim \operatorname{diag}\{\eta, \eta^2\}.$$

Let $s_0 = (2\alpha, \alpha^2)$. Since $\theta \in \operatorname{Aut}_{s_0} G$, $\theta = \gamma_m$ for some $m \in \operatorname{Aut}\mathbb{D}$ such that $m(\alpha) = \alpha$. Therefore $b_\alpha \circ m \circ b_{-\alpha} \in \operatorname{Aut}_0 \mathbb{D}$, and so there exists $\eta \in \mathbb{T}$ such that

$$m = b_{-\alpha} \circ \rho_\eta \circ b_\alpha.$$

Since $m \mapsto \gamma_m$ is an isomorphism,

$$\theta = \gamma_{b_{-\alpha}} \circ \gamma_{\rho_\eta} \circ \gamma_{b_\alpha}.$$

It follows by the chain rule that

$$\begin{aligned}\theta'(s_0) &= X\gamma'_{\rho_\eta}(0,0)X^{-1} \\ &\sim X \operatorname{diag}\{\eta, \eta^2\}X^{-1}\end{aligned}$$

where $X = \gamma'_{b_{-\alpha}}(0,0)$. But $\operatorname{diag}\{\eta, \eta^2\}$ has eigenspaces $\mathbb{C} \oplus 0$ and $0 \oplus \mathbb{C}$ with corresponding eigenvalues η and η^2. Therefore, $\theta'(s_0)$ has eigenspaces $X(\mathbb{C} \oplus 0)$ and $X(0 \oplus \mathbb{C})$ with corresponding eigenvalues η and η^2. We have

$$\gamma_{b_{-\alpha}}(s) = \frac{(2\alpha + (1+|\alpha|^2)s^1 + 2\bar{\alpha}s^2, s^2 + \alpha s^1 + \alpha^2)}{1 + \bar{\alpha}s^1 + \bar{\alpha}^2 s^2}.$$

Hence

$$X = \gamma'_{b_{-\alpha}}(0,0) \sim (1-|\alpha|^2)\begin{bmatrix} 1 & 2\bar{\alpha} \\ \alpha & 1+|\alpha|^2 \end{bmatrix},$$

and therefore

$$X\begin{pmatrix}\mathbb{C}\\0\end{pmatrix} = \mathbb{C}\begin{pmatrix}1\\\alpha\end{pmatrix} = T_{s_0}\mathcal{R}, \quad X\begin{pmatrix}0\\\mathbb{C}\end{pmatrix} = \mathbb{C}\begin{pmatrix}2\bar{\alpha}\\1+|\alpha|^2\end{pmatrix} = T_{s_0}F_{s_0}.$$

Thus $T_{s_0}\mathcal{R}$ and $T_{s_0}F_{s_0}$ are eigenspaces of $\theta'(s_0)$ with corresponding eigenvalues η, η^2 respectively. □

Proposition 1.19. *Let $s_0 = (2\alpha, \alpha^2)$ for some $\alpha \in \mathbb{D}$ and let $m \in \operatorname{Aut}_\alpha \mathbb{D}$. If g is any proper embedding of $\mathbb{D}$ into G such that $g(\mathbb{D}) = F_{s_0}$ and $g(\alpha) = s_0$, then*

$$\gamma_m \circ g = g \circ m \circ m.$$

Proof. Note that

$$\gamma_m \circ R = R \circ m.$$

This equation implies that

$$\gamma_m'(s_0)R'(\alpha) = m'(\alpha)R'(\alpha),$$

which is to say that $m'(\alpha)$ is the eigenvalue of $\gamma_m'(s_0)$ corresponding to the eigenspace $T_{s_0}\mathcal{R}$. Consequently, by Lemma 1.18,

$$\begin{aligned} &m'(\alpha)^2 \text{ is the eigenvalue of } \gamma_m'(s_0) \\ &\qquad \text{corresponding to the eigenspace } T_{s_0}F_{s_0}. \end{aligned} \tag{1.29}$$

Since g is a proper embedding and $\gamma_m(F_{s_0}) = F_{s_0}$, there exists $b \in \operatorname{Aut}\mathbb{D}$ such that

$$\gamma_m \circ g = g \circ b.$$

As $b(\alpha) = \alpha$, this equation implies that

$$\gamma_m'(s_0)g'(\alpha) = b'(\alpha)g'(\alpha),$$

that is, $b'(\alpha)$ is the eigenvalue of $\gamma_m'(s_0)$ corresponding to the eigenspace $T_{s_0}F_{s_0}$. Therefore, statement (1.29) implies that

$$b'(\alpha) = m'(\alpha)^2 = (m \circ m)'(\alpha),$$

Since b, $m \circ m \in \operatorname{Aut}\mathbb{D}$, $b(\alpha) = \alpha = (m \circ m)(\alpha)$, and $b'(\alpha) = (m \circ m)'(z_0)$, it follows that $b = m \circ m$. □

We describe the phenomena described in Propositions 1.18 and 1.19 as the *synchrony property* of G.

2. Royal manifolds

Perhaps the most far-reaching feature of the complex geometry of G is the existence of the special variety $\mathcal{R}$ with the properties described in Proposition 1.3. We formalize these properties in order to characterize G up to isomorphism.

Definition 2.1. *Let Ω be a complex manifold. We say that D is a* royal disc in Ω *if D is a properly embedded analytic disc in Ω and D satisfies the three conditions of Proposition* 1.3, that is,

(1) every automorphism of Ω leaves D invariant,
(2) every automorphism of Ω is uniquely determined by its values on D,
(3) every automorphism of D has an extension to an automorphism of Ω.

A *royal manifold* is an ordered pair (Ω, D) where Ω is a complex manifold and D is a royal disc in Ω.

The following lemma is straightforward.

Lemma 2.2. *If Ω is a complex manifold and $\Lambda : G \to \Omega$ is a biholomorphic map, then $\Lambda(\mathcal{R})$ is a royal disc in Ω and $(\Omega, \Lambda(\mathcal{R}))$ is a royal manifold.*

The next proposition spells out the analog of formula (1.5) on a general royal manifold.

Proposition 2.3. *Let (Ω, D) be a royal manifold. Then $\operatorname{Aut}\Omega$ is isomorphic to $\operatorname{Aut}\mathbb{D}$. Furthermore, if $d : \mathbb{D} \to \Omega$ is a properly embedded analytic disc such that $d(\mathbb{D}) = D$, then there exists a unique isomorphism $\Theta : \operatorname{Aut}\mathbb{D} \to \operatorname{Aut}\Omega$ such that*

$$\Theta(m) \circ d = d \circ m \qquad \text{for all } m \in \operatorname{Aut}\mathbb{D}. \tag{2.1}$$

The counterpart of the commutative diagram (1.6) is

$$\begin{array}{ccccc} \mathbb{D} & \xrightarrow{d} & D & \xrightarrow{\iota_D} & \Omega \\ {\scriptstyle m}\downarrow & & {\scriptstyle \Theta(m)|D}\downarrow & & {\scriptstyle \Theta(m)}\downarrow \\ \mathbb{D} & \xrightarrow{d} & D & \xrightarrow{\iota_D} & \Omega \end{array} \tag{2.2}$$

where ι_D is the injection of D into Ω.

Proof. Fix a properly embedded analytic disc d such that $d(\mathbb{D}) = D$. For each $\tau \in \operatorname{Aut}\Omega$, Condition (i) in Definition 2.1 implies that there exists a function $\varphi_\tau : \mathbb{D} \to \mathbb{D}$ such that

$$\tau \circ d(z) = d \circ \varphi_\tau(z) \qquad \text{for } z \in \mathbb{D}. \tag{2.3}$$

Clearly, since τ is an automorphism of Ω and d is a properly embedded analytic disc, $\varphi_\tau \in \operatorname{Aut}\mathbb{D}$.

If $\tau_1, \tau_2 \in \operatorname{Aut}\Omega$, then for each $z \in \mathbb{D}$ we see using equation (2.3) that

$$\begin{aligned} d(\varphi_{\tau_2 \circ \tau_1}(z)) &= \tau_2 \circ \tau_1 \circ d(z) \\ &= \tau_2(\tau_1 \circ d(z)) \\ &= \tau_2(d \circ \varphi_{\tau_1}(z)) \\ &= d(\varphi_{\tau_2}(\varphi_{\tau_1}(z))) \\ &= d(\varphi_{\tau_2} \circ \varphi_{\tau_1}(z)). \end{aligned}$$

This relation proves that the map $\Psi : \operatorname{Aut}\Omega \to \operatorname{Aut}\mathbb{D}$ given by

$$\Psi(\tau) = \varphi_\tau \tag{2.4}$$

is a homomorphism of automorphism groups.

If $\tau_1, \tau_2 \in \operatorname{Aut}\Omega$ and $\varphi_{\tau_1}(z) = \varphi_{\tau_2}(z)$ for all $z \in \mathbb{D}$, then equation (2.3) implies that $\tau_1(d(z)) = \tau_2(d(z))$ for all $z \in \mathbb{D}$, which is to say

that τ_1 and τ_2 agree on D. Hence, by Condition (ii) in Definition 2.1, $\tau_1 = \tau_2$. This proves that Ψ is injective.

Consider any $b \in \operatorname{Aut}\mathbb{D}$. The map

$$d(z) \mapsto d(b(z)) \in D \qquad \text{for } z \in \mathbb{D}$$

is an automorphism of the complex manifold D. Condition (iii) in Definition 2.1 implies that there exists $\tau \in \operatorname{Aut}\Omega$ such that $\tau(d(z)) = d \circ b(z)$ for all $z \in \mathbb{D}$. But then

$$d(b(z)) = \tau(d(z)) = d(\varphi_\tau(z))$$

for all $z \in \mathbb{D}$, so that $\varphi_\tau = b$. This proves that Ψ is surjective from $\operatorname{Aut}\Omega$ onto $\operatorname{Aut}\mathbb{D}$.

We have shown that Ψ is an isomorphism of groups. In particular, the first assertion of Proposition 2.3 (that $\operatorname{Aut}\Omega$ is isomorphic to $\operatorname{Aut}\mathbb{D}$) is proven. To define an isomorphism Θ satisfying the second assertion of the proposition, let $\Theta = \Psi^{-1}$. Then Θ is an isomorphism from $\operatorname{Aut}\mathbb{D}$ onto $\operatorname{Aut}\Omega$, and equation (2.1) follows from the relation (2.3).

To see that Θ is unique consider $m \in \operatorname{Aut}\mathbb{D}$ and observe that if Θ_1 and Θ_2 are isomorphisms satisfying equation (2.3), then $\Theta_1(m)(d(z)) = \Theta_2(m)(d(z))$ for all $z \in \mathbb{D}$. Since $\Theta_1(m)$ and $\Theta_2(m)$ agree on D, Condition (ii) in Definition 2.1 imples that $\Theta_1(m) = \Theta_2(m)$. As m is arbitrary, $\Theta_1 = \Theta_2$. □

In the light of Proposition 2.3 we adopt the following definition.

Definition 2.4. *Let (Ω, D) be a royal manifold. We say that (d, Θ) is a* concomitant pair for (Ω, D) *if $d : \mathbb{D} \to \Omega$ is a proper analytic embedding, $d(\mathbb{D}) = D$, and $\Theta : \operatorname{Aut}\mathbb{D} \to \operatorname{Aut}\Omega$ is an isomorphism of groups that satisfies, for all $m \in \operatorname{Aut}\mathbb{D}$,*

$$\Theta(m) \circ d = d \circ m$$

as in equation (2.1).

In other words, (d, Θ) is a concomitant pair for (Ω, D) if the diagram (2.2) commutes for every $m \in \operatorname{Aut}\mathbb{D}$.

Remark 2.5. Concomitant pairs are essentially unique in the following sense. If (Ω, D) is a royal manifold and (d_0, Θ_0) is a concomitant pair for (Ω, D), then (d, Θ) is a concomitant pair for (Ω, D) if and only if there exists $b \in \operatorname{Aut}\mathbb{D}$ such that $d = d_0 \circ b$ and $\Theta = \Theta_0 \circ I_b$, where I_b denotes the inner automorphism of $\operatorname{Aut}\mathbb{D}$ defined by $I_b(m) = b \circ m \circ b^{-1}$.

As a companion to Lemma 2.2 we have the following equally straightforward lemma.

Lemma 2.6. *If Ω is a complex manifold, $\Lambda : G \to \Omega$ is a biholomorphic map, $d = \Lambda \circ R$ and*

$$\Theta(m) = \Lambda \circ \gamma_m \circ \Lambda^{-1}, \qquad \text{for } m \in \operatorname{Aut} \mathbb{D}, \tag{2.5}$$

then (d, Θ) is a concomitant pair for $(\Omega, \Lambda(\mathcal{R}))$.

Definition 2.7. *A concomitant pair (d, Θ) for a royal manifold (Ω, D) is* consistent with *a bijective map $\Lambda : G \to \Omega$ if $d = \Lambda \circ R$ and $\Theta(m) = \Lambda \circ \gamma_m \circ \Lambda^{-1}$ for all $m \in \operatorname{Aut} \mathbb{D}$.*

2.1. **Regularity properties of royal manifolds.**

Definition 2.8. *Let (Ω, D) be a royal manifold and (d, Θ) a concomitant pair. We say that (Ω, D) is a* regular *royal manifold if*

(1) *$\Theta : \operatorname{Aut} \mathbb{D} \to \operatorname{Aut} \Omega$ is differentiable;*
(2) *for every $\lambda \in \Omega \setminus D$, the stabilizer of λ in $\operatorname{Aut} \Omega$ is finite, and*
(3) *for every $\lambda \in \Omega \setminus D$, $e'_\lambda(\mathrm{id}_{\mathbb{D}})$ is an invertible real-linear map, where $e_\lambda : \operatorname{Aut} \mathbb{D} \to \Omega$ is defined by*

$$e_\lambda(m) = \Theta(m)(\lambda). \tag{2.6}$$

Remark 2.9. If the complex manifold Ω is isomorphic to a bounded taut domain [18], then Θ is automatically differentiable – indeed, by a theorem of H. Cartan [9], real analytic.

$e'_\lambda(\mathrm{id}_{\mathbb{D}})$ is a real-linear map between real tangent spaces,

$$e'_\lambda(\mathrm{id}_{\mathbb{D}}) : \operatorname{Lie}(\operatorname{Aut} \mathbb{D}) \to T_\lambda \Omega.$$

Conditions (1) to (3) are certainly necessary for Ω to be biholomorphic to G. They do not depend on the choice of concomitant pair for (Ω, D).

The following statement is simple to prove.

Proposition 2.10. *If Ω and Λ are as in Lemma* 2.6 *then $(\Omega, \Lambda(\mathcal{R}))$ is a regular royal manifold.*

There is an analog of Proposition 1.5 for G.

Proposition 2.11. *If (Ω, D) is a regular royal manifold then, for any $\lambda \in \Omega \setminus D$, the map $e_\lambda : \operatorname{Aut} \mathbb{D} \to \operatorname{Orb}_\Omega(\lambda)$ is a local homeomorphism and an N-to-one covering map, where N is the order of the stabilizer group of λ in $\operatorname{Aut} \Omega$.*

Proof. Let (d, Θ) be a concomitant pair for (Ω, D) and let H be the stabilizer of λ in $\operatorname{Aut} \Omega$. By condition (2) in Definition 2.8, H is a finite subgroup of $\operatorname{Aut} \Omega$. For any $m_1, m_2 \in \operatorname{Aut} \mathbb{D}$,

$$e_\lambda(m_1) = e_\lambda(m_2) \Leftrightarrow m_2^{-1} \circ m_1 \in \Theta^{-1}(H).$$

Since Θ is bijective, $|\Theta^{-1}(H)| = N$. It follows that e_λ is an N-to-one map.

To prove that e_λ is a local homeomorphism, consider any point $e_\lambda(\beta)$ of $\mathrm{Orb}_\Omega(\lambda)$, where $\beta \in \operatorname{Aut}\mathbb{D}$. Choose a neighborhood U of $\mathrm{id}_\mathbb{D}$ such that

$$\{m_2^{-1} \circ m_1 : m_1, m_2 \in U\} \cap \Theta^{-1}(H) = \{\mathrm{id}_\mathbb{D}\}.$$

Let V be a compact neighborhood of $\mathrm{id}_\mathbb{D}$ contained in U. Then $\beta \circ V$ is a compact neighborhood of β on which e_λ is injective, and so $e_\lambda | \beta \circ V$ is a homeomorphism onto its range. Thus e_λ is a local homeomorphism. □

Remark 2.12. The natural analog of equation (1.22), to wit,

$$\operatorname{rank}_\mathbb{R} e'_\lambda(\mathrm{id}_\mathbb{D}) = 2 \quad \text{if } \lambda \in D,$$

is not required in Definition 2.8, since the condition holds automatically, as is clear from Proposition 2.14 below.

Proposition 2.13. *Let (Ω, D) be a regular royal manifold with concomitant pair (d, Θ).*

(1) *If $\lambda \in D$ then $\mathrm{Orb}_\Omega(\lambda)$ is a one-dimensional complex manifold properly embedded in Ω.*
(2) *If $\lambda \in \Omega \setminus D$, then $\mathrm{Orb}_\Omega(\lambda)$ is a three-dimensional real manifold properly embedded in Ω.*

In either case,

$$\operatorname{ran} e'_\lambda(\mathrm{id}_\mathbb{D}) = T_\lambda \mathrm{Orb}_\Omega(\lambda). \tag{2.7}$$

Proof. (1) Let $\lambda \in D$. By conditions (1) and (3) in Definition 2.1, $\mathrm{Orb}_\Omega(\lambda) = D$, which is by hypothesis a properly embedded analytic disc in Ω and therefore a one-dimensional complex manifold.

(2) The proof that $\mathrm{Orb}_\Omega(\lambda)$ is a 3-dimensional real manifold for any $\lambda \in \Omega \setminus D$ is almost identical to the proof of the corresponding statement for G in Theorem 1.6, and so we omit it. □

For a domain U in $\mathbb{C}^n$, when necessary we shall write U_r for U considered as a $2n$-dimensional real manifold and U_c for U as a complex manifold. For $p \in U$ the spaces T_pU_r, T_pU_c are respectively the real and complex tangent spaces to U at p. We regard elements of T_pU_r as point derivations at p on the algebra $C^1_p(U)$ of germs at p of real-valued C^1 functions on U_r. Elements of T_pU_c are point derivations at p on the algebra $\mathcal{O}_p(U)$ of germs at p of holomorphic functions on U_c. We express the action of a point derivation δ on a germ g of the appropriate type by the notation $\langle g, \delta \rangle$.

The complexification $(T_pM)_{\mathbb{C}}$ of the real tangent space at p to a real manifold M is the complex vector space comprising the point derivations at p on the complex algebra $C^1_p(M,\mathbb{C})$ of germs at p of *complex*-valued C^1 functions on M. If $\delta \in (T_pM)_{\mathbb{C}}$ then the functional $\operatorname{Re}\delta$ on $C^1_p(M)$ defined by

$$\langle g, \operatorname{Re}\delta\rangle = \operatorname{Re}\langle g + i0, \delta\rangle$$

is a point derivation, that is, a member of T_pM. We also define $\operatorname{Im}\delta \in T_pM$ to be $-\operatorname{Re}(i\delta)$. In the reverse direction, for a tangent vector $\delta \in T_pM$ we denote by $\delta_{\mathbb{C}}$ the complexification of δ, so that, for any complex-valued C^1 function h in a neighborhood of p,

$$\langle h, \delta_{\mathbb{C}}\rangle = \langle \operatorname{Re} h, \delta\rangle + i\langle \operatorname{Im} h, \delta\rangle . \tag{2.8}$$

Then, for $\delta \in (T_pM)_{\mathbb{C}}$, the relation $\delta = (\operatorname{Re}\delta)_{\mathbb{C}} + i(\operatorname{Im}\delta)_{\mathbb{C}}$ holds. Note that, for $\delta \in T_pM$, we have $\operatorname{Re}(\delta_{\mathbb{C}}) = \delta$.

Furthermore, since every holomorphic function on U_c is a $\mathbb{C}$-valued C^1 function on U_r, every tangent vector $\delta \in (T_pU_r)_{\mathbb{C}}$ determines by restriction an element $\delta|\mathcal{O}$ of T_pU_c.

We can summarize the various tangent spaces and their inclusions in the diagram

$$\begin{array}{ccccc} C^1_p(U_r) & \hookrightarrow & C^1_p(U_r,\mathbb{C}) & \hookleftarrow & \mathcal{O}_p(U_c) \\ & & & & \\ T_pU_r & \underset{\operatorname{Re},\operatorname{Im}}{\overset{\cdot_{\mathbb{C}}}{\rightleftharpoons}} & (T_pU_r)_{\mathbb{C}} & \overset{\cdot|\mathcal{O}}{\rightarrow} & T_pU_c \end{array} \tag{2.9}$$

The vector spaces in the bottom row are respectively real of dimension $2n$, complex of dimension $2n$ and complex of dimension n. The composition of $\cdot_{\mathbb{C}}$ and $\cdot|\mathcal{O}$ is a natural real-linear map

$$\kappa : T_pU_r \to T_pU_c, \qquad \text{where } \kappa\delta = \delta_{\mathbb{C}}|\mathcal{O}_p(U_c).$$

For $\delta \in T_pU_r$, the complex tangent vector $\kappa\delta$ satisfies, for $g \in \mathcal{O}_p(U_c)$,

$$\begin{aligned} \langle g, \kappa\delta\rangle &= \langle g, \delta_{\mathbb{C}}\rangle \\ &= \langle \operatorname{Re} g, \delta\rangle + i\langle \operatorname{Im} g, \delta\rangle , \end{aligned} \tag{2.10}$$

the last line by equation (2.8). In terms of the traditional co-ordinates $z^j = x^j + iy^j$ in a neighborhood of p,

$$\kappa\left(\left(\frac{\partial}{\partial x^j}\right)_p\right) = \left(\frac{\partial}{\partial z^j}\right)_p .$$

Therefore κ is surjective, and since both domain and codomain have real dimension $2n$, it follows that κ is a real linear isomorphism.

For $\lambda \in D$ the orbit $\operatorname{Orb}_\Omega(\lambda)$ is the royal disc D, which is a properly embedded analytic disc under the complex structure induced by Ω. Let

the evaluation map e_λ be as in Definition 2.8, so that $e_\lambda(m) = \Theta(m)(\lambda)$. The derivative $e'_\lambda(\mathrm{id}_{\mathbb{D}})$ is then a real-linear map from $\mathrm{Lie}(\mathrm{Aut}\ \mathbb{D})$ to the real tangent space $T_\lambda D_r$, and so if $\kappa : T_\lambda D_r \to T_\lambda D_c$ is the natural embedding of real and complex tangent spaces, then

$$\kappa e'_\lambda(\mathrm{id}_{\mathbb{D}}) : \mathrm{Lie}(\mathrm{Aut}\ \mathbb{D}) \to T_\lambda D_c \tag{2.11}$$

is a real-linear map from a 3-dimensional real space to a 1-dimensional complex space. In fact this map is surjective.

Proposition 2.14. *Let (Ω, D) be a regular royal manifold and let $\lambda \in D$. Then*

$$\operatorname{ran} \kappa e'_\lambda(\mathrm{id}_{\mathbb{D}}) = T_\lambda D_c. \tag{2.12}$$

Proof. Let (d, Θ) be a concomitant pair for (Ω, D).

Since $\lambda \in D$, there exists $z_0 \in \mathbb{D}$ such that $\lambda = d(z_0)$. Consider a tangent vector δ to $\mathrm{Aut}\,\mathbb{D}$ at $\mathrm{id}_{\mathbb{D}}$. We shall calculate $e'_\lambda(\mathrm{id}_{\mathbb{D}})\delta$. For any germ g of real-valued C^1 functions on Ω at λ,

$$\langle g, e'_\lambda(\mathrm{id}_{\mathbb{D}})\delta\rangle = \langle g \circ e_\lambda, \delta\rangle .$$

For $m \in \mathrm{Aut}\,\mathbb{D}$,

$$\begin{aligned} g \circ e_\lambda(m) &= g \circ \Theta(m) \circ d(z_0) \\ &= g \circ d \circ m(z_0). \end{aligned}$$

Hence

$$\langle g, e'_\lambda(\mathrm{id}_{\mathbb{D}})\delta\rangle = \langle g \circ d \circ m(z_0), \delta\rangle ,$$

where $g \circ d \circ m(z_0)$ is understood as a real-valued function of m, with z_0 fixed. Recall the local co-ordinates r, α for $\mathrm{Aut}\,\mathbb{D}$ introduced in equation (1.8). Here we shall write $\alpha = \xi + i\eta$, with $\xi, \eta \in \mathbb{R}$ and shall use the local co-ordinates r, ξ, η for $\mathrm{Aut}\,\mathbb{D}$. Note that $\mathrm{id}_{\mathbb{D}} \in \mathrm{Aut}\,\mathbb{D}$ corresponds to the local co-ordinates $r = \xi = \eta = 0$. By an elementary calculation,

$$\left\langle g \circ d(m_{r,\alpha}(z_0)), \left(\frac{\partial}{\partial r}\right)_{\mathrm{id}_{\mathbb{D}}} \right\rangle = (g \circ d)'(z_0) i z_0,$$

$$\left\langle g \circ d(m_{r,\alpha}(z_0)), \left(\frac{\partial}{\partial \xi}\right)_{\mathrm{id}_{\mathbb{D}}} \right\rangle = (g \circ d)'(z_0)(z_0^2 - 1),$$

$$\left\langle g \circ d(m_{r,\alpha}(z_0)), \left(\frac{\partial}{\partial \eta}\right)_{\mathrm{id}_{\mathbb{D}}} \right\rangle = (g \circ d)'(z_0)(-i)(z_0^2 + 1).$$

Here $(g \circ d)'(z_0)$ is a real linear functional on $T_{z_0}\mathbb{D}_r$. If

$$\delta = \left(\delta_1 \frac{\partial}{\partial r} + \delta_2 \frac{\partial}{\partial \xi} + \delta_3 \frac{\partial}{\partial \eta}\right)_{\mathrm{id}_{\mathbb{D}}}$$

for some real $\delta_1, \delta_2, \delta_3$, then

$$\langle g, e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle = (g\circ d)'(z_0)\left(\delta_1 i z_0 + \delta_2(z_0^2-1) - i\delta_3(z_0^2+1)\right). \quad (2.13)$$

Now we calculate $\kappa e'_\lambda(\mathrm{id}_\mathbb{D}) \in T_\lambda D_c$. To this end consider any $h \in \mathcal{O}_\lambda(D)$. By equation (2.10),

$$\begin{aligned}\langle h, \kappa e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle &= \langle h, (e'_\lambda(\mathrm{id}_\mathbb{D})\delta)_\mathbb{C}\rangle \\ &= \langle \operatorname{Re} h, e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle + i\langle \operatorname{Im} h, e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle.\end{aligned}$$

Thus, by equation (2.13),

$$\begin{aligned}\langle h, \kappa e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle &= \langle \operatorname{Re} h, e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle + i\langle \operatorname{Im} h, e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle \\ &= ((\operatorname{Re} h\circ d)'(z_0) + i(\operatorname{Im} h\circ d)'(z_0))\left(\delta_1 i z_0 + \delta_2(z_0^2-1) - i\delta_3(z_0^2+1)\right) \\ &= (h\circ d)'(z_0)\left(\delta_1 i z_0 + \delta_2(z_0^2-1) - i\delta_3(z_0^2+1)\right).\end{aligned}$$

Since d is only determined up to composition with an automorphism of $\mathbb{D}$, no generality is lost by the assumption that $z_0 = 0$. Hence

$$\langle h, \kappa e'_\lambda(\mathrm{id}_\mathbb{D})\delta\rangle = -(\delta_2 + i\delta_3)(h\circ k)'(0).$$

On the other hand,

$$\left\langle h, d'(0)\left(\frac{d}{dz}\right)_0\right\rangle = \left\langle h\circ d, \left(\frac{d}{dz}\right)_0\right\rangle = (h\circ d)'(0),$$

and therefore

$$\kappa e'_\lambda(\mathrm{id}_\mathbb{D})\delta = -(\delta_2 + i\delta_3)d'(0)\left(\frac{d}{dz}\right)_0.$$

Thus

$$\operatorname{ran}\kappa e'_\lambda(\mathrm{id}_\mathbb{D}) = \mathbb{C}d'(0)\left(\frac{d}{dz}\right)_0 = T_\lambda D_c.$$

□

2.2. **Flat fibrations over royal discs.** In this subsection we shall formalize the consequences for isomorphs of G of the flat fibration of G described in Subsection 1.5.

Definition 2.15. *Let (Ω, D) be a royal manifold. If $\mathcal{E} = \{E_\lambda\}_{\lambda\in D}$ is a family of subsets of Ω indexed by D, then we say that $\mathcal{E}$ is a* flat fibration of Ω over D *if*

(1) *for each $\lambda \in D$, E_λ is a properly embedded analytic disc in Ω such that $E_\lambda \cap D = \{\lambda\}$;*
(2) *$\mathcal{E}$ is a partition of Ω, and*
(3) *if $\theta \in \operatorname{Aut}\Omega$ and $\lambda \in D$, then $\theta(E_\lambda) = E_{\theta(\lambda)}$.*

We say that $(\Omega, D, \mathcal{E})$ *is a* flatly fibered royal manifold *if* (Ω, D) *is a royal manifold and* $\mathcal{E}$ *is a flat fibration of* Ω *over* D*. We define the* flat direction $\lambda^\flat$ *at a point* λ *in* Ω *to be the tangent space at* λ *to* E_μ *where* $\mu \in D$ *and* $\lambda \in E_\mu$*.*

Clearly, if $(\Omega, D, \mathcal{E})$ is a flatly fibered royal manifold then Ω has complex dimension 2.

Note that if (Ω, D) happens to be $(G, \mathcal{R})$ then the definition of the flat direction is consistent with that given earlier in Definition 1.13.

Lemma 2.16. *Let* Ω *be a complex manifold, let* $\Lambda : G \to \Omega$ *be a biholomorphic map, let* $D = \Lambda(\mathcal{R})$ *and let*

$$E_{\Lambda(s)} = \Lambda(F_s) \quad \text{for every } s \in \mathcal{R},$$

where $\{F_s : s \in \mathcal{R}\}$ *is the flat fibration of* G*. Then*

$$\mathcal{E} = \{E_{\Lambda(s)} : s \in \mathcal{R}\}$$

is a flat fibration of the royal manifold (Ω, D) *over* D*, and* $(\Omega, D, \mathcal{E})$ *is a flatly fibered royal manifold.*

Proof. By Lemma 2.2, (Ω, D) is a royal manifold. Since Λ is a bijection from $\mathcal{R}$ to D, we may write $\mathcal{E} = \{E_\lambda : \lambda \in D\}$. Properties (1) and (2) of Definition 2.15 for the sets E_λ follow from the corresponding properties of the sets F_s for G. If $\theta \in \operatorname{Aut}\Omega$ then $\Lambda^{-1} \circ \theta \circ \Lambda \in \operatorname{Aut} G$. Consider any $s \in \mathcal{R}$ and $\lambda = \Lambda(s) \in D$. We have

$$\theta(E_\lambda) = \theta \circ \Lambda(F_s) = \Lambda \circ (\Lambda^{-1} \circ \theta \circ \Lambda)(F_s) = \Lambda(F_{\Lambda^{-1} \circ \theta \circ \Lambda(s)}),$$

the last step by virtue of property (3) for the flat geodesics as a flat fibration of $(G, \mathcal{R})$. Write $\tilde{s} = \Lambda^{-1} \circ \theta(\lambda)$. Now

$$\Lambda(F_{\Lambda^{-1} \circ \theta \circ \Lambda(s)}) = \Lambda(F_{\Lambda^{-1} \circ \theta(\lambda)}) = \Lambda(F_{\tilde{s}}) = E_{\Lambda(\tilde{s})} = E_{\theta(\lambda)}.$$

Hence

$$\theta(E_\lambda) = E_{\theta(\lambda)} \quad \text{for all } \lambda \in D.$$

Thus the partition $\mathcal{E}$ has the property (3) of Definition 2.15, and so $(\Omega, D, \mathcal{E})$ is a flatly fibered royal manifold. □

2.3. **Synchrony in** Ω**.** Lemma 1.19 suggests the following definition concerning the action of $\operatorname{Aut}\Omega$ on a flat fibration.

Definition 2.17. *Let* $(\Omega, D, \mathcal{E})$ *be a flatly fibered royal manifold with concomitant pair* (d, Θ)*, let* $\lambda_0 \in D$ *and let* $\lambda_0 = d(z_0)$ *for some* $z_0 \in \mathbb{D}$*.*

We say that $(\Omega, D, \mathcal{E})$ *is* synchronous at λ_0 *if, for some properly embedded analytic disc* $f : \mathbb{D} \to \Omega$ *such that* $f(z_0) = \lambda_0$ *and* $f(\mathbb{D}) = E_{\lambda_0}$*,*

$$\Theta(m) \circ f = f \circ m \circ m \tag{2.14}$$

for all $m \in \operatorname{Aut}_{z_0} \mathbb{D}$.

Remark 2.18. If $(\Omega, D, \mathcal{E})$ is as in the definition, then the synchrony of $(\Omega, D, \mathcal{E})$ at λ_0 depends neither on the choice of (d, Θ) nor the choice of f.

For suppose $(\Omega, D, \mathcal{E})$ is synchronous at λ_0 with respect to the concomitant pair (d, Θ) and let (f_1, Θ_1) be a second concomitant pair. By Remark 2.5 there exists $b \in \operatorname{Aut} \mathbb{D}$ such that $f_1 = f \circ b$ and $\Theta_1 = \Theta \circ I_b$, where $I_b(m) = b \circ m \circ b^{-1}$ for $m \in \operatorname{Aut} \mathbb{D}$. Let $f_1 = f \circ b$, $z_1 = b^{-1}(z_0)$. Then $d_1(z_1) = \lambda_0 = f_1(z_1)$. Consider $m \in \operatorname{Aut}_{z_1} \mathbb{D}$ and $\zeta \in \mathbb{D}$. Note that $I_b(m) \in \operatorname{Aut}_{z_0} \mathbb{D}$, and therefore, from equation (2.14) with $z = b(\zeta)$,

$$\begin{aligned} \Theta \circ I_b(m)(f \circ b(\zeta)) &= f \circ I_b(m) \circ I_b(m) \circ b(\zeta) \\ &= f \circ b \circ m \circ m(\zeta). \end{aligned}$$

Hence

$$\Theta_1(m)(f_1(\zeta)) = f_1 \circ m \circ m(\zeta).$$

This shows that synchrony at λ_0 does not depend on the choice of concomitant pair.

Nor does it depend on the choice of the map f. For suppose that f_1 is a second properly embedded analytic disc of $\mathbb{D}$ in Ω such that $f_1(z_0) = \lambda_0$ and $f_1(\mathbb{D}) = E_{\lambda_0}$. Then there exists $b \in \operatorname{Aut} \mathbb{D}$ such that $f_1 = f \circ b$ and $b(z_0) = z_0$. Consider any $m \in \operatorname{Aut}_{z_0} \mathbb{D}$ and $\zeta \in \mathbb{D}$. By equation (2.14),

$$\Theta(m) \circ f_1(\zeta) = \Theta(m)\,(f \circ b(\zeta)) = f \circ m \circ m(b(\zeta)).$$

Since $\operatorname{Aut}_{z_0} \mathbb{D}$ is conjugate in $\operatorname{Aut} \mathbb{D}$ to $\operatorname{Aut}_0 \mathbb{D}$, it is an abelian group. Hence

$$\Theta(m) \circ f_1(\zeta) = f \circ b \circ m \circ m(\zeta) = f_1 \circ m \circ m(\zeta),$$

which is the desired relation for f_1.

Remark 2.19. If $(\Omega, D, \mathcal{E})$ is as in the definition, then $(\Omega, D, \mathcal{E})$ is synchronous at a particular $\lambda_0 \in D$ if and only if $(\Omega, D, \mathcal{E})$ is synchronous at λ for every $\lambda \in D$. Consequently, it makes sense to say simply that $(\Omega, D, \mathcal{E})$ *is synchronous.*

For suppose $(\Omega, D, \mathcal{E})$ is synchronous at λ_0 with respect to the concomitant pair (d, Θ) where $\lambda_0 = d(z_0)$, and let $\lambda_1 \in D$.

Suppose $\lambda_1 = d(z_1)$, for $z_1 \in \mathbb{D}$, and $b(z_0) = z_1$, for $b \in \operatorname{Aut} \mathbb{D}$. For every $m \in \operatorname{Aut} \mathbb{D}$, we have $\Theta(m) \circ d = d \circ m$. Hence

$$\Theta(b)(\lambda_0) = \Theta(b) \circ d(z_0) = d(b(z_0)) = d(z_1) = \lambda_1.$$

Let

$$f_1 = \Theta(b) \circ f \circ b^{-1} : \mathbb{D} \to E_{\lambda_1}.$$

Then $f_1(z_1) = \lambda_1 = d(z_1)$. Consider $m \in \operatorname{Aut}_{z_1} \mathbb{D}$ and $\zeta \in \mathbb{D}$. Then

$$\begin{aligned}\Theta(m)(f_1(\zeta)) &= \Theta(m)(\Theta(b)(f \circ b^{-1}(\zeta))) \\ &= \Theta(m \circ b)(f \circ b^{-1}(\zeta)) \\ &= \Theta(b)\Theta(b^{-1} \circ m \circ b)(f(b^{-1}(\zeta))). \qquad (2.15)\end{aligned}$$

Since m fixes $z_1 = b(z_0)$, $b^{-1} \circ m \circ b$ fixes z_0. By assumption, $(\Omega, D, \mathcal{E})$ is synchronous at λ_0 with respect to the concomitant pair (d, Θ). Hence

$$\begin{aligned}\Theta(b^{-1} \circ m \circ b)(f(b^{-1}(\zeta))) &= f \circ (b^{-1} \circ m \circ b) \circ (b^{-1} \circ m \circ b)(b^{-1}(\zeta)) \\ &= f \circ b^{-1} \circ m \circ m(\zeta). \qquad (2.16)\end{aligned}$$

Therefore, by equations (2.15) and (2.16),

$$\begin{aligned}\Theta(m)(f_1(\zeta)) &= \Theta(b) \circ f \circ b^{-1} \circ m \circ m(\zeta) \\ &= f_1 \circ m \circ m(\zeta).\end{aligned}$$

Thus $(\Omega, D, \mathcal{E})$ is synchronous at λ_1 with respect to the concomitant pair (d, Θ).

In view of Remarks 2.18 and 2.19, the following statement follows easily from Lemma 1.19.

Lemma 2.20. *If $(\Omega, D, \mathcal{E})$ is as in Lemma* 2.16*, then $(\Omega, D, \mathcal{E})$ is synchronous.*

2.4. **The sharp direction in Ω.** For a regular royal manifold (Ω, D) we may define the sharp direction just as we did for G in Definition 1.9. By Proposition 2.13, for $\lambda \in \Omega$ the space $T_\lambda \operatorname{Orb}_\Omega(\lambda)$ is either a one-dimensional complex subspace (if $\lambda \in D$) or a 3-dimensional real subspace (if $\lambda \in \Omega \setminus D$) of $T_\lambda \Omega$. We may therefore define the space $\lambda^\sharp$ to be the unique nonzero complex subspace of $T_\lambda \operatorname{Orb}_\Omega(\lambda)$. In either case

$$\lambda^\sharp = T_\lambda \operatorname{Orb}_\Omega(\lambda) \cap i T_\lambda \operatorname{Orb}_\Omega(\lambda).$$

Covariance of the sharp direction under automorphisms is proved in the same way as Proposition 1.10.

Proposition 2.21. *If $\theta \in \operatorname{Aut} \Omega$ and $\lambda \in \Omega$ then*

$$\theta(\lambda)^\sharp = \theta'(\lambda)\lambda^\sharp.$$

Proposition 2.22. *Let $\Lambda : G \to \Omega$ be a biholomorphic map and let (d, Θ) be the concomitant pair for $(\Omega, \Lambda(\mathcal{R}))$ consistent with Λ. If $s \in G$ and $\Lambda(s) = \lambda$, then*

(1) $\Lambda'(s) T_s \operatorname{Orb}_G(s) = T_\lambda \operatorname{Orb}_\Omega(\lambda)$;

(2) $\Lambda'(s)s^\sharp = \lambda^\sharp$.

Moreover, if $s \notin \mathcal{R}$, then $e'_s(\mathrm{id}_\mathbb{D})$ is invertible and

(3) $e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1} = \Lambda'(s)\big|T_s \operatorname{Orb}_G(s)$;
(4) $e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1} : T_s \operatorname{Orb}_G(s) \to T_\lambda \operatorname{Orb}_\Omega(\lambda)$ *is a real linear map whose restriction to $s^\sharp$ is complex linear and maps $s^\sharp$ to $\lambda^\sharp$.*

For $s \in \mathcal{R}$, the real linear map $e'_s(\mathrm{id}_\mathbb{D})$ maps the 3-dimensional space Lie(Aut $\mathbb{D}$) to $T_s\mathcal{R}$, which is 2-dimensional, so we cannot form $e'_s(\mathrm{id}_\mathbb{D})^{-1}$.

Proof. (1) By assumption, $d = \Lambda \circ R$ and $\Theta(m) = \Lambda \circ \gamma_m \circ \Lambda^{-1} \in \operatorname{Aut}\Omega$ for every $m \in \operatorname{Aut}\mathbb{D}$. Hence

$$\begin{aligned} e_\lambda(m) &= \Theta(m)(\lambda) \\ &= \Lambda \circ \gamma_m \circ \Lambda^{-1}(\lambda) \\ &= \Lambda \circ \gamma_m(s) \\ &= \Lambda \circ e_s(m). \end{aligned}$$

That is, $e_\lambda = \Lambda \circ e_s$. Hence

$$\begin{aligned} e'_\lambda(\mathrm{id}_\mathbb{D}) &= \Lambda'(e_s(\mathrm{id}_\mathbb{D}))e'_s(\mathrm{id}_\mathbb{D}) \\ &= \Lambda'(s)e'_s(\mathrm{id}_\mathbb{D}). \end{aligned} \tag{2.17}$$

Therefore

$$\operatorname{ran} e'_\lambda(\mathrm{id}_\mathbb{D}) = \operatorname{ran} \Lambda'(s)e'_s(\mathrm{id}_\mathbb{D}) = \Lambda'(s) \operatorname{ran} e'_s(\mathrm{id}_\mathbb{D}),$$

which is to say (by virtue of equations (1.15) and (2.7)) that

$$T_\lambda \operatorname{Orb}_\Omega(\lambda) = \Lambda'(s)T_s \operatorname{Orb}_G(s).$$

(2) $s^\sharp$ is a nonzero complex subspace of $T_s \operatorname{Orb}_G(s)$. Since $\Lambda'(s)$ is a nonsingular complex linear map, $\Lambda'(s)s^\sharp$ is a nonzero complex linear subspace of $T_\lambda \operatorname{Orb}_\Omega(\lambda)$, hence is $\lambda^\sharp$.
(3) Consider $s \in G\backslash\mathcal{R}$. By Theorem 1.6, the real linear map $e'_s(\mathrm{id}_\mathbb{D})$ has full rank between the 3-dimensional spaces Lie(Aut $\mathbb{D}$) and $T_s \operatorname{Orb}_G(s)$, and so is nonsingular. Hence, $e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1}$ exists and is a real linear map from $T_s \operatorname{Orb}_G(s)$ to $T_\lambda \operatorname{Orb}_\Omega(\lambda)$. By equation (2.17),

$$e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1} = \Lambda'(s) \quad \text{on} \quad T_s \operatorname{Orb}_G(s). \tag{2.18}$$

(4) Since $\Lambda'(s)$ is a complex linear map on $\mathbb{C}^2$, it follows that $e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1}$ is a complex linear map on the complex linear subspace $s^\sharp$ of $\mathbb{C}^2$. By (2) and equation (2.18), $e'_\lambda(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1}s^\sharp = \lambda^\sharp$.

□

2.5. **Sharpness of the action of** $\operatorname{Aut}\Omega$**.** In this section, for a flatly fibered royal manifold $(\Omega, D, \mathcal{E})$, we shall show that sharp action of $\operatorname{Aut}\Omega$, as described in the introduction, is necessary for Ω to be isomorphic to G. In the next subsection we shall show that the condition is also sufficient. We first define sharpness more formally than in the introduction. Recall that, for a flatly fibered royal manifold $(\Omega, D, \mathcal{E})$, we defined the Poincaré parameter $P(\mu)$ for $\mu \in \Omega$ to be the Poincaré distance of μ from λ, where $\lambda \in D$ and $\mu \in E_\lambda$ and the distance is taken in the disc E_λ. That is, if $f : \mathbb{D} \to \Omega$ is a proper analytic embedding with range E_λ and $f(z_0) = \lambda$, $f(z) = \mu$, then

$$P(\mu) \stackrel{\text{def}}{=} \operatorname{arctanh}\left|\frac{z - z_0}{1 - \bar{z}_0 z}\right|. \tag{2.19}$$

It will be convenient to use also the pseudohyperbolic variant $C(\mu)$ of $P(\mu)$, defined for $\mu \in E_\lambda$ to be the pseudohyperbolic distance in E_λ from μ to λ. In other words, if $f \in \Omega(\mathbb{D})$ has range E_λ and $f(z_0) = \lambda$, $f(z) = \mu$, then

$$C(\mu) \stackrel{\text{def}}{=} \left|\frac{z - z_0}{1 - \bar{z}_0 z}\right|. \tag{2.20}$$

Thus P and C are related by the equations

$$\begin{aligned} P(\mu) &= \operatorname{arctanh} C(\mu) \\ &= \tfrac{1}{2} \log \frac{1 + C(\mu)}{1 - C(\mu)}. \end{aligned} \tag{2.21}$$

Remark 2.23. Observe that $P(\cdot)$ and $C(\cdot)$ are invariant under isomorphisms which preserve foliations. If $(\Omega_j, D_j, \mathcal{E}_j)$ is a flatly fibered royal manifold for $j = 1, 2$, if $\Lambda : \Omega_1 \to \Omega_2$ is an isomorphism which maps the leaves of $\mathcal{E}_1$ to those of $\mathcal{E}_2$ and if $\mu \in \Omega_1$ then $C(\mu) = C(\Lambda(\mu))$.

Definition 2.24. *Let $(\Omega, D, \mathcal{E})$ be a regular flatly fibered royal manifold having a concomitant pair (d, Θ). Let $\mu \in \Omega \setminus D$ and let (U, ψ) be a chart in Ω such that $\mu \in U$. We say that* $\operatorname{Aut}\Omega$ *acts sharply at μ with respect to (d, Θ) if*

$$\mathrm{e}^{2P(\mu)} \left(\psi(\Theta(B_{it})(\mu)) - \psi(\mu)\right) = i \left(\psi(\Theta(B_t)(\mu)) - \psi(\mu)\right) + o(t) \tag{2.22}$$

as $t \to 0$ in $\mathbb{R}$.

The condition (2.22) states that the tangents $v_{\mathbf{i}}$ and $v_{\mathbf{1}} \in \mathbb{C}^2$ at $t = 0$ to the curves $\psi(\Theta(B_{\mathbf{i}t})(\mu))$ and $\psi(\Theta(B_t)(\mu))$ in $\psi(U)$ satisfy

$$\mathrm{e}^{2P(\mu)} v_{\mathbf{i}} = \mathbf{i} v_{\mathbf{1}},$$

where **i** is (temporarily, for this sentence) the imaginary unit. This property is clearly independent of the chart ψ since the derivative of any transition function at a point is a complex-linear map.

We need to examine how sharpness depends on the concomitant pair (d, Θ).

Proposition 2.25. *With the notation of Definition* 2.24*, let* (d_1, Θ_1) *(for some* $b \in \operatorname{Aut}\mathbb{D}$*) be the concomitant pair*

$$(d \circ b, \Theta \circ I_b) \quad \text{where } I_b(m) = b \circ m \circ b^{-1} \text{ for } m \in \operatorname{Aut}\mathbb{D}.$$

Let $\mu_1 = \Theta(b)(\mu) \in \Omega \setminus D$*. Then* $\operatorname{Aut}\Omega$ *acts sharply at* μ_1 *with respect to* (d_1, Θ_1) *if and only if* $\operatorname{Aut}\Omega$ *acts sharply at* μ *with respect to* (d, Θ)*.*

Proof. We may choose the chart

$$\psi_1 = \psi \circ \Theta(b)^{-1} \quad \text{on } \Theta(b)(U)$$

at μ_1. For small real t,

$$\begin{aligned} \psi_1(\mu_1) &= \psi \circ \Theta(b)^{-1}((\Theta(b)(\mu)) \\ &= \psi(\mu), \\ \psi_1(\Theta_1(B_{it})(\mu_1)) &= \psi \circ \Theta(b)^{-1} \circ (\Theta(b) \circ \Theta(B_{it}) \circ \Theta(b)^{-1})((\Theta(b)(\mu)) \\ &= \psi(\Theta(B_{it})(\mu)). \end{aligned}$$

These equations, together with the analogous ones with it replaced by t and the fact that $P(\mu_1) = P(\mu)$, imply the statement in the proposition. □

Definition 2.26. *Let* $(\Omega, D, \mathcal{E})$ *be a regular flatly fibered royal manifold having a concomitant pair* (d, Θ)*. Let* $\mu \in \Omega \setminus D$ *and let* (U, ψ) *be a chart in* Ω *such that* $\mu \in U$*. We say that* $\operatorname{Aut}\Omega$ acts sharply on Ω *if* $\operatorname{Aut}\Omega$ *acts sharply with respect to* (d, Θ) *at every point of* $\Omega \setminus D$*.*

Remark 2.27. (1) Propositions 2.5 and 2.25 show that the sharpness of the action of $\operatorname{Aut}\Omega$ on Ω does not depend on the choice of the concomitant pair (d, Θ).
(2) With respect to a fixed concomitant pair, for any $\theta \in \operatorname{Aut}\Omega$, automorphisms act sharply at $\mu \in \Omega \setminus D$ if and only if they act sharply at $\theta(\mu)$. Since every orbit in Ω meets every leaf in $\mathcal{E}$, to conclude that $\operatorname{Aut}\Omega$ acts sharply, it is enough to show that, for some $\lambda \in D$, automorphisms act sharply at every point of $E_\lambda \setminus \{\lambda\}$.

Proposition 2.28. *Let* Ω *be a complex manifold and* $\Lambda : G \to \Omega$ *be a biholomorphic map. There exist a royal disc* D *in* Ω*, a flat fibration* $\mathcal{E}$ *of* Ω *over* D *and a concomitant pair* (d, Θ) *such that* $(\Omega, D, \mathcal{E})$ *is a synchronous regular flatly fibered royal manifold,* (d, Θ) *is consistent with* Λ *and* $\operatorname{Aut}\Omega$ *acts sharply on* $\Omega \setminus D$*.*

Proof. Let $D = \Lambda(\mathcal{R})$. By Lemmas 2.2 and 2.6, (Ω, D) is a royal manifold and there is a concomitant pair (d, Θ) for (Ω, D) such that

$$\Theta(m) \circ \Lambda = \Lambda \circ \gamma_m \tag{2.23}$$

for all $m \in \operatorname{Aut}\mathbb{D}$. Thus (d, Θ) is consistent with Λ. By Proposition 2.10, (Ω, D) is a regular royal manifold. Let $\mathcal{E}$ correspond under Λ to the flat fibration of G. By Lemma 2.20, $(\Omega, D, \mathcal{E})$ is a synchronous regular flatly fibered royal manifold. It remains to show that $\operatorname{Aut}\Omega$ acts sharply on $\Omega \setminus D$.

Consider a point $\mu \in \Omega \setminus D$, say $\mu \in E_\lambda$, for some $\lambda \in D$. Let $s = \Lambda^{-1}(\mu)$. We may assume (by modifying Λ and Θ and utilising Remark 2.27) that s has the form $(0, p)$ for some $p \in (0, 1)$. Then s lies in the flat geodesic F^0, and so $\Lambda^{-1}(E_\lambda) = F^0$. It follows that $\Lambda^{-1}(\lambda) = (0, 0)$ and since isomorphisms preserve the Möbius distance, $p = C(\mu)$.

Let (U, ψ) be a chart at μ. For any $\alpha \in \mathbb{C}$ and all small enough real t, in view of equation (2.23),

$$\begin{aligned}\psi\left(\Theta(B_{t\alpha})(\mu)\right) &= \psi \circ \Lambda \circ \gamma_{B_{t\alpha}}(s) \\ &= \psi(\mu) + (\psi \circ \Lambda)'(s) \frac{\mathrm{d}}{\mathrm{d}t} \gamma_{B_{t\alpha}}(s)\big|_{t=0} + o(t). \end{aligned} \tag{2.24}$$

By Lemma 1.7, with $r = 0$ and $s = (0, p)$,

$$\frac{\mathrm{d}}{\mathrm{d}t} \gamma_{B_{t\alpha}}(s)\big|_{t=0} = -\alpha \begin{pmatrix} 2 \\ 0 \end{pmatrix} - \bar{\alpha} \begin{pmatrix} 2p \\ 0 \end{pmatrix}.$$

Let $A = -2(\psi \circ \Lambda)'(s)$, so that A is a complex-linear map. Taking successively $\alpha = i$ and $\alpha = 1$ in equation (2.24) we obtain

$$\begin{aligned}\psi\left(\Theta(B_{ti})(\mu)\right) - \psi(\mu) &= i(1-p)A(1,0) + o(t), \\ \psi\left(\Theta(B_t)(\mu)\right) - \psi(\mu) &= (1+p)A(1,0) + o(t).\end{aligned}$$

Hence

$$(1+p)\left(\psi\left(\Theta(B_{ti})(\mu)\right) - \psi(\mu)\right) = i(1-p)\left(\psi\left(\Theta(B_t)(\mu)\right) - \psi(\mu)\right) + o(t).$$

Since $p = C(\mu)$, this is to say that $\operatorname{Aut}\Omega$ acts sharply at μ. □

The next statement justifies the terminology of 'sharp action'.

Lemma 2.29. *Let $(\Omega, D, \mathcal{E})$ be a regular flatly fibered royal manifold and suppose that $\operatorname{Aut}\Omega$ acts sharply at a point $\mu \in \Omega \setminus D$. For any $s \in G$ such that $C(s) = C(\mu)$ the map*

$$e'_\mu(\mathrm{id}_\mathbb{D}) e'_s(\mathrm{id}_\mathbb{D})^{-1} : T_s \operatorname{Orb}_G(s) \to T_\mu \operatorname{Orb}_\Omega(\mu) \tag{2.25}$$

maps $s^\sharp$ to $\mu^\sharp$ and is a complex-linear map.

Proof. Let

$$X = e'_\mu(\mathrm{id}_\mathbb{D})e'_s(\mathrm{id}_\mathbb{D})^{-1}.$$

X is a real-linear map from T_sG to $T_\mu\Omega$. We must show that $Xs^\sharp \subseteq \mu^\sharp$ and that X is complex-linear on $s^\sharp$.

We can assume that $s = (0, p)$ where $0 < p < 1$. Clearly

$$p = C(s) = C(\mu).$$

By equation (2.21),

$$\mathrm{e}^{2P(\mu)} = \frac{1+p}{1-p}.$$

The sharpness hypothesis, according to Definition 2.29, is

$$\frac{1+p}{1-p}\left(\psi(\Theta(B_{it})(\mu)) - \psi(\mu)\right) = i\left(\psi(\Theta(B_t)(\mu)) - \psi(\mu)\right) + o(t) \quad (2.26)$$

as $t \to 0$ in $\mathbb{R}$. By Proposition 1.16,

$$s^\sharp = \mathbb{C}\begin{pmatrix}1\\0\end{pmatrix}.$$

We shall use the local co-ordinates $(r, \alpha) \in (-\pi, \pi) \times \mathbb{D}$ for a neighborhood of $\mathrm{id}_\mathbb{D}$ in $\operatorname{Aut}\mathbb{D}$, as in Lemma 1.4. By Lemma 1.7,

$$e'_s(\mathrm{id}_\mathbb{D})(\varphi_1^{-1})'(r, \alpha) = v_{r,\alpha}(0, p) = \begin{pmatrix}-2\alpha - 2p\bar\alpha\\ 2irp\end{pmatrix},$$

which is in $s^\sharp$ if and only if $r = 0$. Thus

$$e'_s(\mathrm{id}_\mathbb{D})^{-1}s^\sharp = (\varphi_1^{-1})'(0 \oplus \mathbb{C}).$$

Moreover, for all $\alpha \in \mathbb{C}$,

$$(\varphi_1^{-1})'(0, \alpha) = e'_s(\mathrm{id}_\mathbb{D})^{-1}\begin{pmatrix}-2\alpha - 2p\bar\alpha\\ 0\end{pmatrix}. \quad (2.27)$$

Note that $m_{0,\alpha} = B_\alpha$ in the notation of equation (0.6). Let ψ be a chart on Ω at μ. For any $\alpha \in \mathbb{C}$, as $t \to 0$ in $\mathbb{R}$,

$$\begin{aligned}\psi\left(\Theta(B_{t\alpha})(\mu)\right) &= \psi(\mu) + \psi'(\mu)e'_\mu(\mathrm{id}_\mathbb{D})(\varphi_1^{-1})'(0, \alpha) + o(t)\\ &= \psi(\mu) + \psi'(\mu)X\begin{pmatrix}-2\alpha - 2p\bar\alpha\\ 0\end{pmatrix} + o(t)\end{aligned}$$

Take in succession $\alpha = 1$ and $\alpha = i$ and use the real-linearity of X to obtain the relations

$$\psi\left(\Theta(B_t)(\mu)\right) - \psi(\mu) = -2(1+p)\psi'(\mu)X(1, 0) + o(t), \quad (2.28)$$

$$\psi\left(\Theta(B_{ti})(\mu)\right) - \psi(\mu) = -2(1-p)\psi'(\mu)Xi(1, 0) + o(t). \quad (2.29)$$

We have

$$\begin{aligned}\psi'(\mu)Xi(1,0) &= -\frac{1}{2(1-p)}\left(\psi(\Theta(B_{ti}(\mu)) - \psi(\mu)\right) + o(t) \\ &\qquad\qquad \text{by equation (2.29)} \\ &= -\frac{1}{2(1+p)}\left(\psi(\Theta(B_t(\mu)) - \psi(\mu)\right) + o(t) \\ &\qquad\qquad \text{by equation (2.26)} \\ &= i\psi'(\mu)X(1,0) + o(t) \quad \text{by equation (2.28).}\end{aligned}$$

Since $\psi'(\mu)$ is an invertible complex-linear map which identifies $T_\mu\Omega$ with $\mathbb{C}^2$, it follows that

$$X(i,0) = iX(1,0). \tag{2.30}$$

The vectors $(1,0)$ and $(i,0)$ span $s^\sharp$ over $\mathbb{R}$, and

$$\operatorname{ran} X \subseteq \operatorname{ran} e'_\mu(\mathrm{id}_\mathbb{D}) \subseteq T_\mu \operatorname{Orb}_\Omega(\mu).$$

Equation (2.30) now shows both that $\operatorname{ran} X \subset \mu^\sharp$ and that X is complex-linear on $s^\sharp$. □

2.6. **A characterization of G.** We have arrived at the main theorem of the paper.

Theorem 2.30. *A complex manifold Ω is isomorphic to G if and only if there exist a royal disc D in Ω and a flat fibration $\mathcal{E}$ of Ω over D such that $(\Omega, D, \mathcal{E})$ is a synchronous regular flatly fibered royal manifold and $\operatorname{Aut}\Omega$ acts sharply on Ω.*

Proof. Necessity is Proposition 2.28. We prove sufficiency. Let (d, Θ) be a concomitant pair for (Ω, D). Choose $z_0 \in \mathbb{D}$ and let $s_0 = R(z_0)$ and $\lambda_0 = d(z_0)$. We shall construct a biholomorphic map $\Lambda : G \to \Omega$ satisfying $\Lambda(s_0) = \lambda_0$. Figure 1 represents the construction.

Choose a properly embedded analytic disc g of $\mathbb{D}$ into G satisfying $g(\mathbb{D}) = F_{s_0}$ and $g(z_0) = s_0$. Choose also a properly embedded analytic disc $f : \mathbb{D} \to \Omega$ such that $f(z_0) = \lambda_0$ and $f(\mathbb{D}) = E_{\lambda_0}$. For $s \in G$ we define $\Lambda(s)$ by the following recipe.

Since each point in G is in a flat geodesic and $\operatorname{Aut} G$ acts transitively on the flat geodesics, we may choose $m \in \operatorname{Aut}\mathbb{D}$ such that $\gamma_m^{-1}(s) \in F_{s_0}$ and hence there exists $z \in \mathbb{D}$ such that

$$s = \gamma_m \circ g(z). \tag{2.31}$$

Let

$$\Lambda(s) = \Theta(m) \circ f(z) \tag{2.32}$$



FIGURE 1. The construction of $\Lambda : G \to \Omega$

Certainly $\Lambda(s) \in \Omega$. To see that this recipe does define Λ as a map from G to Ω, consider $z_1, z_2 \in \mathbb{D}$ and $m_1, m_2 \in \operatorname{Aut}\mathbb{D}$ such that

$$\gamma_{m_1} \circ g(z_1) = \gamma_{m_2} \circ g(z_2). \tag{2.33}$$

We wish to show that

$$\Theta(m_1) \circ f(z_1) = \Theta(m_2) \circ f(z_2). \tag{2.34}$$

Note first that equation (2.33) implies that if $m = m_2^{-1} \circ m_1$, then

$$\gamma_m \circ g(z_1) = g(z_2).$$

Since $g(z_1), g(z_2) \in F_{s_0}$, Lemma 1.14 implies that $m \in \operatorname{Aut}_{z_0}\mathbb{D}$. Consequently, by Lemma 1.19

$$g \circ m \circ m(z_1) = \gamma_m \circ g(z_1) = g(z_2),$$

which implies that

$$z_2 = m \circ m(z_1).$$

By hypothesis, $(\Omega, D, \mathcal{E})$ is synchronous. According to Definition 2.17, it means (since $m \in \operatorname{Aut}_{z_0}\mathbb{D}$) that

$$\Theta(m) \circ f = f \circ m \circ m.$$

Hence

$$\begin{aligned} \Theta(m) \circ f(z_1) &= f \circ m \circ m(z_1) \\ &= f(z_2). \end{aligned}$$

Therefore equation (2.34) is true, and so $\Lambda(s)$ is unambiguously defined.

On taking $m = \mathrm{id}_{\mathbb{D}}$ in equation (2.32) we have

$$\Lambda \circ g(z) = f(z) \tag{2.35}$$

for all $z \in \mathbb{D}$. In particular, $\Lambda(s_0) = \lambda_0$.

Consider any $\upsilon \in \operatorname{Aut} \mathbb{D}$. Since

$$\gamma_\upsilon(s) = \gamma_\upsilon \circ \gamma_m \circ g(z) = \gamma_{\upsilon \circ m} \circ g(z),$$

by the definition (2.32) of Λ,

$$\begin{aligned} \Lambda \circ \gamma_\upsilon(s) &= \Theta(\upsilon \circ m) \circ f(z) \\ &= \Theta(\upsilon) \circ \Theta(m) \circ f(z) \\ &= \Theta(\upsilon) \circ \Lambda(s). \end{aligned}$$

Thus

$$\Lambda \circ \gamma_\upsilon = \Theta(\upsilon) \circ \Lambda \quad \text{ for all } \upsilon \in \operatorname{Aut} \mathbb{D}. \tag{2.36}$$

Now fix a general point $R(z)$ on the royal geodesic $\mathcal{R}$. If $m \in \operatorname{Aut} \mathbb{D}$ is such that $m(z_0) = z$, then

$$\begin{aligned} \Lambda \circ R(z) &= \Lambda \circ R \circ m(z_0) \\ &= \Lambda \circ \gamma_m \circ R(z_0) \qquad \text{by equation (1.5).} \end{aligned}$$

By equation (2.36) and the fact that $s_0 = R(z_0)$,

$$\begin{aligned} \Lambda \circ R(z) &= \Theta(m) \circ \Lambda(s_0) \\ &= \Theta(m)(\lambda_0) \\ &= \Theta(m) \circ d(z_0) \\ &= d \circ m(z_0) \qquad \text{by equation (2.1)} \\ &= d(z). \end{aligned}$$

Thus

$$\Lambda \circ R = d. \tag{2.37}$$

Now fix $z_1 \in \mathbb{D}$ and choose m such that $m(z_0) = z_1$. Since $\gamma_m \circ R(z_0) = R(z_1)$, Lemma 1.14 implies that

$$\gamma_m(F_{R(z_0)}) = F_{R(z_1)},$$

and since $\Theta(m) \circ d(z_0) = d(z_1)$, Condition (3) in Definition 2.15 implies that

$$\Theta(m)(E_{d(z_0)}) = E_{d(z_1)}.$$

Therefore

$$\begin{aligned}\Lambda(F_{R(z_1)}) &= \Lambda \circ \gamma_m(F_{R(z_0)}) \\ &= \Theta(m) \circ \Lambda(F_{R(z_0)}) \\ &= \Theta(m)(E_{d(z_0)}) \\ &= E_{d(z_1)}.\end{aligned}$$

To summarize, we have shown that if $\mathcal{F}$ denotes the partition of G in Lemma 1.12 and $\mathcal{E}$ denotes the partition of Ω in Definition 2.15, then Λ induces a map $\Lambda^\sim : \mathcal{F} \to \mathcal{E}$ given by

$$\Lambda^\sim(F_{R(z)}) = E_{d(z)}.$$

Furthermore, as the map $R(z) \mapsto d(z)$ from $\mathcal{R}$ to D is a bijection, so also is $\Lambda^\sim$.

Consider any point $\lambda \in E_{d(z_1)}$. Then $\Theta(m^{-1})(\lambda) \in E_{\lambda_0}$, and so $\Theta(m^{-1})(\lambda) = f(z)$ for some $z \in \mathbb{D}$. Hence $\lambda = \Theta(m) \circ f(z)$. By equations (2.31) and (2.32), $\lambda = \Lambda \circ \lambda_m \circ g(z)$. Thus $\lambda \in \Lambda(G)$, and so Λ is surjective.

Suppose s_1, $s_2 \in G$ satisfy $\Lambda(s_1) = \Lambda(s_2)$. Since $\Lambda^\sim$ is a bijection, it follows that s_1, s_2 lie in the same flat geodesic in G, say in $F_{R(z_1)}$. Let $m \in \operatorname{Aut}\mathbb{D}$ be such that $m(z_0) = z_1$. We have, for $j = 1, 2$,

$$\gamma_m^{-1}(s_j) \in F_{R \circ m^{-1}(z_1)} = F_{R(z_0)} = F_{s_0}.$$

Hence $\gamma_m^{-1}(s_j) = g(\zeta_j)$ for some $\zeta_1, \zeta_2 \in \mathbb{D}$. By equation (2.32),

$$\Lambda(s_j) = \Theta(m) \circ f(\zeta_j).$$

Hence $\Theta(m) \circ f(\zeta_1) = \Theta(m) \circ f(\zeta_2)$, and therefore $\zeta_1 = \zeta_2$. Thus

$$s_1 = \gamma_m \circ g(\zeta_1) = \gamma_m \circ g(\zeta_2) = s_2.$$

We have shown that $\Lambda : G \to \Omega$ is bijective. Moreover, we can observe that

$$\Lambda | F_{R(z_1)} = \Theta(m) \circ g \circ f^{-1} \circ \gamma_m^{-1} | F_{R(z_1)}. \tag{2.38}$$

There remains to prove that Λ and Λ^{-1} are holomorphic.

We shall first show that Λ is smooth as a mapping between real manifolds by giving a formula for Λ which is clearly differentiable. The assumption that $z_0 = 0$, $g(z) = (0, z)$ and so $s_0 = (0, 0)$ loses no generality. It implies that $F_{s_0} = \{(0, z) : z \in \mathbb{D}\}$.

Consider a point

$$s = (\zeta + \eta, \zeta\eta) \in G$$

for some $\zeta, \eta \in \mathbb{D}$. To evaluate $\Lambda(s)$ we shall choose an automorphism m of $\mathbb{D}$ satisfying $m'(0) > 0$ such that $\gamma_m^{-1}(s) \in F_{s_0}$. To see that this

is possible take $m = B_\alpha$ for some $\alpha \in \mathbb{D}$. Then $m'(0) > 0$. We require $\gamma_{B_\alpha}^{-1}(s) \in F_{s_0}$, which is to say that

$$B_{-\alpha}(\zeta) + B_{-\alpha}(\eta) = 0.$$

Expressing this relation in terms of the components s^1, s^2 of s, we must find $\alpha = \alpha(s) \in \mathbb{D}$ such that

$$s^1 = -\frac{2\alpha}{1+|\alpha|^2} - \frac{2\bar{\alpha}}{1+|\alpha|^2} s^2.$$

Compare this expression with that of the flat co-ordinates for s given in equations (1.25) and (1.26):

$$s^1 = \beta + \bar{\beta} s^2,$$

where

$$\beta = \beta(s) = \frac{s^1 - \bar{s^1} s^2}{1 - |s^2|^2}.$$

One sees that it suffices to choose $\alpha(s)$ such that

$$\beta(s) = -\frac{2\alpha(s)}{1+|\alpha(s)|^2}.$$

A suitable choice of $\alpha(s)$ is

$$\alpha(s) = \frac{-\beta(s)}{1+\sqrt{1-|\beta(s)|^2}}$$

as may readily be checked. Clearly $\beta, \alpha \in \mathbb{D}$ and both β and α are real-analytic functions of s. Moreover

$$\begin{aligned} L(s) \stackrel{\text{def}}{=} \gamma_{B_\alpha}^{-1}(s) &= (0, B_{-\alpha}(\zeta) B_{-\alpha}(\eta)) \\ &= \left(0, \frac{s^2 + \alpha(s) s^1 + \alpha(s)^2}{1 + \overline{\alpha(s)} s^1 + \overline{\alpha(s)}^2 s^2}\right), \end{aligned}$$

which is also real-analytic in s. By the definition of Λ,

$$\Lambda(s) = \Theta(B_{\alpha(s)}) \circ f \circ g^{-1} \circ L(s).$$

The map $s \mapsto B_{\alpha(s)}$ is real-analytic from G to $\operatorname{Aut}\mathbb{D}$. Since the action of $\operatorname{Aut}\mathbb{D}$ on Ω is differentiable, by the regularity assumption on the royal manifold (Ω, D), we conclude that $\Lambda : G \to \Omega$ is differentiable.

Consider $s \in G$ and suppose that $s \in F_{R(z_1)}$. Let $X = \Lambda'(s)$ viewed as a real-linear mapping from $T_s G$ to $T_{\Lambda(s)}\Omega$.

Recall from Definition 1.13 that $s^\flat$ denotes the flat direction at s. Equation (2.38) implies that

$$X(s^\flat) = \Lambda(s)^\flat \text{ and } X|s^\flat \text{ is complex linear.} \tag{2.39}$$

By equations (2.31) and (2.32), for all $z \in \mathbb{D}$ and $m \in \operatorname{Aut}\mathbb{D}$,

$$\Lambda \circ \gamma_m \circ g(z) = \Theta(m) \circ f(z).$$

In view of the definitions (1.3) and (2.6), this equation can be written

$$\Lambda \circ e_{g(z)} = e_{f(z)} : \operatorname{Aut}\mathbb{D} \to \Omega.$$

On differentiating at $\mathrm{id}_{\mathbb{D}}$ we obtain

$$\Lambda' \circ e_{g(z)}(\mathrm{id}_{\mathbb{D}}) e'_{g(z)}(\mathrm{id}_{\mathbb{D}}) = e'_{f(z)}(\mathrm{id}_{\mathbb{D}}) : \operatorname{Lie}(\operatorname{Aut}\mathbb{D}) \to T_{f(z)} \operatorname{Orb}_{\Omega}(f(z)).$$

For $z \neq z_0$, the point $g(z) \notin \mathcal{R}$, and therefore, by Proposition 2.22, $e'_{g(z)}(\mathrm{id}_{\mathbb{D}})$ is invertible, and so

$$\Lambda' \circ g(z) = e'_{f(z)}(\mathrm{id}_{\mathbb{D}}) e'_{g(z)}(\mathrm{id}_{\mathbb{D}})^{-1}.$$

By the hypothesis, $\operatorname{Aut}\Omega$ acts sharply on Ω. By Lemma 2.29, it follows that $\Lambda' \circ g(z)$ maps $g(z)^\sharp$ into $f(z)^\sharp$ and is complex-linear on $g(z)^\sharp$ whenever $g(z) \notin \mathcal{R}$.

Recalling that $X : T_s G \to T_{\Lambda(s)}\Omega$ is real-linear and that (by Proposition 1.15) $s^\flat$ and $s^\sharp$ are linearly independent, we infer from equation (2.39) that $X = \Lambda'(s)$ is complex linear for all $s \in G \setminus \mathcal{R}$. Therefore Λ is analytic on $G \setminus \mathcal{R}$. The restriction of Λ to any co-ordinate plane $P_\zeta \stackrel{\text{def}}{=} \{s \in G : s^1 = \zeta\}$, for $|\zeta| < 2$, is analytic in s^2 except possibly at the sole point $(\zeta, \frac{1}{4}\zeta^2)$ of $P_\zeta \cap \mathcal{R}$ and is continuous on P_ζ. Hence $\Lambda | P_\zeta$ is analytic in s^2. Likewise the restriction of Λ to any of the orthogonal co-ordinate planes is analytic in s^1. Thus Λ is analytic on G. Every bijective holomorphic map between domains has a holomorphic inverse (for example, [20, Chapter 10, Exercise 37]). It follows easily that a bijective holomorphic map between a domain and a complex manifold has a holomorphic inverse. □

3. A characterization of G via flat co-ordinates

Recall from Subsection 1.5 that G is foliated by the sets

$$F^\beta = \{(\beta + \bar{\beta}z, z) : z \in \mathbb{D}\}$$

for $\beta \in \mathbb{D}$ [4, Theorem 2.1]. Thus the map $\eta : \mathbb{D}^2 \to G$ defined by the formula

$$\eta(\beta, z) = (\beta + \bar{\beta}z, z), \qquad \beta, z \in \mathbb{D}, \tag{3.1}$$

is a homeomorphism of $\mathbb{D}^2$ onto G.

We will call β, z the *flat co-ordinates* for points of G. In this section we shall use the variables (β, z) for points in $\mathbb{D}^2$ and the variables (s, p) for points in G, so that

$$s = \beta + \bar{\beta}z, \qquad p = z, \qquad \beta, z \in \mathbb{D}.$$

Flat co-ordinates provide another characterization of domains biholomorphic to G.

The following lemma is a consequence of the Chain Rule.

Lemma 3.1. *If $f = f(s,p)$ is a differentiable function on G, η is defined on $\mathbb{D}^2$ as in equation (3.1) and $\xi = f \circ \eta$, then the following relations hold.*

$$\frac{\partial \xi}{\partial \beta} = \frac{\partial f}{\partial s} + \bar{z}\frac{\partial f}{\partial \bar{s}}, \tag{3.2}$$

$$\frac{\partial \xi}{\partial \bar{\beta}} = z\frac{\partial f}{\partial s} + \frac{\partial f}{\partial \bar{s}}, \tag{3.3}$$

$$\frac{\partial \xi}{\partial z} = \bar{\beta}\frac{\partial f}{\partial s} + \frac{\partial f}{\partial p}, \tag{3.4}$$

$$\frac{\partial \xi}{\partial \bar{z}} = \beta\frac{\partial f}{\partial \bar{s}} + \frac{\partial f}{\partial \bar{p}}. \tag{3.5}$$

Theorem 3.2. *If Ω is a domain in $\mathbb{C}^2$, then Ω is biholomorphic to G if and only if there exists a differentiable homeomorphism $\Xi = (\xi_1, \xi_2)$ from $\mathbb{D}^2$ onto Ω satisfying*

$$\frac{\partial \xi_i}{\partial \bar{\beta}} = z\frac{\partial \xi_i}{\partial \beta}, \qquad i = 1, 2 \tag{3.6}$$

and

$$\frac{\partial \xi_i}{\partial \bar{z}} = 0, \qquad i = 1, 2 \tag{3.7}$$

at all $(\beta, z) \in \mathbb{D}^2$.

Proof. First assume that $F \in \Omega(G)$ is a biholomorphic map of G onto Ω and let $\Xi = F \circ \eta$. Since η is a smooth homeomorphism of $\mathbb{D}^2$ onto G, Ξ is a smooth homeomorphism of $\mathbb{D}^2$ onto Ω.

If we set $F = (f_1, f_2)$ and $\Xi = (\xi_1, \xi_2)$, then f_i is holomorphic and $\xi_i = f_i \circ \eta$ for $i = 1, 2$. Hence, using equations (3.2) and (3.3), we see that

$$\frac{\partial \xi_i}{\partial \bar{\beta}} = z\frac{\partial f_i}{\partial s} = z\frac{\partial \xi_i}{\partial \beta}, \qquad i = 1, 2,$$

which proves that equation (3.6) holds. Also, equation (3.5) implies that the relation (3.7) holds.

Now assume that $\Xi = (\xi_1, \xi_2)$ is a differentiable homeomorphism from $\mathbb{D}^2$ onto Ω satisfying equations (3.6) and (3.7). Define $F = (f_1, f_2)$ by $F = \Xi \circ \eta^{-1}$. Since η is a differentiable homeomorphism of $\mathbb{D}^2$ onto G, it follows that F is a differentiable homeomorphism of G onto Ω. There remains to show that F is holomorphic.

Since $\xi_i = f_i \circ \eta$, we have

$$z\frac{\partial f_i}{\partial s} + \frac{\partial f_i}{\partial \bar{s}} = \frac{\partial \xi_i}{\partial \bar{\beta}} = z\frac{\partial \xi_i}{\partial \beta} \qquad \text{by equations (3.3) and (3.6)}$$

$$= z\left(\frac{\partial f_i}{\partial s} + \bar{z}\frac{\partial f_i}{\partial \bar{s}}\right) \qquad \text{by equation (3.2).}$$

Thus

$$(1 - |z|^2)\frac{\partial f_i}{\partial \bar{s}} = 0$$

on G. Since $|z| < 1$ when $(s, p) \in G$ it follows that

$$\frac{\partial f_i}{\partial \bar{s}} = 0$$

throughout G. Hence f_1, f_2 are holomorphic on G. □

4. Asymmetry of domains

É. Cartan's classification theorem [8] is based on his theory of *symmetric* spaces, in the sense of the first paragraph of the paper. In $\mathbb{C}^2$ and $\mathbb{C}^3$ (but not $\mathbb{C}^4$) *every* bounded homogeneous domain is a symmetric space [8, 14]. In contrast, none of the 'almost homogeneous' domains that we consider is symmetric.

Let us say that a point λ in a domain Ω is a *point of symmetry* of Ω if there exists a holomorphic self-map γ of Ω such that $\gamma \circ \gamma = \mathrm{id}_\Omega$ and λ is an isolated fixed point of γ. Thus a domain is symmetric if every point of the domain is a point of symmetry.

From the fact that the automorphisms of the annulus $\mathbb{A}_q$ are the maps ωz and ωz^{-1} for $\omega \in \mathbb{T}$ (for example, [12, Theorem 6.2]), it is easy to see that the only points of symmetry in $\mathbb{A}_q$ are the points of the unit circle. Hence $\mathbb{A}_q$ is not a symmetric domain.

Proposition 4.1. *Neither the symmetrized bidisc nor the tetrablock contains a point of symmetry.*

Proof. We sketch the proof for the tetrablock; that for the symmetrized bidisc is similar but simpler.

Let E denote the tetrablock defined in equation (0.5). Every orbit in E contains a point of the form $(0, 0, p)$ [26, Theorem 5.2], so it suffices to show that no such point is a point of symmetry. By [1, Theorem 2.2], the tetrablock is foliated by the 'flat geodesics'

$$C_{\beta_1\beta_2} \stackrel{\text{def}}{=} \{(\beta_1 + \bar{\beta}_2 z, \beta_2 + \bar{\beta}_1 z, z) : z \in \mathbb{D}\}$$

where $|\beta_1| + |\beta_2| < 1$. These geodesics are permuted by the automorphisms of E [26, Theorem 5.1]. Moreover the 'royal variety'

$\{x \in E : x^1x^2 = x^3\}$ is invariant under all automorphisms of E (see the proof of [26, Theorem 4.1]).

Consider a holomorphic involution γ of E that fixes $(0, 0, p)$. Then γ fixes the flat geodesic containing $(0, 0, p)$, which is C_{00}. Hence γ fixes the only common point of C_{00} and the royal variety, which is easily seen to be $(0, 0, 0)$. It is shown in [26, Proof of Theorem 4.1, foot of page 766] that an automorphism γ of E fixes $(0, 0, 0)$ if and only if either

$$\gamma(x) = (\omega_1 x^1, \omega_2 x^2, \omega_1\omega_2 x^3) \tag{4.1}$$

or

$$\gamma(x) = (\omega_2 x^2, \omega_1 x^1, \omega_1\omega_2 x^3) \tag{4.2}$$

for some $\omega_1,\ \omega_2 \in \mathbb{T}$.

In the case that γ is of the form (4.1), since γ is an involution, we have $\omega_1^2 = \omega_2^2 = 1$. Thus the four involutions of this form that fix $(0, 0, p)$ have fixed points as in the following table.

ω_1	ω_2	$\gamma(x)$	Fixed points
1	1	x	E
-1	1	$(-x^1, x^2, -x^3)$	$(0, z, 0)$
1	-1	$(x^1, -x^2, -x^3)$	$(z, 0, 0)$
-1	-1	$(-x^1, -x^2, x^3)$	$(0, 0, z)$

where z ranges over $\mathbb{D}$. Hence in the case (4.1), $(0, 0, p)$ is not an isolated fixed point of γ.

In case (4.2),

$$\gamma \circ \gamma(x) = (\omega_1\omega_2 x^1, \omega_1\omega_2 x^2, (\omega_1\omega_2)^2 x^3)$$

and so the involutory property of γ corresponds to the condition $\omega_1\omega_2 = 1$. Hence $\gamma(x) = (\bar{\omega}x^2, \omega x^1, x^3)$ for some $\omega \in \mathbb{T}$. Then the fixed points of γ are the points $(x^1, \omega x^1, x^3)$ in E. Hence $(x^1, \omega x^1, p)$ is a fixed point of γ for all x^1 in a neighborhood of 0, and so $(0, 0, p)$ is not an isolated fixed point of γ. □

References

[1] A. A. Abouhajar, M. C. White and N. J. Young, A Schwarz lemma for a domain related to mu-synthesis, *J. Geometric Analysis* **17** (2007) 717-750.

[2] J. Agler, Z. A. Lykova and N. J. Young, Geodesics, retracts, and the extension property in the symmetrized bidisc, *Memoirs of the American Mathematical Society* **258** (2019) no. 1242, 106pp.

[3] J. Agler, Z. A. Lykova and N. J. Young, Characterizations of some domains via Carathéodory extremals, *J. Geometric Analysis*, DOI 10.1007/s12220-018-0059-6.

[4] J. Agler and N. J. Young, The hyperbolic geometry of the symmetrised bidisc, *J. Geometric Analysis* **14** (2004) 375-403.

[5] J. Agler and N. J. Young, The complex geodesics of the symmetrised bidisc, *Int. J. Math.* **17** (2006) 375-391.
[6] J. Agler and N. J. Young, The magic functions and automorphisms of a domain, *Complex Analysis and Operator Theory* **2** (2008) 383-404.
[7] T. Bhattacharyya, S. Pal and S. Shyam Roy, Dilations of Γ-contractions by solving operator equations, *Advances in Mathematics* **230** (2012) 577-606.
[8] É. Cartan, Sur les domaines bornés homogènes de l'espace de n variables complexes, *Abh. Math. Sen. Univ. Hamburg* **11** (1935) 116-162.
[9] H. Cartan, *Sur les groupes de transformations analytiques*, Hermann et Cie, Paris 1935.
[10] C. Costara, The symmetrized bidisc and Lempert's theorem, *Bull. Lond. Math. Soc.* **36** (2004) 656–662.
[11] M. Cvetic, G. W. Gibbons, H. Lu, C. N. Pope, New cohomogeneity one metrics with spin(7) holonomy, *J. Geom. Phys.* **49** (2004) 350–365.
[12] S. D. Fisher, *Function theory on planar domains*, Wiley, New York, 1983.
[13] A. Edigarian and W. Zwonek, Geometry of the symmetrized polydisc, *Arch. Math. (Basel)* **84**(2005) 364-374.
[14] B. A. Fuks, *Special chapters in the theory of analytic functions of several complex variables*, Translations of Mathematical Monographs **14** (American Mathematical Society, Providence RI, 1965).
[15] K. Grove and W. Ziller, Cohomogeneity one manifolds with positive Ricci curvature, *Inv. Math.* **149** (2002), 619–646.
[16] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Academic Press, New York 1978.
[17] M. Jarnicki and P. Pflug, On automorphisms of the symmetrised bidisc, *Arch. Math. (Basel)* **83** (2004) 264-266.
[18] M. Jarnicki and P. Pflug, *Invariant Distances and Metrics in Complex Analysis*, 2nd Extended Edition, De Gruyter, Berlin, 2013.
[19] L. Kosiński and W. Zwonek, Nevanlinna-Pick problem and uniqueness of left inverses in convex domains, symmetrized bidisc and tetrablock, *J. Geom. Anal.* **26** (2016) 1863-1890.
[20] S. G. Krantz, *Function theory of several complex variables*, Wiley, New York, 1982.
[21] L. Lempert, La métrique de Kobayashi et la représentation des domaines sur la boule, *Bull. Soc. Math. France*, **109** (1981) 427–484.
[22] P. Pflug and W. Zwonek, Description of all complex geodesics in the symmetrized bidisc, *Bull. London Math. Soc.*, **37** (2005) 575–584.
[23] J. Sarkar, Operator theory on symmetrized bidisc, *Indiana Univ. Math. J.* **64** (2015) 847–873.
[24] M. Spivak, *Calculus on Manifolds*, W. A. Benjamin Inc., New York 1965.
[25] M. Trybula, Invariant metrics on the symmetrized bidisc, *Complex Variables and Elliptic Equations* **60** (4) (2015) 559-565.
[26] N. J. Young, The automorphism group of the tetrablock, *J. London Math. Soc.* **77** (2008) 757-770.
[27] N. J. Young, Some analysable instances of mu-synthesis, *Mathematical methods in systems, optimization and control*, Editors: H. Dym, M. de Oliveira, M. Putinar, Operator Theory: Advances and Applications **222** 349–366, Birkhäuser, Basel, 2012.

Index

(Ω, D), 18
$(\Omega, D, \mathcal{E})$, 25
(d, Θ), 20
B_α, 4
F^β, 14
F_s, 14
G, 2
$P(\mu)$, 30
$T_\lambda\Omega$, 4
$\mathbb{D}$, 3
$\Omega(U)$, 5
$\mathbb{T}$, 16
Aut_λ, 16
$\mathcal{V}(s)$, 9
γ_m, 5
$\lambda^\sharp$, 4, 28
$\lambda^\flat$, 25
Lie(Aut $\mathbb{D}$), 9
$\mathbb{A}_q$, 2, 41
$\mathrm{Orb}_U(\lambda)$, 7
ρ_η, 16
e_s, 6
$s^\flat$, 14
$s^\sharp$, 13
$v_{r,\alpha}$, 9

analytic disc
  properly embedded, 5
annulus, 2, 41
automorphism, 3

Cartan
  É., 1, 41
  H., 21
Cartan's theorem, 1
concomitant pair, 20
consistent, 20

domain, 1
  bounded symmetric homogeneous, 1, 41

evaluation map, 6

Figure 1, 34
flat
  direction, 14, 25
  fibration, 15, 25
  geodesic, 13
flat co-ordinates, 39

geodesic
  complex, 3
  flat, 3

homogeneous, 1

Lempert, 2
Lie
  algebra, 9
  group, 6

main theorem, 34

orbit, 2

Poincaré parameter, 3, 30

royal
  disc, 3, 18
  manifold, 18
    flatly fibered, 25
    regular, 21
  variety, 3

sharp
  action, 4, 30
  direction, 4, 13, 28
stabilizer, 16
symmetric, 1, 41
symmetrized bidisc, 2
synchronous, 26
synchrony, 4, 16, 18

tetrablock, 2, 41

Department of Mathematics, University of California at San Diego, CA 92103, USA

School of Mathematics, Statistics and Physics, Newcastle University, Newcastle upon Tyne NE1 7RU, U.K.
Email address: Zinaida.Lykova@ncl.ac.uk

School of Mathematics, Statistics and Physics, Newcastle University, Newcastle upon Tyne NE1 7RU, U.K. *and* School of Mathematics, Leeds University, Leeds LS2 9JT, U.K.
Email address: Nicholas.Young@ncl.ac.uk