



UNIVERSITY OF LEEDS

This is a repository copy of *Utility of Magnetic Resonance Imaging in Diagnosis and Monitoring Enthesitis in Patients with Spondyloarthritis: An OMERACT Systematic Literature Review*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/141081/>

Version: Accepted Version

Article:

Mathew, AJ, Krabbe, S, Kirubakaran, R et al. (4 more authors) (2019) Utility of Magnetic Resonance Imaging in Diagnosis and Monitoring Enthesitis in Patients with Spondyloarthritis: An OMERACT Systematic Literature Review. *Journal of Rheumatology*, 46 (9). pp. 1207-1214. ISSN 0315-162X

<https://doi.org/10.3899/jrheum.181083>

© 2019. All rights reserved. This is an author produced version of a paper published in *Journal of Rheumatology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Table 1: Characteristics of included studies

Author/Year of publication	Study population	Number of participants	Intervention	Comparator	MRI field strength	Sequences	MRI sites	Scoring systems
Cross sectional studies								
Fournie 1997	AS fulfilling Amor criteria and PsA based on seronegative joint disease with psoriasis	8 (5; 3)	MRI	NA	NA	T1W Gd	Anterior chest wall	Qualitative
McGonagle 2002	SpA with plantar fasciitis; Mechanically induced plantar fasciitis	28 (17; 11)	MRI	NA	0.5; 1.5	T1W; SPIR	Plantar fascia	Semi quantitative
Olivieri 2002	SpA with dactylitis	6	MRI	NA	1.5	T1W; T2W FS; GRE-T2W	Finger tendon insertions	Semi quantitative
Tan 2006	DIP joint PsA; DIP joint OA; Healthy subjects	30 (10; 10; 10)	MRI	NA	1.5	T1W; T2W FS; PD; 3DGE; T1W FS Gd	DIP joints	Qualitative
McQueen 2007	PsA	10	MRI	NA	0.6	T1W; T1W Gd; STIR	2nd-5th finger	Semi quantitative
Tan 2007	DIP joint PsA; DIP joint OA; Healthy subjects	30 (10; 10; 10)	MRI	NA	1.5	T1W; T2W FS; PD; 3DGE; T1W FS Gd	DIP joints	Qualitative
Marzo-Ortega 2009	SpA; RA	20 (10; 10)	MRI	NA	1.5	T1W; DCE-MRI; SPIR FS Gd	MCP joints	Qualitative
Maksymowicz 2010	SpA or suspected SpA	35	MRI T1W FS Gd	MRI T2W FS	1.5	T1W; T1W FS; T2W FS; T1W FS Gd	Enthesitis of SIJ ligaments	Semi quantitative
Feydy 2012	SpA; Controls hospitalized low-back pain	75 (51; 24)	MRI	NA	1.5	T1W; STIR	Heel enthesitis	Qualitative
Althoff 2013	SpA	75	MRI	NA	1.5	T1W; STIR	Whole-body	Qualitative
Aydin 2013	SpA with swollen knee	21	MRI	US	1.5	T1W; T2 SPIR; TI SPIR; TI SPIR Gd	Knee entheses	Qualitative
Braum 2013	Suspected inflammatory joint disease	69	MRI	Clinical	1.5	T1W; T1W FS Gd; STIR	Collateral ligaments of finger joints	Qualitative
Paramarta 2014	Knee or ankle arthritis: SpA; RA; crystal arthritis	41 (13; 20; 8)	MRI	NA	1.5	T1W; T2W FS; STIR; T1W FS Gd	Knee and ankle entheses	Qualitative
Ramirez 2014	Greater trochanter pain: SpA/RA/no inflammatory disease	40	MRI	NA	1.5	T1W; T2W FS	Greater femoral trochanter	Qualitative

Godfrin 2004	Entheseal pain at multiple sites	33	MRI	Clinical	1.5	T1W; T2W; T1W FS Gd and/or STIR	Not described	Qualitative
Eshed 2008	SpA with hindfoot pain	27	MRI	NA	0.2; 1.5	T1W; STIR; T1W FS Gd; T1W GRE FS	Heel enthesitis	Qualitative
Karpitscha 2013	AS	10	MRI	NA	1.5	T1W; STIR	Whole-body	Qualitative
Zhen-Guo 2013	AS	58	MRI	CR; Clinical	1.5	T2W; T1W; STIR; T1W FS Gd	Hip	Qualitative
Althof f2016	SpA	41	MRI	Clinical	1.5	T1W; STIR	Whole-body	Qualitative
Marzo-Ortega 2001	SpA	10	MRI	NA	1.5	T1W; T2W FS; T1W FS Gd	Depending on symptoms of each patient	Qualitative
Tan 2004	AS fulfilling modified NY criteria	9	MRI	NA	1.5	T1TSE, STIR	SIJ and spine	Semi quantitative
de Hooge 2013	Chronic back pain	127	MRI T1W FS Gd	MRI STIR	1.5	T1W; T1W FS Gd; STIR	SIJ enthesitis/capsulitis	Qualitative

Randomised controlled trials

Dougados 2010	SpA with heel enthesitis	24	MRI	NA	Not reported	T1W; STIR	Heel enthesitis	Quantitative
Song 2011	SpA	76	MRI	NA	1.5	T1W; STIR	Whole-body	Qualitative
Krabbe 2018	SpA	49	MRI	NA	3	T1W; STIR	Whole-body	Semi quantitative

* AS – ankylosing spondylitis, SpA – spondyloarthritis, PsA – psoriatic arthritis, RA – rheumatoid arthritis, OA – osteoarthritis, MCP – metacarpophalangeal joint, DIP – distal interphalangeal joint, SIJ – sacroiliac joints, T1W – T1 weighted, T2W – T2 weighted, Gd – gadolinium, FS – fat suppressed, SPIR – spectral pre-saturation with inversion recovery, GRE – gradient recalled echo, PD – proton density, 3D-GE – 3D gradient echo, STIR – short tau inversion recovery, DCE – dynamic contrast enhanced, TSE – turbo spin echo, VIBE – volumetric interpolated breath-hold sequence, UTE – quantitative ultrashort echo time, US – ultrasound, CR – conventional radiograph