



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/141086/>

Version: Accepted Version

Article:

Mathew, AJ, Krabbe, S, Eshed, I et al. (2019) The OMERACT MRI in Enthesitis Initiative: Definitions of Key Pathologies, Suggested MRI Sequences and Novel Heel Enthesitis Scoring System (HEMRIS). *Journal of Rheumatology*, 46 (9). pp. 1232-1238. ISSN: 0315-162X

<https://doi.org/10.3899/jrheum.181093>

© 2019 The Journal of Rheumatology. This is an author produced version of a paper published in *Journal of Rheumatology*. Uploaded in accordance with the publisher's self-archiving policy. <https://doi.org/10.3899/jrheum.181093>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Table 3: Exercise 3: Baseline and change single-measure inter-reader ICCs (sum scores), quadratic weighted kappa (individual component scores, per lesion) and mean scores of all readers

<i>BASELINE SCORES</i>							
	Reproducibility (smICC & Kappa)				Range of scores	Reader scores	
	All readers		Subset of readers*			All readers	Subset of readers*
	Median	Mean (Range)	Median	Mean (Range)		**Mean (Range)	Mean (Range)
Inter-reader smICC (total score)	0.83	0.81 (0.57– 0.95)	0.91	0.91 (0.90– 0.94)	0-21	3.55 (0.1 – 13.6)	4.04 (0.25 – 14.5)
<i>Achilles tendon (Kappa – lesion wise)</i>							
Peri-tendon hypersignal	0.64	0.62 (0.29–0.87)	0.79	0.78 (0.68– 0.87)	0-3	0.47 (0 – 2.00)	0.41 (0 – 2.00)
Intra-tendon hypersignal	0.55	0.51 (0.12– 0.89)	0.79	0.81 (0.77– 0.89)	0-3	0.34 (0 – 1.50)	0.33 (0 – 1.50)
Retro-calcaneal bursitis	0.55	0.49 (-0.12-0.93)	0.57	0.60 (0.42– 0.78)	0-3	0.34 (0 – 2.1)	0.40 (0 – 2.5)
Bone marrow edema	0.86	0.86 (0.75– 0.97)	0.89	0.88 (0.84– 0.92)	0-3	0.38 (0 – 3.00)	0.39 (0 – 3.00)
<i>Plantar fascia (Kappa – lesion wise)</i>							
Peri-aponeurosis hypersignal	0.65	0.60 (0.06– 0.89)	0.80	0.80 (0.66– 0.89)	0-3	0.85 (0 – 2.80)	1.01 (0 – 3.00)
Intra-aponeurosis hypersignal	0.55	0.46 (-0.0– 0.93)	0.84	0.84 (0.79– 0.93)	0-3	0.56 (0 – 2.00)	0.80 (0 – 3.00)
Bone marrow edema	0.89	0.89 (0.76– 0.98)	0.87	0.88 (0.81– 0.96)	0-3	0.61 (0 – 2.90)	0.65 (0 – 3.00)
<i>CHANGE SCORES</i>							
Inter-reader smICC (total score)	0.82	0.80 (0.57– 0.92)	0.84	0.85 (0.79– 0.85)	0-21	1.54 (0.1 – 4.9)	1.99 (0 – 6.25)
<i>Achilles tendon (Kappa – lesion wise)</i>							
Peri-tendon hypersignal	0.49	0.47 (0.21– 0.75)	0.50	0.53 (0.42– 0.75)	0-3	0.20 (0 – 0.8)	0.17 (0 – 1.00)
Intra-tendon hypersignal	0.41	0.41 (0.09– 0.63)	0.54	0.52 (0.35– 0.63)	0-3	0.15 (0 – 1.00)	0.17 (0 – 1.25)

Retro-calcaneal bursitis	0.24	0.26 (-0.2– 1.00)	0.45	0.42 (0.15– 0.67)		0-3	0.21 (0 – 0.8)	0.29 (0 – 1.00)
Bone marrow edema	0.52	0.53 (0.30– 0.82)	0.47	0.54 (0.45– 0.76)		0-3	0.14 (0 – 1.1)	0.20 (0 – 1.25)
<i>Plantar fascia (Kappa – lesion wise)</i>								
Peri-aponeurosis hypersignal	0.66	0.61 (0.17– 0.87)	0.77	0.78 (0.70– 0.85)		0-3	0.33 (0 – 1.30)	0.42 (0 – 1.75)
Intra-aponeurosis hypersignal	0.53	0.42 (-0.08-0.80)	0.62	0.64 (0.57– 0.77)		0-3	0.24 (0 – 0.90)	0.40 (0 – 1.25)
Bone marrow edema	0.78	0.77 (0.57– 0.94)	0.81	0.79 (0.69– 0.88)		0-3	0.28 (0 – 1.30)	0.58 (0 – 1.25)

smICCs - single measures intraclass correlation coefficient by two-way random effects, absolute agreement for sum scores (patient level).

* One participating radiologist + three rheumatologist with best individual ICC with other readers for inflammatory pathologies in exercise 2

**Each patient's score was calculated as the mean of all readers. The presented mean and ranges are means/ranges of these values. (Range of readers' mean scores)

Readers: AJM, DG, FG, IE †, MS, MØ ‡, PB, SK ‡, SJP, VF ‡, † Musculoskeletal radiologist. ‡ Three rheumatologists with best individual ICC for inflammatory pathologies in exercise 2.