



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/141245/>

Version: Accepted Version

Proceedings Paper:

Sheese, PE, Walker, KA, Boone, CD et al. (2018) Investigating N₂O produced in the mesosphere - Lower thermosphere and its transport to the middle atmosphere. In: OSA Technical Digest. Fourier Transform Spectroscopy 2018: Light, Energy and the Environment 2018 (E2, FTS, HISE, SOLAR, SSL), 05-08 Nov 2018, Singapore. OSA. Article no: FT5B.2. ISBN: 9781557528209.

<https://doi.org/10.1364/FTS.2018.FT5B.2>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Investigating N₂O produced in the mesosphere – lower thermosphere and its transport to the middle atmosphere

Patrick E. Sheese

University of Toronto, Toronto, Ontario, CANADA

Kaley A. Walker

University of Toronto, Toronto, Ontario, CANADA
e-mail address: kwalker@atmosph.physics.utoronto.ca

Chris D. Boone

University of Waterloo, Waterloo, Ontario, CANADA

Monika Andersson

Finnish Meteorological Institute, Helsinki, FINLAND

Daniel Marsh

National Center for Atmospheric Research, Boulder, UNITED STATES

Christopher Kelly, Wuhu Feng, John Plane, and Martyn Chipperfield

University of Leeds, Leeds, UNITED KINGDOM

Abstract: This paper will describe Atmospheric Chemistry – Experiment Fourier Transform Spectrometer (ACE-FTS) N₂O measurements in the middle atmosphere along with its production mechanisms and long-term variation. Comparisons with model N₂O simulations will also be discussed.

OCIS codes: 010.0280 Remote sensing and sensors; 010.1280 Atmospheric composition

1. Motivation

Nitrous oxide (N₂O) is an important greenhouse gas and contributes to ozone depletion in the Earth's atmosphere. Until recently, it was thought that there were no in situ atmospheric sources of N₂O. However, measurements from the ACE-FTS (Atmospheric Chemistry Experiment – Fourier Transform Spectrometer) instrument show that there is a consistent presence of N₂O in the mesosphere-lower thermosphere (MLT) [1]. These measurements are important for understanding and characterizing the global N₂O budget and its effects on the atmosphere.

2. Introduction

ACE-FTS [2] is a limb-viewing, solar occultation instrument on board the Canadian SCISAT satellite, which was launched into a high inclination orbit in August 2003. It is a high spectral resolution spectrometer that retrieves atmospheric profiles from infrared observations between 750 and 4400 cm⁻¹. One of the 50+ species that ACE-FTS measures is N₂O, which is retrieved between 5 and 95 km with ~3-4 km vertical resolution. The measurements, which span from February 2004 to the present, are made between 85°S to 85°N, with a focus on high latitudes. This study uses the most current version of the ACE-FTS level 2 N₂O data, version 3.5/3.6 [3, 4]. The ACE-FTS data are compared to simulated N₂O data from the Whole Atmosphere Community Climate Model (WACCM) [5], which is run in specified dynamics mode and has been modified to include reactions relevant to N₂O production, including ion chemistry in the D-region.

3. Discussion

Latitudinal cross sections of 2004-2018 zonal mean ACE-FTS N₂O profiles are shown in Figure 1 for January-February and for July-August. A layer of N₂O is observed at high altitudes, around 90 km, that is present at all latitudes. The largest concentrations are observed in the polar winter region on the order of 100 ppbv. At all latitudes, N₂O is indirectly produced via photoelectrons, and at the poles there is an additional source that is dependent on energetic particle precipitation [6].

Occasionally, after strong sudden stratospheric warming events, which often occur in the Arctic winter, strong descent in the polar vortex can bring NO_x and N₂O-rich air down from the upper atmosphere into the upper stratosphere. ACE-FTS has measured N₂O concentrations on the order of 1 ppbv descending from the lower thermosphere down to ~40 km [1]. NO_x, which plays a significant role in O₃ depletion, is produced from N₂O destruction. It is therefore important to understand what role atmospherically produced N₂O plays on NO_x and O₃ throughout the upper stratosphere to lower thermosphere.




Figure 1: Latitudinal cross section of 2004-2018 zonal mean ACE-FTS N₂O profiles for January-February (left) and July-August (right). White sections are regions where no data are available or where mean concentrations are negative.

4. Summary

ACE-FTS is the only instrument that measures N₂O produced in the MLT region. Near 90-95 km, ACE-FTS measures the presence of N₂O concentrations at all latitudes, all seasons, throughout the solar cycle, on the order of 10-100 ppbv. This presentation will detail comparisons between ACE-FTS measurements of N₂O, NO_x, and O₃ with WACCM simulations in order to further the understanding of the role N₂O plays on NO_x and O₃ chemistry in the middle atmosphere.

5. References

- [1] P. E. Sheese et al., "Nitrous oxide in the atmosphere: First measurements of a lower thermospheric source", *Geophys. Res. Lett.*, 43, 2866–2872 (2016).
- [2] P. F. Bernath et al., "Atmospheric Chemistry Experiment (ACE): Mission overview", *Geophys. Res. Lett.*, 32, L15S01 (2005).
- [3] C. D. Boone et al., "Retrievals for the atmospheric chemistry experiment Fourier-transform spectrometer", *Appl. Opt.*, 44, 7218-7231 (2005).
- [4] C. D. Boone et al., Version 3 Retrievals for the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), *The Atmospheric Chemistry Experiment ACE at 10: A Solar Occultation Anthology* (Peter F. Bernath, editor, A. Deepak Publishing, Hampton, Virginia, U.S.A., 2013), 103-127 (2013).
- [5] R. R. Garcia et al., "Simulation of secular trends in the middle atmosphere, 1950–2003", *J. Geophys. Res.*, 112, D09301, (2007).
- [6] C. Kelly et al., "An explanation for the nitrous oxide layer observed in the mesopause region", *Geophys. Res. Lett.*, Accepted, (2018).