



This is a repository copy of *Adhesion and acceptability of novel oral patches in human volunteers*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/142096/>

Version: Accepted Version

---

**Proceedings Paper:**

Colley, H.E. [orcid.org/0000-0003-0053-7468](https://orcid.org/0000-0003-0053-7468), D'Apice, K., Baker, S.R. [orcid.org/0000-0002-2861-451X](https://orcid.org/0000-0002-2861-451X) et al. (5 more authors) (2016) Adhesion and acceptability of novel oral patches in human volunteers. In: Oral Diseases. 13th Biennial Congress of the European Association of Oral Medicine, 15-17 Sep 2016, Torino, Italy. Wiley , p. 19.

<https://doi.org/10.1111/odi.12559>

---

This is the peer reviewed version of the following article: HE Colley, K D'Apice, SR Baker, PV Hatton, ME Santocildes-Romero, MH Thornhill, J Hansen, C Murdoch, Adhesion and acceptability of novel oral patches in human volunteers, (2016), Original Clinical Research. Oral Dis, 22: 14-41, which has been published in final form at <https://doi.org/10.1111/odi.12559>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Adhesion and acceptability of novel oral patches in human volunteers

H.E Colley<sup>1</sup>, K D'Apice<sup>1</sup>, S R Baker<sup>1</sup>, P Hatton<sup>1</sup>, M.E Santocildes-Romero<sup>1</sup>, M.H Thornhill<sup>1</sup>, J Hansen<sup>2</sup>, C Murdoch<sup>1</sup>

<sup>1</sup> *School of Clinical Dentistry, The University of Sheffield, Sheffield, UK*

<sup>2</sup> *DermTreat ApS, Abildgaardsvej 174, 2830, Virum, Denmark*

**Objective:** Oral lesions such as Oral lichen planus (OLP) or Aphthous Ulcers are often characterised by painful or erosive oral lesions. These lesions are often treated with steroid-containing mouthwashes or creams but these preparations can be ineffective due to low drug contact times. We have created an oral adhesive patch using electrospinning technology that is able to firmly adhere to porcine and tissue-engineered oral mucosa for prolonged periods and so may be able to deliver drugs directly to a lesion. In this study we examine the adherence of drug-free oral patches in human volunteers to determine their longevity and acceptability in humans.

**Methods:** Oral patches were placed on the buccal mucosa, lateral tongue or gingiva of healthy human volunteers (with written, informed consent; n=36) for 5 seconds with applied pressure and residence time measured every 10 minutes for up to 2 hours. Volunteers were also asked to complete a questionnaire to provide information on patch performance, comfortableness and acceptability.

**Results:** Oral patches were highly adherent to the human oral mucosal tissues with 100%, 89% and 86% of patches remaining adherent to the gingiva, tongue or buccal mucosa, respectively for 2 hours. 94% of volunteers felt the patches provided good/excellent adherence and over 70% of volunteers felt no irritation whilst wearing the patches. The majority of volunteers (>70%) stated that the patches were comfortable to wear and although 16% reported moderate interference with speech over 70% stated only minor affects on saliva production and swallowing. 97% of volunteers stated that they would be willing to wear the patch twice-a-day to treat an oral lesion if required.

**Conclusion:** Electrospun oral patches were strongly adherent to different sites of the oral mucosal for prolonged periods and, overall, were well tolerated. These patches have great potential to deliver drugs directly to oral lesions.